

UNCLASSIFIED

AD NUMBER
AD003750
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; FEB 1953. Other requests shall be referred to Office of Naval Research, 800 North Quincy Street, Arlington, VA 22217-5660.
AUTHORITY
ONR ltr., 9 Nov 1977

THIS PAGE IS UNCLASSIFIED

Reproduced by

Armed Services Technical Information Agency
DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

AD -

3750

Best Available Copy

UNCLASSIFIED

AD003750

Columbia University
in the City of New York

DEPARTMENT OF CIVIL ENGINEERING



REFLECTION OF FLEXURAL WAVES
AT THE EDGE OF A PLATE

By

T. R. KANE

Office of Naval Research Project NR-064-388

Contract Nonr-266(09)

Technical Report No. 6

CU-7-53-ONR-266(09)-CE

February 1953

ABSTRACT

The reflection of straight-crested flexural waves at the edge of a semi-infinite plate is studied in terms of a two-dimensional plate theory. It is found that, in general, a flexural wave propagated toward the edge at an arbitrary angle of incidence gives rise to three reflected waves: two flexural waves and a shear wave. A number of special cases, involving degenerate forms of these motions, are investigated in detail.

1. Introduction

In this paper Mindlin's [1]¹ equations of flexural motions of plates are used to study the reflection of a straight-crested wave at the edge of a semi-infinite plate. The equations accommodate three modes of motion: two types of flexural waves and a thickness-shear wave. It is found that, in general, all three of these motions are excited upon incidence of any one of them at a free edge. The shear motion, here encountered, is of particular interest; for this motion is absent in the classical theory of plates, whence the applicability of that theory is restricted to a range of frequencies considerably below that corresponding to the first mode of thickness-shear vibration. By the same token, the present theory does not include the higher modes of motion which are to be found in three-dimensional elasticity theory; thus, the solution set forth below may be expected to furnish an adequate description for frequencies which do not materially exceed that of the first thickness-shear mode.

The character of the reflected waves is affected by both the angle of incidence and the ratio of plate thickness to wave-length of the incident wave. Appropriate values of these two parameters give rise to such special cases as waves whose amplitudes decrease exponentially with distance from the edge, vibrations, resonance, disappearance of some of the reflected waves, and, for grazing incidence, complete disappearance of the motion.

Following a resumé of the plate equations in Section 2, straight-crested waves are considered in Section 3. Incident and emergent waves are described in Section 4 and, in terms of these, a formal solution is reached in Section 5. In Sections 6 and 7 various cases of normal and oblique incidence are discussed in detail. Section 8 deals with grazing incidence.

1. Numbers in brackets refer to the Bibliography at the end of the paper.

2. Plate Displacements, Equations of Motion and Stresses

For a plate of thickness h , oriented as in Fig. 1, plate displacements $\psi_x(x, y, t)$, $\psi_y(x, y, t)$ and $w(x, y, t)$ are given by (see [1])

$$\begin{aligned}\psi_x &= \left[(\sigma_1 - 1) \frac{\partial w_1}{\partial x} + (\sigma_2 - 1) \frac{\partial w_2}{\partial x} + \frac{\partial H}{\partial y} \right] e^{i p t} \\ \psi_y &= \left[(\sigma_1 - 1) \frac{\partial w_1}{\partial y} + (\sigma_2 - 1) \frac{\partial w_2}{\partial y} - \frac{\partial H}{\partial x} \right] e^{i p t} \\ w &= (w_1 + w_2) e^{i p t}\end{aligned}\quad (1)$$

The functions $w_1(x, y)$, $w_2(x, y)$ and $H(x, y)$ are governed, respectively, by equations of motion

$$\begin{aligned}(\nabla^2 + \delta_1^2) w_1 &= 0 \\ (\nabla^2 + \delta_2^2) w_2 &= 0 \\ (\nabla^2 + \omega^2) H &= 0\end{aligned}\quad (2)$$

In the above,

$$\begin{aligned}(\sigma_1, \sigma_2) &= (\delta_1^2, \delta_2^2) (R \delta_0^4 - S^{-1})^{-1} (\delta_1^2, \delta_2^2) S \delta_0^4 \\ 2(\delta_1^2, \delta_2^2) &= \delta_0^4 \{ S + R \pm [(S - R)^2 + 4 \delta_0^4]^{1/2} \}\end{aligned}\quad (3)$$

$$(1 - \nu) \omega^2 = 2 (R \delta_0^4 - S^{-1}) \quad (4)$$

$$D \delta_0^4 = \rho p^2 h \quad (5)$$

$$R = \frac{h^2}{12}, \quad S = \frac{D}{2\mu h}$$

$$D = \frac{Eh^3}{12(1-\nu^2)},$$

(6)

ρ is the circular frequency associated with the wave motion. E, ν, ρ are, respectively, Young's Modulus, Poisson's Ratio, and the mass density of the plate material.

Plate bending moments M_x, M_y , shears Q_x, Q_y and twisting moment M_{xy} are given by

$$M_x = D \left(\frac{\partial^2 \psi}{\partial x^2} + \nu \frac{\partial^2 \psi}{\partial y^2} \right)$$

$$M_y = D \left(\frac{\partial^2 \psi}{\partial y^2} + \nu \frac{\partial^2 \psi}{\partial x^2} \right)$$

$$2M_{xy} = D(1-\nu) \left(\frac{\partial^2 \psi}{\partial x \partial y} + \frac{\partial^2 \psi}{\partial y \partial x} \right)$$

(7)

$$Q_x = \kappa^2 \mu h \left(\psi_x + \frac{\partial \psi}{\partial x} \right)$$

$$Q_y = \kappa^2 \mu h \left(\psi_y + \frac{\partial \psi}{\partial y} \right)$$

μ is the modulus of rigidity of the plate material. The definition of κ^2 is given in [1].

3. Straight-Crested Waves

Taking any one of the functions w_1, w_2 , or H to be of the form

$$A \exp(i g x), \quad g > 0$$

and letting the remaining two vanish identically, we find that, in each case, two of the equations of motion are automatically satisfied while the remaining

one requires that

$$f^2 = \begin{cases} \delta_1^2, & W_1 \neq 0, W_2 = H = 0 \\ \delta_2^2, & W_2 \neq 0, W_1 = H = 0 \\ \omega^2, & H \neq 0, W_1 = W_2 = 0 \end{cases} \quad (8)$$

Letting

$$\rho^2 = (fc)^2, \quad c > 0,$$

this leads to [see Equations (3), (4), (5)] three possible types of relationships between phase velocity C and wave number f :

$$\frac{C}{C_s} = \begin{cases} \frac{K}{f} \left[\frac{\Gamma - (\Gamma^2 - 4j^2 RS)^{1/2}}{2K} \right]^{1/2}, & W_1 \neq 0, W_2 = H = 0 \\ \frac{K}{f} \left[\frac{\Gamma + (\Gamma^2 - 4j^2 RS)^{1/2}}{2R} \right]^{1/2}, & W_2 \neq 0, W_1 = H = 0 \\ \frac{K}{f} \left[\frac{2 + j^2(1-\nu)S}{R} \right]^{1/2}, & H \neq 0, W_1 = W_2 = 0 \end{cases} \quad (9)$$

where

$$\Gamma = 1 + j^2(R+S)$$

and C_s is the velocity of shear waves in an infinite medium. A plot of C/C_s versus hf (with $\nu = 1/4$) for each of these cases is shown in Fig. 3. It is seen that $C/C_s \rightarrow 0$ as $hf \rightarrow 0$ for a W_1 wave, while $C/C_s \rightarrow \infty$ as $hf \rightarrow 0$ for W_2 and H waves. In the sequel we shall call W_1 a "slow" flexural wave, W_2 a "fast" flexural wave and H a shear wave. (The fact that the preceding discussion involves waves propagated in the X -direction, does not result in any loss of generality.)

4. Incident and Emergent Waves

Referring the semi-infinite plate to axes x, y, z as shown in Fig. 2 we consider a slow flexural wave² propagated towards the edge of the plate,

$$w_1 = A e^{i\psi}, w_2 = H = 0 \quad (10)$$

where

$$\psi = y \sin \alpha - x \cos \alpha$$

We postulate three emergent waves, propagated respectively towards P', P'', P''' as shown in Fig. 2. These are

a) a slow flexural wave:

$$w_1' = A' e^{i\psi'}, w_2' = H' = 0 \quad (11)$$

b) a fast flexural wave:

$$w_2'' = A'' e^{i\psi''}, w_1'' = H'' = 0 \quad (12)$$

c) a shear wave:

$$H''' = A''' e^{i\psi'''}, w_1''' = w_2''' = 0 \quad (13)$$

where

$$\psi^j = y \sin \alpha^j + x \cos \alpha^j, j = ', ', ''$$

and the α^j are the angles of emergence shown in Fig. 2.

In accordance with Equations (9), these waves are propagated with velocities c^j which depend on their respective wave numbers k^j .

2. The case of a fast incident wave is similar in nature and will not be discussed in the present paper.

The displacements corresponding to these waves are obtained from Equation (1). Denoting the incident wave by w, φ_x and φ_y , and calling the displacement components of the emergent waves $w^j, \varphi_x^j, \varphi_y^j$ ($j = ', ', ''$), we observe that, since the equations of motion (2) are linear, the state of motion given by

$$\begin{aligned} w^* &= w + \sum_j w^j \\ \varphi_x^* &= \varphi_x + \sum_j \varphi_x^j \\ \varphi_y^* &= \varphi_y + \sum_j \varphi_y^j \end{aligned} \quad (14)$$

is a possible state of motion for an infinite plate. For the semi-infinite plate under consideration, three boundary conditions must be satisfied on the edge $x=0$.

5. Boundary Conditions

Plate stresses M_x^*, M_y^* , etc. corresponding to the motion defined by Equations (14) may be computed from Equations (7). To obtain a traction-free edge, we must have, on $x=0$,

$$M_x^* = M_{yx}^* = Q_y^* = 0. \quad (15)$$

In order that these equations be satisfied for all values of the time t and all values of the space variable y , it is necessary that the circular frequencies p and p^j associated with the various waves be identical, i.e.,

$$pc = p^j c^j, \quad (16)$$

and that the angles α and α^j satisfy

$$f \sin \alpha = f^j \sin \alpha^j. \quad (17)$$

Now, the relation between C and f is of the same form as that between C' and f' , both being slow flexural waves. Thus it follows from Equations (16) and (17) that the angle of emergence α' of the slow flexural wave is equal to the angle of incidence α of the incident wave, and that the wave numbers and phase velocities of these two waves are identical. Equations (16) and (17) show, furthermore, that the condition of vanishing traction at the edge of the plate, [Eq. (15)], leads to a determination of the phase velocity, wave number and angle of emergence of each of the postulated emergent waves when the angle of incidence and wave number (or phase velocity) of the incident wave are specified.

The boundary conditions (15) also impose restrictions on the amplitude ratios A^j/A . From Equations (10)-(13), (14) and (7) we get, upon substitution into Equations (15), a system of three non-homogeneous linear algebraic equations governing these amplitude ratios,

$$\sum_j \phi_j^j \frac{A^j}{A} = -\frac{2}{\sqrt{2}} \phi_0' \cos[(2n-1)\frac{\pi}{4}], \quad j = ', ', ', \quad n = 1, 2, 3 \quad (18)$$

where

$$\begin{aligned} \phi_1^j &= M^j \phi^j, \quad j = ', ', ' \\ \phi_1''' &= (1-\nu) f^{\prime\prime\prime} \sin \alpha^{\prime\prime} \cos \alpha^{\prime\prime} \\ \phi_2^j &= M^j \sin 2\alpha^j, \quad j = ', ', ' \\ \phi_2''' &= -f^{\prime\prime\prime} \cos 2\alpha''' \\ \phi_3^j &= S \sigma_0^* f^j (f^j)^{-1} \cos \alpha^j, \quad j = ', ', ' \\ \phi_3''' &= f^{\prime\prime\prime} \sin \alpha''' \end{aligned} \quad (19)$$

with

$$M^j = S\delta_0^2 - (f^j)^2$$

$$\theta^j = \cos^2 \alpha^j + \nu \sin^2 \alpha^j.$$

The analog, in the present theory, to P. Neumann's uniqueness theorem [2] guarantees that the above constitutes the unique solution of the problem. We may say then that a slow incident flexural wave produces, in general, three reflected waves: a slow flexural, a fast flexural, and a shear wave.

6. Normal Incidence

We now proceed to study the character of the reflected waves in terms of the angle of incidence α and the wave number f of the incident wave. We begin by examining the case of normal incidence, i.e., $\alpha = 0$.

The wave numbers f'' and f''' of the fast reflected wave and of the shear wave are given by [see Eqs. (8), (4), (5)]

$$2(f'')^2 = \delta_0^2 \{ S + R - [(S - R)^2 + 4\delta_0^2]^{1/2} \} \quad (20)$$

$$(f''')^2 = 2(1 - \nu)^{-1} (R\delta_0^2 - S^2) \quad (21)$$

Both f'' and f''' vanish when $R\delta_0^2 = 1$, i.e., when $f^2 RS = R + S$. For $f^2 RS < R + S$, f'' and f''' are imaginary while for $f^2 RS > R + S$ both are real. The physical significance of $f^2 RS = 1$ will be discussed when the motion corresponding to that value of the wave number of the incident wave has been determined. We first consider

$$\underline{f^2 RS < R + S}$$

From Eq. (17) we have, for $\alpha = 0$,

$$\alpha' = 0, \quad j = 1, 2, 3$$

The amplitude ratios are found from Eqs. (18):

$$\begin{aligned} \frac{A'}{A} &= \frac{a_1}{A} + i \frac{b_1}{A} \\ \frac{A''}{A} &= \frac{a_2}{A} + i \frac{b_2}{A} \\ \frac{A'''}{A} &= 0 \end{aligned} \quad (22)$$

where

$$\begin{aligned} \frac{a_1}{A} &= \frac{(\bar{f}''')^2 (S\bar{f}_0'' + \bar{f}''')^2 - f^2 (S\bar{f}_0'' - f')^2}{(\bar{f}''')^2 (S\bar{f}_0'' + \bar{f}''')^2 + f^2 (S\bar{f}_0'' - f')^2} \\ \frac{b_1}{A} &= - \frac{2\bar{f}'' f (S\bar{f}_0'' + \bar{f}''') (S\bar{f}_0'' - f')}{(\bar{f}''')^2 (S\bar{f}_0'' + \bar{f}''')^2 + f^2 (S\bar{f}_0'' - f')^2} \\ \frac{a_2}{A} &= - \frac{2\bar{f}''^2 (S\bar{f}_0'' + \bar{f}''') (S\bar{f}_0'' - f')}{\bar{f}''^2 (S\bar{f}_0'' + \bar{f}''')^2 + f^2 (S\bar{f}_0'' - f')^2} \\ \frac{b_2}{A} &= \frac{2\bar{f}'' f (S\bar{f}_0'' - f')^2}{\bar{f}''^2 (S\bar{f}_0'' + \bar{f}''')^2 + f^2 (S\bar{f}_0'' - f')^2} \end{aligned}$$

and

$$\bar{f}'' = -if',$$

so that $\bar{f}''$ is real.

The incident wave is given by [see Eqs. (1) and (10)]

$$\begin{aligned} w &= A \cos f(x+ct) \\ \psi_x &= -AM' f^{-1} \sin f(x+ct) \\ \psi_y &= 0 \end{aligned} \quad (23)$$

The slow reflected wave [Eq. (11)] becomes

$$\begin{aligned}w &= A \cos f(x-ct+l_1) \\ \psi_x' &= -A M f^{-1} \sin f(x-ct+l_1) \\ \psi_y' &= 0\end{aligned}$$

and the fast reflected wave [Eq. (12)] is given by

$$\begin{aligned}w'' &= (a_1^2 + b_1^2)^{1/2} e^{-\tilde{f}'' x} \cos(f'' ct - l_2) \\ \psi_x'' &= (\tilde{f}'')^{-1} (S \delta_0^2 - \tilde{f}''^2) w'' \\ \psi_y'' &= 0\end{aligned}$$

In the above,

$$l_k = \arctan \frac{b_k}{a_k}, \quad k=1,2$$

We see that the slow reflected wave has the same amplitude, wave-length and velocity as the incident wave, but is out of phase with it. The fast reflected "wave" is, in fact, a vibration, the amplitude of which decreases exponentially as the distance from the edge of the plate increases. As $\dot{A}''' = 0$ by Eq. (22), no other waves are reflected.

Now consider

$$\underline{f^2 R S > R + S}$$

Equations (22) are then replaced by

$$\begin{aligned}\frac{A'}{A} &= \frac{f''(S\delta_0^2 - f'^2) + f(S\delta_0^2 - f^2)}{f''(S\delta_0^2 - f'^2) - f(S\delta_0^2 - f^2)} \\ \frac{A''}{A} &= \frac{2(S\delta_0^2 - f^2)f''}{f(S\delta_0^2 - f'^2) - f''(S\delta_0^2 - f^2)} \\ \frac{A'''}{A} &= 0\end{aligned}$$

The incident wave remains unchanged [Equations (23)], the slow reflected wave becomes

$$\begin{aligned} w' &= A' \cos f(x - ct) \\ \psi_x' &= -A' M' f' \sin f(x - ct) \\ \psi_y' &= 0, \end{aligned} \quad (26)$$

the fast reflected wave is given by

$$\begin{aligned} w' &= A'' \cos f''(x - c''t) \\ \psi_x' &= -A'' M'' (f'')' \sin f''(x - c''t) \\ \psi_y'' &= 0, \end{aligned} \quad (27)$$

and the reflected shear wave again vanishes. (The amplitude ratios A'/A and A''/A for this case are plotted versus h_f in Fig. 4.)

The incident wave is seen to give rise to two reflected flexural waves; we may examine the manner in which the energy per unit length of wave-front, per cycle, of the incident wave is distributed to the two reflected waves. Using the expressions for energy given in Reference [4], we find

$$\begin{aligned} \frac{E'}{E} &= \left(\frac{A'}{A}\right)^2 \\ \frac{E''}{E} &= \left(\frac{A''}{A}\right)^2 \left(\frac{f'}{f''}\right)^2 \frac{\bar{E}_0^2 + M''^2}{\bar{E}_0^2 + M'^2}. \end{aligned}$$

Here E , E' and E'' are, respectively, the energy per cycle, per unit length of wave front of the incident wave, for the incident, slow reflected and fast reflected waves. The ratios E'/E and E''/E are plotted versus h_f in Fig. 5.

The motion corresponding to

$$\frac{1}{2}RS = R+S$$

may be found by proceeding to the limit, as $\frac{1}{2}RS \rightarrow R+S$, in either of the preceding cases. By either method it may be verified that an incident wave

$$\begin{aligned} w &= A \cos f(x+ct) \\ \psi_x &= A \sqrt{\frac{R}{S(R+S)}} \sin f(x+ct) \\ \psi_y &= 0 \end{aligned} \quad (28)$$

now gives rise to a slow reflected wave,

$$\begin{aligned} w' &= -A \cos f(x-ct) \\ \psi'_x &= -A \sqrt{\frac{R}{S(R+S)}} \sin f(x-ct) \\ \psi'_y &= 0 \end{aligned} \quad (29)$$

and to a thickness-shear vibration,

$$\begin{aligned} w'' &= 0 \\ \psi''_x &= 2A \sqrt{\frac{S}{R(R+S)}} \sin fct \\ \psi''_y &= 0 \end{aligned} \quad (30)$$

The circular frequency ρ for this vibration is [see Eqs. (3), (5), (6), (8), (9)]

$$\rho = \kappa c_s \frac{\sqrt{12}}{h}$$

Now, the circular frequency, $\bar{\rho}$, of the first antisymmetric mode of thickness-shear vibration of an infinite plate, according to the three-dimensional theory

of elasticity, is

$$\bar{\rho} = \frac{\tau G}{h}$$

Thus it seems appropriate to let

$$\kappa = \frac{T}{\sqrt{2}}$$

so that the thickness-shear vibration in the present case will occur at the frequency predicted by exact theory for an infinite plate of the same thickness.

It is interesting to notice that the motion here being considered, is one of the possible modes of motion of an infinite strip of thickness h and width b (see Reference [3]), provided

$$b = \frac{h n \pi}{[3(R/S+1)]^{1/2}}, \quad n = 1, 2, \dots$$

Hence, for $f R S = R + S$, i.e., for $\rho = \bar{\rho}$, the semi-infinite plate may be regarded as consisting of an infinite number of independently vibrating strips, each with its infinite dimension parallel to the edge of the plate.

To conclude the discussion of normal incidence, we consider two limiting cases:

When the wave-length of the incident wave is large in comparison with the thickness of the plate, we find, by proceeding to the limit in Equations (22)-(25), as $h f \rightarrow 0$,

$$\begin{aligned} w &= A \cos f(x+ct) \\ \psi_x &= A f \sin f(x+ct) \\ \psi_y &= 0 \end{aligned}$$

$$\begin{aligned}
w' &= -A \sin f(x-ct) \\
\psi_x' &= -A f \cos f(x-ct) \\
\psi_y' &= 0
\end{aligned}$$

$$\begin{aligned}
w'' &= A e^{-\delta x} (\cos fct + \sin fct) \\
\psi_x'' &= -A f e^{-\delta x} (\cos fct + \sin fct) \\
\psi_y'' &= 0
\end{aligned}$$

$$\begin{aligned}
w''' &= 0 \\
\psi_x''' &= 0 \\
\psi_y''' &= 0
\end{aligned}$$

These expressions show that the incident wave is reflected without change in amplitude or phase velocity, and that a vibration, confined primarily to a region near the edge of the plate, takes place. The classical theory of plates (whose range of applicability is restricted to the limiting case under consideration) predicts the same results.

For wave-lengths which are very small in comparison with the plate thickness³, we let h_f approach infinity in Equations (26), (27) with the result

$$\begin{aligned}
w &= A \cos f(x+ct), \quad \psi_x = \psi_y = 0 \\
w' &= A \cos f(x-ct), \quad \psi_x' = \psi_y' = 0 \\
w'' &= \psi_x'' = \psi_y'' = 0 \\
w''' &= \psi_x''' = \psi_y''' = 0
\end{aligned}$$

3. The theory is not expected to be good for very short waves. This limiting case is included for the sake of completeness.

The total motion is

$$w^* = 2A \cos \phi x \cos \phi c t$$

which is a standing wave.

7. Oblique Incidence

We have seen that fundamentally different states of motion obtain according as the circular frequency of the incident wave is less than, equal to, or greater than the circular frequency of thickness-shear vibration of an infinite plate. Hence, for $0 < \alpha < \pi/2$ we shall again examine separately the cases $\rho < \bar{\rho}$, $\rho > \bar{\rho}$ and $\rho = \bar{\rho}$.

$$\underline{\rho < \bar{\rho}}$$

As $\rho < \bar{\rho}$ is equivalent to $R S_0^* < 1$, we see from Equations (20) and (21) that ϕ^H and ϕ^N are imaginary.

Setting

$$\tilde{\phi}'' = -i\phi^H, \tilde{\phi}''' = -i\phi^N$$

we get, from Equation (17),

$$\sin \alpha^j = -\frac{c_j}{c} \sin \alpha, \quad j = H, N$$

whence

$$\cos \alpha^j = [1 + \left(\frac{c_j}{c}\right)^2 \sin^2 \alpha]^{1/2} = m^j, \text{ say.}$$

(For all values of α and j we have, as noted earlier, $\phi^H \phi^N = c^2, \alpha^H = \alpha$)

Substituting into the general expressions for the various reflected waves [Equations (11), (12), (13)], we get from Equations (1),

$$\begin{aligned} W' &= A' \exp[i\gamma(x \cos \alpha + \xi)] \\ \psi_x' &= W' M' f' \cos \alpha \\ \psi_y' &= W' M' f' i \sin \alpha \end{aligned}$$

$$\begin{aligned} W'' &= A'' \exp(i\gamma \xi - \tilde{f}'' m'' x) \\ \psi_x'' &= W'' M'' m'' \\ \psi_y'' &= -W'' M'' (\tilde{f}'')' f \sin \alpha \end{aligned} \quad (31)$$

$$\begin{aligned} W''' &= 0 \\ \psi_x''' &= A''' i\gamma \sin \alpha \exp(i\gamma \xi - \tilde{f}''' m''' x) \\ \psi_y''' &= A''' \tilde{f}''' m''' \exp(i\gamma \xi - \tilde{f}''' m''' x) \end{aligned}$$

where

$$\xi = y \sin \alpha - ct$$

These expressions are valid for all α with the possible exception of $\alpha = \pi/2$, i.e., "grazing" incidence. For that case, the solution of Equations (18) is

$$\frac{A'}{A} = -1, \quad A'' = A''' = 0$$

and we get, in place of Equations (31),

$$W' = -W, \psi_x' = -\psi_x, \psi_y' = -\psi_y$$

$$W'' = \psi_x'' = \psi_y'' = 0, j = ", "'$$

so that the entire motion is then [see Eq. (14)]

$$W^* = \psi_x^* = \psi_y^* = 0. \quad (32)$$

This complete disappearance of the motion is analogous to the case of grazing incidence in the reflection of plane waves from the plane boundary of a semi-infinite solid. We shall return to this question later on.

The waves described by Equations (31) are

- ① - a slow flexural wave, reflected at an angle equal to the angle of incidence of the incident wave,
- ② - a fast flexural wave of amplitude decreasing exponentially with departure from the edge, and propagated in a direction parallel to the edge,
- ③ - a shear wave of exponentially decreasing amplitude, propagated along the edge. (See Fig. 6.)

Turning now to the case

$$\mu > \bar{\mu}$$

we note that, from Equations (20) and (21)

$$\frac{\psi''}{\psi} < 1, \frac{\psi'''}{\psi} < 1 \quad \text{and} \quad \frac{\psi''}{\psi} < \frac{\psi'''}{\psi} \quad \text{for all } \psi.$$

(Fig. 8 illustrates these facts for $\mu = 1/4$.)

Equation (17) permits us to construct a table which shows the relationship between α and α' , α'' , α''' in terms of trigonometric functions of these angles.

$\sin \alpha = 0$	$\sin \alpha = 0$	$\sin \alpha' = 0$	$\cos \alpha' = 1$	$\sin \alpha < \sin \alpha' < \sin \alpha''$	$0 < \cos \alpha' < 1$
$0 < \sin \alpha < \frac{\ell''}{\ell}$	$0 < \sin \alpha < \frac{\ell''}{\ell}$	$\sin \alpha < \sin \alpha' < 1$	$0 < \cos \alpha' < 1$	$\sin \alpha < \sin \alpha' < \sin \alpha''$	$0 < \cos \alpha' < 1$
$\sin \alpha = \frac{\ell''}{\ell}$	$\sin \alpha' = \frac{\ell''}{\ell}$	$\sin \alpha' = 1$	$\cos \alpha' = 0$	$\sin \alpha < \sin \alpha' = \frac{\ell''}{\ell} < 1$	$0 < \cos \alpha' < 1$
$\frac{\ell''}{\ell} < \sin \alpha < \frac{\ell'''}{\ell}$	$\frac{\ell''}{\ell} < \sin \alpha < \frac{\ell'''}{\ell}$	$\sin \alpha' > 1$	$\cos \alpha' = \text{imag.}$	$\sin \alpha < \sin \alpha' < 1$	$0 < \cos \alpha' < 1$
$\sin \alpha = \frac{\ell'''}{\ell}$	$\sin \alpha' = \frac{\ell'''}{\ell}$	$1 < \sin \alpha' < \frac{\ell'''}{\ell}$	$\cos \alpha' = \text{imag.}$	$\sin \alpha' = 1$	$\cos \alpha' = 0$
$\frac{\ell'''}{\ell} < \sin \alpha < 1$	$\frac{\ell'''}{\ell} < \sin \alpha < 1$	$\sin \alpha' > 1$	$\cos \alpha' = \text{imag.}$	$1 < \sin \alpha'$	$\cos \alpha' = \text{imag.}$
$\sin \alpha = 1$	$\sin \alpha' = 1$	$1 < \sin \alpha' < \frac{\ell'''}{\ell}$	$\cos \alpha' = \text{imag.}$	$1 < \sin \alpha' = \frac{\ell'''}{\ell}$	$\cos \alpha' = \text{imag.}$

With the help of this table and Equations (10)-(13) we can now describe the motion associated with various angles of incidence.

For $0 \leq \sin \alpha < \frac{\ell''}{\ell}$, the reflected waves are

- ① - a slow flexural wave,
- ② - a fast flexural wave,
- ③ - a shear wave.

These emerge at angles α' , α'' , α''' with

$$\alpha = \alpha' < \alpha''' < \alpha'' < \pi/2$$

as shown in Fig. 7. For instance, with $h\ell = 5$, we have $\ell''/\ell = .302$, $\ell'''/\ell = .593$. Taking $\alpha = 15^\circ$ (so that $\sin \alpha < .302$) we get

$$\alpha' = 15^\circ$$

$$\alpha'' = \arcsin \frac{.259}{.302} = 59.1^\circ$$

$$\alpha''' = \arcsin \frac{.259}{.593} = 25.9^\circ$$

For $\sin \alpha = \frac{\ell''}{\ell}$, the amplitude ratios [see Eqs. (18)] are

$$\frac{A'}{A} = 1, \quad \frac{A''}{A} = \frac{2(5\delta_0^4 - \ell^2)}{\nu(5\delta_0^4 - \ell^2)} \left[\left(\frac{\ell''}{\ell} \right)^2 (1 - \nu) - 1 \right], \quad \frac{A'''}{A} = 0.$$

Thus we have only two reflected waves, (see Fig. 9),

① - a slow flexural wave whose angle of emergence is equal to the angle of incidence of the incident wave,

② - a fast flexural wave propagated along the free edge.

With $h_f = 5$, as in the numerical example above, the critical angle, α_c , for which the shear wave vanishes, is

$$\alpha_c = \arcsin(.302) = 17.6^\circ$$

Letting α increase further, until $\delta'/\delta < \sin \alpha < \delta''/\delta$, we find that the fast flexural wave [Equations (12)] changes in character since $\cos \alpha''$ is now imaginary. The shear wave reappears and we have (Fig. 10)

① - a slow flexural wave,

② - an exponentially decaying fast flexural wave, propagated parallel to the edge,

③ - a shear wave emerging at an angle, α'' , with $\alpha < \alpha'' < \pi/2$.

Another critical value of α is reached when $\sin \alpha = \delta''/\delta$. We then find (Fig. 11)

① - a slow flexural wave,

② - an exponentially decaying fast flexural wave,

③ - a shear wave.

Both ② and ③ are now propagated along the edge. With $h_f = 5$, the critical value for α is 36.4° .

Finally, for $\delta''/\delta < \sin \alpha < 1$, the shear wave also acquires an exponentially decaying amplitude, giving us (Fig. 12)

① - a slow flexural wave,

② - an exponentially decaying fast flexural wave,

③ - an exponentially decaying shear wave.

Grazing incidence, i.e., $\alpha = 90^\circ$, leads to vanishing motion, as it did for $p < \bar{p}$.

To complete the discussion of oblique incidence, we consider the case

$$\underline{p = \bar{p}}$$

As the circular frequency of the incident wave approaches the thickness-shear frequency, the amplitudes of the fast reflected wave and of the reflected shear wave become infinite. The resonance encountered here is due to the fact that reflection at the edge of the plate is equivalent to "forcing" the plate at a frequency equal to a "natural" frequency. For, it may easily be verified that the vibration

$$\begin{aligned} W = \psi_x &= 0 \\ \psi_y &= B_2 e^{i\beta t} \end{aligned}$$

is a motion which satisfies the equations of motion (2) and leaves the edge $x=0$ free of traction.

8. Grazing Incidence

We have seen [Eq. (32)] that the wave motion in a semi-infinite plate disappears as the angle of incidence approaches 90° . A similar situation obtains in the case of waves reflected from the plane boundary of a semi-infinite solid. By application of a suitable limiting process, wave motions for the latter case have recently been found by Goodier and Bishop [4]. A similar limiting process for the present case will now be considered.

Letting

$$\epsilon = \pi/2 - \alpha$$

and neglecting higher powers of ϵ , we have, from Equations (17),

$$\sin \alpha_j = \frac{\epsilon}{f_j} (1 - \frac{\epsilon^2}{2}), \quad j = 1, 2.$$

Substituting into Equations (12), the amplitude ratios are found to be of the form

$$\frac{A'}{A} = -(1 + a'\epsilon), \quad \frac{A''}{A} = a'\epsilon, \quad \frac{A'''}{A} = a''\epsilon \quad (33)$$

Expanding the exponentials which appear in the expressions for displacements, we get from Equations (10)-(13) and Equation (1),

$$\begin{aligned} w &= A(1 - i\epsilon x) \exp(i\epsilon \xi) \\ w' &= A'(1 + i\epsilon x) \exp(i\epsilon \xi) \\ w'' &= A'' \exp(i\epsilon \xi - x f'') \\ w''' &= A''' \exp(i\epsilon \eta - x f''') \end{aligned} \quad (34)$$

where

$$\begin{aligned} \xi &= y - zt \\ f^j &= [c^2 - (f^j)^2]^{1/2}, \quad j = ', ', '''. \end{aligned}$$

These expansions are valid only when higher powers of ϵx can be neglected, that is, for a range of distances from the edge which is small in comparison with the wave-length of the incident wave.

From Equations (14), (32) and (34), the total motion is characterized by the expressions

$$w^* = A\epsilon [-a' - 2i\epsilon x + a'' e^{-f'' x}] \exp(i\epsilon \xi) \quad (35)$$

$$w''' = A\epsilon a''' \exp(i\epsilon \eta - x f'''). \quad (36)$$

If we now permit A to become infinite as ϵ approaches zero, and let the approach to infinity be such that the product $A\epsilon$ remains finite, we see that the first term of Equation (35) represents an incident flexural wave while the second term corresponds to the " P_y " wave found by Goodier and Bishop. The third term and Equation (36) indicate, respectively, a flexural and a shear wave, each propagated along the edge of the plate, and each having an exponentially increasing or decreasing amplitude, according as ρ is less than or greater than $\bar{\rho}$; for $\rho = \bar{\rho}$ we again have resonance.

Acknowledgment

The author wishes to thank Professor R. D. Mindlin of Columbia University both for suggesting this problem, and for his helpful advice in its solution.

Bibliography

1. R. D. Mindlin, "Influence of Rotatory Inertia and Shear on Flexural Motions of Isotropic, Elastic Plates," Journal of Applied Mechanics, Vol. 18 (1951), pp. 31-38.
2. A. E. H. Love, Theory of Elasticity, (Cambridge University Press, London, 1927), 4th Ed., p. 176.
3. R. D. Mindlin, "Thickness-Shear and Flexural Vibrations of Crystal Plates," Journal of Applied Physics, Vol. 22 (1951), pp. 316-323.
4. J. N. Goodier and R. E. D. Bishop, "A Note on Critical Reflections of Elastic Waves at Free Surfaces," Journal of Applied Physics, Vol. 23 (1952), pp. 124-126.




Fig. 1




Fig. 2: Incident and Emergent Wave Normals.




Fig. 3: Relationship between Phase Velocity and Wave Number.




Fig. 4




Fig. 5: Energy Partition - Normal Incidence.




Fig. 6: Oblique Incidence, $p < \bar{p}$.




Fig. 7: Oblique Incidence, $p > \bar{p}$, $0 \leq \sin \alpha < r'/r$.




Fig. 8: Wave Number Ratios.




Fig. 9: Oblique Incidence, $p > \bar{p}$, $\sin \alpha = \gamma'/\gamma$.




Fig. 10: Oblique Incidence, $p > \bar{p}$, $\gamma'\gamma < \sin \alpha < \gamma''/\gamma$.




Fig. 11: Oblique Incidence, $p > \bar{p}$, $\sin \alpha = \bar{v}/v$.




Fig. 12: Oblique Incidence, $p > \bar{p}$, $\bar{v}/v < \sin \alpha < 1$.

DISTRIBUTION LIST

for

Technical and Final Reports Issued Under
Office of Naval Research Project NR-064-388. Contract Nonr-266(C9)

Administrative, Reference and Liaison Activities of ONR

Chief of Naval Research Department of the Navy Washington 25, D.C. Attn: Code 438	(2)	Commanding Officer Office of Naval Research Branch Office 1000 Geary Street San Francisco 24, California	(1)
Director, Naval Research Lab. Washington 25, D.C. Attn: Tech. Info. Officer	(9)	Commanding Officer Office of Naval Research Branch Office	
Technical Library	(1)	1030 Green Street	
Mechanics Division	(2)	Pasadena, California	(1)
Code 3834 (J. P. Walsh)	(1)		
Commanding Officer Office of Naval Research Branch Office 150 Causeway Street Boston 10, Massachusetts	(1)	Contract Administrator, SE Area Office of Naval Research c/o George Washington University 707 22nd Street, N.W. Washington 6, D.C.	(1)
Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, New York	(1)	Officer in Charge Office of Naval Research Branch Office, London Navy No. 100 FPO, New York, N.Y.	(5)
Commanding Officer Office of Naval Research Branch Office 844 N. Rush Street Chicago 11, Illinois	(1)	Library of Congress Washington 25, D.C. Attn: Navy Research Section	(2)

Department of Defense

Other Interested Government Activities

GENERAL		GENERAL (cont.)	
Research and Development Board Department of Defense Pentagon Building Washington 25, D.C. Attn: Library (Code 3D-1075)	(1)	Joint Task Force 3 12 St. & Const. Ave., N.W. (Temp. U) Washington 25, D.C. Attn: Major B. D. Jones	(1)
Armed Forces Special Weapons Project P.O. Box 2610 Washington, D.C. Attn: Col. G. F. Blunda	(1)	ARMY Chief of Staff Department of the Army Research and Development Division Washington 25, D.C. Attn: Chief of Res. and Dev.	(1)

ARMY (cont.)

Office of the Chief of Engineers
Assistant Chief for Works
Department of the Army
Bldg. T-7, Gravelly Point
Washington 25, D.C.
Attn: Structural Branch
(R. L. Bloor) (1)

Office of the Chief of Engineers
Asst. Chief for Military Construction
Department of the Army
Bldg. T-7, Gravelly Point
Washington 25, D.C.
Attn: Structures Branch
(H. F. Carey) (1)

Office of the Chief of Engineers
Asst. Chief for Military Operations
Department of the Army
Bldg. T-7, Gravelly Point
Washington 25, D.C.
Attn: Structures Development Branch
(W. F. Woodlard) (1)

Engineering Research & Development Lab.
Fort Belvoir, Virginia
Attn: Structures Branch (1)

The Commanding General
Sandia Base, P.O. Box 5100
Albuquerque, New Mexico
Attn: Col. Canterbury (1)

Operations Research Officer
Department of the Army
Ft. Lesley J. McNair
Washington 25, D.C.
Attn: Howard Brackney (1)

Office of Chief of Ordnance
Research & Development Service
Department of the Army
The Pentagon
Washington 25, D.C.
Attn: ORDTB (2)

Commanding Officer
Ballistic Research Laboratory
Aberdeen Proving Ground
Aberdeen, Maryland
Attn: Dr. C. W. Lampson (1)

ARMY (cont.)

Commanding Officer
Watertown Arsenal
Watertown, Massachusetts
Attn: Laboratory Division (1)

Commanding Officer
Frankford Arsenal
Philadelphia, Pennsylvania
Attn: Laboratory Division (1)

Commanding Officer
Squier Signal Laboratory
Fort Monmouth, New Jersey
Attn: Components and Materials
Branch (1)

NAVY
Chief of Bureau of Ships
Navy Department
Washington 25, D.C.
Attn: Director of Research (2)

Director
David Taylor Model Basin
Washington 7, D.C.
Attn: Structural Mechanics Div. (2)

Director
Naval Engr. Experiment Station
Annapolis, Maryland (1)

Director
Materials Laboratory
New York Naval Shipyard
Brooklyn 1, New York (1)

Chief of Bureau of Ordnance
Navy Department
Washington 25, D.C.
Attn: Ad-3, Technical Library (1)

Superintendent
Naval Gun Factory
Washington 25, D.C. (1)

Naval Ordnance Laboratory
White Oak, Maryland
RFD 1, Silver Spring, Maryland
Attn: Mechanics Division (2)

Naval Ordnance Test Station
Inyokern, California
Attn: Scientific Officer (1)

NAVY (cont.)

Naval Ordnance Test Station
Underwater Ordnance Division
Pasadena, California
Attn: Structures Division (1)
Physics Division (1)

Chief of Bureau of Aeronautics
Navy Department
Washington 25, D.C.
Attn: TD-41, Technical Library (1)

Naval Air Experimental Station
Naval Air Materiel Center
Naval Base
Philadelphia 12, Pennsylvania
Attn: Head, Aeronautical Materials Laboratory (1)

Chief of Bureau of Yards & Docks
Navy Department
Washington 25, D.C.
Attn: Code P-314 (1)

Officer in Charge
Naval Civil Engr. Research and Eval.
Laboratory
Naval Station
Port Hueneme, California (1)

Superintendent, U.S. Naval Post
Graduate School
Annapolis, Maryland (1)

Commander
U.S. Naval Proving Grounds
Dahlgren, Virginia (1)

AIR FORCES
Commanding General
U.S. Air Forces
The Pentagon
Washington 25, D.C.
Attn: Research & Development Division (1)

Commanding General
Air Materiel Command
Wright-Patterson Air Force Base
Dayton, Ohio
Attn: MOREX-B (E. H. Schwartz) (1)

AIR FORCES (cont.)

Office of Air Research
Wright-Patterson Air Force Base
Dayton, Ohio
Attn: Chief, Applied Mechanics Group (1)

OTHER GOVERNMENT AGENCIES
U.S. Atomic Energy Commission
Division of Research
Washington, D.C. (1)

Argonne National Laboratory
P.O. Box 5207
Chicago 80, Illinois (1)

Director
National Bureau of Standards
Washington, D.C.
Attn: Dr. W. H. Ramberg (1)

U.S. Coast Guard
1300 E Street, N.W.
Washington, D.C.
Attn: Chief, Testing & Development Division (1)

Forest Products Laboratory
Madison, Wisconsin
Attn: L. J. Markwardt (1)

National Advisory Committee for
Aeronautics
1724 F Street, N.W.
Washington, D.C. (1)

National Advisory Committee for
Aeronautics
Langley Field, Virginia
Attn: Dr. E. Lundquist (1)

National Advisory Committee for
Aeronautics
Cleveland Municipal Airport
Cleveland, Ohio
Attn: J. H. Collins, Jr. (1)

U.S. Maritime Commission
Technical Bureau
Washington, D.C.
Attn: Mr. V. Russo (1)

Contractors and Other Investigators
Actively Engaged in Related Research

Professor J. R. Andersen Towne School of Engineering University of Pennsylvania Philadelphia, Pennsylvania	(1)	Dr. G. F. Carrier Graduate Division of Applied Mathematics Brown University Providence, Rhode Island	(1)
Professor Lynn Beedle Fritz Engineering Laboratory Lehigh University Bethlehem, Pennsylvania	(1)	Mrs. Hilda L. Cooper 150 Ravine Avenue Yonkers, New York	(1)
Professor C. B. Biezeno Technische Hoogeschool Nieuw Laan 76 Delft, Holland	(1)	Dr. Antoine E. I. Craya Neyrpic Boite, Postale 52 Grenoble, France	(1)
Professor M. A. Biot 1819 Broadway New York, New York	(1)	Professor J. P. Den Hartog Massachusetts Institute of Technology Cambridge 39, Massachusetts	(1)
Professor R. L. Bisplinghoff Massachusetts Institute of Technology Dept. of Aeronautical Engineering Cambridge 39, Massachusetts	(1)	Dr. Herbert Deresiewicz Dept. of Civil Engineering Columbia University 632 West 125th Street New York 27, New York	(1)
Dr. Hans H. Bleich Dept. of Civil Engineering Columbia University New York 27, New York	(1)	Dr. C. O. Dohrenwend Rensselaer Polytechnic Institute Troy, New York	(1)
Professor J. A. Bogdanoff Furdue University Lafayette, Indiana	(1)	Professor T. J. Dolan Dept. of Theoretical and Applied Mechanics University of Illinois Urbana, Illinois	(1)
Professor B. A. Boley Dept. of Civil Engineering Columbia University New York 27, New York	(1)	Professor Lloyd Donnell Dept. of Mechanics Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)
Professor P. W. Bridgeman Dept. of Physics Harvard University Cambridge, Massachusetts	(1)	Professor D. C. Drucker Brown University Providence, Rhode Island	(1)
Professor D. M. Burmister Dept. of Civil Engineering Columbia University New York 27, New York	(1)	Dr. W. Eckert Watson Scientific Computing Laboratory 612 West 116th Street New York 27, New York	(1)
Dr. V. Cadambe Assistant Director of the National Physical Laboratory of India New Delhi 12, India	(1)	Dr. H. Ekstein Armour Research Foundation Illinois Institute of Technology Chicago 16, Illinois	(1)

Contractors and Other Investigators Actively Engaged in Related Research (cont.)

Engineering Library Columbia University New York 27, New York	(1)	Dr. L. E. Goodman Dept. of Civil Engineering University of Illinois Urbana, Illinois	(1)
Professor E. L. Eriksen University of Michigan Ann Arbor, Michigan	(1)	Dr. R. J. Hansen Massachusetts Institute of Technology Dept. of Civil & Sanitary Engineering Cambridge 39, Massachusetts	(1)
Professor A. C. Eringen Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor R. M. Hermes University of Santa Clara Santa Clara, California	(1)
Dr. W. L. Esmeljer Technische Hoogeschool Nieuwe Laan 76 Delft, Holland	(1)	Professor G. Herrmann Dept. of Civil Engineering Columbia University New York 27, New York	(1)
Professor A. M. Freudenthal Dept. of Civil Engineering Columbia University New York 27, New York	(2)	Professor M. Hetényi Northwestern University The Technological Institute Evanston, Illinois	(1)
Professor B. Fried Washington State College Pullman, Washington	(1)	Professor T. J. Higgins Dept. of Electrical Engineering University of Wisconsin Madison 6, Wisconsin	(1)
Professor K. O. Friedrichs New York University Washington Square New York, New York	(1)	Dr. N. J. Hoff, Head Dept. of Aeronautical Engineering & Applied Mechanics Polytechnic Institute of Brooklyn 99 Livingston Street Brooklyn 2, New York	(1)
Professor M. M. Frocht Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor M. E. Hagan University of Utah Salt Lake City, Utah	(1)
Professor J. M. Garrelts Dept. of Civil Engineering Columbia University New York 27, New York	(1)	Professor D. L. Holl Iowa State College Ames, Iowa	(1)
Dean J. A. Goff University of Pennsylvania Philadelphia, Pennsylvania	(1)	Dr. J. H. Hollomon General Electric Research Labs. 1 River Road Schenectady, New York	(1)
Mr. Martin Goland Midwest Research Institute 4049 Pennsylvania Avenue Kansas City 2, Missouri	(1)	Dr. W. H. Hopmann Dept. of Applied Mechanics John Hopkins University Baltimore, Maryland	(1)
Dr. J. N. Goodier Dept. of Mechanical Engineering Stanford University Stanford, California			

Contractors and Other Investigators Actively Engaged in Related Research (cont.)

Institut de Mathematiques Universite post. fax 55 Skopje, Yugoslavia	(1)	Professor C. T. G. Looney Dept. of Civil Engineering Yale University New Haven, Connecticut	(1)
Professor L. S. Jacobsen Dept. of Mechanical Engineering Stanford University Stanford, California	(1)	Dr. J. L. Lubkin Midwest Research Institute 4049 Pennsylvania Avenue Kansas City 2, Missouri	(1)
Professor Bruce G. Johnston University of Michigan Ann Arbor, Michigan	(1)	Professor J. F. Ludloff School of Aeronautics New York University New York 53, New York	(1)
Professor R. Klotter Stanford University Stanford, California	(1)	Professor J. N. Macduff Rensselaer Polytechnic Institute Troy, New York	(1)
Professor W. J. Krefeld Dept. of Civil Engineering Columbia University New York 27, New York	(1)	Professor C. W. MacGregor University of Pennsylvania Philadelphia, Pennsylvania	(1)
Professor B. J. Lavan Dept. of Materials Engineering University of Minnesota Minneapolis, Minnesota	(1)	Professor Lawrence E. Malvern Dept. of Mathematics Carnegie Institute of Technology Pittsburgh 13, Pennsylvania	(1)
Dr. E. H. Lee Division of Applied Mathematics Brown University Providence, Rhode Island	(1)	Dr. J. H. Marchant Brown University Providence, Rhode Island	(1)
Professor George Lee Rensselaer Polytechnic Institute Troy, New York	(1)	Professor J. Marin Pennsylvania State College State College, Pennsylvania	(1)
Professor J. M. Lessells Massachusetts Institute of Technology Cambridge 39, Massachusetts	(1)	Dr. W. P. Mason Bell Telephone Laboratories Murray Hill, New Jersey	(1)
Library, Engineering Foundation 29 West 39th Street New York, New York	(1)	Professor R. D. Mindlin Dept. of Civil Engineering Columbia University 632 West 125th Street New York 27, New York	(15)
Professor Paul Lieber Dept. of Engineering Rensselaer Polytechnic Institute Troy, New York	(1)	Dr. A. Nadai 136 Cherry Valley Road Pittsburgh 21, Pennsylvania	(1)
Dr. Hsu Lo Purdue University Lafayette, Indiana	(1)	Professor Paul M. Naghdi Dept. of Engineering Mechanics University of Michigan Ann Arbor, Michigan	(1)

Contractors and Other Investigators Actively Engaged in Related Research (cont.)

Professor N. M. Newmark Dept. of Civil Engineering University of Illinois Urbana, Illinois	(1)	Dr. S. Payner Armour Research Foundation Illinois Institute of Technology Chicago 16, Illinois	(1)
Professor Jesse Ormondroyd University of Michigan Ann Arbor, Michigan	(1)	Professor E. Reissner Dept. of Mathematics Massachusetts Institute of Technology Cambridge 39, Massachusetts	(1)
Dr. W. Osgood Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor H. Reissner Polytechnic Institute of Brooklyn 99 Livingston Street Brooklyn 2, New York	(1)
Dr. George B. Pegram Committee on Government Aided Research 313 Low Memorial Library Columbia University New York 27, New York	(1)	Dr. Kenneth Robinson National Bureau of Standards Washington, D.C.	(1)
Dr. R. P. Petersen Director, Applied Physics Division Sandia Laboratory Albuquerque, New Mexico	(1)	Professor M. A. Sadowsky Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)
Mr. R. E. Petersen Westinghouse Research Laboratories East Pittsburgh, Pennsylvania	(1)	Professor M. J. Salvadori Dept. of Civil Engineering Columbia University New York 27, New York	(1)
Dr. A. Phillips School of Engineering Stanford University Stanford, California	(1)	Dr. F. S. Shaw Polytechnic Institute of Brooklyn 99 Livingston Street Brooklyn 2, New York	(1)
Professor Gerald Pickett Dept. of Mechanics University of Wisconsin Madison 6, Wisconsin	(1)	Dr. Daniel T. Sigley Applied Physics Laboratory John Hopkins University 8621 Georgia Avenue Silver Spring, Maryland	(1)
Dr. H. Peritzky General Electric Research Labs. Schenectady, New York	(1)	Dr. C. B. Smith Department of Mathematics Walker Hall University of Florida Gainesville, Florida	(1)
Dr. W. Prager Graduate Division of Applied Mathematics Brown University Providence, Rhode Island	(1)	Professor C. R. Soderberg Massachusetts Institute of Technology Cambridge 39, Massachusetts	(1)
RAND Corporation 1500 4th Street Santa Monica, California Attn: Dr. D. L. Judd	(1)	Professor R. V. Southwell Imperial College of Science and Technology South Kensington London S.W. 7, England	(1)

Contractors and Other Investigators Actively Engaged in Related Research (cont.)

Professor E. Sternberg Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor E. Volterra Rensselaer Polytechnic Institute Troy, New York	(1)
Professor J. J. Stoker New York University Washington Square New York, New York	(1)	Mr. A. M. Wahl Westinghouse Research Laboratories East Pittsburgh, Pennsylvania	(1)
Mr. R. A. Sykes Bell Telephone Laboratories Murray Hill, New Jersey	(1)	Professor C. T. Wang Dept. of Aeronautical Engineering New York University University Heights, Bronx New York, New York	(1)
Professor P. S. Symonds Brown University Providence, Rhode Island	(1)	Dr. R. L. Wegel RFD 2 Peekskill, New York	(1)
Professor J. L. Synge Dublin Institute for Advanced Studies School of Theoretical Physics 64-65 Merrion Square Dublin, Ireland	(1)	Professor E. E. Weibel University of Colorado Boulder, Colorado	(1)
Professor F. K. Teliepsmann Dept. of Aeronautical Engineering New York University University Heights, Bronx New York, New York	(1)	Dr. Alexander Weinstein Institute of Applied Mathematics University of Maryland College Park, Maryland	(1)
Professor S. F. Timoshenko School of Engineering Stanford University Stanford, California	(1)	Professor Dana Young Yale University New Haven, Connecticut	(1)
Dr. C. A. Truesdell Graduate Institute for Applied Mathematics Indiana University Bloomington, Indiana	(1)		
Professor Karl S. Van Dyke Department of Physics Scott Laboratory Wesleyan University Middletown, Connecticut	(1)		
Dr. I. Vigness Naval Research Laboratory Anacostia Station Washington, D.C.	(1)		
Dr. Leonardo Villena Gran Via J. Antonio 6 Madrid, Spain	(1)		