

UNCLASSIFIED

AD NUMBER
AD009911
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; SEP 1951. Other requests shall be referred to Office of Naval Research, 800 North Quincy Street, Arlington, VA 22217-5660.
AUTHORITY
ONR ltr, 26 Oct 1977

THIS PAGE IS UNCLASSIFIED

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

ON THE TWO-CENTER EXCHANGE INTEGRALS*

* This work was assisted by the ONR under Task Order IX of Contract N6601 with The University of Chicago.

K. Rüdenberg
Department of Physics
The University of Chicago
Chicago 37, Illinois

In a recent discussion¹ it was shown that the two-center exchange integrals are most conveniently expressed in terms of certain auxiliary functions, $B_j^{MQ}(\beta)$ and $\phi_{nn}^{ML}(\alpha, \bar{\alpha})$. Of these, the functions ϕ_{nn}^{MQ} are the more complicated ones, and it was felt that a condensed account of the recurrence method of their computation would be useful. In the following this method is therefore presented in the form of an instruction for actual work. At the same time, some minor modifications have been made.

First step: Compute the functions

$$A_0(\alpha), A_1(\alpha), A_2(\alpha), \dots, A_N(\alpha);$$

$$A_0(\bar{\alpha}), A_1(\bar{\alpha}), A_2(\bar{\alpha}), \dots, A_N(\bar{\alpha});$$

$$A_0(\alpha + \bar{\alpha}), A_1(\alpha + \bar{\alpha}), A_2(\alpha + \bar{\alpha}), \dots, A_{N+N}(\alpha + \bar{\alpha});$$

by means of the recurrence procedure

$$A_0(x) = e^{-x}/x, \quad A_n(x) = (1/x) [nA_{n-1}(x) + e^{-x}].$$

For the arguments: $x = 0.25, 0.5, 0.75, \dots$, the value of $A_n(x)$

¹ Cf. K. Rüdenberg, J. Chem. Phys., 19, 1459 (1951).

can be taken from Kotani's Table.

The values of the highest indices, i.o., N , $\bar{N}$, are determined by the recurrence procedure which follows.

Second step: Compute the functions

$$G_0(\alpha), G_1(\alpha), G_2(\alpha), \dots, G_N(\alpha);$$

$$G_0(\bar{\alpha}), G_1(\bar{\alpha}), G_2(\bar{\alpha}), \dots, G_{\bar{N}}(\bar{\alpha});$$

$$G_0(\alpha + \bar{\alpha}), G_1(\alpha + \bar{\alpha}), G_2(\alpha + \bar{\alpha}), \dots, G_{N+\bar{N}}(\alpha + \bar{\alpha});$$

by means of the recurrence procedure

$$G_0(x) = \frac{1}{2}e^{-x} \left[C + \log 2x - e^{2x} \text{Ei}(-2x) \right],$$

$$G_1(x) = \frac{1}{2}e^{-x} \left[C + \log 2x + e^{2x} \text{Ei}(-2x) \right],$$

$$G_n(x) = G_{n-2}(x) - A_{n-2}(x).$$

For the argument values: $x = 0.25, 0.5, 0.75, \dots$, the functions $f_0(0, x)$, $f_0(1, x)$, of Kotani's Table can be used in order to calculate G_0 and G_1 according to

$$G_0(x) = x f_0(0, x), \quad G_1(x) = x f_0(1, x) - f_0(0, x).$$

Third step: Compute the functions

$$\phi_{n\bar{n}}^{00}(\alpha, \bar{\alpha}) = \phi_{n\bar{n}}(\alpha, \bar{\alpha}),$$

for $n = 0, 1, 2, \dots, N$; $\bar{n} = 0, 1, 2, \dots, \bar{N}$, in the following way. First draw the skeleton of rules of the table shown in Fig. 1

(the diagonal lines preferably in a different color) and enter the values for all functions except the functions $\phi_{n\bar{n}}$. Then compute and enter successively the latter using the recurrence formulae:

$$\begin{aligned} \phi_{n\bar{n}} = (1/\alpha\bar{\alpha}) \left[\bar{\alpha}n \phi_{n-1,\bar{n}} + e^{-\bar{\alpha}} g_n(\alpha) \right. \\ \left. + \alpha\bar{n} \phi_{n,\bar{n}-1} + e^{-\alpha} g_{\bar{n}}(\bar{\alpha}) \right. \\ \left. - n\bar{n} \phi_{n-1,\bar{n}-1} - g_{n+\bar{n}}(\alpha + \bar{\alpha}) \right]. \end{aligned}$$

If $n = 0$ or $\bar{n} = 0$, or if both are zero, then the terms on the right hand side which are multiplied by zero have simply to be omitted.

Each $\phi_{n\bar{n}}$ is obtained by one operation (on a calculating machine permitting accumulative multiplication) following a simple geometrical path in the table.

Fourth step: Compute the functions $A_{n\bar{n}}(\alpha, \bar{\alpha})$. First draw the skeleton of rules of Figure 2 and enter the values for all functions except the functions $A_{n\bar{n}}$. Then compute and enter successively the latter using the recurrence formulae:

$$A_{n\bar{n}} = (1/\alpha + \bar{\alpha}) \left[nA_{n-1,\bar{n}} + \bar{n}A_{n,\bar{n}-1} + A_n(\alpha) A_{\bar{n}}(\bar{\alpha}) \right].$$

For $n = 0$ or $\bar{n} = 0$, or both $= 0$, the terms containing the factor zero have to be omitted. Each $A_{n\bar{n}}$ is obtained by one operation (on a calculating machine permitting accumulative multiplication) following a simple geometrical path in the table.

Fifth step: Compute the functions $\phi_{n\bar{n}}^{01}(\alpha, \bar{\alpha})$ and $\phi_{n\bar{n}}^{02}(\alpha, \bar{\alpha})$ by means of the recurrence formulae:

$$\phi_{n\bar{n}}^{01}(\alpha, \bar{\alpha}) = \phi_{n+1,\bar{n}+1}^{00}(\alpha, \bar{\alpha}) - A_n(\alpha)A_{\bar{n}}(\bar{\alpha}) - A_{n\bar{n}}(\alpha, \bar{\alpha}),$$

$$\begin{aligned} 4 \phi_{n\bar{n}}^{02}(\alpha, \bar{\alpha}) = & 9 \phi_{n+1,\bar{n}+1}^{01}(\alpha, \bar{\alpha}) + \phi_{n\bar{n}}^{00}(\alpha, \bar{\alpha}) \\ & + 3 \left[A_n(\alpha)A_{\bar{n}+1}(\bar{\alpha}) + A_{n+1}(\alpha)A_{\bar{n}}(\bar{\alpha}) \right] \\ & - 3 \left[\phi_{n+2,\bar{n}}^{00}(\alpha, \bar{\alpha}) + \phi_{n,\bar{n}+2}^{00}(\alpha, \bar{\alpha}) \right] \\ & - 3 \left[A_n(\alpha)A_{\bar{n}}(\bar{\alpha}) + A_{n\bar{n}}(\alpha, \bar{\alpha}) \right]. \end{aligned}$$

Sixth step: Compute the functions $\phi_{n\bar{n}}^{03}$, $\phi_{n\bar{n}}^{04}$, ..., $\phi_{n\bar{n}}^{0l}$

by means of the recurrence formulae:

$$9 \phi_{n\bar{n}}^{03} = 25 \phi_{n+1, \bar{n}+1}^{02} + 9 \phi_{n\bar{n}}^{01} + 5 \phi_{n+1, \bar{n}+1}^{00}$$

$$- 15 \left[\phi_{n+2, \bar{n}}^{01} + \phi_{n, \bar{n}+2}^{01} \right],$$

$$\begin{aligned} l^2(2l-5) \phi_{n\bar{n}}^{0,l} &= (2l-5)(2l-1)^2 \phi_{n+1, \bar{n}+1}^{0,l-1} \\ &+ \left[(2l-5)(l-1)^2 + (2l-1)(l-2)^2 \right] \phi_{n\bar{n}}^{0,l-2} \\ &+ (2l-1)(2l-5)^2 \phi_{n+1, \bar{n}+1}^{0,l-3} \\ &- (2l-1)(2l-3)(2l-5) \left[\phi_{n+2, \bar{n}}^{0,l-2} + \phi_{n, \bar{n}+2}^{0,l-2} \right] \\ &- (2l-1)(l-3)^2 \phi_{n\bar{n}}^{0,l-4}. \end{aligned}$$

The second formula is valid for all $l \geq 4$.

Seventh step: Compute the functions $\phi_{n\bar{n}}^{Ml}(\alpha, \bar{\alpha})$ for $M = 1, 2,$

... by means of the recurrence formulae:

$$\phi_{n\bar{n}}^{M+1,l} = \phi_{n+1, \bar{n}+1}^{M,l} - \frac{(l+M)}{(2l+1)} \phi_{n\bar{n}}^{M,l-1} - \frac{(l+1-M)}{(2l+1)} \phi_{n\bar{n}}^{M,l+1}.$$

		$e^{-\alpha} G_0(\bar{\alpha})$	$e^{-\alpha} G_1(\bar{\alpha})$	$e^{-\alpha} G_2(\bar{\alpha})$	$e^{-\alpha} G_3(\bar{\alpha})$	...
$(\alpha \bar{\alpha})$		0	1α	2α	3α	...
$e^{-\alpha} G_0(\alpha)$	0	ϕ_{00}	ϕ_{01}	ϕ_{02}	ϕ_{03}	...
$e^{-\alpha} G_1(\alpha)$	$1\bar{\alpha}$	ϕ_{10}	ϕ_{11}	ϕ_{12}	ϕ_{13}	...
$e^{-\alpha} G_2(\alpha)$	$2\bar{\alpha}$	ϕ_{20}	ϕ_{21}	ϕ_{22}	ϕ_{23}	...
$e^{-\alpha} G_3(\alpha)$	$3\bar{\alpha}$	ϕ_{30}	ϕ_{31}	ϕ_{32}	ϕ_{33}	...
.	.	.	.	.	.	.
.	.	.	.	.	.	.

Fig. 1

$G_0(\alpha + \bar{\alpha})$
$G_1(\alpha + \bar{\alpha})$
$G_2(\alpha + \bar{\alpha})$
$G_3(\alpha + \bar{\alpha})$
$G_4(\alpha + \bar{\alpha})$
$G_5(\alpha + \bar{\alpha})$
$G_6(\alpha + \bar{\alpha})$
.
.
.

		$A_0(\bar{\alpha})$	$A_1(\bar{\alpha})$	$A_2(\bar{\alpha})$	$A_3(\bar{\alpha})$	...
$(\alpha + \bar{\alpha})$		0	1	2	3	...
$A_0(\alpha)$	0	A_{00}	A_{01}	A_{02}	A_{03}	...
$A_1(\alpha)$	1	A_{10}	A_{11}	A_{12}	A_{13}	...
$A_2(\alpha)$	2	A_{20}	A_{21}	A_{22}	A_{23}	...
$A_3(\alpha)$	3	A_{30}	A_{31}	A_{32}	A_{33}	...
.	.	.	.	.	.	.
.	.	.	.	.	.	.
.	.	.	.	.	.	.

Fig. 2