

UNCLASSIFIED

AD _____

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION ALEXANDRIA, VIRGINIA

DOWNGRADED AT 3 YEAR INTERVALS:
DECLASSIFIED AFTER 12 YEARS
DCD DIR 5200.10



UNCLASSIFIED

THIS REPORT HAS BEEN DECLASSIFIED
AND CLEARED FOR PUBLIC RELEASE.

DISTRIBUTION A
APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

40 No. 10 113
ASTIA FILE COPY

OFFICE OF NAVAL RESEARCH

Task Contract N5ori-07822

Project Designation No. NR-356-096

Technical Report No. 18

CYCLIC POLYOLEFINS. XXIX. CYCLOOCTATETRAENE DERIVATIVES FROM
COPOLYMERIZATION AND SIDE CHAIN MODIFICATION.

by

Arthur C. Cope and Ronald M. Pike

(Prepared for publication in the
Journal of the American Chemical Society)

Massachusetts Institute of Technology
Department of Chemistry
Cambridge 39, Mass.

March 23, 1953

[Contribution from the Department of Chemistry,
Massachusetts Institute of Technology]

CYCLIC POLYOLEFINS. XXIX. CYCLOOCTATETRAENE DERIVATIVES FROM
COPOLYMERIZATION AND SIDE CHAIN MODIFICATION¹

(1) Supported in part by the Office of Naval Research under Contract
N5ori-07822, Project Designation NR-356-096.

By Arthur C. Cope and Ronald M. Pike

Received _____

γ -Cyclooctatetraenyl-n-propyl alcohol (I), γ -cyclooctatetraenyl-n-propyl cyanide (V), cyclooctatetraenylmethylcarbinol (VIII), cyclooctatetraenyl methyl ketone (IX) and cyclooctatetraenyldimethylcarbinol (XI) have been prepared by copolymerization of acetylene with functionally substituted acetylenes. N, N-Dimethyl γ -cyclooctatetraenyl-n-propylamine (IV) and γ -cyclooctatetraenyl-n-propyl cyanide (V) were prepared by displacement reactions of the *p*-toluenesulfonate III of the alcohol I, and δ -cyclooctatetraenyl-n-butylamine (VI) and γ -cyclooctatetraenyl-n-butyric acid (VII) by reduction and saponification, respectively, of the cyanide V. The structures of several of the new cyclooctatetraene derivatives were established by conversion to, or preparation from, known cyclooctatetraenes: The primary alcohol I was oxidized to β -cyclooctatetraenylpropionic acid (II); cyclooctatetraenylmethyl-

carbinol (VIII) was oxidized with sodium hypobromite to cyclooctatetraene-carboxylic acid; cyclooctatetraenyl methyl ketone (IX) was prepared by oxidation of the carbinol VIII; and cyclooctatetraenyldimethylcarbinol (XI) was prepared from methyl cyclooctatetraenecarboxylate and methylmagnesium iodide.

This paper reports the preparation of additional mono-substituted cyclooctatetraenes by two routes; copolymerization of acetylene with monosubstituted acetylenes, and chemical reactions of functional groups in the side chains of cyclooctatetraenes prepared by copolymerization.

New cyclooctatetraenes prepared by copolymerization were the following:

γ -cyclooctatetraenyl-n-propyl alcohol (I) (8%) from 4-pentyn-1-ol;
 γ -cyclooctatetraenyl-n-propyl cyanide (V) (11%) from 5-hexynenitrile;
cyclooctatetraenyldimethylcarbinol (VIII) (11%) from 3-buten-2-ol; cyclooctatetraenyl methyl ketone (IX) (1.5%) from 3-buten-2-one; cyclooctatetraenyldimethylcarbinol (XI) (13%) from 2-methyl-3-buten-2-ol. These compounds were isolated and purified by a combination of steam distillation, fractionation and chromatography on silica gel described previously. Physical


(2) A. C. Cope and D. F. Ruge, This Journal, 75, 0000 (1953).

evidence for their structures was provided by their infrared spectra (Figs. 1 and 2), which had the characteristics previously noted for monosubstituted cyclooctatetraenes.² Chemical evidence for the structure of γ -cyclo-

cyclooctatetraenyl-n-propyl alcohol (I) was obtained by oxidation with chromic acid to the known ² crystalline β -cyclooctatetraenylpropionic acid (II) in 37% yield. Oxidation of cyclooctatetraenylmethylcarbinol (VIII) provided evidence for its structure and also for the structure of the ketone IX. Oxidation with chromic acid or Oppenauer oxidation with acetone and aluminum tert.-butoxide yielded cyclooctatetraenyl methyl ketone (IX) (also prepared by copolymerization), while oxidation with sodium hypobromite formed the known ³ crystalline cyclooctatetraenecarboxylic acid. The structure of

(3) A. C. Cope, M. Burg and S. W. Fenton, ibid., 74, 173 (1952).

cyclooctatetraenyldimethylcarbinol (XI) was verified by an independent synthesis from methyl cyclooctatetraenecarboxylate ³ and methylmagnesium iodide.



- I, R = (CH₂)₃OH
- II, R = (CH₂)₂COOH
- III, R = (CH₂)₃OSO₂C₆H₄CH₃(-p)
- IV, R = (CH₂)₃N(CH₃)₂
- V, R = (CH₂)₃CN
- VI, R = (CH₂)₄NH₂
- VII, R = (CH₂)₃COOH
- VIII, R = CH(OH)CH₃
- IX, R = $\begin{array}{c} \text{CCH}_3 \\ \parallel \\ \text{O} \end{array}$
- X, R = $\begin{array}{c} \text{CCH}_3 \\ \parallel \\ \text{NOH} \end{array}$
- XI, R = C(OH)(CH₃)₂

γ -Cyclooctatetraenyl-n-propyl p-toluenesulfonate (III) was prepared from the alcohol I and p-toluenesulfonyl chloride in pyridine and used as an intermediate in the preparation of other cyclooctatetraenes in the manner previously reported for the next lower homolog.² Reaction of the p-toluenesulfonate III with dimethylamine yielded N, N-dimethyl γ -cyclooctatetraenyl-n-propylamine (IV) (54%) while with potassium cyanide γ -cyclooctatetraenyl-n-propyl cyanide (V) (74%) was formed, identical in properties with V prepared by copolymerization. Reduction of the cyanide V with lithium aluminum hydride yielded δ -cyclooctatetraenyl-n-butylamine (VI) (39%), and saponification of V yielded γ -cyclooctatetraenyl-n-butyric acid (VII) (69%). The acid VII was isolated as a yellow liquid that formed a crystalline salt with quinine. After recrystallization of the salt to constant melting point and optical rotation, the acid was regenerated by treatment with 1% hydrochloric acid at 0°, and proved to be optically inactive.

Reactions that were used successfully in the course of this work to modify the side chains of substituted cyclooctatetraenes that had not been demonstrated previously to be applicable without disruption of the cyclooctatetraene ring included the following: oxidation of a primary alcohol to an acid with chromic acid; oxidation of a secondary alcohol to a ketone with chromic acid or by the Oppenauer method; sodium hypobromite oxidation of a methyl carbinol to an acid; Grignard synthesis of a tertiary alcohol from a carboxylic ester. Types of functionally substituted acetylenes that were used successfully for the first time in the preparation of substituted cyclooctatetraenes by copolymerization with acetylene were an acetylenic ketone and an acetylenic nitrile.

Experimental ⁴

-
- (4) Melting points are corrected and boiling points are uncorrected. We are indebted to Dr. S. M. Nagy and his associates for analyses, and for the infrared spectra, which were determined with a Baird Double Beam Infrared Recording Spectrometer, Model B, fitted with a sodium chloride prism.
-

Procedures that have been described previously for the copolymerization of substituted acetylenes with acetylene and purification of the resulting substituted cyclooctatetraenes by chromatography on silica gel were used. ²

γ -Cyclooctatetraenyl- n -propyl Alcohol (I). - 4-Pentyn-1-ol was prepared from 120.5 g. of tetrahydrofurfuryl chloride and sodamide in liquid ammonia in a yield of 65.5 g. (78%), b.p. 70-71° (29 mm.), n_D^{25} 1.4443. ⁵ Copolymerization of 20 g. of 4-pentyn-1-ol with acetylene by

-
- (5) G. Eglinton, E. R. H. Jones and M. C. Whiting, J. Chem. Soc., 2873 (1952); Org. Syntheses, 33, 000 (1953).
-

the general procedure described previously ² resulted in an absorption of 1200 p.s.i. of acetylene and an increase in weight of 171 g. in 8 hours at 84-92°. After purification by chromatography on silica gel ² and fractionation through a semimicro column, the yield of the pure alcohol I was 3.5 g. (8%), b.p. 91-91.5° (0.15 mm.), n_D^{25} 1.5426, d_4^{25} 1.0244.

Anal. Calcd. for $C_{11}H_{14}O$: C, 81.48; H, 8.64. Found:

C, 81.36; H, 8.85.

β -Cyclooctatetraenylpropionic Acid (II) was obtained by treating a solution of 0.50 g. of the alcohol I in 40 ml. of acetone (distilled from potassium permanganate) with 8 ml. of a solution prepared from 23.3 g. of

chromium trioxide, 19.7 ml. of concentrated sulfuric acid and 30 ml. of water.⁶

-
- (6) Based on an oxidation procedure described by R. G. Curtis, I. Heilbron, E.R.H. Jones and G. F. Woods, J. Chem. Soc., 0000 (1953).
-

The oxidizing agent was added dropwise with stirring and cooling to keep the reaction temperature below 25°, and the mixture was allowed to stand for 30 minutes. Water (40 ml.) was added and the mixture was extracted thoroughly with ether. The extracts were combined, washed with water, and extracted with 10% sodium hydroxide solution. The alkaline extracts were acidified with hydrochloric acid and re-extracted with ether. The extracts were dried over sodium sulfate, concentrated under reduced pressure after addition of benzene, and the yellow acid II that crystallized from the residue on cooling was recrystallized from hexane; yield 0.22 g. (37%), m.p. and mixed m.p. with an authentic sample,² 58.8-59.8°.

γ -Cyclooctatetraenyl-n-propyl p-toluenesulfonate (III). -

A solution of 4.0 g. of the alcohol I in 6 ml. of pyridine was cooled in an ice bath, and 6.87 g. of p-toluenesulfonyl chloride was added in one portion. The mixture was allowed to stand in the ice bath for 3.5 hours, with frequent agitation by swirling. Water (20 ml.) was added, and after 30 minutes the mixture was extracted twice with ether, and the extracts were washed with 10% hydrochloric acid, sodium bicarbonate solution and water. The ether solution was dried over magnesium sulfate, concentrated under reduced pressure, and the liquid residue of the crude p-toluenesulfonate III was evacuated at 16 mm. and room temperature for 15 minutes; yield 7.47 g. (96%).

N, N-Dimethyl γ -Cyclooctatetraenyl-n-propylamine (IV). -

A solution of 3.0 g. of the p-toluenesulfonate III in 15 ml. of benzene was saturated with dimethylamine and allowed to stand overnight at room temperature. Ether (20 ml.) was added, and the solution was washed with water and dried over magnesium sulfate. Concentration of the solution and fractionation of the residue through a semimicro column yielded 0.98 g. (54%) of the amine IV, b.p. 50-55° (0.01 mm.), n_D^{25} 1.5158-1.5164. An analytical sample had b.p. 55° (0.01 mm.), n_D^{25} 1.5164, d_4^{25} 0.9234.

Anal. Calcd. for $C_{12}H_{18}N_2$: C, 82.48; H, 10.12; N, 7.40.

Found: C, 82.47; H, 10.07; N, 7.17.

N, N-Dimethyl γ -Cyclooctatetraenyl n-propylamine Picrate was

prepared by adding the amine IV to a saturated solution of picric acid in ether, and recrystallized to a constant melting point of 103.5-104° from 95% ethanol.

Anal. Calcd. for $C_{15}H_{23}N_4O_7$: C, 54.53; H, 5.30; N, 13.39.

Found: C, 54.23; H, 5.35; N, 13.55.

γ -Cyclooctatetraenyl-n-propyl Cyanide (V). - (a). A solution

of 1.32 g. of the p-toluenesulfonate III and 0.75 g. of potassium cyanide in 10 ml. of 80% ethanol was heated under reflux on a steam bath for 2 hours. The solution was poured into 25 ml. of water, and extracted with ether. The ether extracts were dried over magnesium sulfate, concentrated, and the residue was fractionated through a semimicro column, yielding 0.73 g. (74%) of the nitrile V, b.p. 95-103.5° (1 mm.), n_D^{25} 1.5263-1.5290. A redistilled analytical sample had b.p. 103.5-104.5° (0.5 mm.), n_D^{25} 1.5300, d_4^{25} 0.9911.

Anal. Calcd. for $C_{12}H_{13}N$: C, 84.21; H, 7.60; N, 8.19.

Found: C, 84.27; H, 7.81; N, 8.06.

(b). The nitrile V also was prepared by the copolymerization of 10 g. of 5-hexynenitrile (b.p. 64-65° at 10 mm., n_D^{25} 1.4389, prepared from 4-pentyn-1-yl p-toluenesulfonate and sodium cyanide⁷) with acetylene

(7) G. Eglington, E. R. H. Jones and M. C. Whiting, J. Chem. Soc., 0000 (1953).

at 86-95°, with acetylene absorption of 1020 p.s.i. (134 g.) in 11 hours, and purified by chromatography on silica gel. The yield of V with physical properties and infrared spectrum identical within experimental error with the sample described above that was prepared from the p-toluenesulfonate III was 2.1 g. (11%).

α -Cyclooctatetraenyl-n-butylamine (VI). - A solution of 1.23 g. of the nitrile V in 15 ml. of dry ether was added slowly with stirring to 0.31 g. of lithium aluminum hydride in 20 ml. of ether in a nitrogen atmosphere. The mixture was heated under reflux for 1 hour, cooled, and 3 ml. of water was added cautiously, followed by 35 ml. of 10% hydrochloric acid. After extraction with ether (extracts discarded), the solution was made basic with 10% sodium hydroxide and re-extracted with ether. The extracts were dried over magnesium sulfate, concentrated, and the residue was distilled through a semimicro column, yielding 0.49 g. (39%) of the amine VI, b.p. 82-83° (0.7 mm.), n_D^{25} 1.5330-1.5334. An analytical sample had b.p. 83° (0.7 mm.), n_D^{25} 1.5334, d_4^{25} 0.9535.

Anal. Calcd. for $C_{12}H_{17}N$: C, 82.31; H, 9.71; N, 8.00.

Found: C, 82.16; H, 9.92; N, 7.90.

δ -Cyclooctatetraenyl-n-butylamine Hydrochloride was prepared

by passing hydrogen chloride into a solution of 0.246 g. of the amine VI in 10 ml. of dry ether. The hydrochloride was obtained in a yield of 0.184 g. (59%) after recrystallization from methanol-ether to a constant melting point of 110.5-111.5°.

Anal. Calcd. for $C_{12}H_{18}ClN$: C, 68.07; H, 8.57; N, 6.62.

Found: C, 68.05; H, 8.72; N, 6.63.

γ -Cyclooctatetraenyl-n-butyric Acid (VII). - The nitrile

V (1.9 g.) was heated under reflux for 12 hours with a solution of 3.0 g. of sodium hydroxide in 50 ml. of water. The solution was cooled, extracted with ether (extracts discarded), and acidified with 10% hydrochloric acid. The acid VII was extracted with three 20-ml. portions of ether, which were dried over magnesium sulfate and concentrated. Fractionation of the residue through a semimicro column yielded 1.45 g. (69%) of the acid VII, b.p. 95-97° (0.019 mm.), n_D^{25} 1.5310. The acid failed to crystallize.

Anal. Calcd. for $C_{12}H_{14}O_2$: C, 75.77; H, 7.41. Found:

C, 75.25; H, 7.45.

The Quinine Salt of γ -Cyclooctatetraenyl-n-butyric Acid was

prepared by adding a solution of 0.15 g. of the acid VII in 1 ml. of ether to a solution of 0.299 g. of quinine hydrate in 50 ml. of ether. The solution was concentrated to 10 ml. and cooled overnight at 5°. The light yellow crystalline salt (0.325 g., 80%) was crystallized twice from ether to a constant

melting point of 101-102°, $[\alpha]_D^{27} -114.8^\circ$ (l. = 1, c 0.95 in absolute ethanol).

Anal. Calcd. for $C_{22}H_{33}N_2O_4$: C, 74.68; H, 7.44; N, 5.44.

Found: C, 74.45; H, 7.56; N, 5.56.

The acid VII that was regenerated from the quinine salt with 1% hydrochloric acid at 0° and immediately extracted with carbon tetrachloride was optically inactive.

Cyclooctatetraenylmethylcarbinol (VIII). - 3-Butyn-2-ol

(20 g., Farchan Research Laboratories) was copolymerized with acetylene by the general procedure described previously.² Acetylene absorption amounted to 1080 p.s.i., with a weight increase of 102 g. during 10.5 hours at 84-90°. Separation of the products by the usual procedure, including chromatography on silica gel, yielded 36 g. of cyclooctatetraene and 4.56 g. (11%) of the alcohol VIII (eluted from the silica gel with ether); b.p. 80-85° (0.9 mm.), $n_D^{25} 1.5455-1.5472$. A redistilled analytical sample had b.p. 70° (0.4 mm.), $n_D^{25} 1.5460$, $d_4^{25} 1.0385$.

Anal. Calcd. for $C_{10}H_{12}O_2$: C, 81.04; H, 8.10. Found: C, 81.10; H, 8.16.

Cyclooctatetraenecarboxylic Acid was obtained by adding 0.50 g. of the alcohol VIII dropwise with stirring at a temperature below 10° to the sodium hypobromite prepared from 1.9 g. of sodium hydroxide in 16 ml. of water and 2.42 g. of bromine. The mixture was stirred at room temperature for 12 hours, extracted with ether (extracts discarded), acidified with 6 N sulfuric acid, and re-extracted with ether. The extracts were washed with

water, dried over sodium sulfate, and concentrated under reduced pressure. Sublimation of the residue by heating at 0.5 mm. yielded 0.183 g. (37%) of cyclooctatetraenecarboxylic acid, m.p. 69-75°, which after recrystallization from ether had m.p. and mixed m.p. with an authentic sample ³ of 71.5-72.8°.

Cyclooctatetraenyl Methyl Ketone (IX). - (a). To a solution of 3.5 g. of the carbinol VIII in 20 ml. of acetone was added dropwise with stirring at 0° and in a nitrogen atmosphere during 25 minutes a solution of 1.43 g. of chromium trioxide and 1.5 ml. of sulfuric acid in 5 ml. of water. After the addition was completed, the mixture was stirred for 2.5 hours at 0° and poured into 100 ml. of ice water. The mixture was extracted with ether, and the extracts were washed with water and dried over sodium sulfate. Concentration and distillation of the residue through a semimicro column yielded 1.69 g. (53%) of the ketone IX, b.p. 65-68.5° (0.55 mm.), n_D^{25} 1.5470-1.5518, which was purified through the oxime.

Cyclooctatetraenyl Methyl Ketoxime (X) was prepared by heating a solution of 0.35 g. of the ketone IX and 0.17 g. of hydroxylamine hydrochloride in 5 ml. of pyridine and 5 ml. of absolute ethanol on a steam bath under reflux for 15 hours. The solution was concentrated and water was added to the crystalline residue, which was collected on a filter and washed with water. The yield of the oxime was 0.1 g. (85%) after recrystallization from methanol; m.p. 150-151°.

Anal. Calcd. for $C_{10}H_{11}NO$: C, 74.53; H, 6.83; N, 8.70.

Found: C, 74.35; H, 6.89; N, 8.48.

The ketone IX was regenerated from 0.60 g. of the oxime, which was heated under reflux for 3.5 hours with 0.78 g. of pyruvic acid and 0.38 g. of sodium acetate in 5 ml. of acetic acid and 2 ml. of water.⁸

(8) Based on a procedure described by E. B. Hershberg, J. Org. Chem. 13, 542 (1948).

The solution was poured into 20 ml. of water and extracted with two 20-ml. portions of ether. The extracts were washed with saturated sodium bicarbonate solution, water, and dried over sodium sulfate, and concentrated. Benzene was added and distilled under reduced pressure to complete removal of the water, and the residue was fractionated through a semimicro column. The yield of IX was 0.31 g. (57%), b.p. 73-74° (1.2 mm.), n_D^{25} 1.5519-1.5549. A redistilled analytical sample had b.p. 74° (1.2 mm.), n_D^{25} 1.5548, d_4^{25} 1.0319. A number of analyses of several samples of the ketone were slightly low in carbon, but it is believed to be pure because the physical properties (including infrared spectra) of the samples regenerated under nitrogen from the analytically pure ketoxime X were identical.

Anal. Calcd. for $C_{10}H_{10}O$: C, 82.20; H, 6.90. Found:
C, 81.43; H, 6.98.

The ketone IX also was prepared in poorer (18-30%) yield by Oppenauer oxidation of the carbinol VIII with acetone and aluminum tert. butoxide in benzene.

Cyclooctatetraenyl Methyl Ketone 2,4-Dinitrophenylhydrazone

was prepared by adding a solution of the ketone in 95% ethanol at room temperature to a slight excess of a solution prepared from 0.4 g. of 2,4-dinitrophenylhydrazine, 2 ml. of concentrated sulfuric acid, 3 ml. of water and 10 ml. of 95% ethanol. The derivative was recrystallized from a mixture of ethanol and ethyl acetate; m.p. 202-203° (dec.).

Anal. Calcd. for $C_{16}H_{14}N_4O_6$: C, 58.89; H, 4.33; N, 17.18.

Found: C, 58.72; H, 4.50; N, 16.98.

(b). Copolymerization of 19 g. of 3-buten-2-one⁹ with

(9) E. R. H. Jones, K. Bowden, I. M. Heilbron, and B. C. L. Weedon, J. Chem. Soc., 39 (1946).

acetylene resulted in the absorption of 1160 p.s.i. (117 g.) of acetylene in 11 hours at 85-90°. Purification of the product by the general procedure described previously, including chromatography on silica gel, yielded 28.3 g. of cyclooctatetraene and 0.60 g. (1.5%) of IX, b.p. 85-90° (1 mm.), identified by m.p. and mixed m.p. of the 2,4-dinitrophenylhydrazone with the derivative described above (201-202° dec.).

Cyclooctatetraenyldimethylcarbinol (XI). - Copolymerization of 20 g. of 2-methyl-3-buten-2-ol (Farchan Research Laboratories) with acetylene resulted in the absorption of 1030 p.s.i. of acetylene (153 g.) in 8.5 hours at 85-88°. Purification by the usual procedure,² including chromatography on silica gel, yielded 37 g. of cyclooctatetraene and 3.0 g.

(13%) of the carbinol XI, b.p. 73-75.5° (0.5 mm.), n_D^{25} 1.5370-1.5382.

A redistilled analytical sample had b.p. 74.5-75.5° (1.3 mm.), n_D^{25} 1.5370, d_4^{25} 1.1024.

Anal. Calcd. for $C_{11}H_{14}O$: C, 81.48; H, 8.64. Found:

C, 81.49; H, 8.79.

An authentic sample of the carbinol XI was prepared by adding methyl cyclooctatetraenecarboxylate ³ (1.28 g.) in 10 ml. of dry ether to the methylmagnesium iodide prepared from 0.36 g. of magnesium and 2.27 g. of methyl iodide in 20 ml. of dry ether. The magnesium salts were decomposed with aqueous ammonium chloride and the ether solution was washed with water, dried over magnesium sulfate, and concentrated. Distillation of the residue through a semimicro column yielded 0.81 g. of crude XI, b.p. 73-75° (2.5 mm.), n_D^{25} 1.5369-1.5371. The infrared spectrum of the product showed a strong carbonyl band, so it was purified by chromatography on a column containing 25 g. of neutral alumina with activity 3.¹⁰

(10) H. Brockman and E. Schodder, Ber., 74, 73 (1945).

Elution with 300 ml. of benzene followed by concentration and distillation yielded 0.49 g. of recovered methyl cyclooctatetraenecarboxylate, b.p. 76° (2.3 mm.), n_D^{25} 1.5371, identified by the infrared spectrum. Elution with 300 ml. of ether followed by concentration and distillation through a semimicro column yielded 0.21 g. of XI, b.p. 74.5° (3.0 mm.), n_D^{25} 1.5357, which had an infrared spectrum identical with XI prepared by copolymerization, except for the presence of weak bands at 5.83 and 6.5 μ due to a small amount of impurity.




Fig. 1. - Infrared absorption spectra: curve 1, δ -cyclooctatetraenyl-n-propyl alcohol (I); curve 2, N, N-dimethyl- δ -cyclooctatetraenyl-n-propylamine (IV); curve 3, γ -cyclooctatetraenyl-n-propyl cyanide (V); curve 4, δ -cyclooctatetraenyl-n-butylamine (VI). The spectra were determined without solvent in a 0.025 mm. cell.




Fig. 2. - Infrared absorption spectra: curve 1, γ -cyclooctatetraenyl-*n*-butyric acid (VII); curve 2, cyclooctatetraenylmethylcarbinol (VIII); curve 3, cyclooctatetraenyl methyl ketone (IX); curve 4, cyclooctatetraenyldimethylcarbinol (XI). Curve 1 was determined with solutions (100 mg./ml.) in carbon tetrachloride in the region 2-8 μ and in carbon disulfide in the region 8-16 μ in a 0.10 mm. cell. Curves 2-4 were determined without solvent in a 0.025 mm. cell.

DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Addressee</u>
2	Commanding Officer Office of Naval Research Branch Office 150 Causeway Street Boston, Massachusetts
1	Commanding Officer Office of Naval Research Branch Office 844 North Rush Street Chicago 11, Illinois
1	Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, New York
1	Commanding Officer Office of Naval Research Branch Office 1000 Geary Street San Francisco 9, California
1	Commanding Officer Office of Naval Research Branch Office 1030 N. Green Street Pasadena 1, California
2	Officer-in-Charge Office of Naval Research Branch Office Navy Number 100 Fleet Post Office New York, New York
9	Director, Naval Research Laboratory Washington 25, D. C. Attention: Technical Information Officer
4	Chief of Naval Research Office of Naval Research Washington 25, D. C. Attention: Chemistry Branch
1	Research and Development Board Pentagon, Room 3D1041 Washington 25, D. C. Attention: Technical Reference Section

DISTRIBUTION LIST

Page 2

<u>No. of Copies</u>	<u>Addressee</u>
1	Dr. Ralph G. H. Siu, Research Director General Laboratories QM Depot 2800 S. 20th Street Philadelphia 45, Pennsylvania
1	Dr. Warren Stubblebine, Research Director Chemical and Plastics Section, RDB-MPD Quartermaster General's Office Washington 25, D. C.
1	Dr. A. Stuart Hunter, Tech. Director Research and Development Branch MPD Quartermaster General's Office Washington 25, D. C.
1	Dr. A. G. Horney Wright Air Development Center Wright-Patterson Air Force Base Dayton, Ohio Attention: WCRRS-4
1	Dr. A. Weissler Department of The Army Office of the Chief of Ordnance Washington 25, D. C. Attention: ORDTB-PS
1	Research and Development Group Logistics Division, General Staff Department of The Army Washington 25, D. C. Attn: Dr. W. T. Read, Scientific Adviser
2	Director, Naval Research Laboratory Washington 25, D. C. Attention: Chemistry Division
2	Chief of the Bureau of Ships Navy Department Washington 25, D. C. Attention: Code 340
2	Chief of the Bureau of Aeronautics Navy Department Washington 25, D. C. Attention: Code TD-4
2	Chief of the Bureau of Ordnance Navy Department Washington 25, D. C. Attention: Code Rexd
1	Dr. H. A. Zahl, Tech. Director Signal Corps Engineering Laboratories Fort Monmouth, New Jersey

DISTRIBUTION LIST

Page 3

No. of Copies

Addressee

6

Mr. J. H. Heald
Library of Congress
Navy Research Section
Washington 25, D. C.

1

U.S. Naval Radiological Defense Laboratory
San Francisco 24, California
Attention: Technical Library

1

Naval Ordnance Test Station (Inyokern)
China Lake, California
Attention: Head, Chemistry Division

1

Office of Ordnance Research
2127 Myrtle Drive
Durham, North Carolina

1

Technical Command
Chemical Corps
Chemical Center, Maryland

1

U.S. Atomic Energy Commission
Research Division
Washington 25, D. C.

1

U.S. Atomic Energy Commission
Chemistry Division
Brookhaven National Laboratory
Upton, New York

1

U.S. Atomic Energy Commission
Library Branch, Tech. Information, ORE
P. O. Box E
Oak Ridge, Tennessee