

AD NO. 10251
ADP FILE COPY

ODS HOLD OCEANOGRAPHIC INSTITUTION

NO. 10251, MAR 1961, 1/15.

WOODS HOLE OCEANOGRAPHIC INSTITUTION

Woods Hole, Massachusetts

Reference No. 53-24

THE SURFACE OUTCROP OF THE
GULF STREAM FRONT

By

William S. von Arx
and
William S. Richardson

Technical Report
Submitted to the Office of Naval Research
Under Contract N6onr-27701 (NR-083-004)

April 1953

Approved for Distribution


Director

THE SURFACE OUTCROP OF THE GULF STREAM FRONT

William S. von Arx
and
William S. Richardson

During 26-27 February 1953, an attempt was made to fly along the surface expression of the Gulf Stream front between Miami, Florida, and 70° W longitude. On this flight, as during the earlier flight made in November 1952 (Science, 1953), the Stommel-Parson airborne radiation thermometer was used together with photographic and visual observations to determine the position of the frontal outcrop on the sea surface. It was possible to follow phenomena apparently related to the Gulf Stream front throughout most of the distance between Miami and 70° W, but it was found that both the visual and thermometric evidence of the front was discontinuous. Figure 1 shows the manner in which the frontal outcrop was broken. When approaching a discontinuity, one had the experience that both the temperature gradient at the surface and the change of sea state and sea color coincident with the temperature change grew fainter as observations were pursued downstream until in each case none of these clues were sufficiently pronounced to direct the airplane further. It was learned in November 1952 that the frontal phenomena generally reappeared nearer shore. This was found to be so in every case on the more recent flight. The surface expressions of the Gulf Stream front are therefore considered to be

a succession of traceable surface evidences arranged on echelon in a manner similar in many ways to the very large-scale pattern of the multiple Gulf Stream proposed by Fuglister, 1952, from evidence obtained by Bathythermograph in the upper 200 m layer.

At present it is difficult to understand how the surface evidence obtained from the air may be related to the meander structure such as that observed from shipboard (Fuglister and Worthington, 1951). It is quite possible, for example, that the discontinuities observed from the airplane are only superficial phenomena. It is also possible, however, that the relatively wide spacing between successive Bathythermograph observations and successive crossings by ships have led to a smoother representation of the frontal structure than is actually the case.

Observation from the airplane has also indicated a fine structure which interrupts the continuity of the frontal outcrop at intervals from 1/2 to 10 miles. When the weather is clear it is possible to fly across these individual irregularities without losing track of the larger-scale trend. Figure 2 shows some of these smaller features as they were observed near Cape Hatteras on 27 February 1953.

The validity of visual and radiant signals from the sea surface as a means for tracing the frontal outcrop of the Gulf Stream is open to some question. It is known, for example, that the steepest temperature gradients at the




FIG.2

Appearance of the fine structure at the frontal outcrop seen off Cape Hatteras near $35^{\circ}00'N$, $75^{\circ}00'W$ on 27 Feb 1953 from an altitude of 1500 feet. View toward Northeast. Wind force 3 (approx) from Southwest. Warm water on the right. Original exposure on Kodachrome.

Unusually marked change in sea state noted at the frontal outcrop on 27 Feb 1953 near $35^{\circ}45'N$, $74^{\circ}20'W$ from an altitude of 1500 feet. View toward Northeast. Wind force about 4 from Southwest. Warm water on right, presumably moving with the wind. Original exposure on Kodachrome.

sea surface are usually found a few miles shoreward of the steepest horizontal gradients 100 to 200 m below the surface in winter and may be as much as 50 miles or more shoreward of the deeper temperature gradients in summer. Therefore, the evidence of the radiation thermometer alone is probably most reliable during the winter months. Visible evidence consists of a number of factors which include a change in the water color, a change in the sea state due to the horizontal shear of the surface currents, and possibly, accumulations of sargassum or, in calm weather, characteristic patterns of surface slicks. Slicks alone are unreliable indices of the front but when accompanied with a change in water color it is usually found that a strong temperature gradient is also associated. In rough weather the slicks are no longer visible but the sea state may change abruptly across the frontal outcrop (especially when near force 3) together with a change of water color and temperature. Therefore, the flights along the front are guided by an association of visible signs and evidence of an abrupt temperature change. From a study of Bathythermograph sections through Slope water it seems possible that these clues may remain useful from Cape Hatteras eastward during June and possibly early July before surface heating destroys the correlation of temperature gradients with the other physical factors. A trustworthy check has still to be made of the relationship between the frontal outcrop detected from the air with that established from shipboard.

REFERENCES

- Fuglister, F. C. and L. V. Worthington, 1951: Some results of a multiple ship survey of the Gulf Stream. *Tellus* 3 (1) 1-14.
- Fuglister, F. C., 1952: Multiple currents in the Gulf Stream. *Tellus* 3 (4) 230-233.
- Stommel, Henry, W. S. von Arx, D. Parson, and W. S. Richardson, 1953: Rapid aerial survey of Gulf Stream with camera and radiation thermometer. *Science* (in press).
- von Arx, W. S., 1952: Notes on the surface velocity profile and horizontal shear across the width of the Gulf Stream. *Tellus* 4 (3) 211-214.

Technical Report Distribution List
ONR Project NR-083-004

- 1 -

<u>Address</u>	<u>Copies</u>
Commanding Officer, Air Force Cambridge Research Center, 230 Albany Street, Cambridge 39, Massachusetts	1
Chief, Bureau of Ships, Department of the Navy, Washington 25, D. C. Attn: Code 847	2
Chief of Naval Research, Department of the Navy, Washington 25, D. C. Attn: Code 416	3
Commanding General, Research and Development Division, Department of the Air Force, Washington, D. C.	1
Commanding Officer, Branch Office of Naval Research, 150 Causeway Street, Boston 14, Massachusetts	1
Commanding Officer, Branch Office of Naval Research, 346 Broadway, New York 13, New York	1
Commanding Officer, Branch Office of Naval Research, 1000 Geary Street, San Francisco 9, California	1
Director, Lamont Geological Observatory, Torrey Cliff, Palisades, New York	1
Director, Naval Research Laboratory, Washington 25, D. C. Attn: Technical Services	9
Hydrographer, U. S. Navy Hydrographic Office, Washington 25, D. C. Attn: Division of Oceanography	8
Naval War College, Newport, Rhode Island	1
Chief, Bureau of Yards and Dock, Department of the Navy, Washington 25, D. C.	1
Director, Chesapeake Bay Institute, Box 426A, RFD 2, Annapolis, Maryland	1
Commandant, U. S. Coast Guard, 1300 E Street, N. W., Washington, D. C. Attn: Aerology and Oceanographic Section	1
Commanding General, Research and Development Division, Department of the Army, Washington, D. C.	1
Commanding Officer, Branch Office of Naval Research, 844 North Rush Street, Chicago, Illinois	1

Technical Report Distribution List
CNR Project NR-083-004

- 2 -

<u>Address</u>	<u>Copies</u>
Commanding Officer, Branch Office of Naval Research, 1030 East Green Street, Pasadena 1, California	1
Commanding Officer, Naval Ordnance Laboratory, White Oak, Silver Spring 19, Maryland	1
Director, U. S. Coast and Geodetic Survey, Department of Commerce, Washington 25, D. C.	1
Head, Department of Oceanography, Texas A and M, College Station, Texas	1
National Research Council, 2101 Constitution Avenue, Washington 25, D. C. Attn: Committee on Undersea Warfare	1
Assistant Naval Attache for Research, American Embassy, Navy Number 100, Fleet Post Office, New York, New York	2
Director, Scripps Institution of Oceanography, La Jolla, California	2
Director, U. S. Navy Electronics Laboratory, San Diego 52, California Attn: Codes 550, 552	2
Director, Marine Laboratory, University of Miami, Coral Gables 34, Florida	1
Chief of Naval Research, Department of the Navy, Washington 25, D. C. Attn: Code 466	1
Director, Narragansett Marine Laboratory, Kingston, Rhode Island	1
U. S. Fish and Wildlife Service, Fort Crockett, Galveston, Texas	1
Director, Hawaii Marine Laboratory, University of Hawaii, Honolulu, T. H.	1
British Joint Services Mission, Main Navy Building, Washington 25, D. C.	3
Research and Development Board, National Military Establish- ment, Washington 25, D. C. Attn: Committee on Geophysics and Geography	1
Director, Department of Engineering, University of California Berkeley, California	1

Technical Report Distribution List
ONR Project NR-083-004

- 3 -

<u>Address</u>	<u>Copies</u>
Head, Department of Oceanography, University of Washington Seattle 5, Washington	1
The Oceanographic Institute, Florida State University, Tallahassee, Florida	1
U. S. Fish and Wildlife Service, PO Box 3830, Honolulu, T.H.	1
Department of Conservation, Cornell University, Ithaca, New York. Attn: Dr. J. C. Ayers	1
Director, U. S. Fish and Wildlife Service, Department of the Interior, Washington 25, D. C. Attn: Dr. L. A. Walford	2
U. S. Army Beach Erosion Board, 5201 Little Falls Road, N. W. Washington 16, D. C.	1
Allan Hancock Foundation, University of Southern California, Los Angeles 7, California	1
U. S. Fish and Wildlife Service, 450B Jordan Hall, Stanford University, Stanford, California	1
Head, Department of Oceanography, Brown University, Providence, Rhode Island	1
Department of Zoology, Rutgers University, New Brunswick, New Jersey. Attn: Dr. H. H. Haskin	1
U. S. Fish and Wildlife Service, South Atlantic Offshore Fishery Investigations, c/o Georgia Game and Fish Comm. P. O. Box 312, Brunswick, Georgia	1