

AD No. 14 972
ASTIA FILE COPY

EQUILIBRIUM CONSTANTS IN THE REACTIONS

3-METHYLHEXANE = 2-METHYLHEXANE

AND

2,3-DIMETHYLPENTANE = 2,4-DIMETHYLPENTANE

By Robert L. Burwell, Jr. and Lucien G. Maury

Technical Report No. 5

to

THE OFFICE OF NAVAL RESEARCH

Project Number' NR 055-199

Contract N7 onr-54006

at

Northwestern University

under the direction of Robert L. Burwell, Jr.

June 1, 1953

This work was not actually supported by the Office of Naval Research in regard to salaries since Mr. Maury had been awarded a fellowship during his last year. However, this report is submitted since it followed directly from work of the project, employed compounds synthesized under the project and would ordinarily have been carried out as part of the project.

ABSTRACT

Equilibrium constants have been measured for the reactions, 3-methylhexane = 2-methylhexane, and 2,3-dimethylpentane = 2,4-dimethylpentane. The values were determined by direct equilibration employing sulfuric acid at 60°, chlorosulfonic acid at -33.4° and an equimolar mixture of the two at 0°.


The results are:

3MH = 2MH				23MP = 24MP			
temp.°	x_{2MH}^{eq}	K	ΔF°	x_{2MH}^{eq}	K	ΔF°	
60	0.595	1.47	-255	0.596	1.48	-260	
0	.628	1.69	-285	.706	2.40	-475	
-33.4	.655	1.90	-306	.728	2.68	-470	

Sulfuric acid and chlorosulfonic acid specifically interconvert methylalkanes.^{2,3,4} Change of chain branching as in the conversion

-
- (2) R.L. Burwell, Jr. and G.S. Gordon, III, *J. Am. Chem. Soc.*, 70, 3128 (1948); Gordon and Burwell, *ibid.*, 71, 2355 (1949).
 - (3) R.L. Burwell, Jr., Robert B. Scott, Lucien G. Maury and Allen S. Hussey, Technical Report No. 2 of this project.
 - (4) R.L. Burwell, Jr., Lucien G. Maury and Robert B. Scott, Technical Report No. 4 of this project.
-

of the 2- and 3-methylhexane pair to the 2,3- and 2,4-dimethylpentane pair occurs much more slowly. It, thus, seemed desirable to test the applicability of this discovery to the determination of the equilibrium constants of the following two reactions by direct equilibration.



Voorthuijsen⁵ determined the equilibrium constants of these

-
- (5) J.J.B. Voorthuijsen, *Rec. trav. chim.*, 66, 323 (1947).
-

reactions by direct equilibration of all heptanes by aluminum chloride at 20°. The more specific nature of the reaction with sulfuric acid offered hope of increased accuracy. The equilibrium constants have accordingly been measured at 60°, 0° and -33.4°.

Experimental

The preparation of the hydrocarbons has been described.³ The infrared absorption spectra were indistinguishable from those of the Bureau of Standards standard samples and no ultraviolet absorption was exhibited at wave lengths longer than 2200 Å.

As isomerizing agents, 96.04% sulfuric acid was employed at 60°, an equimolar mixture of this acid and freshly distilled Eastman Kodak

Practical Grade Chlorosulfonic acid was employed at 0°, and the chlorosulfonic acid alone was employed at -33.4°. Thermostating was by a water bath at 60°, by an ice-water mixture at 0° and by refluxing ammonia at -33.4°.

The value of the equilibrium constant was first approximated from the relative initial rates of isomerization of the two hydrocarbons involved by the relation $K = k_1/k_{-1}$. Mixtures of known composition on either side of the estimated equilibrium composition were prepared and subjected to the isomerizing conditions. According to the observed changes in composition, the estimated equilibrium composition was revised and new mixtures prepared. These were then subjected to isomerizing conditions. By this means, it was possible to employ short times of contact and relatively mild condition and, thus, to minimize side reactions.

Reaction flasks and stirring procedures have been described.⁸ To avoid preferential vapor loss of one component, ratios of gas to liquid volume were kept low, never exceeding 15 to 1.

The flask and acid were brought to temperature before the hydrocarbon mixture was added. After "emulsification" by magnetic stirring for the desired time, stirring was stopped and a sample of the hydrocarbon layer was pipetted into a cold 30% solution of potassium hydroxide. The hydrocarbon layer was then removed and stored over solid potassium hydroxide before analysis. Precautions were taken throughout to prevent change in composition of the hydrocarbon sample after removal from the thermostated reactor either by isomerization or preferential vapor loss.

Analysis.- An infrared spectrometric technique was employed³ using a pair of matched cells and substitution against a standard mixture. As appropriate, this was 59.18 mole% 2-methylhexane and 40.82 mole% 3-methylhexane or 62.13 mole% 2,4-dimethylpentane and 37.87 mole% 2,3-dimethylpentane. In the first system, measurements were made at 7.43, 8.63, 10.43, 11.09 and 12.98 μ ons; in the second, at 8.68, 9.02, 9.96, 10.46, 12.38 and 12.78.

In the methylhexane system, a plot of $\log \% \text{ transmission vs. } \lambda$ 2-methylhexane was prepared at each of the five wave lengths. The standard mixture was always in Cell A and the transmission was always adjusted to read 100% with Cell A in the beam. The plots were prepared from readings with the various synthetic mixtures prepared for isomerization runs in Cell B. Average deviations from smooth curves was less than 0.1%. Analyses could be performed on synthetic mixtures giving agreeing results to this accuracy at all wave lengths. With products of isomerization runs, particularly those extending over longer periods of time, there was some spread in the values determined for each wave length owing to side reactions.

Results and Discussion

By exposing mixtures of hydrocarbons in the vicinity of the equilibrium mixture to isomerizing conditions for relatively short times, the position of the steady state has been located with minimum interference by side reactions. We feel that the position of the steady state, so located, is accurate to within a few tenths of 1%. The experimental data are shown in Tables I and II.

However, as Bates has pointed out,⁶ in the presence of side

(6) J.R. Bates, J. Am. Chem. Soc., 68, 511 (1946).

Table I

The 2-Methylhexane-3-Methylhexane Equilibrium

Temp. °	Reaction Time, Hours	Reactant Mixture, % 2-Methylhexane	Product Mixture, Av. % 2MH	Average Deviation%
60	1.00	56.7	58.0	0.4
	1.00	60.4	60.0	.3
	.50	59.2	59.5	.1
	.50	59.8	59.6	.3
0	.67	65.6	64.6	.8
	.33	63.5	62.8	.4
	.25	62.7	62.8	.2
	.67	60.5	62.0	.4
-33.4	1.00	65.3	65.5	.7
	.50	65.7	65.5	.3
	1.00	65.4	65.5	.2

Table II

The 2,3-Dimethylpentane-2,4-Dimethylpentane Equilibrium

Temp. °	Reaction Time, Hours	Reactant Mixture, %2,4-Dimethylpentane	Product Mixture, Av. % 2,4MH	Average Deviation%
60.	.16	60.2	60.0	0.1
	.16	59.5	59.6	.2
	.16	59.8	59.7	.2
0.	.16	70.5	70.6	.1
	.16	70.8	70.6	.1
	.16	70.6	70.6	.1
-33.4	.16	74.2	73.0	.2
	.16	71.4	72.2	.2
	.16	72.7	72.8	.1
	.16	73.1	72.9	.1

reactions, the steady state does not necessarily correspond to the equilibrium composition. Since the interconversion of 2- and 3-methylhexane proceeds about 10 times as fast as the formation of dimethylpentanes,⁵ the error introduced into the equilibrium constant is small.⁵ Furthermore, at 60°, at any rate, if the mechanism advanced⁵ is correct, no error is introduced. The small spread in the analyses at different wave lengths (Tables I and II) would indicate that no serious error was introduced by side reactions.

Computed equilibrium compositions and constants are presented in Table III where they are compared with the values of Voorthuijsen,⁵

Table III
Equilibrium Results

Temp. °	3MH = 2MH ^a			23MP = 24MP ^b		
	X_{2MH}^{eq}	K	$\Delta F_{cal.}^{\circ}$	X_{24MP}^{eq}	K	ΔF°
60 ^c	0.595	1.47	-255	0.596	1.48	-260
0 ^c	.628	1.69	-285	.706	2.40	-475
-33.4 ^c	.655	1.90	-306	.728	2.68	-470
25 ^d		1.35	-180		1.90	-380
20 ^e		1.63	-284		1.50	-236
60 ^f		1.05	-29		0.25	+921
0 ^f		1.27	-130		.29	+668
-33.4 ^f		1.49	-191		.33	+525

^a 2MH and 3MH are 2- and 3-methylhexane, respectively.

^b 23MP and 24MP are 2,3- and 2,4-dimethylpentane, respectively.

^c Results of this investigation.

^d Voorthuijsen (ref. 5).

^e Roebuck and Evering (ref. 7).

^f Rossini, et al. (ref. 8).

the recent values of Roebuck and Evering,⁷ and the computed values

(7) A.K. Roebuck and B.L. Evering, *ibid.*, 75, 1631 (1953). These values were obtained by direct equilibration with sulfuric acid.

of Rossini.⁸

 (8) American Petroleum Institute, Research Project 44 at the National Bureau of Standards, "Selected Values of Properties of Hydrocarbons." F.R. Rossini, E.J. Prosen and K.S. Pitzer, J. Res. Nat. Bureau of Standards. 27, 529 (1941). The values of ΔF° for the vapor state were corrected by the Antoine Equation from the first cited reference to the liquid state assuming Raoult's Law.

In terms of the experimental procedures employed, the experimental results of previous investigators should not be taken as diverging seriously from our results since the deviation in K corresponds to a deviation in X^{eq} of but a few %. Similarly the computed results of Rossini⁸ may be taken as in quite satisfactory agreement for the equilibrium between the methylhexanes. The discrepancy in the case of the dimethylpentanes is larger and amounts to about 1000 cal. in terms of ΔF° .

Within the accuracy of the data, the values of ΔF° for the methylhexane reaction are representable by $\Delta H^\circ = -436$ cal. and $\Delta S^\circ = +0.55$ cal./deg. Standard states are the pure liquids.

The rate of change of ΔF° with temperature is too large to permit an accurate computation of values of ΔH° and ΔS° from the data for the dimethylpentanes. ΔS° changes sign in the vicinity of 0° at which temperature ΔH° has a value of about -475 cal.

Evanston, Illinois

DISTRIBUTION LIST FOR TECHNICAL REPORTS

Northwestern Univ. - ISOMERIZATION AND EXCHANGE IN HYDROCARBONS - Contract
N7onr-45006

NR-056-199

<u>No. of Copies</u>	<u>Addresses</u>	<u>No. of Copies</u>	<u>Addresses</u>
1	Commanding Officer Office of Naval Research Br. Office 150 Causeway Street Boston, Massachusetts	1	Research and Development Board Pentagon, Room 3d1041 Washington 25, D. C. Attn: Technical Reference Sec.
2	Commanding Officer Office of Naval Research Br. Office 844 North Rush Street Chicago 11, Illinois	1	Dr. Ralph G.H. Siu, Res. Dir. General Laboratories, QM Depot 2800 South 20th Street Philadelphia 45, Pennsylvania
1	Commanding Officer Office of Naval Research Br. Office 346 Broadway New York 13, New York	1	Dr. Warren Stubblebine, Research Director Chemical and Plastics Section RDB-MPD Quartermaster General's Office Washington 25, D. C.
1	Commanding Officer Office of Naval Research Br. Office 1000 Geary Street San Francisco 9, California	1	Dr. A. Stuart Hunter, Technical Director Res. and Dev. Branch MPD Quartermaster General's Office Washington 25, D. C.
1	Commanding Officer Office of Naval Research Br. Office 1030 North Green Street Pasadena 1, California	1	Dr. A. G. Horney Wright Air Development Center Wright-Patterson Air Force Base Dayton, Ohio Attn: WCRRS -4
2	Officer-in-Charge Office of Naval Research Br. Office Navy Number 100 Fleet Post Office New York, New York	1	Dr. A. Weissler Department of the Army Office of the Chief of Ordnance Washington 25, D. C. Attn: ORDTB-PS
9	Director, Naval Research Laboratory Washington 25, D. C. Attn: Technical Information Off.	1	Research and Development Group Logistics Division, Gen. Staff Department of the Army Washington 25, D. C. Attn: Dr. W.T. Read, Scientific Adviser
4	Chief of Naval Research Office of Naval Research Washington 25, D. C. Attn: Chemistry Branch		
2	Director, Naval Research Laboratory Washington 25, D. C. Attn: Chemistry Division		

DISTRIBUTION LIST FOR TECHNICAL REPORTS - page 2

Northwestern Univ. - ISOMERIZATION AND EXCHANGE IN HYDROCARBONS - Contract
F7onr-45006

NR-056-199

<u>No. of Copies</u>	<u>Addresses</u>	<u>No. of Copies</u>	<u>Addresses</u>
2	Chief of the Bureau of Ships Navy Department Washington 25, D. C. Attention: Code 340	1	Technical Command Chemical Corps Chemical Center, Maryland
2	Chief of the Bureau of Aeronautics Navy Department Washington 25, D. C. Attention: Code Td-4	1	U.S. Atomic Energy Commission Chemistry Division Brookhaven National Laboratory Upton, New York
2	Chief of the Bureau of Ordnance Navy Department Washington 25, D. C. Attention: Code Rexd	1	U.S. Atomic Energy Commission Research Division Washington 25, D. C.
5	Mr. J. H. Heald Library of Congress Nav. Research Section Washington 25, D. C.	1	U.S. Atomic Energy Commission Library Br., Tech. Info., ORE P. O. Box E Oak Ridge, Tennessee
1	Dr. H. A. Zahl, Tech. Director Signal Corps Engineering Labs. Fort Monmouth, New Jersey	1	Dr. Henry Eyring University of Utah Salt Lake City 1, Utah
1	U.S. Naval Radiological Def. Lab San Francisco 24, California Attention: Technical Library	1	Dr. M. S. Newman Department of Chemistry Ohio State University Columbus 10, Ohio
1	Naval Ordnance Test Station Inyokern CHINA LAKE, California Attn: Head, Chemistry Division	1	Dr. P. D. Bartlett Department of Chemistry Harvard University Cambridge, Massachusetts
1	Office of Ordnance Research 2127 Myrtle Drive Durham, North Carolina	1	Dr. D. J. Cram University of California Los Angeles 7, California
1	Dr. Louis P. Hammett Department of Chemistry Columbia University New York 27, New York	1	Dr. J. D. Danforth Division of Natural Sciences Grinnell College Grinnell, Iowa