

Reproduced by

Armed Services Technical Information Agen
DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

AD -

17504

UNCLASSIFIED

AD No. 17504
ASTIA FILE COPY

Technical Report No. 4
THE FINITE STURM-LIOUVILLE TRANSFORM

By

A. Cemal Eringen

to

Office of Naval Research
Department of the Navy

Contract N7onr-32909

Department of Mechanics
Illinois Institute of Technology
Technology Center
Chicago, Illinois

1 August 1953

Acknowledgment

The author is deeply indebted
to Mr. Thomas King for checking the
analysis.

TABLE OF CONTENTS

	Page
Acknowledgment	1
Table of Contents	11
1. Introduction.	1
2. The Sturm-Liouville Expansion	2
3. Finite Sturm-Liouville Transform	4
4. Transform of Lf	5
5. Solution of Some Partial Differential Equations	6
6. Solutions ϕ_0 and χ_0 of Some Special Equations	9
7. Application	11
References	17

The Finite Sturm-Liouville Transform

By

A. Cemal Eringen
Illinois Institute of Technology

1. Introduction

It is known that certain partial differential equations can be solved with the use of particular types of definite integrals having appropriate kernels. The choice of kernel depends on the type of boundary value problem. The solution obtained by these transforms is direct in the sense that it contains the boundary values in the solution. This, of course, is lacking in the classical approach. Finite Fourier transforms are of this type [1].

Recently Tranter [2], used a Legendre polynomial as a kernel. Scott [3] following a similar approach used a Jacobi polynomial as a kernel which extends the result of [2].

It is the purpose of the present paper to extend and unify all such special transforms. Thus we employ a kernel which may be determined to suit each particular type of problem. The transform can be employed to solve a wide class of linear second order partial differential equations.

In this paper we deal only with those transforms whose intervals are finite. Thus, the results obtained here are particularly useful for finite domains, or domains which are finite in one direction. An extension to infinite domains and singular cases will be made in a later paper.

2. The Sturm-Liouville Expansion

A second order, linear, homogeneous differential equation containing an arbitrary parameter λ has the general form:

$$Mv = p_0 v_{,yy} + p_1 v_{,y} + (p_2 + \lambda p_3)v = 0 \quad (1)$$

$$a_0 \leq y \leq b_0$$

where $p_1 = p_1(y)$; v and y are the dependent and real independent variables, respectively. Indices after comma represent differentiation, i.e.: $\phi_{,y} = d\phi/dy$.

Equation (1) can be transformed into the canonical form:

$$Lu = \lambda u, \quad L = q(x) - \frac{d^2}{dx^2}, \quad a \leq x \leq b \quad (2)$$

where:

$$\left. \begin{aligned} u &= \theta(x)v, \quad x = \int (p_3/p_0)^{1/2} dy \\ a &= \int^{a_0} (p_3/p_0)^{1/2} dy, \quad b = \int^{b_0} (p_3/p_0)^{1/2} dy \\ q(x) &= \frac{\theta_{,xx}(x)}{\theta(x)} - \phi(x), \quad \theta(x) = (p_3 k^2/p_0)^{1/4} \\ \phi(x) &= p_2/p_3, \quad k = \exp \int (p_1/p_0) dy \end{aligned} \right\} \quad (3)$$

It is simpler to work with (2).

We consider two solutions $\phi(x, \lambda)$ and $\chi(x, \lambda)$ of (2) satisfying the boundary conditions:

$$\left. \begin{aligned} \phi(a, \lambda) &= \sin \alpha, \quad \phi_{,x}(a, \lambda) = -\cos \alpha \\ \chi(b, \lambda) &= \sin \beta, \quad \chi_{,x}(b, \lambda) = -\cos \beta \end{aligned} \right\} \quad (4)$$

Both solutions are unique if $q(x)$ is real, continuous everywhere in (a, b) and has finite limits for $x = a$ and $x = b$

(see [4], p.6). We can build up $\phi(x, \lambda)$ and $\chi(x, \lambda)$ as a linear combination of two independent solutions $\phi_0(x, \lambda)$ and $\chi_0(x, \lambda)$ of (2). For let:

$$\left. \begin{aligned} \omega_0(\lambda)\phi(x, \lambda) &= \phi_0(x, \lambda)[\chi_0(a, \lambda)\cos\alpha + \chi_{0,x}(a, \lambda)\sin\alpha] \\ &\quad - \chi_0(x, \lambda)[\phi_0(a, \lambda)\cos\alpha + \phi_{0,x}(a, \lambda)\sin\alpha] \\ \omega_0(\lambda)\chi(x, \lambda) &= \phi_0(x, \lambda)[\chi_0(b, \lambda)\cos\beta + \chi_{0,x}(b, \lambda)\sin\beta] \\ &\quad - \chi_0(x, \lambda)[\phi_0(b, \lambda)\cos\beta + \phi_{0,x}(b, \lambda)\sin\beta] \\ \omega_0(\lambda) &= W(\phi_0, \chi_0) = \phi_0(x, \lambda)\chi_{0,x}(x, \lambda) - \phi_{0,x}(x, \lambda)\chi_0(x, \lambda) \end{aligned} \right\} (5)$$

Functions $\phi(x, \lambda)$ and $\chi(x, \lambda)$ defined by (5) satisfy (2) and (4). Here $W(u, v)$ is the Wronskian, and can be shown to be independent of x .

Let λ_n be a root of $W(\phi, \chi) = 0$. Then we have:

$$\omega(\lambda_n) = W(\phi_n, \chi_n) = \phi_n \chi_{n,x} - \phi_{n,x} \chi_n = 0$$

or

$$\phi_{n,x}/\phi_n = \chi_{n,x}/\chi_n$$

where:

$$\phi_n = \phi(x, \lambda_n), \quad \text{and} \quad \chi_n = \chi(x, \lambda_n).$$

Hence the integration gives:

$$\chi(x, \lambda_n) = k_n \phi(x, \lambda_n) \quad (6)$$

Consequently $\phi(x, \lambda_n)$ and $k_n \phi(x, \lambda_n)$ satisfy all of the boundary conditions (4).

It is also easy to show that $\phi(x, \lambda_n)$ is an orthogonal set. For, in view of (2), (4) and (6) we have:

$$(\lambda_m - \lambda_n) \int_a^b \phi_m \phi_n dx = \int_a^b (\phi_m L \phi_n - \phi_n L \phi_m) dx = W(\phi_m, \phi_n) = 0.$$

$$(\lambda_m \neq \lambda_n)$$

Hence we can write:

$$\int_a^b \phi(x, \lambda_n) \phi(x, \lambda_m) dx = N^2 \delta_{mn} \quad (7)$$

where δ_{mn} is the Kronecker delta (equal to zero for $m \neq n$, one for $m=n$)

It can be shown also that all roots of $\omega(\lambda) = 0$ are real and distinct (see [4], pp 11-12).

From (5) and (6) it follows that:

$$k_n = \frac{\phi_0(b, \lambda_n) \cos \beta + \phi_{0,x}(b, \lambda_n) \sin \beta}{\phi_0(a, \lambda_n) \cos \alpha + \phi_{0,x}(a, \lambda_n) \sin \alpha}$$

$$= \frac{\chi_0(b, \lambda_n) \cos \beta + \chi_{0,x}(b, \lambda_n) \sin \beta}{\chi_0(a, \lambda_n) \cos \alpha + \chi_{0,x}(a, \lambda_n) \sin \alpha} \quad (8)$$

If now $f(x)$ is an integrable function over (a, b) and if $a < x < b$, we have the Sturm-Liouville expansion of $f(x)$:

$$f(x) = \sum_{n=0}^{\infty} [k_n / \omega_{\lambda}(\lambda_n)] \phi(x, \lambda_n) \int_a^b \phi(y, \lambda_n) f(y) dy \quad (9)$$

For the convergence of (9) to $f(x)$ see [4] pp 12-15 or [5] pp 275-276.

3. Finite Sturm-Liouville Transform

Let $f(x)$ be a real continuous and integrable function of x , in the interval (a, b) . We define the finite Sturm-Liouville transform $S\{f\} = F(\lambda_n)$ associated with system (2) and (4) by:

$$F(\lambda_n) = S\{f\} = \int_a^b f(y) \psi(y, \lambda_n) dy \quad (10)$$

where:

$$\psi(y, \lambda_n) = [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} \phi(y, \lambda_n) \quad (11)$$

The inversion theorem now follows from (9):

$$f(x) = \sum_{n=0}^{\infty} F(\lambda_n) \psi(x, \lambda_n) \quad (12)$$

Here functions $\psi(y, \lambda_n)$ are orthonormal, i.e.:

$$\int_a^b \psi(x, \lambda_n) \psi(x, \lambda_m) dx = \delta_{mn} \quad (13)$$

This is readily seen from (7) and (11).

Equations (10) and (12) are basic for the solution of some partial differential equations.

4. Transform of Lf

We are now going to prove that:

$$\begin{aligned} S_x \{ Lf \} &= S_x \{ q(x)f - \partial^2 f / \partial x^2 \} = B_f(\lambda_n, t) \\ &+ \lambda_n F(\lambda_n, t) \end{aligned} \quad (14)$$

where:

$$\begin{aligned} B_f(\lambda_n, t) &= [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} \left\{ -f(b, t) \frac{\cos \beta}{k_n} \right. \\ &\left. - f_{,x}(b, t) \frac{\sin \beta}{k_n} + f(a, t) \cos \alpha + f_{,x}(a, t) \sin \alpha \right\} \end{aligned} \quad (15)$$

is a function depending on the boundary values of $f(x, t)$, $f_{,x}(x, t)$ at $x = a, b$. Subscript f implies that the function B contains boundary values of the function f . Here S_x represents the finite Sturm-Liouville transform taken with respect to x :

$$S_x \{ \phi(x,t) \} = \bar{\phi}(\lambda_n, t) \quad (16)$$

To prove (14) we multiply Lf by $\psi(x, \lambda_n)$ and integrate between (a, b) , that is, by definition we have:

$$S_x \{ Lf \} = \int_a^b [q(x)f(x,t) - \partial^2 f / \partial x^2] \psi(x, \lambda_n) dx \quad (17)$$

Integrating the second term in the integrand of (17) twice by part we obtain:

$$S_x \{ Lf \} = -[f_{,x} \psi - \psi_{,x} f]_a^b + \int_a^b f(x,t) L\psi(x, \lambda_n) dx \quad (18)$$

If we now use (4) and (6) in the first term of the right hand side and (2) inside the integrand we obtain (14).

5. Solution of Some Partial Differential Equations

a) Let us consider a parabolic partial differential equation

$$Mv + p_4 v_{,t} = 0 \quad (19)$$

where Mv is defined by (1), $p_i = p_i(y, t)$ are known and $v(y, t)$ is the unknown dependent variable.

The use of transformation (3) reduces (19) to:

$$u_{,xx} + [\lambda - p(x, t)]u + (p_4/p_3)u_{,t} = 0 \quad (20)$$

where:

$$p(x, t) = q(x, t) - p_4 p_0 (k/2p_0)^2 \frac{\partial}{\partial t} (p_0/p_3 k^2) \quad (21)$$

Here $q(x, t)$, $k(t)$ and $\lambda(t)$ are given by (3) with t as a parameter. If $\phi(x, t, \lambda_n)$ and $\psi(x, t, \lambda_n)$ are now obtained as before except with $q(x)$ replaced by $p(x, t)$, t a parameter,

and p_4/p_3 a function of t alone, then (20), with the use of the finite Sturm-Liouville transform, can be reduced to an ordinary differential equation in t . For, let $p_4/p_3 = r(t)$, then application of the transform to (20) gives:

$$-B_u(\lambda_n, t) + (\lambda - \lambda_n)\bar{u}(\lambda_n, t) + r(t)\bar{u}_{,t}(\lambda_n, t) = 0 \quad (22)$$

Equation (22) is of first order. The complete solution is:

$$\begin{aligned} \bar{u}(\lambda_n, t) = & C(\lambda_n) \exp\left(\int^t \frac{\lambda_n - \lambda}{r} dt\right) + \left[\exp\left(\int^t \frac{\lambda_n - \lambda}{r} dt\right)\right] \cdot \\ & \int^t (B_u/r) \left[\exp\left(-\int^t \frac{\lambda_n - \lambda}{r} dt\right)\right] dt \end{aligned} \quad (23)$$

Now, inversion theorem (12) gives:

$$u(x, t) = \sum_{n=0}^{\infty} \bar{u}(\lambda_n, t) \psi_n(x, t, \lambda_n) \quad (24)$$

Therefore, the solution of (19) is effected. We must, however, remember that at least one of each pair of boundary values $u(a, t)$, $u_{,x}(a, t)$ and $u(b, t)$, $u_{,x}(b, t)$ must be known. By selecting α and β properly in $B_u(\lambda_n, t)$ we can make the terms containing the unknown pair zero. Of course any knowledge of the boundary values of u leading to the complete evaluation of B_u is sufficient for the solution.

b) A second order linear partial differential equation containing a parameter λ has the general form:

$$\begin{aligned} Nv + a_4 v_{,z} + a_5 v_{,\tau} + (a_6 + \lambda a_7) v &= 0 \\ Nv = a_1 v_{,zz} + 2a_2 v_{,z\tau} + a_3 v_{,\tau\tau} \end{aligned} \quad (25)$$

where $a_1 = a_1(z, \tau)$. If the equation is of elliptic type we have:

$$a_1 a_3 - a_2^2 > 0 \quad (26)$$

Consider now an elliptic equation (25). Let $\xi(z, \tau) = \text{const.}$ and $\eta(z, \tau) = \text{const.}$, respectively be the solutions of the following differential equations:

$$\begin{aligned} d\tau/dz &= \zeta_1(z, \tau) & d\tau/dz &= \zeta_2(z, \tau) \\ \left. \begin{matrix} \zeta_1 \\ \zeta_2 \end{matrix} \right\} &= -\frac{a_2}{a_1} \pm 1 \left[\frac{a_3}{a_1} - \left(\frac{a_2}{a_1} \right)^2 \right]^{1/2} \end{aligned} \quad (27)$$

If we set:

$$2y = \xi + \eta, \quad 2it = \xi - \eta \quad (28)$$

equation (25) can be transformed into the canonical form [6]:

$$v_{,yy} + p_1 v_{,y} + (p_2 + \lambda p_3) v + p_4 v_{,t} + v_{,tt} = 0 \quad (29)$$

where:

$$\begin{aligned} p_1 &= \frac{2}{\beta} p_y, \quad p_2 = 2a_6/\beta, \quad p_3 = 2a_7/\beta \\ p_4 &= \frac{2}{\beta} p_t, \quad \beta = a_1(y_{,z}^2 + t_{,z}^2) + 2a_2(y_{,z}y_{,\tau} + t_{,z}t_{,\tau}) \\ &\quad + a_3(y_{,\tau}^2 + t_{,\tau}^2) \end{aligned} \quad (30)$$

Here operator P is defined by:

$$P\phi = N\phi + a_4\phi_{,z} + a_5\phi_{,\tau} \quad (31)$$

and $N\phi$ is given by the second of (25).

Using the transformation (3) with $p_0 = 1$, (29) becomes:

$$u_{,xx} + (\lambda - p)u + r_1 u_{,t} + r_0 u_{,tt} = 0 \quad (32)$$

where:

$$\begin{aligned} p &= q(x,t) - mp_4g_{,t} - mg_{,tt}, \quad r_0 = p_3^{-1} \\ r_1 &= mp_4g + 2mg_{,t}, \quad g = (p_3k^2)^{-1/4}, \\ m &= (k^2/p_3^3)^{1/4} \end{aligned} \quad (33)$$

In the special case where r_1 , and r_0 are functions of t alone equation (32) can be transformed into an ordinary differential equation by applying the finite Sturm-Liouville transform.

Hence:

$$r_0 \bar{u}_{,tt} + r_1 \bar{u}_{,t} + (\lambda - \lambda_n) \bar{u}(\lambda_n, t) = B_u(\lambda_n, t) \quad (34)$$

We can now either use transformation (3) with respect to t , with r_0 , r_1 replacing p_0 , p_1 and $p_2 = 0$, $p_3 = 1$ and find the solution of (34) and then invert it or we can solve (34) directly for $\bar{u}(\lambda_n, t)$ and then (24) gives $u(x, t)$.

It must be remembered that this method has the advantage that it contains the boundary values of the function within the solution, consequently, the difficulty of satisfying boundary conditions is eliminated. The method is also valid in the cases where the usual technique of separation of variables fail.

6. Solutions ϕ_0 and χ_0 of Some Special Equations

Below are given two independent solutions ϕ_0 and χ_0 of some special differential equations (D. E.) of mathematical physics. Their canonical forms (2) will be given by only determining u , q , and λ .

a) Bessel Functions

$$v_{,yy} + \frac{1}{y} v_{,y} + (s^2 - \frac{v^2}{y^2})v = 0 \quad (\text{D. E.})$$

$$u = x^{1/2}v, \quad y = x, \quad q = (v^2 - \frac{1}{4})x^{-2}, \quad \lambda = s^2$$

$$\phi_0(x, \lambda) = x^{1/2} J_v(xs), \quad \chi_0(x, \lambda) = x^{1/2} Y_v(xs)$$

Interval (a,b) should not contain $x = 0$.

b) Spherical Harmonics

$$[(1-y^2)v_{,y}]_{,y} + [v(v+1) - \frac{\mu^2}{1-y^2}]v = 0 \quad (\text{D. E.})$$

$$u = v \cos^{1/2}x, \quad y = \sin x, \quad q = \frac{\mu^2}{\cos^2 x} - \frac{1}{4} \tan^2 x - \frac{1}{2},$$

$$\lambda = v(v+1), \quad \phi_0(x, \lambda) = \cos^{1/2}x \cdot P_v^\mu(\sin x),$$

$$\chi_0(x, \lambda) = \cos^{1/2}x \cdot Q_v^\mu(\sin x)$$

Interval (a,b) should not contain $(-1, 1)$.

c) Hermite Polynomials

$$v_{,yy} - yv_{,y} + nv = 0 \quad (\text{D. E.})$$

$$u = e^{-x^2/4}v, \quad y = x, \quad q = -\frac{1}{2} + \frac{x^2}{4}, \quad \lambda = n,$$

$$\phi_0(x, \lambda) = e^{-x^2/4} \text{He}_n(x), \quad \chi_0(x, \lambda) = e^{-x^2/4} \text{he}_n(x)$$

d) Tschebyscheff Polynomials

$$(1-y^2)v_{,yy} - yv_{,y} + n^2v = 0 \quad (\text{D. E.})$$

$$u = v, \quad y = \sin x, \quad q = 0, \quad \lambda = n^2$$

$$\phi_0(x, \lambda) = T_n(\sin x) = \cos n(\sin^{-1}y),$$

$$\chi_0(x, \lambda) = U_n(\sin x) = \sin n(\sin^{-1}y)$$

e) Mathieu Functions

$$u_{,xx} + (\lambda - 2h^2 \cos 2x)u = 0 \quad (D. E.)$$

$$\phi_0(x, \lambda) = ce_{2n}(x), \quad ce_{2n+1}(x), \quad se_{2n}(x), \quad s_{2n+1}(x)$$

$$\chi_0(x, \lambda) = ce_{2n}^{(2)}(x), \quad ce_{2n+1}^{(2)}(x), \quad se_{2n}^{(2)}(x)$$

$$s_{2n+1}^{(2)}(x)$$

($n = 0, 1, 2, \dots$ except for se_{2n} only $n = 1, 2, \dots$)

$\phi_0(x, \lambda)$ is periodic.

f) Wittaker Functions

$$v_{,yy} + \left(-\frac{1}{4} + \frac{k}{y} + \frac{\frac{1}{4} - \mu^2}{y^2}\right)v = 0 \quad (D. E.)$$

$$u = (x/2)^{1/2}v, \quad y = x^2/4, \quad q = \frac{16\mu^2 - 17}{4x^2} + \frac{x^2}{16}$$

$$\phi_0(x, \lambda) = (x/2)^{1/2} W_{k, \mu}(x^2/4), \quad \chi_0(x, \lambda) = (x/2)^{1/2}$$

$$W_{-k, \mu}(-x^2/4) \quad (*)$$

Interval (a, b) should not contain $x = 0$.

7. Application

As an illustration we solve a heat conduction problem which may be looked upon as a mathematical model of volcanos. The problem also has application in the exhaust parts of jet engines. As far as I know the problem has not previously been solved.

* Here comma does not mean differentiation.

Consider the conical shell enclosed by two coaxial cones having the same apex O , and two concentric spheres having O as their center (Fig. 1).




Fig. 1

In polar coordinates the inner and outer conical surfaces are given by $\theta = \theta_0$ and $\theta = \theta_1$, and the end surfaces by $r = r_0$ and $r = r_1$.

Problem: Determine the steady temperature distribution within the shell under the general axi-symmetric boundary conditions in temperature. This is the general Dirichlet problem for the domain under consideration. The differential equation and boundary conditions (B. C.) are given below:

$$\begin{aligned} \text{(D. E.) } \Delta V &= r^{-2}(r^2 V_{,r})_{,r} + r^{-2}[(1-y^2)V_{,y}]_{,y} = 0 \\ y &= \cos \theta, \quad r_0 < r < r_1, \quad \theta_0 < \theta < \theta_1 \end{aligned} \quad (35)$$

$$\begin{aligned} \text{(B. C.) } V &= V_0(r) \quad \text{for } \theta = \theta_0 \\ V &= V_1(r) \quad \text{for } \theta = \theta_1 \end{aligned} \left. \vphantom{\begin{aligned} V &= V_0(r) \\ V &= V_1(r) \end{aligned}} \right\} r_0 < r < r_1 \\ \\ V &= V_3(\theta) \quad \text{for } r = r_0 \\ V &= V_4(\theta) \quad \text{for } r = r_1 \end{aligned} \left. \vphantom{\begin{aligned} V &= V_3(\theta) \\ V &= V_4(\theta) \end{aligned}} \right\} \theta_0 < \theta < \theta_1 \quad (36)$$

Here $V(r, y)$ is the temperature function, r and θ are the polar coordinates.

Solution: We can exclude the second term of (D. E.) (35) if we use a finite Sturm-Liouville transform associated with the Legendre equation:

$$[(1-y^2)v_{,y}]_{,y} + v(v+1)v = 0 \quad (37)$$

In view of (6, b) with $\mu = 0$ we find that if we select $y = \sin x$, $u = v \cos^{1/2} x$ we can transform (37) to canonical form leading to solutions:

$$\begin{aligned} \phi_0(x, \lambda) &= \cos^{1/2} x \cdot P_v(\sin x), \quad \chi_0(x, \lambda) = \cos^{1/2} x \\ &\cdot Q_v(\sin x), \quad \lambda = v(v+1) \end{aligned} \quad (38)$$

where P_v and Q_v are Legendre functions of the first and second kind, respectively. In view of (14), when we apply the transform to (35) the second term gives $-v_n(v_n+1) - B_v(r, \lambda_n)$. Now B_v contains four arbitrary functions which must be specified on the surfaces $\theta = \theta_0$ and $\theta = \theta_1$. The terms containing the derivatives of v are not given. Thus if we

select $\alpha = \beta = 0$ these terms drop out, leaving the terms containing $v_0(r)$ and $v_1(r)$, which are given. Hence $\phi(x, \lambda)$ and $\omega_0(\lambda)$ of (5) become:

$$\begin{aligned} \phi(x, \lambda) &= \cos^{1/2} x_0 \cos^{1/2} x [P_v(\sin x) Q_v(\sin x_0) \\ &\quad - P_v(\sin x_0) Q_v(\sin x)] , \quad x_0 = \frac{\pi}{2} - \theta_0 , \\ x_1 &= \frac{\pi}{2} - \theta_1 , \quad \omega_0(\lambda) = 1 \end{aligned} \quad (39)$$

By (11) and (8) we have:

$$\begin{aligned} \psi(x, \lambda_n) &= [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} \phi(x, \lambda_n) , \\ k_n &= \frac{\cos^{1/2} x_1 P_{v_n}(\sin x_1)}{\cos^{1/2} x_0 P_{v_n}(\sin x_0)} \end{aligned} \quad (40)$$

Calculation of $\omega(\lambda) = \phi \chi_x - \chi \phi_x$ gives

$$\begin{aligned} \omega(\lambda) &= \cos^{1/2} x_0 \cos^{1/2} x_1 [P_v(\sin x_0) Q_v(\sin x_1) \\ &\quad - P_v(\sin x_1) Q_v(\sin x_0)] \end{aligned} \quad (41)$$

Consequently, the roots $\lambda_n = v_n(v_n+1)$ of $\omega(\lambda) = 0$ satisfy the following equation

$$\frac{P_v(\sin x_1)}{P_v(\sin x_0)} = \frac{Q_v(\sin x_1)}{Q_v(\sin x_0)} \quad (42)$$

Therefore $\psi(x, \lambda_n)$ is completely determined.

After transforming (35) with $y = \sin x$ and $u = v \cos^{1/2} x$ we apply the finite Sturm-Liouville transform. That is, we multiply the equation by $\psi(x, \lambda_n)$ and integrate between x_0 and x_1 . The result is:

$$(r^2 \bar{V}_{,r})_{,r} - v_n(v_n+1) \bar{V} = B_V(r, \lambda_n) \quad (43)$$

$$B_V(r, \lambda_n) = [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} [V_0(r) - k_n^{-1} V_1(r)]$$

Equation (43) is an ordinary differential equation of Euler type whose solution can be found by variation of parameters.

Hence:

$$\begin{aligned} \bar{V}(r, v_n) &= C_1(v_n) r^{v_n} + C_2(v_n) r^{-v_n-1} + F(r, v_n) \\ F(r, v_n) &= \int^r \frac{B_V(\rho, \lambda_n)}{1+2v_n} (r^{v_n} \rho^{-v_n-1} - r^{-v_n-1} \rho^{v_n}) d\rho \end{aligned} \quad (44)$$

Let the transforms of $V_3(\theta)$ and $V_4(\theta)$ be $\bar{V}_3(v_n)$ and $\bar{V}_4(v_n)$.

C_1 and C_2 will then be determined from the remaining conditions:

$$\begin{aligned} \bar{V} &= \bar{V}_3(v_n) \quad , \quad \text{for } r = r_0 \\ \bar{V} &= \bar{V}_4(v_n) \quad , \quad \text{for } r = r_1 \end{aligned} \quad \theta_0 < \theta < \theta_1$$

This gives two linear equations for C_1 and C_2 whose solutions are:

$$\begin{aligned} C_1(v_n) &= \frac{[\bar{V}_3(v_n) - F(r_0, v_n)] r_1^{-v_n-1} - [\bar{V}_4(v_n) - F(r_1, v_n)] r_0^{-v_n-1}}{r_0^{v_n} r_1^{-v_n-1} - r_1^{v_n} r_0^{-v_n-1}} \\ C_2(v_n) &= \frac{[\bar{V}_4(v_n) - F(r_1, v_n)] r_0^{v_n} - [\bar{V}_3(v_n) - F(r_0, v_n)] r_1^{v_n}}{r_0^{v_n} r_1^{-v_n-1} - r_1^{v_n} r_0^{-v_n-1}} \end{aligned} \quad (45)$$

Hence, $\bar{V}(r, v_n)$ is completely determined. The inversion theorem (12) now gives:

$$V(r, x) = \sum_n \bar{V}(r, v_n) \psi(x, \lambda_n) \quad (46)$$

where the summation is extended over all roots of (42).

The analysis given above is formal. It can, however, be made rigorous by showing that the solution satisfies both the differential equation and the boundary conditions.

It may be worth while to remark that with the use of the method of separation of variables the solution of the above problem would have been difficult and lengthy, if not impossible.

References

1. Churchill, R. V., Modern Operational Mathematics in Engineering, McGraw-Hill Book Company, 1944, p. 267-291.
2. Tranter, C. J., "Legendre Transform", Quarterly Journal of Mathematics, (Oxford), (2), 1, 1950, 1-8.
3. Scott, E. J., "Jacobi Transform", Quarterly Journal of Mathematics, (Oxford), (4), 13, 1953, 36-40.
4. Titchmarsh, E. C., Eigenfunction Expansion Associated with Second Order Differential Equations, Oxford University Press, 1946.
5. Ince, E. L., Ordinary Differential Equations, Dover Publications, 1944.
6. Tamarkin, J. D. and Feller, W., Partial Differential Equations, Brown University Notes, June 23-Sept. 13, 1941. 1-5.

Contributing Personnel:

Dr. A. C. Eringen, Research Associate Professor
of Mechanics

Mr. T. King, Staff Computer

Respectfully submitted,

A. C. Eringen

A. C. Eringen

William R. Osgood

William R. Osgood
Chairman, Department
of Mechanics

Distribution List

I: Administrative, Reference and Liaison Activities of ONR

Chief of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 438 (2) Code 432 (1) Code 466(via Code 108) (1)	Library of Congress Washington 25, D. C. Attn: Navy Research Section (2)
Director, Naval Research Lab. Washington 25, D. C. Attn: Tech. Info. Officer (9) Technical Library (1) Mechanics Division (2)	Office of Naval Research Branch Office The John Crerar Library Building Tenth Floor, 86 E. Randolph St. Chicago 1, Illinois (2)

II: Department of Defense and other interested Government Activities

(A) GENERAL

Research and Development Board
Department of Defense
Pentagon Building
Washington 25, D. C.
Attn: Library(Code 3D-1075) (1)

Armed Forces Special Weapons
Project
P.O. Box 2610
Washington, D. C.
Attn: Col. G.F.Blunda (2)

Joint Task Force 3
12 St. and Const. Ave.,N.W.
(Temp. U)
Washington 25, D. C.
Attn: Major B. D. Jones (1)

(B) ARMY

Chief of Staff
Department of the Army
Research and Development Div.
Washington 25, D. C.
Attn: Chief of Res. and Dev. (1)

Office of the Chief of Engineers
Assistant Chief for Works
Department of the Army
Bldg. T-7, Gravelly Point
Washington 25, D. C.
Attn: Structural Branch
(R. L. Bloor) (1)

Engineering Research and
Development Laboratory
Fort Belvoir, Virginia
Attn: Structures Branch (1)

Distribution List (Cont.)

Office of the Chief of Engineers
Asst. Chief for Military
Construction
Department of the Army
Bldg. T-3, Gravelly Point
Washington 25, D. C.
Attn: Structures Branch
(M. F. Carey) (1)
Protective Construction
Branch (I.O.Thorley) (1)

Office of the Chief of Engineers
Asst. Chief for Military
Operations
Department of the Army
Bldg T-7, Gravelly Point
Washington 25, D. C.
Attn: Structures Development
Branch (W.F.Woollard) (1)

U. S. Army Waterways Experiment
Station
P. O. Box 631
Halls Ferry Road
Vicksburg, Mississippi
Attn: Col. H. J. Skidmore (1)

The Commanding General
Sandia Base, P.O.Box 5100
Albuquerque, New Mexico
Attn: Col. Canterbury (1)

Operations Research Officer
Department of the Army
Ft. Lesley J. McNair
Washington 25, D. C.
Attn: Howard Brackney (1)

Office of Chief of Ordnance
Office of Ordnance Research
Department of the Army
The Pentagon Annex No. 2
Washington 25, D. C.
Attn: ORDTB-PS (1)

Ballistics Research Laboratory
Aberdeen Proving Ground
Aberdeen, Maryland
Attn: Dr. C. W. Lampson (1)

(C) NAVY

Chief of Naval Operations
Department of the Navy
Washington 25, D. C.
Attn: OP-31 (1)
OP-363 (1)

Chief of Bureau of Ships
Department of the Navy
Washington 25, D. C.
Attn: Director of Research (2)
Code 423 (1)
Code 442 (1)
Code 421 (1)

Director, David Taylor Model Basin
Department of the Navy
Washington 7, D. C.
Attn: Code 720, Structures
Division (1)
Code 740, Hi-Speed
Dynamics Div. (1)

Commanding Officer
Underwater Explosion Research Div.
Code 290
Norfolk Naval Shipyard
Portsmouth, Virginia (1)

Commander
Portsmouth Naval Shipyard
Portsmouth, N. H.
Attn: Design Division (1)

Director, Materials Laboratory
New York Naval Shipyard
Brooklyn 1, New York (1)

Chief of Bureau of Ordnance
Department of the Navy
Washington 25, D. C.
Attn: Ad-3, Technical Library (1)
Roc, P.H. Girouard (1)

Naval Ordnance Laboratory
White Oak, Maryland
RFD 1, Silver Spring, Maryland
Attn: Mechanics Division (1)
Explosive Division (1)
Mech. Evaluation Div. (1)

Distribution List (Cont.)

(C) NAVY

Commander
U.S. Naval Ordnance Test Station
Inyokern, California
Post Office - China Lake, Calif.
Attn: Scientific Officer (1)

Naval Ordnance Test Station
Underwater Ordnance Division
Pasadena, California
Attn: Structures Division (1)

Chief of the Bureau of Aeronautics
Department of the Navy
Washington 25, D. C.
Attn: TD-41, Technical Library (1)

Chief of Bureau of Ships
Department of the Navy
Washington 25, D. C.
Attn: Code P-314 (1)
Code C-313 (1)

Officer in Charge
Naval Civil Engr. Research and
Evaluation Laboratory
Naval Station
Port Hueneme, California (1)

Superintendent
U.S. Naval Postgraduate School
Annapolis, Maryland (1)

(D) AIR FORCES

Commanding General
U.S. Air Forces
The Pentagon
Washington 25, D. C.
Attn: Res. and Develop. Div. (1)

Deputy Chief of Staff, Operations
Air Targets Division
Headquarters, U.S. Air Forces
Washington 25, D. C.
Attn: AFOIN-T/PV (1)

Flight Research Laboratory
Wright-Patterson Air Force Base
Dayton, Ohio
Attn: Chief, Applied Mechanics
Group (1)

(E) OTHER GOVERNMENT AGENCIES

U.S. Atomic Energy Commission
Division of Research
Washington, D. C. (1)

Director, National Bureau of
Standards
Washington, D. C.
Attn: Dr. W. H. Ramberg (1)

Forest Products Laboratory
Madison 5, Wisconsin (1)

Supplementary Distribution List

Professor Lynn Beedle
Fritz Engineering Laboratory
Lehigh University
Bethlehem, Pennsylvania (1)

Professor R. L. Bisplinghoff
Dept. of Aeronautical Engineering
Massachusetts Institute of Tech.
Cambridge 39, Massachusetts (1)

Professor G. F. Carrier
Graduate Division of Applied
Mathematics
Brown University
Providence, R. I. (1)

Professor R. J. Dolan
Dept. of Theoretical and Applied
Mechanics
University of Illinois
Urbana, Illinois (1)

Distribution List (Cont.)

Professor Hans Bleich Dept. of Civil Engineering Columbia University Broadway at 117th St. New York 27, New York	(1)	Professor L. S. Jacobsen Department of Mechanical Eng. Stanford University Stanford, California	(1)
Professor B. A. Boley Dept. of Aeronautical Eng. Ohio State University Columbus, Ohio	(1)	Dr. W. H. Hoppmann Department of Applied Mechanics Johns Hopkins University Baltimore, Maryland	(1)
Professor Lloyd Donnell Department of Mechanics Illinois Institute of Tech. Technology Center Chicago 16, Illinois	(1)	Professor George Lee Department of Mathematics Rensselaer Polytechnic Institute Troy, New York	(1)
Professor B. Fried Dept. of Mechanical Engineering Washington State College Pullman, Washington	(1)	Professor Glen Murphy, Head Department of Theoretical and Applied Mechanics Iowa State College Ames, Iowa	(1)
Mr. Martin Goland Midwest Research Institute 4049 Pennsylvania Avenue Kansas City 2, Missouri	(1)	Professor N. M. Newmark Department of Civil Engineering University of Illinois Urbana, Illinois	(1)
Dr. J. N. Goodier School of Engineering Stanford University Stanford, California	(1)	Professor Jesse Ormondroyd University of Michigan Ann Arbor, Michigan	(1)
Professor M. Hetenyi Walter P. Murphy Professor Northwestern University Evanston, Illinois	(1)	Dr. R. P. Peterson, Director Applied Physics Division Sandia Laboratory Albuquerque, New Mexico	(1)
Dr. N. J. Hoff, Head Department of Aeronautical Engineer- ing and Applied Mechanics Polytechnic Institute of Brooklyn 99 Livingston Street Brooklyn 2, New York	(1)	Dr. A. Phillips School of Engineering Stanford University Stanford, California	(1)
Dr. J. H. Hollomon General Electric Research Laboratories 1 River Road Schenectady, New York	(1)	Dr. W. Prager, Chairman Graduate Division of Applied Mathematics Brown University Providence 12, R. I.	(1)

Distribution List (Cont.)

Dr. S. Raynor Armour Research Foundation Illinois Institute of Technology Chicago 16, Illinois	(1)	Professor E. Sternberg Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)
Professor E. Reissner Department of Mathematics Massachusetts Institute of Tech. Cambridge 39, Massachusetts	(1)	Professor F. K. Teichmann Department of Aeronautical Engineering New York University University Heights, Bronx New York, N. Y.	(1)
Professor M. A. Sadowsky Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor C. T. Wang Department of Aeronautical Engineering New York University University Heights, Bronx New York, N. Y.	(1)
Professor Paul Lieber Department of Aeronautical Eng. Rensselaer Polytechnic Institute Troy, New York	(1)	Project File	(2)
Professor J. E. Stallmeyer Talbot Laboratory Department of Civil Engineering University of Illinois Urbana, Illinois	(1)	Project Staff	(5)
		For possible future distribution by the University	(10)
Dr. C. B. Smith College of Arts and Sciences Department of Mathematics Walker Hall University of Florida Gainesville, Florida	(1)		
Professor J. R. Andersen Towne School of Engineering University of Pennsylvania Philadelphia, Pennsylvania	(1)		
Commander U.S. Naval Proving Ground Dahlgren, Virginia	(1)		