

Reproduced by

Armed Services Technical Information Agency DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

AD -

17972

UNCLASSIFIED

On the Singularity at a Concentrated Load
Applied to a Curved Surface

By

E. Sternberg and R. A. Eubanks

A Technical Report to the
Office of Naval Research
Department of the Navy
Washington, D. C.

Contract N7onr-32906

Project No. NR 035-302

Department of Mechanics
Illinois Institute of Technology
Chicago, Illinois

November 1, 1953

On the Singularity at a Concentrated Load Applied to a Curved Surface*

By

E. Sternberg and R. A. Eubanks
Illinois Institute of Technology

Summary

This paper deals with the singularity at the point of application of a concentrated load acting perpendicular to a curved boundary of an elastic body. In a neighborhood of the point of application, the boundary is assumed to be representable by a sufficiently smooth arbitrary surface of revolution, the axis of which coincides with the load axis. In the event the surface is locally analytic, it is shown that the singularity is identical with that appropriate to a concentrated load applied normal to a plane boundary if and only if the curvature of the meridian of the surface vanishes at the load point. The required modified singularity for the case of non-vanishing curvature is determined in closed form to the extent where the residual surface tractions are finite and continuous.

Introduction

The singularities encountered in concentrated force problems of the theory of elasticity require special and separate treatment if one is to arrive at practically useful representations of the solution to such problems. Indeed, in order to assure results which are amenable to a complete numerical evaluation, it is essential to determine in closed form the relevant singularities at least to the extent where the residual problem is characterized by finite and continuous surface tractions. This process

*The results presented in this paper were obtained in the course of an investigation conducted under Contract N7onr-32906 with the Office of Naval Research, Department of the Navy, Washington, D. C.

was carried out in a previous paper by F. Rosenthal and one of the present authors [1]¹ in connection with the problem of the sphere under concentrated loads. In this particular case it was found that the singularity at a concentrated normal force is not identical with the well known singularity corresponding to a concentrated load applied perpendicular to a plane boundary, and that certain supplementary singularities have to be introduced to effect a reduction of the problem to one obeying the foregoing regularity requirements.

The investigation just cited suggests more general questions: Under what circumstances does Boussinesq's solution [2]² for the semi-infinite medium bounded by a plane and acted on by a concentrated surface load, supply the complete singularity appropriate to a concentrated load applied to a curved boundary? What are the supplementary singularities needed in the event the Boussinesq singularity fails to yield a regular residual problem? It is the purpose of the present paper to deal with these questions on the assumption that the load is perpendicular to the boundary and that the boundary, in a neighborhood of the load point, is representable by an arbitrary, sufficiently smooth, surface of revolution, the axis of which coincides with the load axis.

The usual conditions imposed on the solution to a problem involving concentrated surface forces, are as follows: (a) it must satisfy the field equations of the theory of elasticity throughout the body under consideration;³ (b) it must satisfy the boundary conditions for distributed

¹Numbers in brackets refer to the bibliography at the end of the paper.

²See also [3], p. 191.

³It is assumed here that concentrated forces at interior points, and otherwise singular body force distributions, are absent.

tractions;⁴ (c) it must possess a singularity at each point of application of a concentrated load such that the resultant of the tractions on any surface enclosing the load point, and lying wholly in the region occupied by the medium, tends to the corresponding prescribed concentrated load in the limit as the surface is contracted to the load point. As emphasized in [1], these three conditions, although necessary, represent an incomplete formulation of the problem and fail to characterize the solution uniquely. A unique characterization is reached by considering the modified problem in which each of the concentrated loads is replaced with an arbitrary continuous distribution of surface tractions over finite surface elements surrounding the load points. The solution to the concentrated force problem is then uniquely defined as the limit of the solution to the modified problem as the surface elements are shrunk to the load points while the resultants of the distributed tractions are made to approach the given concentrated forces.

This definition, which is analogous to Kelvin's definition through a limit process of a concentrated force at an interior point,⁵ is natural both on theoretical and on physical grounds; its usefulness ultimately depends on, and is confirmed by, experimental evidence, such as that supplied by Frocht and Guernsey [4]. The solution to a concentrated force problem which is in accord with the definition adopted here, automatically satisfies the three necessary conditions listed earlier. On the other hand, as demonstrated in [1], there exist pseudo-solutions which meet the three conditions cited but fail to agree with the foregoing limit criterion.

⁴If only concentrated loads are present, the solution must leave the boundary free from surface tractions.

⁵See, for example, [3], art. 130, p. 183.

The validity of Boussinesq's solution for the half-space under a concentrated load is readily established by applying the preceding definition of a concentrated surface force, say, to Cerruti's solution⁶ for the semi-infinite body bounded by a plane and subjected to distributed surface tractions. The pseudo-solutions to concentrated force problems discussed in [1] exhibit a common property: their singularities at the load points are of a higher order than that of the Boussinesq singularity. On the other hand, it is clear from similarity considerations that the singularity at a concentrated load applied to a curved surface must remain dominated by the Boussinesq singularity; it cannot be stronger. This observation leads to an additional necessary requirement applicable to the solution of concentrated force problems: (d) the order of the singularity at each load point must not exceed the order r^{-2} of the Boussinesq singularity.⁷

The treatment of the general problem to be considered presently will be based on conditions (a), (b), (c), and (d). In the special instance of the sphere under radial concentrated loads, the correctness of the solution so obtained was verified analytically through a limit process. The question as to whether the four necessary conditions stated always assure a unique solution, which thus coincides with the limit solution defined previously, is apparently not easy to dispose of with complete generality. Until a general affirmative answer is established, solutions to concentrated force problems conforming to the four conditions listed remain subject to individual verification through appropriate limit processes.

⁶See [3], art. 164-166.

⁷ r here is the distance from the load point.

We now turn to the determination of a sequence of singular harmonic functions which is needed in the subsequent analysis. It is hoped that this part of the paper may prove useful beyond the present application.

A Sequence of Harmonic Functions

If (x, y, z) denote Cartesian coordinates, the spherical coordinates (r, θ, γ) are defined through the mapping,

$$\left. \begin{aligned} x &= r \sin \theta \cos \gamma, & y &= r \sin \theta \sin \gamma, & z &= r \cos \theta \\ 0 \leq r < \infty, & & 0 \leq \theta \leq \pi, & & 0 \leq \gamma \leq 2\pi. \end{aligned} \right\} (1)$$

With the notation,

$$p = \cos \theta, \quad \bar{p} = \sin \theta, \quad (2)$$

we have

$$\rho = (x^2 + y^2)^{1/2} = r\bar{p}, \quad z = rp \quad (3)$$

for the relations between the spherical coordinates and the cylindrical coordinates (ρ, γ, z) . Laplace's equation $\nabla^2 H = 0$, in the axisymmetric case, for which $H = H(r, p)$, becomes

$$H_{rr} + \frac{2}{r} H_r + \frac{\bar{p}^2}{r} H_{pp} - \frac{2p}{r^2} H_p = 0. \quad (4)^8$$

Equation (4) admits the product solutions,⁹

$$H(r, p) = \left[r^n \text{ or } r^{-n-1} \right] \left[P_n(p) \text{ or } Q_n(p) \right] \quad (n = 0, 1, 2, \dots), \quad (5)$$

where P_n and Q_n are the Legendre polynomials and the Legendre functions of the second kind, respectively.

⁸Subscripts attached to functions which originally bear no subscripts, denote partial differentiation with respect to the argument indicated.

⁹For a comprehensive treatment of spherical harmonics, see [5].

The interior spherical harmonics $r^n P_n(p)$ are regular throughout the finite space whereas the exterior harmonics $r^{-n-1} P_n(p)$ vanish at infinity and, with increasing n , possess progressively stronger singularities at the origin. If we adopt the notation,

$$H_{-n}(r, p) = r^{-n} P_{n-1}(p) \quad (n = 1, 2, \dots), \quad (6)$$

then

$$H_{-n-1} = \frac{-1}{n} \frac{\partial}{\partial z} H_{-n}, \quad (7)$$

so that the sequence of exterior harmonics H_{-n} may be generated through successive differentiations with respect to z of the first order exterior harmonic $H_{-1} = r^{-1}$.

For future reference we recall that $P_n(p)$ satisfies Legendre's equation,

$$\frac{d}{dp} (\bar{p}^2 P_n') + n(n+1) P_n = 0 \quad (8)^{10}$$

and cite the recursion formulas,

$$\left. \begin{aligned} P_{-n-1} &= P_n \\ (2n+1) p P_n' &= (n+1) P_{n+1} + n P_{n-1} \\ \bar{p}^2 P_n' &= n P_{n-1} - n p P_n' \end{aligned} \right\} \quad (9)$$

Furthermore, we record the special values

$$\left. \begin{aligned} P_{2n}(0) &= (-1)^n \frac{2n}{2^{2n} (\underline{n})^2}, & P_{2n+1}(0) &= 0; \\ P_{2n}'(0) &= 0, & P_{2n+1}'(0) &= (2n+1) P_{2n}(0) \end{aligned} \right\} \quad (10)$$

and

$$P_n(1) = 1, \quad P_n'(1) = \frac{n(n+1)}{2}. \quad (11)$$

¹⁰ The argument of P_n is henceforth assumed to be p and $P_n' \equiv \frac{dP_n}{dp}$.

The Legendre functions of the second kind admit the representation,

$$Q_n(p) = \frac{1}{2} P_n(p) \ln \frac{1+p}{1-p} - W_{n-1}(p) \quad (n = 0, 1, 2, \dots) \quad (12)$$

in which

$$\left. \begin{aligned} W_{n-1} &= \frac{2n-1}{1 \cdot n} P_{n-1} + \frac{2n-5}{3 \cdot (n-1)} P_{n-3} + \frac{2n-9}{5 \cdot (n-2)} P_{n-5} + \dots \\ &= \sum_{m=1}^n \frac{1}{m} P_{m-1} P_{n-m}, \quad W_{-1} = 0. \end{aligned} \right\} (13)$$

Since $Q_n(p)$ has a logarithmic singularity at $p = \pm 1$, the harmonics $r^n Q_n(p)$ and $r^{-n-1} Q_n(p)$ in (5) are singular along the entire z -axis.

In the problem under consideration we shall have need for a sequence of axisymmetric harmonic functions which is regular in the half-space $z \geq 0$ with the exception of the origin $r = 0$, where it shall possess singularities, progressively weaker than that of $H_{-1} = r^{-1}$. Such a set of potentials may be constructed through successive integrations with respect to z of the first order exterior harmonic H_{-1} . We thus seek a sequence of functions $H_n(r, p)$ which satisfies the following requirements:

$$\nabla^2 H_n = 0, \quad (14)$$

$$\frac{\partial}{\partial z} H_n = n H_{n-1} \quad (n = 1, 2, \dots), \quad \frac{\partial}{\partial z} H_0 = H_{-1} = \frac{1}{r}. \quad (15)^{11}$$

Through a process of induction we are led to

$$\left. \begin{aligned} H_n(r, p) &= r^n T_n(p) \quad (n = 0, 1, 2, \dots) \\ T_n(p) &= P_n(p) L - V_n(p), \end{aligned} \right\} (16)$$

where

$$L = \ln r (1 + p) \equiv \ln (r + z), \quad (17)$$

¹¹The coefficient n in the first of (15) is introduced for later convenience. Note that (15) is analogous to (7).

and

$$\left. \begin{aligned} V_n &= (3c_n - 2c_{2n}) P_n - 2 \sum_{m=0}^{n-1} \frac{(-1)^{n-m} (2m+1) P_m}{(n-m)(n+m+1)}, \\ V_0 &= 0, \end{aligned} \right\} (18)$$

provided,

$$c_n = \sum_{m=1}^n \frac{1}{m} \quad (n = 1, 2, \dots). \quad (19)$$

In order to verify that the functions H_n , defined by (16) to (19) have the properties (14) and (15), we first observe that a lengthy computation, based on (18), (19), and (9), yields the recursion relations,

$$\left. \begin{aligned} \bar{p}^2 V_n' + npV_n - nV_{n-1} &= P_n, \\ n(nV_n - pV_n' + V_{n-1}') &= P_n' - 2P_{n-1}', \\ (n = 1, 2, \dots), \end{aligned} \right\} (20)$$

which, in turn, imply the differential equation,

$$\frac{d}{dp} (\bar{p}^2 V_n') + n(n+1)V_n = 2(P_n' - P_{n-1}') \quad (n = 0, 1, 2, \dots). \quad (21)$$

From (16), (20), (21), (9), we have

$$\left. \begin{aligned} \bar{p}^2 T_n' + npT_n - nT_{n-1} &= -pP_n, \\ n(nT_n - pT_n' + T_{n-1}') &= 2P_{n-1}' - pP_n', \\ (n = 1, 2, \dots), \end{aligned} \right\} (22)^{12}$$

$$\frac{\partial}{\partial p} (\bar{p}^2 T_n') + n(n+1)T_n = -(2n+1)P_n, \quad (n = 0, 1, 2, \dots). \quad (23)$$

With the aid of (16), (22), (23), and (4), Equations (14), (15) now readily follow. For future purposes we list the special values,

¹²Note that $T_n' \equiv \frac{\partial T_n}{\partial p}$.

$$\left. \begin{aligned} V_{2n}(0) &= c_n P_{2n}(0), & V_{2n+1}(0) &= \left[(2n+1) P_{2n}(0) \right]^{-1} \\ V'_{2n}(0) &= P_{2n}(0) - \left[P_{2n}(0) \right]^{-1}, & V'_{2n+1}(0) &= (2n+1) c_n P_{2n}(0), \end{aligned} \right\} (24)$$

which are obtained by induction with the aid of (18), (19), (22), (9), and (10).

According to (12), (13), (18),

$$V_n(p) - (-1)^n V_n(-p) = 2W_{n-1}(p) \quad (25)$$

and (12), (16), (17), (25) yield

$$H_n(r, p) - (-1)^n H_n(r, -p) = 2r^n Q_n(p). \quad (26)$$

This establishes the connection between the harmonic functions H_n , which are no longer product solutions of Laplace's equation, and the spherical harmonics of the second kind given in (5). Finally, we record explicitly the first three members of the aggregate $H_n(r, p)$:

$$\left. \begin{aligned} H_0 &= \ln r(1+p), & H_1 &= r \left[p \ln r(1+p) - 1 \right], \\ H_2 &= \frac{r^2}{2} \left[(3p^2 - 1) \ln r(1+p) - p^2 - 3p + 1 \right]. \end{aligned} \right\} (27)$$

Sequences of Singular Solutions

The sequence of harmonic functions established previously may be used to generate certain sequences of singular solutions of the field equations of elasticity theory. In the absence of body forces, and in case of torsionless axisymmetry about the z -axis, the general solution of the displacement equations of equilibrium may be written as,¹³

$$2G \left[u_\rho, u_\gamma, u_z \right] = \text{grad} (\phi + z\psi) - \left[0, 0, 4(1-\nu)\psi \right] \quad (28)$$

and

$$\nabla^2 \phi(\rho, z) = \nabla^2 \psi(\rho, z) = 0, \quad (29)$$

¹³See [2], [6].

where $[u_\rho, u_\gamma, u_z]$ are the cylindrical components of displacement, whereas G and ν designate the shear modulus and Poisson's ratio, respectively.

We define two sequences of particular solutions¹⁴ $[A_n]$ and $[B_n]$ in terms of their generating stress functions as follows:

$$\left. \begin{aligned} [A_n] \dots \phi &= H_n(r, p), & \psi &= 0 & (n = 0, 1, 2, \dots) \\ [B_n] \dots \phi &= 0, & \psi &= H_n(r, p) & (n = 0, 1, 2, \dots) \end{aligned} \right\} (30)$$

in which H_n is defined by (16) to (19). The spherical components of displacement u_r, u_θ and the spherical stress components $\sigma_r, \sigma_\theta, \sigma_\gamma, \tau_{r\theta}$, belonging to (28), (29), were given explicitly in [7].¹⁵ Substitution of (30) into (4) to (7) of Reference [7], and use of (22), (23), (8), (9), yields the following representations of $[A_n], [B_n]$ in spherical coordinates.

$[A_n]$:

$$\left. \begin{aligned} 2Gu_r &= r^{n-1}(nT_n + P_n), & 2Gu_\theta &= -\bar{p} r^{n-1} T_n' \\ \sigma_r &= r^{n-2} [n(n-1)T_n + (2n-1)P_n] \\ \sigma_\theta &= r^{n-2} [-n^2T_n + pT_n' - 2nP_n] \\ \sigma_\gamma &= r^{n-2} [nT_n - pT_n' + P_n] \\ \tau_{r\theta} &= -\bar{p} r^{n-2} [(n-1)T_n' + P_n'] \end{aligned} \right\} (31)$$

¹⁴ Throughout this paper capital letters in brackets denote either the displacement vector-field or the stress tensor-field of a solution of the field equations, and equality, addition, as well as multiplication by a scalar, are to be interpreted accordingly.

¹⁵ $u_\gamma, \tau_{\gamma r}, \tau_{\gamma\theta}$ vanish identically in view of the assumed rotational symmetry.

$[B_n]$:

$$\left. \begin{aligned} 2Gu_r &= pr^n \left[(n-3+4\nu) T_n + P_n \right] \\ 2Gu_\theta &= \bar{p}r^n \left[(3-4\nu) T_n - pT'_n \right] \\ \sigma_r &= r^{n-1} \left[n(n-3+2\nu)pT_n + (2n-3+2\nu)pP_n - 2\nu\bar{p}^2T'_n \right] \\ \sigma_\theta &= r^{n-1} \left\{ -n(n+2\nu)pT_n - 2(n+\nu)pP_n + \left[1 - (3-2\nu)\bar{p}^2 \right] T'_n \right\} \\ \sigma_y &= r^{n-1} \left\{ (1-2\nu)p(nT_n + P_n) - \left[1 - (1-2\nu)\bar{p}^2 \right] T'_n \right\} \\ \tau_{r\theta} &= \bar{p}r^{n-1} \left[(1-2\nu)(nT_n + P_n) - (n-2+2\nu)pT'_n - pP'_n \right] \end{aligned} \right\} \quad (32)$$

Let $Z(r)$ be the resultant force of the tractions to which a stress field with rotational symmetry about the z -axis gives rise on a hemisphere centered at $r = 0$, lying wholly in the region $z \geq 0$, and having an outer normal which is directed toward the origin. If $Z(r)$ is positive when its sense is that of the positive z -axis, it was shown in [8]¹⁶ that

$$Z(r) = 2\pi r \left\{ \left[\frac{\partial \phi}{\partial r} \right]_{p=0} + 2(1-\nu)r \int_0^1 \frac{\partial \psi}{\partial r} dp \right\}, \quad (33)$$

where $\phi(r,p)$ and $\psi(r,p)$ are the corresponding generating stress functions. Applying (33) to (30), we find, with the aid of (9), (10), (11), (16), (17), (23), and (24), the values of $Z(r)$ for $[A_n]$ and $[B_n]$.

$$\left. \begin{aligned} [A_{2n}]: \quad Z(r) &= \frac{2\pi(-1)^n \lfloor 2n \rfloor r^{2n}}{2^{2n}(\lfloor n \rfloor)^2} \left[2n \ln r + 1 - 2nc_n \right] \\ [A_{2n+1}]: \quad Z(r) &= - \frac{2\pi(-1)^n 2^{2n}(\lfloor n \rfloor)^2 r^{2n+1}}{\lfloor 2n \rfloor} \end{aligned} \right\} \quad (34)$$

¹⁶See Equations (31), (32) of Reference [8].

$$\begin{aligned}
 [B_{2n}]: \quad Z(r) &= \frac{4\pi(1-\nu)(-1)^n 2^{2n} (\underline{n})^2 r^{2n+1}}{2n+1} \\
 [B_{2n-1}]: \quad Z(r) &= \frac{-4\pi(1-\nu)(-1)^n \underline{2n+1} r^{2n}}{2^{2n} (\underline{n})^2} [2n \ln r + 1 - 2nc_n]
 \end{aligned}
 \quad \left. \vphantom{\begin{aligned} [B_{2n}]: \\ [B_{2n-1}]: \end{aligned}} \right\} \begin{array}{l} (34) \\ (Cont.) \end{array}$$

It follows from (34) that the aggregate of solutions defined by the linear combination,

$$\begin{aligned}
 [E_n] &= 2(1-\nu)[A_{n+1}] + (n+1)[B_n] \\
 (n &= 0, 1, 2, \dots)
 \end{aligned}
 \quad \left. \vphantom{[E_n]} \right\} (35)$$

is self-equilibrated in the sense that $Z(r) = 0$ for $[E_n]$. For later convenience, we record here the displacements and stresses of the first two members of the sequence $[E_n]$, which also appear in [1].

$$\begin{aligned}
 [E_0]: \\
 2Gu_r &= -(1-2\nu)pL + (3-2\nu)p - 2(1-\nu) \\
 2Gu_\theta &= \bar{p} \left[(1-2\nu)L - (3-2\nu) \frac{p}{1+p} \right] \\
 \sigma_r &= -\frac{1}{r} \left[(1-2\nu)p + 2\nu \right] \\
 \sigma_\theta &= \frac{1}{r(1+p)} \left[(1-2\nu)p^2 - 2p \right] \\
 \sigma_y &= \frac{1}{r(1+p)} \left[(1-2\nu)p - 2 \right] \\
 \tau_{r\theta} &= \frac{\bar{p}}{r(1+p)} \left[(1-2\nu)p - 1 \right]
 \end{aligned}
 \quad \left. \vphantom{\begin{aligned} [E_0]: \\ 2Gu_r \\ 2Gu_\theta \\ \sigma_r \\ \sigma_\theta \\ \sigma_y \\ \tau_{r\theta} \end{aligned}} \right\} (36)$$

$$\begin{aligned}
 [E_1]: \\
 2Gu_r &= r \left\{ 2 \left[\nu(1+p^2) - \bar{p}^2 \right] L + (3-\nu)p^2 - 2(1+\nu)p + 1 - \nu \right\} \\
 2Gu_\theta &= \bar{p}r \left\{ -2(1+\nu)pL + \frac{1}{1+p} \left[(-3+\nu)p^2 - (1-3\nu)p - 2(1-2\nu) \right] \right\} \\
 \sigma_r &= -2(1+\nu)(\bar{p}^2 L + p) + (5+\nu)p^2 - 1 + \nu \\
 \sigma_\theta &= -2(1+\nu)p^2 L + \frac{p}{1+p} \left[-(5+\nu)p^2 - (3-\nu)p + 6 \right] \\
 \sigma_y &= -2(1+\nu)L + \frac{1}{1+p} \left[(-3+\nu)p + 1 - \nu \right] \\
 \tau_{r\theta} &= -2(1+\nu)p\bar{p}L + \frac{\bar{p}}{1+p} \left[-(5+\nu)p^2 - (3-\nu)p + 2 \right]
 \end{aligned}
 \quad \left. \vphantom{\begin{aligned} [E_1]: \\ 2Gu_r \\ 2Gu_\theta \\ \sigma_r \\ \sigma_\theta \\ \sigma_y \\ \tau_{r\theta} \end{aligned}} \right\} (37)$$

Solutions $[A_n]$, $[B_n]$, $[E_n]$ ($n = 0, 1, 2, \dots$), with increasing n , possess progressively weaker singularities along the portion $z \leq 0$ of the z -axis, and are otherwise regular in the finite space; for $n = 0$ these solutions vanish as $z \rightarrow \infty$ so far as the stresses are concerned.

Construction of the Singularity due to a Concentrated Surface Force on a Curved Surface

Boussinesq's solution $[2]$ ¹⁷ corresponding to a concentrated load of magnitude $|Q|$ applied at $r = 0$ normal to the plane boundary of a medium occupying the half-space $z \geq 0$, appears as

$$[S_0] = -\frac{Q}{2\pi} \left\{ (1 - 2\nu)[A_0] + [B_{-1}] \right\}, \quad (38)^{18}$$

where $[A_0]$ is defined by (30) and

$$[B_{-1}] \dots \phi = 0, \quad \psi = H_{-1} = r^{-1}. \quad (39)$$

The displacements and stresses of $[S_0]$ follow.

$[S_0]$:

$$\left. \begin{aligned} u_r &= \frac{Q}{4\pi Gr} [4(1 - \nu)p - 1 + 2\nu] \\ u_\theta &= \frac{Q\bar{p}}{4\pi Gr} \left[\frac{1 - 2\nu}{1 + p} - 3 + 4\nu \right] \\ \sigma_r &= \frac{-Q}{2\pi r^2} [2(2 - \nu)p - 1 + 2\nu], & \sigma_\theta &= \frac{Q(1 - 2\nu)p^2}{2\pi r^2(1 + p)} \\ \sigma_y &= \frac{Q(1 - 2\nu)(p^2 + p - 1)}{2\pi r^2(1 + p)}, & \tau_{r\theta} &= \frac{Q(1 - 2\nu)p\bar{p}}{2\pi r^2(1 + p)} \end{aligned} \right\} (40)$$

Evidently, $Z(r) = Q$ for $[S_0]$, as is readily verified by means of (38), (39), (30), and (33).

¹⁷See also [3], p. 191.

¹⁸If $Q > 0$ the force has the sense of the positive z -axis.

We now turn to a concentrated load acting perpendicular to a curved boundary (Figure 1). Let the point of application of the load of magnitude $|Q|$ be the origin and again let $Q > 0$ if the load acts in the positive z -direction. Suppose that the boundary, in a neighborhood of the origin, coincides with a surface of revolution Σ , the semi-meridian Γ of which admits the representation,

$$\left. \begin{aligned} \Gamma: z = f(\rho) \quad (0 \leq \rho \leq \epsilon) \\ f(0) = f'(0) = 0, \end{aligned} \right\} (41)$$

and let $f(\rho)$ be at least four times continuously differentiable.¹⁹ Since $f'(0) = 0$, Σ possesses a uniquely defined tangent plane²⁰ at its vertex $r = 0$. The function $f(\rho)$ has a Taylor expansion of the form,

$$f(\rho) = a_2 \rho^2 + a_3 \rho^3 + \dots \quad (42)$$

so that

$$a_2 = \frac{1}{2} f''(0) = \frac{k_0}{2}, \quad a_3 = \frac{1}{6} f'''(0) = \frac{k'_0}{6} \quad (43)$$

where $k(\rho)$ is the curvature of Γ and

$$k_0 = k(0), \quad k'_0 = \left[\frac{dk}{d\rho} \right]_{\rho=0}. \quad (44)$$

With a view toward examining the nature of the singularity at the load point $r = 0$, we first determine the normal and shearing tractions σ and τ (Figure 1) induced by the Boussinesq solution $[S_0]$ on the arc Γ . To this end we observe by means of the law of stress transformation and elementary geometric considerations that along Γ ,

¹⁹Note that the regularity restrictions on f imply the same degree of smoothness for Σ only if the (right-hand) derivatives of odd order of $f(\rho)$ at $\rho = 0$ vanish. Otherwise, the even extension of $f(\rho)$ exhibits discontinuities in its derivatives of odd order at the origin.

²⁰The subsequent analysis is not valid if $r = 0$ is a conical point of Σ .

$$\left. \begin{aligned} \sigma &= \frac{\sigma_r + \sigma_\theta}{2} + \frac{\sigma_r - \sigma_\theta}{2} \cos 2(\beta + \theta) - \tau_{r\theta} \sin 2(\beta + \theta) \\ \tau &= \frac{\sigma_r - \sigma_\theta}{2} \sin 2(\beta + \theta) + \tau_{r\theta} \cos 2(\beta + \theta) \end{aligned} \right\} (45)$$

in which β is the inclination of the tangent of Γ . Evidently,

$$\cos 2\beta = \frac{1 - (f')^2}{1 + (f')^2}, \quad \sin 2\beta = \frac{2f'}{1 + (f')^2}. \quad (46)$$

Recalling that

$$r = \rho \left[1 + \left(\frac{z}{\rho} \right)^2 \right]^{1/2}, \quad (47)$$

we reach by means of (3), (41), (42), the expansions,

$$\left. \begin{aligned} \frac{1}{r} &= \frac{1}{\rho} \left(1 - \frac{a_2^2}{2} \rho^2 - a_2 a_3 \rho^3 + \dots \right) \\ p &= a_2 \rho + a_3 \rho^2 + \left(a_4 - \frac{a_2^3}{2} \right) \rho^3 + \dots \\ \bar{p} &= 1 - \frac{a_2^2}{2} \rho^2 - a_2 a_3 \rho^3 + \dots, \end{aligned} \right\} (48)$$

valid on Γ . From (46), (48), and (2) follows

$$\left. \begin{aligned} \cos 2(\beta + \theta) &= -1 + 2a_2^2 \rho^2 + 8a_2 a_3 \rho^3 + \dots \\ \sin 2(\beta + \theta) &= -2a_2 \rho - 4a_3 \rho^2 - 6(a_4 - a_2^3) \rho^3 + \dots \end{aligned} \right\} (49)$$

Substitution of (49) and (40) into (45), after a lengthy routine computation, and final use of (43), yields

$$\left. \begin{aligned} [S_0]: \\ \sigma(\rho) &= \frac{Q(1 - 2\nu) k_0^2}{2\pi} + o(1) = O(1) \\ \tau(\rho) &= \frac{Q}{8\pi} \left[\frac{-4(1 - 2\nu)k_0}{\rho} + (5 - 4\nu)k_0^2 - 2(1 - 2\nu)k_0' \right] + o(1) \end{aligned} \right\} (50)$$

where $o(1)$ and $O(1)$, respectively, denote functions of ρ which tend to zero or to a finite limit as ρ approaches zero.

If $[S]$ designates the complete solution to the concentrated force problem under consideration, we may write,

$$[S] = [S_0] + [R_0], \quad (51)$$

in which $[R_0]$ stands for the solution to a "residual problem", the characterization of which is implicit in (51). We shall suppose, without loss in generality, that the boundary of the medium is free from distributed tractions on Σ . Furthermore, we shall agree to call $[R_0]$ "regular" on Σ if the residual problem on Σ is governed by finite and continuous surface tractions. This requires for $[R_0]$,

$$\sigma(\rho) = O(1), \quad \tau(\rho) = o(1). \quad (52)^{21}$$

Equations (50), (51) now permit the following conclusions. $[R_0]$ is regular on Σ if and only if the curvature k and the rate of change of curvature k' of the semi-meridian Γ of Σ both vanish at the load point. We note parenthetically that k'_0 is zero automatically if the even extension of Γ has at least a continuous rate of change of curvature.²² In the event that the even extension of $f(\rho)$ is an analytic function, so that Σ is an analytic surface, the surface tractions of $[R_0]$ are analytic on Σ if and only if $k_0 = 0$. In this case, and in this case only, the Boussinesq singularity constitutes the entire singularity of $[S]$ at the load point in question.

In order to effect a reduction of the problem to a residual problem which is regular on Σ when k_0 and k'_0 are not both zero, we need to introduce supplementary singular solutions, in addition to $[S_0]$. Moreover, in accordance with conditions (c) and (d) stated in the Introduction, the

²¹Observe that $\tau(\rho) = O(1)$ is insufficient since it admits a discontinuity in the surface shearing tractions of $[R_0]$ at the vertex of Σ .

²²See footnote No. 19.

supplementary singularities sought must be self-equilibrated and of a lower order than r^{-2} . This leads us to solutions $[E_n]$ defined by (35) and listed explicitly for $n = 0, 1$ in (36), (37).

The local behavior of $[E_0]$, $[E_1]$ on Γ , in the vicinity of $\rho = 0$, is established by a process which is strictly analogous to that followed in the derivation of (50). We may, therefore, give the results of these computations directly.

$[E_0]$:

$$\sigma(\rho) = -2k_0 + o(1), \quad \tau(\rho) = \frac{1}{\rho} - (1 - 2\nu)k_0 + o(1)$$

$[E_1]$:

$$\sigma(\rho) = o(1), \quad \tau(\rho) = -2 + o(1)$$

(53)

(54)

It is apparent from (53), (54) that τ of $[E_0]$ still becomes infinite at $\rho = 0$ whereas τ of $[E_1]$ displays merely a finite discontinuity at the origin; the normal tractions σ of both of these solutions are again finite and continuous on Σ , as in the case of $[S_0]$.

We now set

$$\begin{aligned} [S] &= [S_2] + [R_2] \\ [S_2] &= [S_0] + \lambda_0[E_0] + \lambda_1[E_1], \end{aligned} \quad (55)$$

and ask for the values of the coefficients λ_0 , λ_1 which assure the regularity in the sense of (52) of the solution $[R_2]$ to the new residual problem. In view of (50), (53), (54) this leads to

$$\begin{aligned} \lambda_0 &= \frac{(1 - 2\nu) Qk_0}{2\pi} \\ \lambda_1 &= \frac{Q}{16\pi} \left[(1 + 12\nu - 16\nu^2)k_0^2 - 2(1 - 2\nu)k_0' \right] \end{aligned} \quad (56)$$

supplementary singularities sought must be self-equilibrated and of a lower order than r^{-2} . This leads us to solutions $[E_n]$ defined by (35) and listed explicitly for $n = 0, 1$ in (36), (37).

The local behavior of $[E_0]$, $[E_1]$ on Γ , in the vicinity of $\rho = 0$, is established by a process which is strictly analogous to that followed in the derivation of (50). We may, therefore, give the results of these computations directly.

$$\left. \begin{aligned} [E_0]: \\ \sigma(\rho) = -2k_0 + o(1), \quad \tau(\rho) = \frac{1}{\rho} - (1 - 2\nu)k_0 + o(1) \end{aligned} \right\} (53)$$

$$\left. \begin{aligned} [E_1]: \\ \sigma(\rho) = o(1), \quad \tau(\rho) = -2 + o(1) \end{aligned} \right\} (54)$$

It is apparent from (53), (54) that τ of $[E_0]$ still becomes infinite at $\rho = 0$ whereas τ of $[E_1]$ displays merely a finite discontinuity at the origin; the normal tractions σ of both of these solutions are again finite and continuous on Σ , as in the case of $[S_0]$.

We now set

$$\left. \begin{aligned} [S] &= [S_2] + [R_2] \\ [S_2] &= [S_0] + \lambda_0 [E_0] + \lambda_1 [E_1], \end{aligned} \right\} (55)$$

and ask for the values of the coefficients λ_0 , λ_1 which assure the regularity in the sense of (52) of the solution $[R_2]$ to the new residual problem. In view of (50), (53), (54) this leads to

$$\left. \begin{aligned} \lambda_0 &= \frac{(1 - 2\nu) Q k_0}{2\pi} \\ \lambda_1 &= \frac{Q}{16\pi} \left[(1 + 12\nu - 16\nu^2) k_0^2 - 2(1 - 2\nu) k_0' \right] \end{aligned} \right\} (56)$$

With $[S_2]$ determined as in (55), (56), the surface tractions of $[R_2]$ are finite and continuous on Σ . Alternatively, $[S]$ admits the representation,

$$\left. \begin{aligned} [S] &= [S_1] + [R_1] \\ [S_1] &= [S_0] + \lambda_0 [E_0], \end{aligned} \right\} (57)$$

where λ_0 again has the value appearing in (56). The surface tractions of $[R_1]$ are finite but exhibit a discontinuity in τ at the origin.

It is important to emphasize that the surface tractions of $[R_2]$ on Σ , although finite and continuous, are by no means analytic even if Σ is analytic. This is due to the fact that the derivatives with respect to ρ of σ and τ in $[E_1]$, unlike the corresponding derivatives belonging to $[S_0]$, $[E_0]$, possess a logarithmic singularity²³ at $\rho = 0$. Consequently, the singularity of $[S_2]$, regardless of the analyticity of Σ , does not represent the entire singularity inherent in $[S]$ at the load point.

The sequence of singular solutions $[S_0]$, $[S_1]$, $[S_2]$ thus gives rise to a sequence of solutions $[R_0]$, $[R_1]$, $[R_2]$ to the complementary residual problems, which are characterized by progressively increasing regularity in the corresponding boundary conditions. The process of successive "regularization" of the residual problem can be continued in an obvious manner by recourse to the remaining members of the sequence of self-equilibrated singular solutions $[E_n]$ defined in (35).

If Σ is a portion of a sphere of radius r_0 , lying in the half-space $z \geq 0$, then $k_0 = r_0^{-1}$, $k'_0 = 0$, and the results obtained here reduce to those given in [1] in connection with the particular problem of the

²³The functions designated by $o(1)$ in (54) contain a term, $g(\rho) = p \ln r(1 + p)$.

sphere under concentrated loads. Whereas the general questions raised in the Introduction have been answered within the limitations placed on the relative orientation of the load and on the local character of the boundary, it should be pointed out that certain difficulties in the large may arise in the application of the results to specific concentrated force problems. Thus, the representations (51), (55), (57) lead to serious complications if the line of action of a concentrated load intersects the boundary of the medium on opposite sides of the tangent plane at the point of application of the load. Here $[S_0]$, $[S_1]$, $[S_2]$, each of which is singular along the negative z-axis, give rise to line singularities in the interior of the body.²⁴ Finally, if the region occupied by the medium is not bounded, the representation (55) becomes inadmissible in view of the behavior of $[E_1]$ at infinity.

Bibliography

- [1] "The elastic sphere under concentrated loads", by E. Sternberg and F. Rosenthal, Journal of Applied Mechanics, vol. 19, no. 4, 1952, p. 413.
- [2] "Application des potentiels à l'étude de l'équilibre et du mouvement des solides élastiques", by J. Boussinesq, Gauthier-Villars, Paris, 1885.
- [3] "A treatise on the mathematical theory of elasticity", by A.E.H.Love, fourth edition, Dover Publications, New York, N.Y., 1944.
- [4] "A special investigation to develop a general method for three-dimensional photoelastic stress analysis", by M.M.Frocht and R.Guernsey, Jr., NACA Technical Note 2822, December 12, 1952.
- [5] "The theory of spherical and ellipsoidal harmonics", by E. W. Hobson, University Press, Cambridge, England, 1931.

²⁴Consider, for example, a concentrated load applied to the inner boundary of a spherical shell.

- [6] "On the stress-function approaches of Boussinesq and Timpe to the axisymmetric problem of elasticity theory", by E. Sternberg, R. A. Eubanks, and M. A. Sadowsky, Journal of Applied Physics, vol. 22, no. 9, 1951, p. 1121.
- [7] "On the axisymmetric problem of elasticity theory for a region bounded by two concentric spheres", by E. Sternberg, R. A. Eubanks, and M. A. Sadowsky, Proceedings, First U. S. National Congress of Applied Mechanics, The American Society of Mechanical Engineers, New York, 1952.
- [8] "Stress concentration due to a hemispherical pit at a free surface", by R. A. Eubanks, Technical Report to the ONR, Contract N7omr-32906, January 1, 1953. To appear in a future issue of the Journal of Applied Mechanics.




Figure 1.

DISTRIBUTION LIST - PROJECT NO. NR 035-302

Administrative, Reference, and Liaison Activities

Chief of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 438 (2) : Code 432 (1)	Commanding Officer Office of Naval Research John Crerar Library Building 86 E. Randolph St. Chicago 11, Illinois (2)
Head, Mechanics Branch Office of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 438 (15)	Commanding Officer Office of Naval Research Branch Office 801 Donahue Street San Francisco 24, California (1)
Director, Naval Research Lab Washington 25, D. C. Attn: Tech. Info. Officer (9) : Technical Library (1) : Mechanics Division (2) : Code 3834 (J.P. Walsh) (1)	Commanding Officer Office of Naval Research Branch Office 1030 Green Street Pasadena, California (1)
Commanding Officer Office of Naval Research Branch Office 195 Summer Street Boston 10, Massachusetts (1)	Officer in Charge Office of Naval Research Branch Office, London Navy No. 100 FPO, New York, New York (5)
Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, New York (1)	Library of Congress Washington 25, D. C. Attn: Navy Research Section (2)

Department of Defense and Other Interested Government Activities

GENERAL

Research & Development Board Department of Defense Pentagon Building Washington 25, D. C. Attn: Library (Code 3D-1075) (1)	Armed Forces Special Weapons Project P. O. Box 2610 Washington, D. C. Attn: Lt. Col. G. F. Blunda (1)
	Joint Task Force 3 12th St. & Constitution Ave., NW (Temple U.) Washington 25, D. C. Attn: Major B. D. Jones (1)

ARMY

Chief of Staff
Department of the Army
Research & Development Division
Washington 25, D. C.
Attn: Chief of Res. & Dev. (1)

Office of the Chief of Engineers
Assistant Chief for Works
Department of the Army
Bldg. T-7, Gravelly Point
Washington 25, D. C.
Attn: Structural Branch
(R. L. Bloor) (1)

Office of the Chief of Engineers
Asst. Chief for Military Construction
Department of the Army
Bldg. T-7, Gravelly Point
Washington 25, D. C.
Attn: Structures Branch
(H. F. Carey) (1)

Office of the Chief of Engineers
Asst. Chief for Military Operations
Department of the Army
Bldg. T-7, Gravelly Point
Washington 25, D. C.
Attn: Structures Development
Branch (W. F. Woolard) (1)

Engineering Research & Development
Laboratory
Fort Belvoir, Virginia
Attn: Structures Branch (1)

Office of Chief of Ordnance
Research & Development Service
Department of the Army
The Pentagon
Washington 25, D. C.
Attn: ORDTB (2)

Commanding Officer
Watertown Arsenal
Watertown, Massachusetts
Attn: Laboratory Division (1)

NAVY

Chief of Bureau of Ships
Navy Department
Washington 25, D. C.
Attn: Director of Research (2)
: Code 440 (1)
: Code 421 (1)

Director
David Taylor Model Basin
Washington 7, D. C.
Attn: Structural Mech. Div. (1)
: Dr. E. Wenk (1)

Director
Naval Engr. Experiment Station
Annapolis, Maryland (1)

Director
Materials Laboratory
New York Naval Shipyard
Brooklyn 1, New York (1)

Chief of Bureau of Ordnance
Navy Department
Washington 25, D. C.
Attn: Ad-3, Technical Library (1)

Superintendent
Naval Gun Factory
Washington 25, D. C. (1)

Naval Ordnance Laboratory
White Oak, Maryland
RFD 1, Silver Spring, Maryland
Attn: Mechanics Division (2)

Naval Ordnance Test Station
Inyokern, California
Attn: Scientific Officer (1)

Naval Ordnance Test Station
Underwater Ordnance Division
Pasadena, California
Attn: Structures Division (1)

Chief of Bureau of Aeronautics
Navy Department
Washington 25, D. C.
Attn: TD-41, Tech. Library (1)
: DE-22, C. W. Hurley (1)

NAVY (Cont'd)

Naval Air Experimental Station
Naval Air Materiel Center
Naval Base
Philadelphia 12, Pennsylvania
Attn: Head, Aeronautical
Materials Laboratory (1)

Chief of Bureau of Yards & Docks
Navy Department
Washington 25, D. C.
Attn: Code P-314 (1)

Officer in Charge
Naval Civil Engr. Research
and Eval. Laboratory
Naval Station
Port Hueneme, California (1)

Superintendent
U. S. Naval Post Graduate School
Annapolis, Maryland
Attn: Library (1)

AIR FORCES

Office of Air Research
Administrative Office
Hq Air Research & Development
Command
P. O. Box 1395
Baltimore 3, Maryland (RDMPE)(15)

Commanding General
U. S. Air Forces
The Pentagon
Washington 25, D. C.
Attn: Research & Development
Division (1)

Commanding General
Air Materiel Command
Wright-Patterson Air Force Base
Dayton, Ohio
Attn: MCREX-B (E.H.Schwartz) (1)

Office of Air Research
Wright-Patterson Air Force Base
Dayton, Ohio
Attn: Chief, Applied Mechanics
Group (1)

OTHER GOVERNMENT AGENCIES

U. S. Atomic Energy Commission
Division of Research
Washington, D. C. (1)

Argonne National Laboratory
P. O. Box 5207
Chicago 80, Illinois (1)

Director
National Bureau of Standards
Washington, D. C.
Attn: Dr. W. H. Ramberg (2)

U. S. Coast Guard
1330 E Street, NW
Washington, D. C.
Attn: Chief, Testing & Development
Division (1)

Forest Products Laboratory
Madison, Wisconsin
Attn: L. J. Markwardt (1)

National Advisory Committee
for Aeronautics
1724 F Street, NW
Washington, D. C. (1)

National Advisory Committee
for Aeronautics
Langley Field, Virginia
Attn: Mr. E. Lundquist (1)
: Dr. B. Budiansky (1)

National Advisory Committee
for Aeronautics
Cleveland Municipal Airport
Cleveland, Ohio
Attn: J. H. Collins, Jr. (1)

U. S. Maritime Commission
Technical Bureau
Washington, D. C.
Attn: Mr. V. Russo (1)

Contractors and Other Investigators Actively Engaged in Related Research

Professor J. R. Andersen Towne School of Engineering University of Pennsylvania Philadelphia, Pennsylvania (1)	Dr. Francis H. Clauser, Chairman Dept. of Aeronautics School of Engineering Johns Hopkins University Baltimore 18, Maryland (1)
Mr. T. T. Arai Armour Research Foundation Technology Center Chicago 16, Illinois (1)	Professor Sisir Chandra Das Chandernagore College Chandernagore West Bengal, <u>INDIA</u> (1)
Professor Lynn Beedle Fritz Engineering Laboratory Lehigh University Bethlehem, Pennsylvania (1)	Professor J. P. Den Hartog Massachusetts Institute of Technology Cambridge 39, Massachusetts (1)
Professor C. B. Biezeno Technische Hoogeschool Nieuwe Laan 76 Delft, <u>HOLLAND</u> (1)	Dr. Herbert Deresiewicz Dept. of Civil Engineering Columbia University 632 West 125th Street New York 27, New York (1)
Professor M. A. Biot 1819 Broadway New York, New York (1)	Dr. C. O. Dohrenwend Rensselaer Polytechnic Institute Troy, New York (1)
Professor R. L. Bisplinghoff Massachusetts Institute of Technology Dept. of Aeronautical Engrg. Cambridge 39, Massachusetts (1)	Professor T. J. Dolan Dept. of Theoretical and Applied Mechanics University of Illinois Urbana, Illinois (1)
Dr. Hans. H. Bleich Dept. of Civil Engineering Columbia University New York 27, New York (1)	Professor Lloyd Donnell Dept. of Mechanics Illinois Institute of Technology Technology Center Chicago 16, Illinois (1)
Professor J. A. Bogdanoff Purdue University Lafayette, Indiana (1)	Professor D. C. Drucker Brown University Providence, Rhode Island (1)
Professor B. A. Boley Dept. of Civil Engineering Columbia University New York 27, New York (1)	Dr. H. Ekstein Armour Research Foundation Technology Center Chicago 16, Illinois (1)
Professor F. W. Bridgeman Dept. of Physics Harvard University Cambridge, Massachusetts (1)	Engineering Library Columbia University New York 27, New York (1)
Dr. G. F. Carrier Harvard University Cambridge, Massachusetts (1)	

Professor E. L. Eriksen University of Michigan Ann Arbor, Michigan	(1)	Professor J. N. Goodier Dept. of Mechanical Engineering Stanford University Stanford, California	(1)
Professor A. C. Eringen Purdue University Lafayette, Indiana	(1)	Professor L. E. Goodman Dept. of Civil Engineering University of Illinois Urbana, Illinois	(1)
Dr. W. L. Esmeijer Technische Hoogeschool Nieuwe Laan 76 Delft, <u>HOLLAND</u>	(1)	Professor A. E. Green Dept. of Mathematics Stephenson Building King's College Newcastle-upon-Tyne 2 <u>ENGLAND</u>	(1)
Professor R. A. Eubanks Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor H. J. Greenberg Dept. of Mathematics Carnegie Institute of Technology Pittsburgh, Pennsylvania	(1)
Professor K. Federhofer Technische Hochschule Graz, <u>AUSTRIA</u>	(1)	Professor George Handelman Dept. of Mathematics Carnegie Institute of Technology Pittsburgh, Pennsylvania	(1)
Professor A. M. Freudenthal Dept. of Civil Engineering Columbia University New York 27, New York	(1)	Professor R. J. Hansen Dept. of Civil & Sanitary Engrg. Massachusetts Institute of Technology Cambridge 39, Massachusetts	(1)
Professor B. Fried Washington State College Pullman, Washington	(1)	Professor R. M. Hermes University of Santa Clara Santa Clara, California	(1)
Professor K. O. Friedrichs New York University Washington Square New York, New York	(1)	Professor G. Herrmann Dept. of Civil Engineering Columbia University New York 27, New York	(1)
Professor M. M. Frocht Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor M. Hetényi The Technological Institute Northwestern University Evanston, Illinois	(1)
Professor Hilda Geiringer Department of Mathematics Wheaton College Norton, Massachusetts	(1)	Professor T. J. Higgins Dept. of Electrical Engineering University of Wisconsin Madison 6, Wisconsin	(1)
Dean J. A. Goff University of Pennsylvania Philadelphia, Pennsylvania	(1)		
Mr. Martin Goland Midwest Research Institute 4049 Pennsylvania Avenue Kansas City 2, Missouri	(1)		

Professor N. J. Hoff, Head
Dept. of Aeronautical Engineering
& Applied Mechanics
Polytechnic Institute of Brooklyn
99 Livingston Street
Brooklyn 2, New York (1)

Professor M. B. Hogan
University of Utah
Salt Lake City, Utah (1)

Professor D. L. Holl
Iowa State College
Ames, Iowa (1)

Dr. J. H. Hollomon
General Electric Research Labs
1 River Road
Schenectady, New York (1)

Professor W. H. Hoppmann
Dept. of Applied Mechanics
The Johns Hopkins University
Baltimore, Maryland (1)

Institu de Mathematiques
Université
post. fah 55
Skoplje, YUGOSLAVIA (1)

Professor L. S. Jacobsen
Dept. of Mechanical Engineering
Stanford University
Stanford, California (1)

Professor Bruce G. Johnston
University of Michigan
Ann Arbor, Michigan (1)

Professor K. Klotter
Stanford University
Stanford, California (1)

Professor W. J. Krefeld
Dept. of Civil Engineering
Columbia University
New York 27, New York (1)

Professor B. J. Lazan
Dept. of Materials Engineering
University of Minnesota
Minneapolis, Minnesota (1)

Dr. E. H. Lee
Division of Applied Mathematics
Brown University
Providence, Rhode Island (1)

Professor George Lee
Rensselaer Polytechnic Institute
Troy, New York (1)

Professor J. M. Lessells
Massachusetts Institute of
Technology
Cambridge 39, Massachusetts (1)

Library, Engineering Foundation
29 West 39th Street
New York 18, New York (1)

Professor Paul Lieber
Dept. of Engineering
Rensselaer Polytechnic Institute
Troy, New York (1)

Dr. Hsu Lo
Purdue University
Lafayette, Indiana (1)

Professor C. T. G. Looney
Dept. of Civil Engineering
Yale University
New Haven, Connecticut (1)

Dr. J. L. Lubkin
Midwest Research Institute
4049 Pennsylvania Avenue
Kansas City 2, Missouri (1)

Professor J. F. Ludloff
School of Aeronautics
New York University
New York 53, New York (1)

Professor J. N. Macduff
Rensselaer Polytechnic Institute
Troy, New York (1)

Professor C. W. MacGregor
University of Pennsylvania
Philadelphia, Pennsylvania (1)

Professor Lawrence E. Malvern
Dept. of Mathematics
Carnegie Institute of Technology
Pittsburgh 13, Pennsylvania (1)

Dr. J. H. Marchant
Brown University
Providence, Rhode Island (1)

Professor J. Marin
Pennsylvania State College
State College, Pennsylvania (1)

Professor E. Masur
Illinois Institute of Technology
Technology Center
Chicago 16, Illinois (1)

Mr. E. L. McDowell
Armour Research Foundation
Technology Center
Chicago 16, Illinois (1)

Professor R. D. Mindlin
Dept. of Civil Engineering
Columbia University
632 West 125th Street
New York 27, New York (1)

Dr. A. Nadai
136 Cherry Valley Road
Pittsburgh 21, Pennsylvania (1)

Professor Paul M. Naghdi
Dept. of Engineering Mechanics
University of Michigan
Ann Arbor, Michigan (1)

Professor N. M. Newmark
Dept. of Civil Engineering
University of Illinois
Urbana, Illinois (1)

Professor Jesse Ormondroyd
University of Michigan
Ann Arbor, Michigan (1)

Professor W. Osgood
Illinois Institute of Technology
Technology Center
Chicago 16, Illinois (1)

Dr. R. P. Petersen
Director, Applied Physics Division
Sandia Laboratory
Albuquerque, New Mexico (1)

Mr. R. E. Peterson
Westinghouse Research Laboratories
East Pittsburgh, Pennsylvania (1)

Dr. A. Phillips
School of Engineering
Stanford University
Stanford, California (1)

Professor Gerald Pickett
Dept. of Mechanics
University of Wisconsin
Madison 6, Wisconsin (1)

Dr. H. Poritsky
General Electric Research Labs
Schenectady, New York (1)

Professor W. Frager
Graduate Div. of Applied Mathematics
Brown University
Providence, Rhode Island (1)

RAND Corporation
1500 Fourth Street
Santa Monica, California
Attn: Dr. D. L. Judd (1)

Professor E. Reissner
Dept. of Mathematics
Massachusetts Institute of Technology
Cambridge 39, Massachusetts (1)

Professor H. Reissner
Polytechnic Institute of Brooklyn
99 Livingston Street
Brooklyn 2, New York (1)

Dr. Kenneth Robinson
National Bureau of Standards
Washington, D. C. (1)

Dr. Felix Rosenthal
Armour Research Foundation
Technology Center
Chicago 16, Illinois (1)

Professor M. A. Sadowsky
Dept. of Mechanics
Rensselaer Polytechnic Institute
Troy, New York (1)

Professor M. G. Salvadori
Dept. of Civil Engineering
Columbia University
New York 27, New York (1)

Dr. F. S. Shaw
Polytechnic Institute of Brooklyn
99 Livingston Street
Brooklyn 2, New York (1)

Dr. Daniel T. Sigley
Applied Physics Laboratory
The Johns Hopkins University
8621 Georgia Avenue
Silver Spring, Maryland (1)

Professor C. B. Smith
Department of Mathematics
Walker Hall
University of Florida
Gainesville, Florida (1)

Professor C. R. Soderberg
Massachusetts Institute of
Technology
Cambridge 39, Massachusetts (1)

Professor I. S. Sokolnikoff
Dept. of Mathematics
University of California at
Los Angeles
Los Angeles 24, California (1)

Professor R. V. Southwell
Imperial College of Science and
Technology
South Kensington
London SW 7, ENGLAND (1)

Professor E. Sternberg
Illinois Institute of Technology
Technology Center
Chicago 16, Illinois (10)

Professor J. J. Stoker
New York University
Washington Square
New York, New York (1)

Professor P. S. Symonds
Brown University
Providence, Rhode Island (1)

Professor J. L. Synge
Dublin Institute for Advanced
Studies
School of Theoretical Physics
64-65 Merrion Square
Dublin, IRELAND (1)

Professor F. K. Teichmann
Dept. of Aeronautical Engineering
New York University
University Heights, Bronx
New York, New York (1)

Professor S. P. Timoshenko
School of Engineering
Stanford University
Stanford, California (1)

Professor C. A. Truesdell
Graduate Institute for Applied
Mathematics
Indiana University
Bloomington, Indiana (1)

Professor Karl S. Van Dyke
Department of Physics
Scott Laboratory
Wesleyan University
Middletown, Connecticut (1)

Dr. I. Vigness
Naval Research Laboratory
Anacostia Station
Washington, D. C. (1)

Professor E. Volterra
Rensselaer Polytechnic Institute
Troy, New York (1)

Mr. A. M. Wahl
Westinghouse Research Laboratories
East Pittsburgh, Pennsylvania (1)

Dr. Conrad Wan
3422 Shorecraft Drive
Dallas, Texas (1)

Professor C. T. Wang
Dept. of Aeronautical Engineering
New York University
University Heights, Bronx
New York, New York (1)

Professor E. E. Weibel
University of Colorado
Boulder, Colorado (1)

Professor Dana Young
Yale University
New Haven, Connecticut (1)

Professor Alexander Weinstein
Institute of Applied Mathematics
University of Maryland
College Park, Maryland (1)