

ON THE MIESCHER-BAER EMISSION BANDS IN THE FAR ULTRAVIOLET REGION[†]

Yoshio Tanaka[‡]
Laboratory of Molecular Structure and Spectra
Department of Physics
The University of Chicago
Chicago 37, Illinois

DATA
13
AD 019144
In studying a high-current discharge (of He + enriched N¹⁵O or ordinary NO), Miescher and Baer¹ found a new system of bands in the region from 1,368 to 1,680Å. These bands are single-headed and degraded toward the red. The vibrational frequency of the lower state is close to the frequency of the X¹Σ_g⁺ state of the N₂ molecule as well as to that of the A²Σ⁺ state of the NO molecule. They did not find any alternation of intensity in the rotational structure, nor did they observe the heads of N₂¹⁵ isotope bands. Hence, they were unable to decide whether this system arose from the N₂ or the NO molecule.

With a view to determine experimentally the origin of these bands, we tried a high-current discharge of NO, (He + NO), N₂ and (He + N₂), each with and without a condenser. The 84-cm normal-incidence vacuum spectrograph was used.

We could not observe the bands in the condensed discharge of any of the above-mentioned gases, nor in the uncondensed discharges of N₂ and (He + N₂). We can, however, observe them in the uncondensed discharge of either pure NO or, much stronger, in the case of (He + NO). The results of the measurements are shown in Table I together with those of Miescher and Baer. In the present work, the system has been greatly extended towards shorter wavelengths, down to about 1,200Å. Some of the bands come close to and are confused with the Lyman-Birge-Hopfield bands of N₂, which appear quite strong in the region below 1,500Å. We can, nevertheless, discriminate between them by comparison with the picture of the pure N₂ spectrum because, as observed by Miescher and Baer, this new system has no intensity alternation in the rotational structure.

[†]This work was assisted by the Office of Naval Research under Task Order IX of Contract N6ori-20 with The University of Chicago.

[‡]Present address: Geophysics Research Division, Air Force Cambridge Research Center, 230 Albany Street, Cambridge 39, Massachusetts.

¹R. Miescher and P. Baer, Nature 169, 581 (1952).

Y. TANAKA

TABLE I

v'	v''	ν observed (cm^{-1}) present	I	ν (cm^{-1}) Miescher and Baer
0	6	59513.9	1	59517
0	5	61702.5	2	61704
0	4	63908.8	4	63910
0	3	66149.4	6	66161
0	2	68431.3	6	68437
1	2	69983.3	3	69999
1	1	70745.4	3	70751
1	1	72307.6	6	72310
6	4	72585.8	1	
0	0	73083.9	3	73087
2	1	73819.8	1	
1	4	73874.9	1	
1	0	74642.5	3	
2	0	76158.0	1	
6	2	77121.8	1	
5	0	77636.1	6	
5	1	78134.2	1	
4	3	78361.9	4	
4	0	79072.0	6	
5	1	79455.9	4	
5	0	80488.1	1	
5	2	80645.2	2	$\omega_e' = 1599.6\text{cm}^{-1}$
6	1	80690.1	1	
7	0	81768.8	3	$\omega_e' \chi_e' = 20.6\text{cm}^{-1}$
8	1	81887.3	0	
8	0	83042.0	2	$\omega_e'' = 2366.5\text{cm}^{-1}$
8	0	84209.8	2	
9	0	85284.9	1	$\omega_e'' \chi_e'' = 14.6\text{cm}^{-1}$
10	0	86214.3	1	

It seems probable that this system is due to NO or NO^+ rather than N_2 . The reasons for this conclusion are the following. First of all, no intensity alternation is observed. Secondly, as pointed out by Miescher and Baer, there are no corresponding isotope bands of N_2^{15} observed. This new system appears only in the silent discharge of (He + NO) or NO and are not observed in the N_2 or (He + N_2) spectra. If we assume that the lower state of this system is $\text{A}^2\Sigma^+$ of NO, then the upper state is about 14.54eV above the $v = 0$ level of the ground state ($\text{X}^2\Pi$) of NO. Tanaka² has found several progressions of absorption bands in this region but none fit with the upper state of the present bands.

There is the possibility that this system is due to the NO^+ ion. As Mulliken³ has pointed out, the ground state of NO^+ is $1\Sigma^+$, having the electron configuration

²Y. Tanaka, Sci. Pap. Inst. Phys. Chem. Res., Tokyo 39, 456 (1942).

³R. S. Mulliken, Rev. Mod. Phys. 4, 73 (1932).

MIESCHER-BAER EMISSION BANDS

(kk) $\sigma^2\sigma^2\pi^4\sigma^2$, and both the dissociation energy and the vibration frequency of this state will be larger than that of the ground state of NO because the outermost antibonding electron π of the NO is missing in NO^+ . We may then also expect a $^1\Pi$ excited state in NO^+ analogous to the upper state of the L-B-H bands of N_2 which falls at about 8.55eV from the ground state ($^1\Sigma_g^+$). The (0,0) band of the new system is 9.06eV, and both the ω_e' and ω_e'' values of this system agree closely with those of the $^1\Pi_g - ^1\Sigma_g^+$ system of N_2 . Therefore, it seems quite probable that this system is due to transitions from the upper $^1\Pi$ state to the ground $^1\Sigma^+$ state of NO^+ . Experimentally, this system appears far stronger in the case of (He + NO) compared with pure NO. This also seems to support the idea suggested above.

In conclusion the author wishes to express his sincere gratitude to Professors R. S. Mulliken and J. R. Platt for their cooperation in this work, and to Mr. M. Schagrin for his assistance in the laboratory.
