

# Armed Services Technical Information Agency

Because of our limited supply, you are requested to return this copy WHEN IT HAS SERVED YOUR PURPOSE so that it may be made available to other requesters. Your cooperation will be appreciated.

**AD**

**36998**

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Reproduced by  
**DOCUMENT SERVICE CENTER**  
KNOTT BUILDING, DAYTON, 2, OHIO

**UNCLASSIFIED**

AD NO. 36998

ASTIA FILE COPY



THE PENNSYLVANIA STATE COLLEGE

*School of Chemistry and Physics*

STATE COLLEGE PENNSYLVANIA

TECHNICAL REPORT ON CONTRACT

N6 ONR - 269, T. O. III

by

J. G. Aston  
Director, Cryogenic Laboratory

July 26, 1954

(Contribution from the College of Chemistry and Physics of the  
Pennsylvania State University.)

THE THERMODYNAMIC PROPERTIES AND CONFIGURATION OF TRIMETHYLHYDRAZINE  
IDENTIFICATION AND CORRECTION FOR A MAJOR IMPURITY<sup>1</sup>

---

(1) This work was carried out under Contract N6onr-269 T.O. III of the O.N.R.

---

By J. G. Aston, T. P. Zolki and J. L. Wood

The heat capacities of trimethylhydrazine have been measured from 12 to 294°K. The triple point ( $201.24 \pm 0.01^\circ\text{K.}$ ), the heat of fusion ( $2267 \pm 1.5$  cal./mole), and the heat of vaporization at  $292.16^\circ\text{K.}$  ( $7949 \pm 7$  cal./mole) have been measured. The vapor pressures have been measured between 257 and  $287^\circ\text{K.}$ , and the data found to be represented by the equation  $\log_{10} p = -2614.506/T - 6.873235 \log T + 28.06810$ . The entropy of the ideal gas at  $292.16^\circ\text{K.}$  and 1 atm. was found to be  $79.45 \pm 0.14$  e.u. This value has been compared with that calculated from spectroscopic and molecular data to show that only one rotational isomer is present. The entropies of hydrazine and its mono and dimethyl derivatives have been reviewed and shown to be consistent with barriers and isomer composition closely similar to those of the analogous hydrocarbons.

A method is given for identifying and correcting for the presence of a major impurity.

Rotation about the nitrogen - nitrogen bond in trimethylhydrazine should give rise to isomerism similar to that found in hydrazine,<sup>2</sup> methylhydrazine<sup>3a,b</sup> and the

---

(2) W.G. Penney and G.B.B.M. Sutherland, Trans. Faraday Soc., 30, 902 (1934); J. Chem. Phys., 2, 492 (1934).

(3)(a) J.G. Aston, H.L. Fink, G.J. Janz and K.E. Russell, THIS JOURNAL, 73, 1939 (1951);  
(b) G.J. Janz and K.E. Russell, J. Chem. Phys., 17, 1352 (1949).

---

dimethylhydrazines,<sup>4a,b</sup> in which a skew form is favored. This paper presents the results

- 
- (4)(a) J. G. Aston, G. J. Janz and K.E. Russell, THIS JOURNAL, 73, 1943 (1951);  
 (b) J. G. Aston, J. L. Wood and T. P. Zolki, ibid., 75, 6202 (1953),
- 

of a calorimetric investigation of trimethylhydrazine and a comparison of the third law entropy with that from spectroscopic and molecular data, together with a discussion of the probable configuration of the molecule.

### Experimental

The Trimethylhydrazine Sample. - Trimethylhydrazine was prepared by the method of Class and Aston,<sup>5a,b</sup> purified by distillation through a column of glass helices with

- 
- (5)(a) J.B. Class and J.G. Aston, ibid., 73, 2359 (1951);  
 (b) J.B. Class, J.G. Aston and T.S. Oakwood, ibid., 75, 2937 (1953).
- 

approximately 100 theoretical plates, and further purified by fractional melting<sup>6</sup> in a

- 
- (6) J.G. Aston and S.V.R. Mastrangelo, Anal. Chem., 22, 636 (1950),
- 

glass apparatus. Unfortunately the sample still contained 2.6 mole per cent impurity, 2.0 mole per cent of which was proved to be unsym-dimethylhydrazine. To have effected further purification would have required a prohibitively large sample. However, since the amount and nature of the major impurity was known, this was not necessary.

Heat Capacity Measurements. - The heat capacities of a sample of 48.924 g. (0.66001 moles) were measured from 12 to 294°K. in calorimeter G, which was used in the work on unsym-dimethylhydrazine.<sup>4b</sup> The results are listed in Table I. Smoothed values at integral temperatures are given in Table II.

The mean deviation of the experimental points from the best curve is 1% below 20°K., 0.02% for the solid above 20°K., and less than 0.01% for the liquid.

Table I

## HEAT CAPACITY OF TRIMETHYLHYDRAZINE

Mol. wt. 74.126;  $0^{\circ}\text{C.} = 273.16^{\circ}\text{K.}$ ; 1 cal. = 4.1833 int. joules.

| Temp., $^{\circ}\text{K.}$ | $C_p$<br>cal.deg. $^{-1}$ mole $^{-1}$ | Temp., $^{\circ}\text{K.}$ | $C_p$<br>cal.deg. $^{-1}$ mole $^{-1}$ |
|----------------------------|----------------------------------------|----------------------------|----------------------------------------|
| 12.871                     | 0.78                                   | 61.819                     | 10.38                                  |
| 14.703                     | 1.08                                   | 67.604                     | 11.26                                  |
| 16.984                     | 1.44                                   | 71.902                     | 11.87                                  |
| 20.261                     | 2.18                                   | 76.787                     | 12.59                                  |
| 22.558                     | 2.70                                   | 81.461                     | 13.30                                  |
| 25.317                     | 3.32                                   | 83.802                     | 13.64                                  |
| 28.431                     | 4.05                                   |                            |                                        |
|                            |                                        | 91.721                     | 14.65                                  |
| 24.257                     | 3.11                                   | 96.865                     | 15.26                                  |
| 26.013                     | 3.50                                   | 101.818                    | 15.87                                  |
| 28.343                     | 4.02                                   | 106.605                    | 16.44                                  |
| 31.092                     | 4.67                                   | 111.244                    | 16.93                                  |
| 36.019                     | 5.78                                   | 116.172                    | 17.54                                  |
| 39.809                     | 6.54                                   | 121.360                    | 18.13                                  |
| 43.901                     | 7.27                                   |                            |                                        |
| 49.228                     | 8.28                                   | 111.288                    | 16.91                                  |
| 53.808                     | 9.08                                   | 117.484                    | 17.71                                  |
| 58.931                     | 9.91                                   | 123.492                    | 18.33                                  |
| 65.235                     | 10.92                                  | 129.312                    | 18.99                                  |
| 71.889                     | 11.86                                  | 134.960                    | 19.66                                  |

Table I, contd.

| Temp., °K. | $C_p - 1_{\text{mole}}^{-1}$<br>cal.deg. |
|------------|------------------------------------------|
| 140.450    | 20.24                                    |
| 145.771    | 20.77                                    |
| 152.025    | 21.36                                    |
| 158.173    | 21.91                                    |
| 164.499    | 22.90 premelt                            |
| 205.378    | 43.34 liq.                               |
| 211.466    | 43.63                                    |
| 217.003    | 43.84 <sub>5</sub>                       |
| 220.797    | 43.93                                    |
| 226.183    | 44.05 <sub>5</sub>                       |
| 232.625    | 44.13                                    |
| 239.012    | 44.16                                    |
| 248.831    | 44.33                                    |
| 257.122    | 44.45                                    |
| 263.387    | 44.26                                    |
| 269.590    | 44.37                                    |
| 275.737    | 44.42                                    |
| 281.838    | 44.42 <sub>5</sub>                       |
| 287.890    | 44.44                                    |
| 293.881    | 44.45                                    |

Table II

## HEAT CAPACITY OF TRIMETHYLHYLRAZINE AT INTEGRAL TEMPERATURES

Mol. wt. 74.126;  $0^{\circ}\text{C.} = 273.16^{\circ}\text{K.}$ ; 1 cal. = 4.1833 int. joules.

| Temp.,<br>$^{\circ}\text{K}$ | $C_p$<br>cal.deg. $^{-1}$ mole $^{-1}$ | Temp.,<br>$^{\circ}\text{K}$ | $C_p$<br>cal.deg. $^{-1}$ mole $^{-1}$ | Temp.,<br>$^{\circ}\text{K}$ | $C_p$<br>cal.deg. $^{-1}$ mole $^{-1}$ |
|------------------------------|----------------------------------------|------------------------------|----------------------------------------|------------------------------|----------------------------------------|
| 13                           | 0.80                                   | 80                           | 13.08                                  | 195                          | 24.225                                 |
| 14                           | 0.965                                  | 85                           | 13.80                                  | 200                          | 24.42                                  |
| 15                           | 1.125                                  | 90                           | 14.45                                  | 201.24                       | 24.47 cryst.                           |
| 16                           | 1.295                                  | 95                           | 15.04                                  | Fusion                       |                                        |
| 17                           | 1.47                                   | 100                          | 15.65                                  | 201.24                       | 43.12 liq.                             |
| 18                           | 1.65                                   | 105                          | 16.235                                 | 205                          | 43.315                                 |
| 19                           | 1.85                                   | 110                          | 16.80                                  | 210                          | 43.56                                  |
| 20                           | 2.07                                   | 115                          | 17.39                                  | 215                          | 43.77                                  |
| 21                           | 2.305                                  | 120                          | 17.975                                 | 220                          | 43.93                                  |
| 22                           | 2.555                                  | 125                          | 18.54                                  | 225                          | 44.03                                  |
| 23                           | 2.80                                   | 130                          | 19.105                                 | 230                          | 44.10                                  |
| 24                           | 3.04                                   | 135                          | 19.67                                  | 235                          | 44.145                                 |
| 25                           | 3.265                                  | 140                          | 20.195                                 | 240                          | 44.18                                  |
| 30                           | 4.41                                   | 145                          | 20.695                                 | 245                          | 44.27                                  |
| 35                           | 5.56                                   | 150                          | 21.17                                  | 250                          | 44.345                                 |
| 40                           | 6.57                                   | 155                          | 21.63                                  | 255                          | 44.42                                  |
| 45                           | 7.48                                   | 160                          | 22.055                                 | 260                          | 44.40                                  |
| 50                           | 8.43                                   | 165                          | 22.45                                  | 265                          | 44.27                                  |
| 55                           | 9.28                                   | 170                          | 22.82                                  | 270                          | 44.375                                 |
| 60                           | 10.09                                  | 175                          | 23.155                                 | 275                          | 44.415                                 |
| 65                           | 10.89                                  | 180                          | 23.47                                  | 280                          | 44.425                                 |
| 70                           | 11.60                                  | 185                          | 23.75                                  | 285                          | 44.435                                 |
| 75                           | 12.325                                 | 190                          | 24.00                                  | 290                          | 44.44                                  |
|                              |                                        |                              |                                        | 292.16                       | 44.445                                 |

Due to the presence of a large amount of impurity in the sample, premelting was observed at and above  $160^{\circ}\text{K}$ . Measurements were taken in the entire premelting region, and the heat capacity values listed in Table II between  $160^{\circ}\text{K}$ . and the melting point are based on a combination of correction and extrapolation. Values of the heat capacity of the solid near the melting point may be in error by several tenths of a per cent.

The Melting Point, the Impurity, and its Effect. Equilibrium temperatures of solid and liquid trimethylhydrazine were observed with increasing fractions of the sample melted. The data are summarized in Table III. The solid-insoluble impurity was found to be 2.6 mole per cent, and the calculated melting point of pure trimethylhydrazine is  $201.24 \pm 0.01^{\circ}\text{K}$ .

The intuition that the major part of the impurity was unsym-dimethylhydrazine was quantitatively confirmed by the thermal data in the premelting region. Therefore it is possible to correct exactly for the effect of the major impurity provided there is perfect solution; this is proved below.

From the heat of fusion of unsym-dimethylhydrazine<sup>4b</sup> and that of trimethylhydrazine, the eutectic temperature of the system trimethylhydrazine-unsym-dimethylhydrazine is calculated to be  $185.0^{\circ}\text{K}$ ., provided that Raoult's law is obeyed; the calculated eutectic composition is 0.609 mole per cent trimethylhydrazine and 0.391 mole per cent unsym-dimethylhydrazine; the heat of fusion of one mole of eutectic mixture is 2320 cal., assuming perfect solution.

A strong inflection was observed in the apparent heat capacity ( $\Delta E/\Delta T$ ) curve of the sample in the vicinity of  $180^{\circ}\text{K}$ .. From the calorimetric data enthalpies were calculated at the end of each energy addition. The solid upper curve of Figure 1 shows the heat capacities derived from these enthalpies. The maximum in the heat capacities occurs at  $182.2 \pm 0.1^{\circ}\text{K}$ . Using this value in conjunction with the calculated properties of the system trimethylhydrazine - unsym-dimethylhydrazine, it is calculated that the lowering in eutectic temperature corresponds to a third constituent present to the extent of 9.5

Table III

## MELTING POINT OF TRIMETHYLHYDRAZINE

| Temp., °K.       | 1/Fraction melted |
|------------------|-------------------|
| 193.540          | 8.301             |
| 197.236          | 4.350             |
| 197.743          | 3.778             |
| 199.036          | 2.394             |
| 199.659          | 1.715             |
| 200.031          | 1.313             |
| 200.32 (extrap.) | 1                 |
| 201.24 (extrap.) | 0                 |

mole per cent of the eutectic, or that the bulk sample contains 2.1 mole per cent unsym-dimethylhydrazine and 0.5 mole per cent of another impurity or impurities.

This conclusion may be checked in the following manner: In an excess of trimethylhydrazine, 0.391 moles of unsym-dimethylhydrazine are equivalent to one mole of eutectic mixture, and therefore the sample should contain 5.4 mole per cent eutectic. Now, at 182.2°K.,  $f = 1 = Q/(2320 N)$ , where  $f$  is the fraction of eutectic that has melted,  $N$  is the mole fraction of eutectic in the bulk sample, and  $Q$  is the area between the apparent heat capacity curve and the extrapolated true heat capacity curve (the lower curve of Figure 1) of the solid up to 182.2°K. The value of  $Q$  was found to be 126 cal./mole, and hence  $N = 0.054$  or 5.4 mole per cent, in agreement with the expected value.

A final argument in support of the contention that the major part of the impurity is unsym-dimethylhydrazine is provided by an examination of the fraction of sample melted at various temperatures up to 182.2°K. The fractions of eutectic melted may be obtained directly from the thermal data by making use of the assumed composition of the eutectic together with computed amount of other impurities to ascertain the specific heat of fusion. The results are given in the first two columns of Table IV. It is easily verified that the values of  $1/f$  listed in the first column are a linear function of the temperature. The value of  $T_0$  obtained by extrapolation to  $1/f = 0$  agrees with the calculated eutectic temperature well within the limits of experimental error. Another relation between  $1/f$  and  $T$  may be obtained by putting a straight line through the points ( $1/f = 0$ ,  $T = 185.0$ ) and ( $1/f = 1$ ,  $T = 182.2$ ) where  $f$  is again the fraction of eutectic that has melted. This procedure is solely based upon the hypothesis that unsym-dimethylhydrazine is present in the sample (which implies that  $T_0 = 185.0^\circ\text{K.}$ ), and is obviously independent of any assumptions regarding the quantities of unsym-dimethylhydrazine in the eutectic or other material in the sample. Equilibrium temperatures taken from this theoretical line are listed in the last column of Table IV. The calculated and observed temperatures must, of course, agree by hypothesis at  $1/f = 1$ , but the agreement at other values of  $1/f$  serves to confirm the assumptions that were made.

The dashed curve of Figure 1 shows the heat capacities calculated by using all the assumptions.

Table IV

## FRACTIONS OF EUTECTIC MELTED

| 1/Fraction melted | T., °K.<br>observed | T., °K.<br>calculated |
|-------------------|---------------------|-----------------------|
| 1.89              | 179.4               | 179.7                 |
| 1.57              | 180.3               | 180.6                 |
| 1.40              | 181.0               | 181.1                 |
| 1.21              | 181.5               | 181.6                 |
| 1.10              | 181.9               | 181.9                 |
| 1                 | 182.2               | (182.2)               |
| 0                 | 185.4 (extrap.)     | 185.0                 |

The values of the specific heat of unsym-dimethylhydrazine are so close to those of trimethylhydrazine at any temperature that no correction is necessary for the presence of 2.1 mole per cent of the former in the sample. Corrections of 0.5% are required in the heat of fusion and in the heat of vaporization, but since these corrections are known to about 3% of their value, their application introduces negligible uncertainty. However, the possibility remains that all results of this paper may be in error by 0.1% to 0.2% due to the presence of 0.5 mole per cent of the additional 0.5% impurities.

The Heat of Fusion. - Three independent determinations of the heat of fusion of trimethylhydrazine were made, and the resulting values are presented in Table.V. The usual corrections <sup>7</sup> have been applied to the heat input.

---

(7) J. G. Aston and G. J. Szasz, THIS JOURNAL, 69, 3108 (1947)

---

The entropy of mixing due to impurities is calculated to be 0.26 e.u. This is automatically excluded from the entropy of fusion when the heat capacities are corrected for premelting and all the heat due to melting assumed to be put in at the true melting point. On the other hand, if the melting phenomenon be treated in such a way that all latent heat is counted as sensible heat (i.e. the measured heat capacities are used directly and the main part of fusion is treated as a heat capacity over the range 1° below the melting point), the entropy of the liquid at the melting point does include the entropy of mixing.

The entropy of the liquid at the melting point calculated by the former method is 39.06 e.u. (not corrected for the slightly different heat of fusion of unsym-dimethylhydrazine), and consequently the second method should result in a value of 39.32 e.u. The actual value found by the sensible heat method is 39.34 e.u. As far as the authors are aware this is the first time attention has been called to the question of entropy of mixing where there is impurity and how this is eliminated.

Table V

## HEAT OF FUSION OF TRIMETHYLHYDRAZINE

Mol. wt. 74.126;  $0^{\circ}\text{C.} = 273.16^{\circ}\text{K.}$ ; 1 cal. = 4.1833 int. joules; m.p.  $200.32^{\circ}\text{K.}$

| Temp. interval, $^{\circ}\text{K.}$ | Heat input,<br>cal./mole | $\int \text{CpdT}$<br>cal./mole | Premelting,<br>cal./mole | $\Delta H$ fusion,<br>cal./mole |
|-------------------------------------|--------------------------|---------------------------------|--------------------------|---------------------------------|
| 164.082 - 203.940                   | 4220.0                   | 1943.5                          | 1.1                      | 2277.6                          |
| 162.796 - 203.298                   | 4108.1                   | 1827.9                          | 0.0                      | 2280.2                          |
| 161.420 - 202.111                   | 4088.4                   | 1808.9                          | 0.0                      | 2279.5                          |

Mean  $\Delta H = 2279.1 \pm 1.5$

Mean  $\Delta H$ , corrected for the different heat of fusion  
of 2.1 mole per cent unsym-dimethylhydrazine = 2267.1

The Vapor Pressures. - The vapor pressure of trimethylhydrazine at 292.16°K. is required for the evaluation of the entropy of compression of the gas at that temperature. Measurements were made as for methylhydrazine <sup>3a</sup>. Observed values of the vapor pressure at various temperatures are listed in column 2 of Table VI. These were fitted to the equation

$$\log p = - 2621.916/T - 6.920951 \log T + 28.20910 \quad (1)$$

Deviations of the experimental values from this equation are tabulated in column 3. Use was made of the vapor pressure equation for unsym-dimethylhydrazine <sup>4b</sup> to correct the observed pressures for the presence of 2.1 mole per cent of that substance. Corrected values are listed in the last column, and are summarized by the equation

$$\log p = - 2614.506/T - 6.873235 \log T + 28.06810 \quad (2)$$

The values of  $\Delta C_p$  calculated from equation (1) or equation (2) agree with that estimated from the spectroscopic and molecular data for the gas along with the measured heat capacities of the liquid to about 6 cal., which is in accord with the consistency of the data. As a check on internal consistency, the value of the heat of vaporization at 292.16°K. calculated from the corrected vapor pressures as given by equation (2) is appended at the foot of Table VII for comparison with the experimental value corrected for impurity.

In spite of the thermodynamic consistency of the corrected as well as the uncorrected vapor pressures, their reliability is open to serious doubt because of the presence of 0.5 mole per cent of unknown impurity or impurities in the sample. The vapor pressure at 274°K. was lowered by 0.8 mm. after about one third of the sample had been distilled off, whereas the calculated drop due to enrichment of the residue in the less volatile unsym-dimethylhydrazine only amounts to 0.04 mm.

The Heat of Vaporization. - Measurements were made as for methylhydrazine <sup>3a</sup>, except that a grooved stopcock was used instead of a valve to maintain the temperature constant. The results are presented in Table VII. Due account was taken of the temperature variation

Table VI

## VAPOR PRESSURES OF TRIMETHYLHYDRAZINE

$$0^{\circ}\text{C.} = 273.16^{\circ}\text{K.}$$

| T, $^{\circ}\text{K.}$ | P <sub>obs.</sub> , mm. | P <sub>calcd.</sub> - | Corrected<br>P |
|------------------------|-------------------------|-----------------------|----------------|
|                        |                         | P <sub>obs.</sub>     |                |
| 256.939                | 21.20                   | 0.00                  | 21.34          |
| 263.791                | 32.52                   | + 0.01                | 32.73          |
| 269.058                | 44.41                   | 0.00                  | 44.67          |
| 274.222                | 59.38                   | + 0.02                | 59.72          |
| 279.568                | 79.19                   | - 0.01                | 79.59          |
| 279.576                | 79.25                   | - 0.03                | 79.63          |
| 286.986                | 115.44                  | - 0.01                | 115.98         |
| 286.995                | 115.51                  | - 0.02                | 116.05         |
| 292.16                 | 148.05 (extrap.)        |                       | 148.72         |

Table VII

## HEATS OF VAPORIZATION OF TRIMETHYLHYDRAZINE

Mol. wt. 74.126;  $0^{\circ}\text{C.} = 273.16^{\circ}\text{K.}$ , 1 cal. = 4.1833 int. joules

| Mole<br>vaporized | Mean temp. of<br>vaporization,<br>$^{\circ}\text{K}$ | $\Delta H$<br>real gas,<br>$T, ^{\circ}\text{K.}$ | $\Delta H$<br>real gas<br>$292.16^{\circ}\text{K}$<br>cal./mole |
|-------------------|------------------------------------------------------|---------------------------------------------------|-----------------------------------------------------------------|
| 0.047673          | 292.525                                              | 7984                                              | 7991                                                            |
| .065766           | 292.614                                              | 7990                                              | 7999                                                            |
| .030584           | 292.801                                              | 7978                                              | 7991                                                            |
| .032602           | 292.703                                              | 7976                                              | 7987                                                            |

Average 7992  $\pm$  7

Value corrected for presence of 2.1 mole  
per cent unsym-dimethylhydrazine

7949

Calculated from equation (2) and estimated  
critical data,  $T_c = 500^{\circ}\text{K.}$ ,  $P_c = 46 \text{ atm.}$ ,  
(Berthelot correction = 62 cal.)

7910

of the heat of vaporization of unsym-dimethylhydrazine in the correction for the presence of 2.1 mole per cent of that substance in the sample.

Since these determinations involved the use of stopcocks, they were made after all other measurements had been completed.

The Entropy from Calorimetric Data. - The calculation of the entropy of trimethylhydrazine from the calorimetric data is summarized in Table VIII. The correction for gas imperfection was made with the estimated critical constants given in Table VII in conjunction with the Berthelot equation.

While most of the work on trimethylhydrazine was in progress, this laboratory underwent extensive structural alterations in the course of which the room temperature could not be maintained much above 20°C. In order to avoid condensation of material, it was necessary to abandon the usual standard temperature of 298.16°K. in reporting results.

The Entropy from Spectroscopic and Molecular Structure Data. - The possible forms of trimethylhydrazine, in order of decreasing probability, are the outer form, the inner form, and the trans form, each corresponding to a sector of  $2\pi/3$  for rotation about the N - N bond. Each form exhibits a pair of optical isomers. The inner form has the greatest clustering of methyl groups; the nomenclature is that used for methylhydrazine.<sup>3b</sup>

The entropy was calculated for the outer form, assuming all angles to be tetrahedral and using the following bond distances:<sup>8</sup>

---

(8) W. H. Beamer, ibid., 70, 2979 (1948).

---

N-N, 1.45 Å.; C-N, 1.47 Å.; C-H, 1.09 Å.; N-H, 1.04 Å. The product of the principal moments of inertia is  $6.12 \times 10^{-114}$  g. cm.<sup>2</sup> The reduced moments are  $23.826 \times 10^{-40}$  g. cm.<sup>2</sup> for the methylamino group,  $5.055 \times 10^{-40}$  g. cm.<sup>2</sup> for the methyl group of the methylamino group and the methyl group farthest removed from the methylamino group, and  $5.150 \times$

Table VIII

## ENTROPY OF TRIMETHYLHYDRAZINE FROM THERMAL DATA

|                                                                    | E.u./mole                            |
|--------------------------------------------------------------------|--------------------------------------|
| 0 - 15°K., Debye function, $\theta = 141$ , six degrees of freedom | $0.368 \pm 0.02$                     |
| 15°K. - 155°K.                                                     | $21.311 \pm 0.02$                    |
| 155°K. - 201.24°K., graphical                                      | $6.055 \pm 0.00$                     |
| Fusion, 2267.1/201.24                                              | $11.266 \pm 0.01$                    |
| 201.24°K. - 292.16°K., graphical                                   | <u><math>16.445 \pm 0.02</math></u>  |
| Entropy of liquid at 292.16°K.                                     | $55.445 \pm 0.07$                    |
| Vaporization, 7949/292.16                                          | <u><math>27.208 \pm 0.03</math></u>  |
| Entropy real gas 292.16°K., 148.7 mm.                              | $82.653 \pm 0.10$                    |
| Entropy ideal gas at 292.16°K., 148.7 mm.                          | $82.689 \pm 0.11$                    |
| Compression, $R \ln 148.7/760$                                     | - <u><math>3.241 \pm 0.03</math></u> |
| Entropy ideal gas at 292.16°K., 760 mm.                            | $79.45 \pm 0.14$                     |

$10^{-40}$  g. cm.<sup>2</sup> for the third methyl group. They were calculated by the method of Pitzer.<sup>9a,b,c</sup>

- 
- (9)(a) Pitzer and Gwinn, *J. Chem. Phys.*, 10, 428 (1942);  
 (b) K. S. Pitzer, *ibid.*, 14, 239 (1946);  
 (c) J. E. Kilpatrick and K. E. Pitzer, *ibid.*, 17, 1064 (1949).
- 

The frequencies used in the calculation of the vibrational entropy contribution were 307,414, 436, 498, 712, 783, 888, 1007, 1044. (2), 1134 (2), 1168 (2), 1180, 1270, 1398 (3), 1477 (6), 2858 (4), 2970 (2), 3082, 3092 (2), 3405 cm.<sup>-1</sup> <sup>10</sup>

- 
- (10) E.R. Shull, J.L. Wood, J.G. Aston and D.H. Rank, *ibid.*, 22, 0000 (1954).
- 

The entropy calculation is summarized in Table IX. The choice of barriers of 3850 ± 150 cal./mole hindering all internal rotations leads to agreement of the calculated entropy of the outer form with the experimental value. This is consistent with the result for the hydrocarbon analogue, isopentane, which apparently exists in the C<sub>1</sub> form with an average barrier of about 3200 cal./mole, but with methyl barriers as high as 3620 cal./mole. <sup>11</sup> For completeness the data on the liquid have been extrapolated, and a calculation made based on the spectroscopic data and the derived barrier to yield the entropies of the

- 
- (11) Scott, McCullough, Williamson and Waddington, *THIS JOURNAL*, 73, 1707 (1951).
- 

liquid and the ideal vapor at 298.16°K and 1 atm. The values are respectively 56.34 and 80.00 e.u./mole.

#### Discussion

The potential barrier hindering rotation about the nitrogen-nitrogen bond in hydrazine <sup>12</sup> has the same magnitude as the methyl barrier in ethane.

- 
- (12) Scott, Oliver, Gross, Hubbard and Huffman, *ibid.*, 71, 2293 (1949).
-

Table IX

ENTROPY OF TRIMETHYLHYDRAZINE FROM MOLECULAR DATA AT 292.16°K.

|                                            | <u>outer form</u><br>E.u. |
|--------------------------------------------|---------------------------|
| S translation                              | 38.723                    |
| S vibration                                | 4.897                     |
| S external rotation                        | 25.555                    |
| S mixing optical isomers ( $R \ln 2$ )     | 1.377                     |
|                                            | <hr/>                     |
|                                            | 70.552                    |
| S int. rot. Me ( $V = 3850$ )              | 1.877                     |
| S int. rot. Me                             | 1.862                     |
| S int. rot. Me                             | 1.862                     |
| S int. rot. $\text{NHCH}_3$ ( $V = 3850$ ) | 3.292                     |
|                                            | <hr/>                     |
| Sum                                        | 79.44                     |
| Observed                                   | 79.45 $\pm$ 0.14          |

For methylhydrazine, several choices for the barrier hindering internal rotation of the methyl group were considered: <sup>3a</sup> A barrier of 2800 cal./mole if the molecule be entirely in the outer form; 3200 cal./mole for a mixture of 96 mole per cent outer form and 4 mole per cent inner form; 3500 cal./mole for a mixture of 92 mole per cent outer form and 8 mole per cent inner form. This <sup>latter</sup> corresponds to an energy difference of 1450 cal./mole. An assignment of 3500 cal./mole agrees within limits of experimental error with the methyl barrier of the hydrocarbon analogue, propane (3400 cal./mole).

The case of sym-dimethylhydrazine will be considered later in some detail. Regardless of the interpretation of experimental results, the choice of methyl barriers in sym-dimethylhydrazine of 3000 cal./mole is compatible with the value for its hydrocarbon analogue, butane (3300 cal./mole). <sup>13</sup>

---

(13) K. S. Pitzer and J. E. Kilpatrick, Chem. Revs., 39, 435 (1946).

---

Only one assignment is possible for the methyl barriers of unsym-dimethylhydrazine in order to obtain agreement between the calculated and the observed entropy, namely 3700 cal./mole, <sup>4b</sup> which corresponds to the existence of the molecule in the skew "gauche" form and agrees within limits of experimental error with the value of the methyl barriers in the hydrocarbon analogue, isobutane (3620 cal./mole). <sup>13</sup>

The results for trimethylhydrazine show so much similarity to those for isopentane as to suggest a revision of the conclusions that were drawn from an entropy study of sym-dimethylhydrazine. <sup>4a</sup> There will be three forms, outer-outer, inner-outer, and inner-inner, both of the latter corresponding to the gauche configuration of butane. Denoting an inner interaction energy by the letter i, and a gauche interaction energy by the letter g, the three forms (oo, io and ii) of sym-dimethylhydrazine have the interaction energies

$$\underline{oo} = 0, \underline{io} = i + g, \underline{ii} = 2i + g.$$

Adopting Pitzer's value <sup>14</sup> of 800 cal./mole for the energy difference between potential

---

(14) K. S. Pitzer, THIS JOURNAL, 63, 2413 (1941).

---

minima (gauche interaction energy), the inner interaction energy required to obtain agreement with the observed entropy is 400 cal./mole, and sym-dimethylhydrazine exists as a mixture of approximately 75 mole per cent outer-outer form, 20 mole per cent inner-outer form and 5 mole per cent inner-inner form. It has not been possible to relate any of the lines in the spectrum to these forms.<sup>15</sup> Strong hydrogen bonding in the liquid

---

(15) D.W.E. Axford, G.J. Janz and K.E. Russell, J. Chem. Phys., 19, 704 (1951).

---

would make a study of the temperature dependence difficult. A determination of the gaseous heat capacities of all the methyl-substituted hydrazines is obviously the best way to verify the conclusions.

It should be noted that the inner interaction energy for sym-dimethylhydrazine is due to the effect of the interaction of one hydrogen atom with the inner methyl group, the other part of the interaction being absorbed in the 300 cal./mole gauche interaction. Therefore, twice the value of 400 cal./mole or 800 cal./mole is to be compared with the value of 1450 cal./mole found for the energy difference between the two form of methylhydrazine. If the value of 800 cal./mole be adopted for this inner interaction, and using a methyl barrier of 3500 cal./mole, the value obtained for the entropy of methylhydrazine is 67.06 e.u., compared with the experimental value of  $66.61 \pm 0.2$  e.u. at 298.16°K. This corresponds to 21 mole per cent of the inner form and to a dipole moment of 1.76 D. The measured value is  $1.68 \pm 0.14$  D.<sup>16</sup>

---

(16) West and Killingsworth, ibid., 6, 1 (1938).

---

Acknowledgement. - We wish to thank Dr. J. B. Class for preparing the trimethylhydrazine sample, Mr. H. L. Pinch for help with the measurements and L. F. Shultz for the preparation of liquid hydrogen and technical maintenance of facilities.

STATE COLLEGE, PENNA.




Figure I