

UNCLASSIFIED

**A
D 86021**

Armed Services Technical Information Agency

Reproduced by
DOCUMENT SERVICE CENTER
KNOTT BUILDING, DAYTON, 2, OHIO

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address: **Armed Services Technical Information Agency, Document Service Center, Knott Building, Dayton 2, Ohio.**

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED

AD No. 86021

ASTIA FILE COPY

U.S. AIR FORCE
Project **RAND** **FC**
RESEARCH MEMORANDUM



This is a working paper. It may be expanded, modified, or withdrawn at any time. The views, conclusions, and recommendations expressed herein do not necessarily reflect the official views or policies of the U. S. Air Force.

The **RAND** *Corporation*
SANTA MONICA • CALIFORNIA

U. S. AIR FORCE

PROJECT RAND

RESEARCH MEMORANDUM

ON MULTI-STAGE GAMES WITH IMPRECISE PAYOFF

Richard Bellman

RM-1337

9 September 1954

Assigned to _____

ASTIA (17)

This is a working paper. It may be expanded, modified, or withdrawn at any time. The views, conclusions, and recommendations expressed herein do not necessarily reflect the official views or policies of the United States Air Force.

The **RAND** Corporation

1700 MAIN ST. • SANTA MONICA • CALIFORNIA

Summary: It is shown that under certain natural conditions the play of a multi-stage game is to a great extent independent of the payoff function.

ON MULTI-STAGE GAMES WITH IMPRECISE PAYOFF

Richard Bellman

§1. Introduction

There is a large class of situations of economic and military significance which can be considered to be multi-stage games. In many of these situations, the payoff function is easily determined; in others, it is a matter of difficulty to determine a suitable criterion.

The purpose of this note is to show, in a heuristic fashion, that in many cases the optimal play is independent of the precise form of the payoff, provided only that this payoff possesses certain intuitive properties.

§2. Description of the Multi-stage Game

Let us consider a zero-sum multi-stage game where each play is determined by the game matrix $A = (a_{ij})$. Let initially the first player possess a quantity x of resources and the second player a quantity y . Since the game is zero sum, the state of the game at any time is described by x .

Defining a suitable criterion function, let $f(x)$ represent the value of the game to the first player.

Then, $f(x)$ satisfies the functional equation

$$\begin{aligned} f(x) &= \text{Min}_q \text{Max}_p \sum_{I,J} p_i q_j f(x + a_{ij}) \\ &= \text{Max}_p \text{Min}_q \sum_{I,J} p_i q_j f(x + a_{ij}) \end{aligned} \quad (2.1)$$

(see [1], [3]). The quantities p_i and q_j will, in general, depend upon x .

§3. Assumptions Concerning $f(x)$

Let us now assume that x and y are large compared to a_{ij} . In other words, the state of the system is only slightly disturbed by any one play of the game. Furthermore, let us assume that the value of the matrix A is zero, which is to say, it is on the average a fair game. Otherwise, the play is relatively trivial.

Finally, we assume that it pays to start with a larger initial resource. Then

$$f'(x) > 0 \quad (3.1)$$

§4. Heuristic Conclusion

Under these assumptions, we wish to show plausibly but not rigorously that the p_i and q_j are approximately independent of x and $f(x)$. This means that, under these assumptions, on each

play the players attempt to maximize the single-stage return, the ordinary expected value.

Let us write

$$f(x + a_{1j}) \approx f(x) + a_{1j}f'(x) \quad (4.1)$$

Then, from (2.1),

$$f(x) \approx \text{Min}_q \text{Max}_p \sum_{i,j} p_i q_j [f(x) + a_{1j}f'(x)] \quad (4.2)$$

or

$$f(x) \approx f(x) \sum_{i,j} p_i q_j + \text{Min}_q \text{Max}_p f'(x) \sum_{i,j} a_{1j} p_i q_j \quad (4.3)$$

whence, since $f'(x) \neq 0$,

$$0 \approx \text{Min}_q \text{Max}_p \sum_{i,j} a_{1j} p_i q_j \quad (4.4)$$

$$\approx \text{Max}_p \text{Min}_q \sum_{i,j} a_{1j} p_i q_j$$

§5. Nonzero-sum Games

Let us consider a two-person, multi-stage, nonzero-sum game where the first and second players have, respectively, the game matrices

$$A = (a_{ij}), \quad B = (b_{ij}) \quad (5.1)$$

and initially the amounts x and y , respectively.

Let $f(x,y)$ be some criterion function, such as probability of survival, assumed to satisfy the conditions

$$f_x > 0, \quad f_y < 0 \quad (5.2)$$

and assume that

$$\sum_{i,j} a_{ij} p_i q_j, \quad \sum_{i,j} b_{ij} p_i q_j < 0 \quad (5.3)$$

a game of attrition.

Then $f(x,y)$ satisfies the functional equation

$$\begin{aligned} f(x,y) &= \text{Max}_p \text{Min}_q \left[\sum_{i,j} p_i q_j f(x+a_{ij}, y+b_{ij}) \right], \quad x,y > 0 \\ &= \text{Min}_q \text{Max}_p \left[\dots \right] \\ &= 1, \quad x > 0, \quad y < 0 \\ &= 0, \quad x < 0, \quad y > 0 \\ &= 1/2, \quad x = y = 0 \quad (\text{for the sake of completeness}) \end{aligned} \quad (5.4)$$

Assume as above that x and y are large compared to a_{ij} and b_{ij} .

Then

$$f(x+a_{ij}, y+b_{ij}) \approx f(x,y) + a_{ij} f_x + b_{ij} f_y \quad (5.5)$$

Equation (5.4) then yields

$$0 \approx \text{Max}_p \text{Min}_q \left[f_x \sum_{i,j} a_{ij} p_i q_j + f_y \sum_{i,j} b_{ij} p_i q_j \right] \quad (5.6)$$

$$\approx \text{Min}_q \text{Max}_p \left[\dots \right]$$

or

$$\frac{f_x}{f_y} \approx \text{Min}_p \text{Max}_q \left[\frac{\sum a_{ij} p_i q_j}{\sum b_{ij} p_i q_j} \right] \quad (5.7)$$

$$\approx \text{Max}_p \text{Min}_q \left[\frac{\sum a_{ij} p_i q_j}{\sum b_{ij} p_i q_j} \right]$$

This shows that the single-stage play is approximately governed by the criterion function

$$K(p,q) = \frac{\sum a_{ij} p_i q_j}{\sum b_{ij} p_i q_j} \quad (5.8)$$

That min-max = max-min in (5.7) is a result due to von Neumann. An elegant short proof based on the usual min-max theorem will be found in [4].

We thus have a rationale for the play of large classes of two-person nonzero-sum games.

BIBLIOGRAPHY

1. Bellman, R. and J. LaSalle, On Nonzero-sum Games and Stochastic Processes, The RAND Corporation, RM-212, August 1949.
2. ——— and D. Blackwell, On a Particular Nonzero-sum Game, The RAND Corporation, RM-250, September 1949.
3. ———, An Introduction to the Theory of Dynamic Programming, The RAND Corporation, R-245, June 1953.
4. Shapley, L., "Stochastic Games," Proc. Nat. Acad. Sci., Vol. 39, No. 10, October 1953.