

UNCLASSIFIED

**A
D 218020**

Armed Services Technical Information Agency

**ARLINGTON HALL STATION
ARLINGTON 12 VIRGINIA**

**FOR
MICRO-CARD
CONTROL ONLY**

1 OF 2

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

DISCLAIMER NOTICE

**THIS DOCUMENT IS BEST QUALITY
PRACTICABLE. THE COPY FURNISHED
TO DTIC CONTAINED A SIGNIFICANT
NUMBER OF PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

218020
AD NO. 218020
ASTIA FILE COPY

HEADQUARTERS
QUARTERMASTER RESEARCH & ENGINEERING COMMAND
U S ARMY

TECHNICAL REPORT
EP-III

ASTIA
JUL 24 1959
1100R B

CLIMATIC ANALOGS OF FORT GREELY, ALASKA AND
FORT CHURCHILL, CANADA IN NORTH AMERICA

FC
BAC



QUARTERMASTER RESEARCH & ENGINEERING CENTER
ENVIRONMENTAL PROTECTION RESEARCH DIVISION

MAY 1959

NATICK, MASSACHUSETTS

HEADQUARTERS
QUARTERMASTER RESEARCH & ENGINEERING COMMAND, US ARMY
OFFICE OF THE COMMANDING GENERAL
NATICK, MASSACHUSETTS

Major General Andrew T. McNamara
The Quartermaster General
Department of the Army
Washington 25, D. C.

Dear General McNamara:

This report, "Climatic Analogs of Fort Greely and Fort Churchill in North America", compares climatic conditions throughout northern North America with those at two Department of the Army cold-weather test sites.

The analysis of individual climatic elements shows large areas of close analogy with each test site. Combinations of elements (e.g., composite of winter and summer temperatures with mean annual precipitation) however, show much smaller areas of analogy.

The distributions mapped in this study provide a basis for determining the degree to which each test site is climatically representative of the North American Arctic and Subarctic. The study will aid test and design personnel in estimating regional suitability of end items based upon performance trials conducted at either test site. Furthermore, areas of more severe climatic stress, shown in this report, should be considered when clothing and equipage are being designed for use in the extreme cold.

Sincerely yours,

C. G. Calloway
C. G. CALLOWAY
Major General, USA
Commanding

1 Incl
EP-111

HEADQUARTERS
QUARTERMASTER RESEARCH & ENGINEERING COMMAND, US ARMY
Quartermaster Research & Engineering Center
Natick, Massachusetts

ENVIRONMENTAL PROTECTION RESEARCH DIVISION

Technical Report

EP-111

CLIMATIC ANALOGS OF FORT GREELY, ALASKA, AND
FORT CHURCHILL, CANADA, IN NORTH AMERICA

Andrew D. Hastings, Jr.
Geographer

REGIONAL ENVIRONMENTS RESEARCH BRANCH

Prepared for the R&D Project "Military Evaluation of Geographic Areas"
(8-70-09-400), U. S. Army Engineer Waterways Experiment Station
Vicksburg, Mississippi

Project Reference:
7-83-01-005

May 1969

Foreword

This report is one of a series designed to determine the extent of world areas having climates similar to those of the various Department of the Army test sites. The present study compares climatic conditions at Fort Greely, Alaska, and Fort Churchill, Manitoba, Canada, with those of other parts of northern North America as well as certain western islands of Europe (Iceland, Jan Mayen, the Faeroes, and the British Isles). Data are plotted as far south as 45° North latitude.

The series of climatic analog studies, prepared at the request of the U.S. Army Waterways Experiment Station, Corps of Engineers, Vicksburg, Mississippi, is part of a program for evaluating the suitability and representativeness of all Department of the Army test sites. They constitute the Quartermaster Corps contribution to the Corps of Engineers project on "Military Evaluation of Geographic Areas."

AUSTIN HENSCHKE, Ph.D.
Chief
Environmental Protection Research
Division

Approved:

CARL L. WHITNEY, Lt Col, QMC
Commanding Officer
QM R&E Center Laboratories

J. FRED OESTERLING, Ph.D.
Acting Scientific Director
QM Research & Engineering Command

Contents

	<u>Page</u>
Abstract	iv
1. Introduction	1
2. Major physical features	1
3. Climate	4
4. Analysis of climatic elements	6
5. Analysis of composite maps	16
6. Tables of monthly climatic values for selected stations	18
7. Limitations of methods	18
8. Bibliography	33
9. Maps	43
Map Errata	44

Abstract

Fort Greely, Alaska, and Fort Churchill, Manitoba, Canada, are used by the Armed Forces of the United States as cold-weather field laboratories where physiological performance studies are conducted and effectiveness of clothing and equipment for cold climate operations is evaluated. This study undertakes to indicate the extent to which the climatic environment at each of the test sites is representative of arctic and sub-arctic areas elsewhere in North America.

Twelve climatic elements are mapped and analogous areas are indicated. Ranges of analogy vary with the element being measured, but for most elements the southern limit of close analogy for both test sites lies north of the 45° parallel (roughly the northern tier of states in the United States). Areas of analogy are designated on the maps by yellow areas for Fort Greely and blue areas for Fort Churchill. Except for absolute minimum temperature, Fort Churchill has the colder environment of the two sites; hence the more northerly zone of cold analogy shown is usually that of Fort Churchill. Areas of analogy for both test sites tend to extend farthest southward over midcontinent regions.

Composite areas of analogy for multiple elements are also mapped. Areas analogous to Fort Churchill for combined coldest month temperatures, snow depth, and windchill are found primarily in the Hudson Bay Lowland. For Fort Greely, analogous areas for the above combination of elements are confined to the immediate vicinity of Fort Greely, northwestern Alaska, southern Manitoba, and small areas on the coast of Greenland.

Areas of composite analogy with Fort Greely for winter and summer temperature and mean annual precipitation occur in the Tanana Valley of Alaska, southern Yukon Territory, and in small scattered areas of western Alaska. Fort Churchill analogy for the same combination of elements occurs in a 500-mile-wide salient stretching westward from Hudson Bay and a smaller patch in interior northern Quebec.

CLIMATIC ANALOGS OF FORT GREELY, ALASKA, AND FORT CHURCHILL, CANADA, IN NORTH AMERICA

1. Introduction

This study was prepared in answer to a request from the Office, Chief of Engineers, for a climatic comparison of the arctic and subarctic regions of the world with Fort Greely, Alaska, and Fort Churchill, Manitoba, Canada. It is the second of two reports, each treating one-half of the Northern Hemisphere north of 45° N. latitude; the first covered Eurasia. The land areas covered in this, the western portion, include North America, and certain western islands of Europe (Iceland, Jan Mayen Island, the Faeroe Islands, and the British Isles).

Only those climatic elements which seem to best define stress factors in cold environments were selected for the study. For each selected element a map (or maps) has been prepared showing areas having conditions considered analogous to those at Fort Greely and Fort Churchill. Information from some of these maps has been consolidated on composite maps to show areas of coincidence of analogous elements.

The climatic conditions at Fort Greely and Fort Churchill have been summarized and evaluated by the Environmental Protection Research Division in the following publications:

Handbook of Fort Churchill Environment, Technical Report EP-4 (25)

Handbook of Big Delta*, Alaska, Environment, Technical Report EP-5 (23)

Data from these and other studies listed in the bibliography furnished the bases for comparison in this report.

2. Major physical features (Fig. 2)

Physiographically, the arctic and subarctic North American mainland have a certain basic uniformity. The terrain of the interior is relatively low and gentle and bordering uplands lie on or near the east and west coasts.

The Hudson Bay Lowland and its northward extension into the Arctic Archipelago form a broad depression with Hudson Bay in the center. This is a country of boggy tundra which lies, for the most part, close to sea level.

* Renamed Fort Greely in August, 1955

Fort Churchill is located within this area on the west shore of Hudson Bay at the mouth of the Churchill River; the meteorological station is on a low moraine with the Bay to the north and relatively level muskeg to the south.

Except in the extreme insular north, this lowland is bordered by the Canadian Shield (Laurentian Upland). Here the term "upland" can be somewhat misleading. Although isolated elevations within this region exceed 3,000 feet, the transition from lowland to shield topography is gradual in many areas and much of the upland averages no more than 300 feet above sea level. Basically the shield is distinctive because of its old crystalline bedrock rather than its elevation.

In the District of Keewatin the Shield extends north of the forested subarctic as a moss-covered barren. South and southeastward it becomes a subdued upland covered by taiga forest. The taiga is a diverse wilderness of dense spruce forests, muskeg, lakes, barren knobs, glacial debris, and patternless water courses. Tree line is a relatively sharp break in many instances; where the forest ends, tundra begins, with scattered outlying trees in protected spots.

A nearly continuous fringe of lowlands surrounds the Shield country from the lower Mackenzie Valley (Mackenzie Lowland) in the northwest down through the northern United States (Central Lowlands and Great Lakes Lowland) and on around to Hudson Strait in northern Quebec. This lowland is prominently marked by a succession of some of the largest lakes on the continent: Great Bear, Great Slave, Athabaska, Winnipeg, Lake of the Woods, and the Great Lakes.

Southwestward the lowland slopes gently upward to reach over 4,000 feet in the Great Plains. The rise into the Franklin and Mackenzie Mountains in the northwest and into the Appalachian Plateau and related uplands of the southeast is far more abrupt.

The major highlands of the continent rise beyond the belt of circumferential lowlands and transitional plains. In the west, the great ranges of the Rocky Mountains stretch from the Mackenzie Mountains near the Beaufort Sea to well south of the area in the United States shown on the maps in this report. In the east, lower and more discontinuous mountains rim the Atlantic Coast of the continent from the Appalachian Mountains to northern Ellesmere Island. Highest summits of the eastern ranges average little more than 5,000 feet; those of the Rockies are well over twice as high. West of the Rockies lies a fairly continuous band of intermontane plateaus extending from the Columbia Plateau in the United States to the Tanana Hills of Alaska.

Along the western margin of the continent, coastal ranges form a high, nearly continuous wall from the Aleutian Chain to Mexico. Alaskan coastal mountains are heavily mantled with icecaps and valley glaciers, with tongues of the glaciers often terminating in the coastal waters. The Malaspina and Bering Glaciers fan out on the coastal plain of southern Alaska to form two of the largest piedmont glaciers in the world. Glaciers and permanent snow fields are common at higher elevations as far south as southern British Columbia. Even the Cascades in Oregon support a few glaciers.

Icefields are rare in the Rocky Mountains north of the Yukon-British Columbia boundary because the coastal ranges effectively impede the passage of moisture-bearing winds from the west. In southern Alberta and British Columbia, significant icefields do occur mainly as a result of considerably higher elevations and a rather broad system of transverse valleys through the coastal ranges which give access to the westerly winds.

Southern Alaska is dominated by the extensively glaciated Alaska Range and its eastern extension, the Wrangell Mountains. North of these ranges the valley of the Yukon River and its tributary, the Tanana River, drain westward from Yukon Territory. Fort Greely is situated in the Tanana Valley at the point where the Delta River joins the valley, about halfway between the Canadian border and the confluence of the Tanana and the Yukon rivers. The elevation at Fort Greely is 1,274 feet. Within 40 miles to the south, glaciated summits around Mount Hayes rise to over 13,000 feet.

In the central region of Alaska, forested plains penetrate far inland along major streams. These river plains are separated by clusters of dissected uplands such as the Kilbuck Mountains, the Kuskokwim Range, and the mountains in the area bordering Bering Strait.

North of the central valleys rise the barren summits of the Brooks Range. Poleward from there, the northern coastal lowland, or Arctic Shelf, reaches across all of northern Alaska and southwestward along the Bering Strait.

Of Greenland's total area of 840,000 square miles, over 705,000 are blanketed with a great ice sheet rising from valley glaciers on or near the coasts to an interior plateau level of about 10,000 feet. Although mountain peaks are prominent (some over 11,000 feet) along the east coast, the overall aspect of the island is one of a uniformly domed icecap bordered by a relatively narrow fringe of fiordal coast.

Iceland is a volcanic island which averages about 2,500 feet above sea level with ice-capped highlands nearly 7,000 feet high in the southeast. Very narrow lowlands are confined mainly to the southwest and north coasts.

Ireland is largely deforested lowland. Small upland areas occur principally along the northwest and southeast coasts. On the island of Great Britain, rolling, low topography characterizes most of England, but the Pennine Chain, with elevations of about 1,500 feet, forms a prominent north-south spine in the northern interior. The Cambrian Mountains of Wales average about 2,000 feet, topped by Munt Snowdon, 3,560 feet. The Highlands of Scotland have many elevations above 3,000 feet. Ben Nevis, the highest peak, is over 4,400 feet and has a permanent snowfield in the Great Corrie of Allt-a-Mhullin.

The eastern tip of Siberia and the northwest coast of France, which both appear in the map area, are discussed in the Eurasian counterpart of this study.(28)

3. Climate

Arctic North America is an area of extremely cold winters, cool or cold summers, and scanty precipitation, most of which falls as snow, usually heaviest in autumn. These conditions are most extreme in interior Greenland where mean annual temperatures average about -20°F and summer temperatures seldom, if ever, rise above freezing. Included in the Arctic climatic region as used here, are the eastern Arctic Shelf of Alaska, most of the Arctic Archipelago and Keewatin Barrens, and practically all of Greenland. Southern Greenland coasts have cool to cold winters and cool summers with moderate precipitation on the southwest coast and heavy precipitation in the southeast. Most of the precipitation in the south falls as snow, with a winter maximum. Most coastal locations on the southeast side of Baffin Island have conditions similar to those of the southwest Greenland littoral. The arctic region extends southward in a broad salient across the Keewatin and eastern Mackenzie District barrens.

In the Great Bear Lake area of Mackenzie District and along the northwest coast of Alaska, winters are somewhat less severe and summers slightly warmer, but precipitation remains scanty. Throughout the Brooks Range of Alaska and in some valleys of southern Yukon Territory in the Whitehorse-Teslin area, precipitation averages less than 10 inches per year, winters are very cold, and summers are generally warm.

A large central continent area, including the shores of Hudson Bay and the Ungava section of northern Quebec, has very cold winters, cool summers, and light precipitation. Precipitation is concentrated in late summer or early autumn. The area receives 10 to 20 inches of precipitation annually, 30 to 50 percent of which falls as snow. The site at Fort Churchill lies within this area and may be regarded as typical.

In most of the southern interior of Canada east of the Rocky Mountains and northwestward through southern Mackenzie District, central Yukon,

interior Alaskan valleys, and the Gulkana Basin, there is a zone of very cold winters, warm summers, and light to moderate precipitation. This is the region in which Fort Greely is located. Annual precipitation, usually with a midsummer maximum, is lightest in Alaska and northwestern Canada and gradually increases southeastward to over 30 inches in central Quebec. Some places in the Yukon and the southern Prairie Provinces receive less than 40 inches of snow annually, but this too increases eastward, until it is over 200 inches per year in parts of Quebec and Newfoundland.

On either side of the Rocky Mountains in British Columbia and Alberta are foothill areas which are somewhat warmer in winter than the wide cross-continent zone immediately to the north and east. Summers in this area are warm and precipitation is light to moderate, depending upon local exposure.

The western mountains themselves have more precipitation, very cold winters, and warm summer days with cool nights. Snowfall increases rapidly with elevation and some stations at high elevations in the Rocky Mountains receive nearly 400 inches annually.

Montana and the Dakotas have cold winters and hot summers, with light precipitation reaching its maximum in early summer. Eastward through the Great Lakes to the St. Lawrence Valley, New England, and the Maritime Provinces, winters are cool or cold, summers are warm, and the precipitation is varied both in time of maximum and intensity. In the Great Lakes vicinity, precipitation is moderate and falls most heavily in spring and autumn. Nearly all of southern Quebec and all except the very southernmost coasts of the Maritimes receive between 30 and 50 inches of precipitation with a maximum in late summer, including at least 100 inches of snow in most areas.

The coasts of New England and the southeastern Maritimes have cool winters, warm summers, frequent fogs, and 40 to 60 inches of precipitation, with a tendency toward a winter or late autumn maximum.

The Pacific Coast littoral of British Columbia, Washington, and Oregon has mild, damp winters, warm, humid summers, and very heavy precipitation falling mostly as rain with a maximum in winter. Certain places on the British Columbian coast receive over 200 inches of precipitation.

Western Alaskan coasts have moderate precipitation, heaviest in autumn, 40 to 60 inches of snowfall, cold to very cold winters and cool summers. The Aleutian Islands have cool winters and summers, frequent fogs and high wind, and heavy precipitation. East of Kodiak, the Alaskan coast has heavy precipitation and cool winters, but summers are warmer here. Very heavy snowfall occurs in the Alaskan Panhandle, with some locations receiving over 400 inches per year. This is also generally true of the British Columbian coastal ranges and the Cascade Mountains of Washington and Oregon. East of the Cascades, the interior valleys have cool winters, hot, dry summers, and scanty precipitation.

Most of Iceland has moderate precipitation, usually in the form of rain or sleet, falling mostly in winter. Only the southern littoral zone receives heavy precipitation. Cool winters and cool summers prevail over all but the highest, glaciated parts of the island.

Moderate to heavy precipitation falls in the British Isles, mostly as rain, reaching its greatest intensity in late autumn and winter but with no dry season at any time of year. Like the coast of British Columbia, the British Isles have mild winters and warm summers.

4. Analysis of climatic elements

The maps included in this study cover certain basic climatic elements considered essential to an understanding of the environment of the area: elements which reflect stress on men and materiel in the Arctic and Subarctic.

Close analogy of any climatic element to that of the test sites is shown on the maps by means of station symbols and colored overprints, blue for Fort Churchill and yellow for Fort Greely. Overlap of these colors in areas analogous to both sites produces a green tone on the maps. Degree of analogy for each observation station is indicated by station symbols, which are defined on the map legends. Numerical values of each element are given beside the appropriate station symbol.

Figure 1 shows the weather stations from which data were used in the various map analyses. Figure 2 is a reference map of major physiographic regions, with mountains and icecaps differentiated in color.

On the individual climatic maps, only those stations actually used in a particular analysis are shown. They vary in density of distribution from map to map, depending upon the availability of data.

The relative reliability of the position of isopleths is indicated as follows: solid lines indicate fair to good reliability and dashed lines, poor reliability. Lines indicating poor reliability appear in mountainous areas, the uninhabited Arctic, and other remote places for which adequate meteorological data are not available. The complex influence of elevation is evidenced only in a general way where such effect is actually reflected by station data. No attempt was made to differentiate reliability of lines on the composite maps, Figures 14, 15, and 16.

Additional comments concerning determination of analogous ranges and details of evaluation are given in the following discussion of separate climatic elements. The associated map figure is indicated beside element subtitle.

a. Mean Temperature, Coldest Month (Fig. 3)

Fort Churchill	-19°F (January)
Fort Greely	-5°F (January)

Standard deviation was considered in determining the analogous ranges of mean monthly temperature. A range of mean temperatures, within which the average temperature of a single January could be expected to occur in 50 percent of the years, was regarded as a reasonable basis for delimiting close analogy.

Fort Churchill has a standard deviation of 7 F° from the mean temperature of its coldest month. The 50 percent expectancy level falls within 14.9 F° of the long-term monthly mean. Fort Greely, with a standard deviation of approximately 9 F°, establishes 50 percent expectancy within 16.3 F° of the long-term mean. The average range then, between the two sites, is 15.6 F°. Previous analog studies have used a range of ± 5 F° for temperature. In this case precedent agreed so closely with standard deviation determinations that the more convenient 5 F° increment was adopted. Thus, a range of ± 5 F° from mean was considered "closely analogous". The next 5 F° above and below this range were regarded as "semianalogous". This range was used for all temperature maps.

For stations with means at either limit of analogy, but with standard deviations differing from those of the test sites, the probability of their means for a given year being within the 10-degree range would decrease at stations with higher standard deviations and increase at stations with lower standard deviations.

Areas analogous to the two test sites appear separate on the map because the difference between the mean temperatures at the sites exceeds 10 F degrees. The areas form broad, undulating belts dipping far south over midcontinent, and almost encircling Greenland. Fort Greely analogy reaches as far as northernmost United States in the Montana Rockies, northern North Dakota, and Minnesota. The cold nucleus of north-central Canada and the arctic islands is much colder than either test site.

The coldest month is January in most areas throughout interior Alaska, Canada, and the United States. December is coldest in a narrow area along the northwest flanks of the Alaska Range. February is generally the coldest month in Greenland and the Arctic Archipelago, except on Melville Sound,

between Baffin Bay and the Beaufort Sea. In many maritime locations seasons are delayed, February being the coldest month, August the warmest. The Canadian Maritime Provinces, the Great Lakes region, and most of the coast of Labrador are also coldest in February. In certain locations in the Bering Strait, the north side of the Aleutian Peninsula, and the southeast coast of Iceland, the coldest period occurs as late as March. The coldest month is also February in parts of the coastal area of the British Isles and western North America.

b. Mean Daily Minimum Temperature, Coldest Month (Fig. 4)

Fort Churchill	-27°F (January)
Fort Greely	-13°F (January)

With minor variations, the mean daily minimum temperature map for the coldest month shows essentially the same distribution of analogous areas as the mean monthly temperature map. One of these exceptions is the small coastal area in Northern Greenland, which is analogous to Fort Churchill in respect to mean temperature, but has mean daily minimum temperatures too cold for analogy.

c. Absolute Minimum Temperature (Fig. 5)

Fort Churchill	-57°F (January)
Fort Greely	-65°F (January)

The area of close analogy with Fort Churchill extends from the Beaufort Sea to Wyoming through 30 degrees of latitude. Fort Greely analogy occurs in interior locations encircling the coldest spots of North America. These cold spots, with temperatures too low for close analogy to Fort Greely, are found in central Alaska and the Yukon to the Mackenzie Valley, the Peace River Valley of British Columbia and Alberta, interior Greenland and Ellesmere Island, and in localized spots around Copper Center, Alaska; Prince Albert, Saskatchewan; Rogers Pass, Montana; and Iroquois Falls, Ontario. The lowest temperature on record for North America, exclusive of Greenland, is -81°F at Snag, Yukon. The Greenland icecap stations, North Ice and Central Station, have each recorded -87°F during different 2-year periods.

d. Mean Windchill, Coldest Month (Fig. 6)

Fort Churchill	1820 Kg Cals/m ² /hr (January)
Fort Greely	1630 " " " " (January)

Windchill is, in a sense, a measure of human discomfort, since it is the cooling effect of moving air on the human body. It is based upon an empirical formula determined by Siple and Passel (Antarctica in 1940) (53)

from experiments with the freezing rate of water in a plastic cylinder. The formula is used to compute kilogram calories of heat loss per square meter of exposed skin surface per hour for each Centigrade degree of temperature difference between the skin and the air. It is written: $H = (\sqrt{100V + 10.45} - V) (33 - T)$, where H = Windchill in Kg Cal/m²/hr, V = wind velocity in meters per second, and T = ambient temperature in degrees Centigrade.

This formula is best applied to simultaneous wind and temperature measurements from hourly records, but such records have not been available in sufficient numbers to map. For this reason monthly mean values of temperature and windspeed were used in this study. Calculations by Arnold Court (20) indicate that the use of such mean values gives results reasonably comparable with those using hourly values. Because of the scarcity of good summarized wind data for northern Canada, however, most of the isopleths on Figure 6 are classed as having poor reliability.

The mean windspeed of the coldest month at Fort Greely is 17 miles per hour. A range from 15 to 19 miles per hour was considered closely analogous. For Fort Churchill, with a mean windspeed of 15 miles per hour, for the coldest month, a range from 13 to 17 miles per hour was used. These ranges were plotted on a windchill nomogram (inset on Fig. 6) together with the corresponding mean temperature ranges (Fig. 3) to derive the range of windchill analogy. Close analogy for windchill then falls within a range of plus or minus 130 Kilogram Calories from the mean windchill at the test sites.

It will be noted that the windchill map shows a predominance of isopleths which show poor reliability. This is due to the difficulty in obtaining satisfactory wind data in suitable quantities for Canadian stations.

Fort Greely analogy extends southward to the south end of Lake Winnipeg in Manitoba. The southern limit of Fort Churchill analogy agrees roughly with the northern tree line across the mainland.

e. Mean Cloudiness, Coldest Month (Fig. 7)

Fort Churchill	4.7 tenths of sky cover (January)
Fort Greely	5.5 tenths of sky cover (January)

During the coldest months, cloudiness is greatest in the maritime areas, particularly over the Pacific coast and the Gulf of St. Lawrence. Inland, cloudiness is less, especially in the north-central area of greatest cold. Widespread areas are analogous to Fort Greely, smaller areas are analogous to less cloudy Fort Churchill. Except for a small area in eastern Maine, the northern United States is too cloudy to be analogous to Fort Churchill.

Greenland presents a different picture. Inland from the coast, cloudiness decreases rather sharply to less than Fort Churchill analogy. Still farther inland, according to the brief records available, cloudiness increases over the high interior of the icecap to values great enough for close analogy with Fort Greely.

f. Mean Snow Depth, Month of Greatest Depth (Fig. 8)

Fort Churchill	25.1 inches (March)
Fort Greely	17.0 inches (February)

Snow depth is so dependent upon a number of other climatic elements that its distribution is highly varied. Mean depths for the month of maximum accumulation range from none along parts of the Oregon-Washington coast to more than 20 feet in the Cascade Mountains a hundred miles to the east. Except for high elevations, only two areas record notably deep snow cover exceeding the limit of Fort Churchill analogy (i.e., more than 30 inches). One is located in central Ontario and the other includes all of Labrador, most of Quebec north of the 50th parallel, and the mountains of eastern Baffin Land.

Fort Churchill analogy occurs in most mountainous regions, except at the highest elevations, and along the southern coast of Greenland. It also appears in the widespread areas over the interior of the continent. One of these reaches from interior Alaska to Great Bear and Great Slave Lakes. The other area stretches from the inland side of Great Slave Lake south and east to the Appalachians and the Gulf of St. Lawrence, skirting Lake Huron and encompassing Hudson Bay.

As a rule, mean snow depths grow shallower on all sides of areas which are analogous to Fort Churchill. These peripheral zones are analogous to Fort Greely.

In Greenland, precipitation patterns over the interior are little known. It might be presumed that snowfall would be relatively light on the interior icecap because of persistent low temperatures. Seasonal snow accumulations in this case should be correspondingly small, as indicated on the map. However, evidence from snow strata measurements made in the central icecap (26) indicates that snowfall there may be greater than shown on Figure 8, which is based upon standard observations. If this is true, the actual area of snow depth analogy on central Greenland may resemble the pattern of cloudiness analogy shown on Figure 7 (i.e., additional areas of analogy on the high interior of the icecap). This information would also modify Figures 9 and 13.

g. Mean Snowfall, Month of Greatest Snowfall (Fig. 9)

Fort Churchill	10.3 inches (November)
Fort Greely	7.0 inches (January)

Fort Churchill and Fort Greely have only moderate mean snowfall even during the snowiest months. Analogous conditions prevail in most of the lowland and high prairie country of western Canada and eastern Alaska, as well as littorals in most of Greenland, western Alaska, and British Columbia. On the other hand, most of eastern Canada, the Rocky Mountains, and the western interior of Alaska have substantially heavier snowfall than either Fort Greely or Fort Churchill.

Much of north-central Canada has relatively light snowfall and is analogous to Fort Greely but not to the slightly snowier Fort Churchill. Relatively light snowfall is also characteristic of parts of the semi-arid western Great Plains and intermontane plateaus of the United States and Canada.

It is difficult to generalize about Greenland because of the scarcity of data. (See closing paragraph, subsection "f." above)

h. Mean Temperature, Warmest Month (Fig. 10)

Fort Churchill	54°F (July)
Fort Greely	59°F (July)

Mean temperatures for the warmest month range from over 75°F in northern United States to below 10°F in central Greenland. The warmest month is generally July in interior and far northern areas, and August near the coasts south of 60° North latitude. Also, a broad salient extending west of Hudson Bay to Great Slave Lake is warmest in August.

Mean warmest month temperatures analogous to those of Fort Greely occur in a broad belt across Alaska and Canada and in the British Isles, mostly north of the 45th parallel. The area of Fort Churchill analogy lies north of the 50th parallel except at high elevations and in limited coastal areas, where it extends farther south. It is limited on the north by cold coasts from the Bering Strait in Alaska to the Hudson Strait in Quebec and Labrador. None of the Arctic Archipelago is warm enough for analogy.

Most of Iceland and the bords of fiords in southwest Greenland are analogous to Fort Churchill alone. The icecaps and north coasts of the two islands are too cold for analogy.

1. Mean Daily Maximum Temperature, Warmest Month (Fig. 11)

Fort Churchill	64°F (July)
Fort Greely	69°F (July)

This map presents essentially the same picture as the mean warmest month temperature map (Fig. 10), except that the southern limits do not extend as far south. The most striking departure from Figure 10 is the presence of an area too warm for analogy to either test site. This area extends from the Central Lowlands of the United States to the Mackenzie Valley on the Northwest Territories.

In contrast to North America, a greater area of the British Isles is analogous to Fort Churchill in mean daily maximum than in mean temperature during the warmest month.

J. Absolute Maximum Temperature

Fort Churchill	96°F (July)
Fort Greely	90°F (May)

No map has been prepared for absolute maxima because these temperatures are not considered to be critical in evaluating arctic and subarctic environments. Table VI contains selected data for this element.

Northwest and central parts of the Arctic Archipelago and the south-east coast of Baffin Island report maximum temperatures below 60°F. Interior Alaska near the Yukon border, on the other hand, has registered 100°F, and temperatures above 120°F have been recorded in central North Dakota.

A 10-degree range of close analogy for Fort Churchill, 91° to 101°F, includes most of the Canadian Maritime Provinces and New England, most of Quebec, Ontario, Manitoba, northern Saskatchewan, Yukon Territory, central Alaska, and British Columbia. The interior plateaus of British Columbia have absolute maxima too hot for Fort Churchill analogy.

Fort Greely analogy, 85° to 95°F, overlaps the Fort Churchill range and extends farther coastward everywhere and up into the mountains in the higher ranges of Alaska and the Yukon Territory.

Records for the interior of the Greenland icecap indicate that maximum temperatures seldom, if ever, rise above the freezing point, but in coastal Greenland they range between 50° and 75°F even on the northern littoral.

K. Mean Cloudiness, Warmest Month (Fig. 12)

Fort Churchill	6.4 tenths of sky cover (July)
Fort Greely	6.4 tenths of sky cover (July)

This map is unique in the series in that the cloudiness values and ranges of analogy for both sites are identical; therefore, analogy is shown by the green tint indicating 100 percent overlap. Cloudiest skies in summer are found over the Aleutian Islands, the Arctic Archipelago westward along the cold coasts to the Bering Sea, the coasts north and east of the Gulf of Alaska, central Quebec and Labrador, and interior high valleys of central British Columbia. Least sky cover occurs over interior United States, the southern part of the Prairie Provinces, and interior lowlands from Alberta to the Yukon Valley in Alaska.

The area of analogy stretches, with few exceptions, from coast to coast and north-south from the Arctic Ocean to northern United States.

The general cloudiness pattern in Greenland for the coldest month, as shown in Figure 7, is repeated in the warmest month. Cloud cover decreases rapidly inland from analogy on the coasts to below analogy and increases again to analogous and greater semi-analogous amounts over the central plateau. The central plateau of Greenland and the west coast of Iceland fall within the analogous range, as do all of the British Isles except the cloudier parts of northern Scotland and western Ireland.

1. Mean Annual Precipitation (Fig. 13)

Fort Churchill	14.4 inches
Fort Greely	11.4 inches

The lowlands of western Canada and central Alaska, parts of the Rockies, the Hudson Bay - Baffin Island areas, and interior Iceland are characterized by rather low precipitation such as occurs in Fort Churchill and Fort Greely. Southeast Canada, the Pacific coast and adjacent ranges, and the south coast of Greenland are much wetter than the two referenced stations.

At the other extreme are the northern parts of Canada, Alaska, and Greenland, with much less precipitation than Fort Greely and Fort Churchill.

Areas analogous only to Fort Churchill occupy the wetter margins of analogy on the map, while those analogous only to Fort Greely occupy the drier margins.

Precipitation in North America is commonly associated with cyclonic activity. Maximum precipitation falls in summer throughout the northern and central regions, from interior Alaska to interior Greenland, including both Fort Churchill and Fort Greely. The maximum occurs in autumn (usually October) along the Gulf of Alaska, Baffin Bay, the St. Lawrence Valley, and east shore of Hudson Bay. In Iceland, the British Isles, and southern and northeastern Greenland, the precipitation maximum occurs a little later in the season, usually November. Winter precipitation maxima prevail in the northwest states, British Columbia, and the Canadian Maritime provinces, but in the east, maxima occur about a month later than in the west.

There is a spring rainfall maximum in a narrow belt of the Rocky Mountain foothills in Alberta. The Great Lakes Region has two peaks of precipitation, the first occurring in May or June and the second in September. Inland New England and the intermontane plateaus of western Canada have a relatively even distribution of precipitation throughout the year.

Data were unavailable for the interior of Greenland (see closing paragraph subsection "f." above).

m. Humidity

Maps of mean dewpoint are not included in this series since dewpoint temperatures, particularly during winter, are generally not much lower than dry bulb temperatures throughout most of the study area. Table XIII shows mean relative humidity at selected stations. No data were available for Fort Greely. Fairbanks, 100 miles west-northwest, has been substituted to provide an approximation of the conditions at Fort Greely.

In winter, mean dewpoint temperatures in most places in the Arctic and Subarctic are about 2 to 5°F below the mean dry-bulb temperatures. The difference is only slightly greater in the prairies of northern United States. During the warmest month, dewpoints average about 10°F lower than mean temperatures in continental locations and only about 2 to 5°F lower in coastal areas. Many arctic and subarctic areas have low absolute humidity, but very high relative humidity during the cold months (e.g., Fort Churchill and Aklavik). Certain marine stations in the northwest have lower relative humidity in winter than in summer (e.g., Barrow and Nome). However, in lower latitudes there is a tendency for relative humidity to be higher in winter than in summer (e.g., Vancouver, B.C. and Gander, Newfoundland). In extremely cold areas of the Arctic it is not uncommon for relative humidity to exceed 100 percent, with resulting condensation producing a diffuse ice fog.

n. Mean Number of Days with Fog (Fig. 13A)

Fort Churchill	13 days per annum
Fort Greely	8 days per annum

Figure 13A and Table XIV show mean frequency of days with fog wherein visibility is restricted to about one mile or less. Data for only dense fog (visibility 1/4 mile or less) were available for most stations in the United States and certain interior stations in Alaska. A comparison with nearby stations in similar situations indicates that the total frequency of occurrence of fog for these Alaskan stations would not be appreciably different from that for dense fog alone, so the data are undifferentiated on the map. In the United States, comparison of data did not indicate a similar relationship, and fog was therefore not mapped for the United States.

Radiation fogs resulting from radiative cooling near the ground are the most widespread type in the study area. Such fogs occur most frequently at temperatures of -40°F or colder, in the form of ice fogs. This phenomenon is rather common during cold months in the Arctic. Ice fog and accompanying smoke fog from motor exhaust often obscure the immediate

vicinity of landing fields for several hours following a few successive aircraft landings or take-offs. Ice fog for the month of maximum occurrence (February) averages 108 hours at Fort Churchill while at Fort Greely the maximum monthly average is about 57 hours in January.

Summer is the season of greatest fog frequency along most arctic and subarctic coasts from Alaska to Iceland and as far south as the Queen Charlotte Islands on the Pacific Coast and Nova Scotia on the Atlantic Coast. However, along the poleward shores of the Arctic Archipelago from a point east of Herschel Island, Yukon, to northern Greenland, autumn is the season with the greatest mean number of days with fog. Summer is the foggiest season along the eastern flanks of the Alberta Rockies, in the Great Lakes Region, and on the southern coasts of the British Isles.

Throughout a broad inland area from the Tanana Valley and Brooks Range of Alaska south to Oregon and southeast along the margin of the Canadian Shield to the St. Lawrence Valley, fog is most frequent in autumn. Autumn is also foggiest on the icecap in southern Greenland and the highlands of Iceland. Winter is the foggiest season in most of the central Alaskan and Yukon valleys and plateau country, high plains and plateaus of British Columbia, the Prairie Provinces, northern Plains States, the upper St. Lawrence and Ottawa River basins, central Greenland, and most of the British Isles.

Localities with spring fog maxima are less extensive and less numerous. Notable among these smaller areas are the Foxe basin, northern Ontario, central Quebec, southern shores of the Gulf of St. Lawrence, and parts of the north and east coasts of Greenland.

The map shows areas having very high and very low fog frequency. Those with very high fog occurrence (60 days or more per year) are likely to fall in one of three classifications. The most widespread of these is the high elevation type where the normal cloud cover is frequently low enough to obscure the sun. Highest parts of the western ranges, the Baffin highlands, and high icecaps of Ellesmere Island, Greenland, and Iceland belong to this category.

The second type occurs mainly in scattered coastal situations where summer sea fogs predominate. Notable examples of this type are found in northern and western Alaska, east Greenland, and parts of the Canadian Maritime Provinces.

The third type of locality having frequent fog is found in the vicinity of major settlements in very cold areas where combustion of large quantities of fuel in winter produces numerous condensation nuclei. This condition accounts for relatively frequent ice fogs around a townsite, although a few miles away a very low fog frequency may prevail.

Areas of very low annual fog incidence are found at relatively low elevations in interior locations. The interior valleys of Alaska, the valleys, plains, and low shield country of central Canada, heads of fiords in Greenland, and north and west coasts of Scotland and Ireland have less than 10 days of fog per year. Fort Greely, with 8 days, falls within this frequency range and Fort Churchill, with 13 days, is only slightly foggier.

5. Analysis of composite maps

Figures 14 through 16 synthesize several of the foregoing maps to show analogous areas of coincidence among multiple climatic elements. Temperatures were chosen to serve for the basic zones of analogy. These zones were called: (a) "winter temperature analogy", the areal coincidence of mean and mean daily minimum temperatures for the coldest month, and (b) "summer temperature analogy", the areal coincidence of mean and mean daily maximum temperatures for the warmest month. Over these basic analogous zones were superimposed analogous areas of additional stress elements (i.e., snow depth, windchill, and precipitation) in various combinations. Certain lines were generalized.

a. Composite Analogous Areas, Coldest Month (Fig. 14)

Because the ranges of coldest month temperature analogy for the two test sites do not overlap, it was possible to draw the coldest month composite for both sites on the same map. Areas of analogous snow depth and windchill were placed over the winter temperature analogy for both Fort Greely and Fort Churchill. Yellow indicates Fort Greely and blue indicates Fort Churchill temperature analogy, with black line patterns designating snow depth and windchill analogy for both sites.

Zones that are completely analogous (with respect to temperature, snow depth, and windchill) to the respective test sites are small and widely dispersed. For Fort Churchill, the most extensive area is in the vicinity of the site itself, on both sides of Hudson Bay. Other patches at the entrance to Foxe Basin, east of Coppermine, and on the southern Greenland icecap, are less reliably defined, since they do not include stations for corroboration.

Complete analogy for Fort Greely is narrowly confined to the immediate vicinity of Fort Greely, northwestern Alaska, the south end of Lake Winnipeg, and scattered patches bordering the icecap of southern Greenland. The combination of temperature and snow depth analogy is most widespread in the western half of the continent; coincidence of temperature and windchill analogies is more common in the east.

b. Composite Analogous Areas, Fort Greely (Fig. 15)

Seasonal and year-round analogies are shown in Figure 15. Areas of analogous winter and summer temperatures are represented in blue and yellow diagonal lines respectively. Overlaps of the two give the appearance of a green pattern. Areas of analogous mean annual precipitation are shown by black stipple pattern, only where analogous temperatures occur.

Combined analogy (winter and summer temperatures and precipitation) for Fort Greely occurs in three areas: (1) adjacent to Fort Greely along the north flank of the Alaska Range and southward through Broad Pass into the Gulkana Basin, (2) a small patch straddling the lower Yukon River south of its confluence with the Koyukuk, and (3) the largest of the three areas, southern Yukon Territory.

The most widespread area of temperature-precipitation analogy occurs in summer. It includes central Alaska and most of the Yukon Territory stretching southeastward in a broad band through the larger Canadian lakes to Hudson Bay and northern Ontario. Other isolated spots are found primarily in high valleys from Alaska's Gulkana Basin to the state of Wyoming.

Combined winter and summer temperature analogy appears in parts of lowland Alaska, plateau valleys of northern British Columbia and southern Yukon, scattered Canadian prairie locations near lakes, and in a sizable portion of the Laurentian Upland stretching from central Ontario to southern Labrador.

Analogous areas of precipitation in combination with winter temperatures occur in widely-dispersed, small spots from Alaska to the Greenland icecap.

c. Composite Analogous Areas, Fort Churchill (Fig. 16)

This map was prepared in the same manner as Figure 15. Combined winter and summer temperature and precipitation analogy for Fort Churchill occurs in only three locations. The principal area extends east from northern Saskatchewan to include the west shore of Hudson Bay from James Bay nearly to Chesterfield Inlet. The other locations (but with no verifying stations) are the central Ungava Peninsula of northern Quebec, and the lowland just west of Great Bear Lake.

Summer temperatures and precipitation are jointly analogous in the interior plateau valleys of northern British Columbia and southern Yukon, the western lowlands of Alaska, and other scattered small areas.

Analogy of combined winter temperatures and precipitation occurs in a narrow strip from central Yukon to northern Manitoba and along the northeast shores of Hudson Bay and southeastern Baffin Island.

Combined summer and winter temperatures are analogous in parts of the Brooks Range of Alaska, in a broad swath stretching from the Mackenzie Delta along the eastern shores of the larger lakes of the Shield margin to northern Saskatchewan and in a large part of the Laurentian Upland of north-central Quebec.

6. Tables of monthly climatic values for selected stations

Table I is a list of stations with elevations and map coordinates for which climatic data are presented in the subsequent tables.

Tables II through XIII show mean values of climatic elements for all months for forty of the stations selected from Table I to illustrate comparative climatic regimes among representative parts of the study area. These elements are the same as used in the map series with the addition of Table VI, Absolute Maximum Temperature and Table XIII, Mean Relative Humidity.

Table XIV is a frequency table showing Mean Number of Days with fog. Some stations (as indicated) show only dense fog, although most figures represent fogs of all densities affording less than about 1 mile horizontal surface visibility.

While the maps were prepared on a seasonal basis for warmest or coldest months (except Fig. 13 and Fig. 17) the tables provide information for making year-round regional comparisons.

7. Limitations of methods

The following limiting factors may affect the reliability of the accompanying maps.

(a) Many parts of the study area have few or no climatic stations. There are practically no representative data for most zones of high elevation, such as the Alaskan and western Cordilleran ranges and the ice-capped highlands of Ellesmere Island, Baffin Island, and Iceland.

(b) Many far northern stations have very short periods of record, many obtained 20 to 50 years ago. It was not possible to obtain data from comparable periods of record in every case.

(c) The map analyses necessitated broad comparisons between large, diverse areas and two localized climatic stations.

(d) Times and frequencies of daily meteorological observations vary from one locality or political unit to another.

(e) Station exposure and elevation are not always representative of very large surrounding areas for which inferences had to be made.

Table I: LOCATION AND ELEVATION OF STATIONS

Country (Area)	Climatic Station	Elevation (feet)	Lat. (N)		Long. (W)	
			(°)	(')	(°)	(')
<u>Alaska</u>	Barrow	22	71	18	156	47
	Bethel	10	60	47	161	43
	Dutch Harbor	72	53	53	166	32
	Fairbanks	440	64	51	147	43
	FORT GREELY	1,274	64	00	145	44
	Juneau	72	58	18	134	24
	Kodiak	21	57	45	152	31
	Nome	13	64	30	165	26
<u>British Isles</u>	London (Kew), England	18	51	28	0	19
	Stornoway, Hebrides	79	58	11	6	21
<u>Canada</u> (Western Provinces)	Calgary, Alta.	3,540	51	02	114	01
	Embarras, Alta.	775	58	12	111	23
	FORT CHURCHILL, MAN.	115	58	47	94	17
	Fort Vermilion, Alta.	950	58	23	116	03
	Keg River, Alta.	1,402	57	47	117	50
	Prince George, B. C.	2,218	53	54	122	40
	The Pas, Man.	890	53	49	101	15
	Vancouver, B. C.	45	49	17	123	05
	(Eastern Provinces)					
	Charlottetown, P. E. I.	186	46	14	63	07
	Gander, Nfld.	582	48	57	54	34
	Goose Bay (Lab.), Nfld.	144	53	20	60	25
	Kapuskasing, Ont.	752	49	25	82	28
	Lake Doré (Ombougamau), Que.	1,234	49	54	74	18
<u>(Yukon & N.W. Territories)</u>	Moose Factory, Ont.	29	51	14	80	31
	Moosonee, Ont.	34	51	16	80	34
	Nitchequon, Que.	1,690	53	12	70	
	Port Harrison, Que.	66	58	27	78	08
	Quebec, Que.	296	46	48	71	13
	Sioux Lookout, Ont.	1,227	50	07	91	54
	Aklavik, Mackenzie	30	68	14	135	00
	Alert, Franklin	205	82	30	62	20
	Arctic Bay, Franklin	36	73	00	85	18
	Baker Lake, Keewatin	30	64	18	96	05
	Coppermine, Mackenzie	13	67	47	115	15
	Dawson, Yukon	362	64	04	139	29
	Fort Simpson, Mackenzie	415	61	52	121	21
	Mould Bay, Franklin	50	76	17	119	28
	Resolution Island, Franklin	127	61	18	64	53
<u>Greenland</u>	Jakobshavn	102	69	13	51	02
	Mygghukta	10	73	29	21	34
	Nanortalik	23	60	10	45	17
	Narsarsuaq	88	61	11	45	25
	Scoresbysund	56	70	29	21	58
	Søndre Strømfjord	187	67	00	50	48
<u>Iceland</u>	Thule	121	76	33	68	49
	Reykjavik	131	63	49	22	43
	U. S. A.					
	Devils Lake, N. D.	1,471	48	07	98	52
	Helena, Mont.	3,893	46	36	112	00
	Sault Ste. Marie, Mich.	721	46	28	84	22

TABLE II. MEAN TEMPERATURE (°F)*

Station	Ice Rec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
<u>Alaska</u>														
Barrow	30	-15	-18	-15	0	19	34	40	38	31	17	1	-10	10
Bethel	30	7	9	12	27	40	53	55	53	45	32	17	7	30
Dutch Harbor	25	32	22	34	36	41	46	51	51	49	42	36	33	40
<u>FORT GREENGLASS</u>														
Juneau	9	-5	2	12	28	46	56	59	54	44	27	5	-4	27
Kodiak	45	28	30	34	41	48	54	52	55	50	43	31	42	42
Koon	38	30	32	33	37	43	49	54	55	50	42	35	31	41
Nome	74	3	6	9	20	34	46	50	49	42	29	16	8	26
<u>British Isles</u>														
London	41	41	41	43	47	55	58	61	62	58	51	44	41	50
Stornoway	52	40	40	41	44	48	52	55	55	52	47	43	40	46
<u>Canada (West)</u>														
Calgary	59	13	17	26	40	50	56	62	60	51	42	28	19	38
<u>FORT CHURCHILL</u>														
Fort Vermilion	30	-19	-17	-6	14	30	43	51	52	42	27	6	-11	18
Prince George	30	-2	-4	11	32	49	57	62	58	43	34	11	-5	29
The Pas	27	-13	18	30	40	49	56	60	59	50	41	29	16	38
Timmins	39	-2	-2	11	33	48	59	65	61	49	35	17	1	31
Vancouver	61	36	39	43	48	54	60	64	63	57	50	43	39	49
<u>Charlottetown (East)</u>														
Charlottetown	65	18	17	26	36	43	58	62	65	58	48	36	25	42
Gender	14	19	18	24	33	44	52	62	61	54	44	34	25	39
Goose Bay	10	0	4	16	28	40	51	61	58	50	38	24	9	30
Moose Factory	40	-6	-3	10	24	41	54	61	59	50	38	21	4	30
Hitchewon	10	-11	-6	6	21	36	49	55	55	45	34	17	-1	25
Port Harrison	26	-11	-16	-6	11	28	39	47	47	41	31	17	-1	19
Quebec	72	10	12	23	37	51	62	67	64	56	44	30	16	39
St. John's	23	-3	2	15	32	48	59	65	62	52	38	21	3	33
<u>Alaskan (Territories)</u>														
Alert	25	-18	-17	-9	9	31	49	56	50	38	20	-3	-17	16
Arctic Bay	6	-28	-29	-27	-8	11	32	39	33	16	-3	-15	-23	-1
Baker Lake	13	-21	-26	-16	-4	19	36	43	41	29	14	-4	-16	8
Copernicus	6	-30	-32	-15	-1	19	34	48	49	38	20	-2	-16	9
Denton	30	-19	-20	-14	1	22	39	49	47	36	19	4	-16	12
Fort Simpson	53	-18	-11	5	29	46	57	60	55	43	26	1	-13	23
Waula Bay	42	-18	-11	1	26	45	56	62	57	46	29	4	-12	24
Waula Bay	8	-30	-33	-25	1	31	43	48	46	34	20	-2	-15	1
<u>Resolution Island</u>														
Resolution Island	22	0	-1	7	15	27	40	46	37	38	29	21	8	21
<u>Jakobshavn</u>														
Jakobshavn	52	0	-3	3	15	31	40	46	44	35	25	17	9	22
<u>Greenland</u>														
Naortalik	42	23	24	26	31	37	41	43	42	35	35	29	25	33
Scorebyund	12	4	5	14	27	37	42	42	39	34	20	14	9	21
Thule	6	-11	-15	-12	-2	22	36	41	38	27	14	5	-10	11
<u>Iceland</u>														
Reykjavik	40	30	30	31	36	43	49	52	50	46	39	34	30	39
<u>U.S.A.</u>														
Devils Lake	47	3	7	22	39	52	62	68	66	56	43	25	10	38
<u>U.S.A.</u>														
Malena	73	25	24	32	44	52	60	68	66	56	46	33	25	44
Sault Ste. Marie	64	15	13	23	38	49	59	64	63	56	45	32	21	40

*Values for coldest and warmest months are underlined.

TABLE III. MEAN DAILY MINIMUM TEMPERATURE (°F)*

Station	Rec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Barrow	25-28	-23	-24	-22	-8	14	29	34	33	27	12	-6	-18	4
Bethel	17-16	-1	0	3	18	31	35	46	45	36	24	11	0	21
Dutch Harbor	22-24	23	27	28	31	35	39	45	46	43	37	32	29	35
FORT GREENLY	9	-13	-7	1	18	36	49	44	44	36	20	-2	-13	18
Juneau	43-47	24	26	29	34	40	46	50	49	45	39	31	27	37
Kodiak	20-22	26	27	28	32	37	44	47	48	44	38	31	26	36
Kona	34-35	-4	-2	1	12	28	38	44	43	36	24	10	1	20
London	34	36	32	37	40	46	51	55	54	50	44	39	36	41
Stordomy	22	36	35	35	42	47	49	50	47	42	39	32	37	41
Calgary	59	2	6	14	27	36	43	47	45	37	29	17	9	26
FORT CHURCHILL	30	-27	-25	-16	4	22	34	43	43	34	20	-2	-19	9
Fort Vermilion	30	-19	-16	-3	19	36	44	49	45	35	24	3	-14	17
Prince George	27	3	6	18	27	34	42	44	43	36	20	21	-8	26
The Pas	39	-18	-13	-2	21	26	48	54	50	38	26	8	-8	20
Vancouver	11	32	33	36	40	46	50	54	53	49	44	38	35	43
Charlottetown	65	10	9	19	30	39	50	58	51	41	31	31	18	34
Gander	14	12	11	17	26	35	43	52	52	46	37	28	19	32
Goose Bay	10	-8	-5	6	19	32	42	52	49	42	31	18	2	23
Moose Factory	40	-18	-15	-4	15	30	42	49	48	40	30	14	-5	19
Nitchequon	10	-31	-18	-6	10	27	40	48	47	39	28	11	-9	16
Port Harrison	26	-22	-24	-14	3	22	32	39	40	36	26	12	-8	12
Quebec	72	2	4	15	29	41	52	57	54	48	37	24	9	31
Sioux Lookout	23	-11	-9	4	23	37	48	54	51	43	32	16	-4	24
Aklavik (Territories)	22	-26	-24	-17	-2	22	40	47	42	32	15	-9	-24	8
Alert	5	-35	-37	-36	-17	7	27	33	29	11	-10	-21	-30	-7
Arctic Bay	13	-28	-31	-25	-14	12	30	36	35	25	9	-10	-22	1
Baker Lake	6	-40	-35	-19	-11	10	27	38	40	30	7	-13	-23	5
Copparmine	30	-26	-28	-22	-4	15	32	42	40	31	14	-12	-23	5
Dawson	53	-26	-19	-6	16	34	45	47	42	33	20	-4	-19	13
Fort Simpson	42	-27	-21	-11	14	33	45	50	46	36	21	-3	-20	14
Mould Bay	7	-37	-32	-32	-15	8	27	34	30	16	-6	-21	-32	-6
Resolution Island	15	-6	-2	2	10	23	30	33	34	32	26	18	4	17
Jacobshavn	50	-7	-10	-5	6	25	35	40	38	30	20	11	4	16
Nanortalik	47	17	15	17	25	33	40	39	35	30	24	17	27	27
Scoresbysund	12	-3	-3	-4	6	20	31	36	33	29	15	7	2	14
Thule	6	-20	-23	-21	-11	16	30	36	33	21	6	-3	-17	3
Reykjavik	11	21	30	30	33	38	44	47	46	40	35	30	29	26
Devils Lake	47	-6	-2	13	29	40	51	56	53	44	32	17	2	27
Helena	73	11	15	23	33	41	48	54	53	44	35	24	16	33
Sault Ste. Marie	64	7	1	14	29	39	48	53	53	47	38	27	14	31

*Values for coldest month are underlined.

TABLE IV. MEAN DAILY MAXIMUM TEMPERATURE (°F).

Station	Yrs Rec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
<u>Barrow</u> <u>Alaska</u>	25-29	-10	-11	-7	8	25	40	47	44	35	22	6	-5	16
<u>Bellevue</u>	16-18	16	17	22	36	50	63	53	61	53	38	24	15	28
<u>Dutch Harbor</u>	22-24	37	36	39	41	46	52	58	59	54	47	41	38	46
<u>FORT GREENLY</u>	9	3	11	23	38	56	66	69	63	52	34	13	4	36
<u>Junesau</u>	44-48	32	35	39	47	55	62	62	62	56	48	40	35	48
<u>Kodiak</u>	20-22	35	37	39	43	48	56	60	60	56	48	40	33	46
<u>Kona</u>	33-35	11	13	17	27	41	53	56	55	48	35	21	14	32
<u>London</u> <u>British Isles</u>	34	45	46	50	54	63	68	71	70	65	58	49	46	57
<u>Stornoway</u>	52	44	44	46	49	54	58	60	60	58	52	48	45	51
<u>Calgary</u> <u>Canada</u> <u>(West)</u>	59	24	28	37	53	63	69	76	74	64	54	38	29	51
<u>FORT CHAMPLAIN</u>	30	-11	-8	4	24	38	52	64	62	49	34	13	-3	26
<u>Fort Vermilion</u>	30	2	8	25	45	62	70	76	72	60	44	20	5	41
<u>Prince George</u>	27	23	31	42	54	64	70	73	74	65	52	38	25	51
<u>The Pas</u>	39	1	10	24	45	60	70	72	72	66	45	25	9	41
<u>Vancouver</u>	41	40	44	50	57	63	69	74	72	66	56	42	27	57
<u>Charlottetown</u> <u>(East)</u>	65	26	26	33	43	56	66	73	72	65	54	42	31	49
<u>Gander</u>	14	26	26	31	40	53	62	72	70	62	51	40	31	47
<u>Goose Bay</u>	10	8	14	25	37	49	61	71	70	67	59	45	31	40
<u>Moose Factory</u>	40	6	10	23	38	52	67	73	73	61	47	29	14	41
<u>Kitchiquon</u>	20	0	5	18	32	44	58	64	62	52	40	23	7	34
<u>Port Harrison</u>	26	-8	-9	3	20	34	46	54	54	46	36	23	6	25
<u>Quebec</u>	72	18	20	31	44	61	72	76	73	64	51	36	22	47
<u>Sioux Lookout</u>	23	5	13	26	43	60	69	76	72	60	45	27	10	42
<u>Alert</u>	22	-10	-9	0	19	40	58	66	58	44	25	3	-8	24
<u>Arctic Bay</u>	5	-20	-22	-20	-1	20	37	45	39	21	3	-8	-16	6
<u>Baker Lake</u>	13	-14	-19	-8	5	27	42	51	47	34	19	1	-9	15
<u>Coppermine</u>	6	-23	-26	-8	6	27	40	57	59	43	26	6	-10	16
<u>Dawson</u>	30	-11	-13	-6	10	29	45	53	53	41	25	3	-9	19
<u>Fort Simpson</u>	53	-12	-4	17	42	59	70	71	68	53	33	7	-7	33
<u>Fort St. John</u>	42	-10	0	13	38	56	68	74	68	56	36	11	-4	34
<u>Ward Bay</u>	7	-23	-27	-17	1	19	36	43	38	25	7	-8	-19	6
<u>Resolute Island</u>	19	5	4	12	20	30	37	42	38	32	25	14	25	25
<u>Jakobshavn</u> <u>Greenland</u>	32	8	6	14	24	38	46	51	50	41	31	22	15	29
<u>Narsarsuaq</u>	47	35	38	36	38	42	47	47	48	44	40	35	34	40
<u>Narsarsuaq</u>	12	12	12	14	22	30	43	45	45	39	25	20	16	27
<u>Scoresbyund</u>	6	-2	-7	3	8	20	33	41	44	34	21	13	-2	18
<u>Thule</u>	11	36	39	39	43	43	54	57	55	49	43	39	38	45
<u>Baytharvik</u> <u>Iceland</u>	47	13	17	32	51	64	73	81	79	68	54	34	20	49
<u>Devils Lake</u> <u>U. S. A.</u>	73	28	32	42	55	63	71	81	80	68	56	42	33	54
<u>Sault Ste. Marie</u>	64	22	22	32	46	59	70	77	72	55	52	38	27	48

Values for warmest months are underlined.

TABLE V. ABSOLUTE MINIMUM TEMPERATURE (°F)•

Station	Yrs Rec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Barrow	35	-53	-56	-52	-42	-18	8	22	20	4	-19	-40	-55	-56
Etah	32	-52	-47	-36	-25	-5	28	30	30	18	-5	-31	-44	-52
Dutch Harbor	17-25	10	2	9	13	20	30	36	35	29	12	16	10	5
PORT GREENLY	9	-65	-60	-39	-37	-1	32	35	-2	9	-15	-42	-62	-65
Juneau	90	-15	-15	-5	13	24	32	35	36	29	13	-1	-21	-21
Kodiak	42-49	-9	-3	2	5	20	32	35	34	26	7	-3	-12	-12
Nome	39	-47	-42	-38	-30	-11	3	24	16	-4	-39	-42	-47	-47
London	65	9	11	17	26	30	37	43	41	31	25	20	11	9
British Isles	52	11	12	13	16	26	31	32	33	32	28	20	12	11
Calgary Canada (West)	59	-48	-49	-34	-14	12	26	31	28	8	-8	-31	-45	-49
PORT CHURCHILL	61	-57	-52	-52	-26	-14	13	22	25	15	-17	-57	-57	-57
Fort Vermilion	32	-78	-66	-52	-38	-20	18	26	22	6	-21	-44	-72	-78
Prince George	30	-58	-52	-31	-12	15	24	29	25	6	-4	-39	-56	-58
The Pas	39	-54	-54	-43	-21	13	25	32	22	16	-11	-13	-51	-54
Vancouver	30	-2	8	15	27	33	36	43	39	30	21	10	8	2
Charlottetown (East)	30	-27	-23	-12	8	22	32	42	40	32	23	5	-17	-27
Gender	14	-13	-16	-14	7	22	28	36	30	31	22	6	-5	-16
Goose Bay	9	-32	-35	-32	-13	10	30	38	32	20	11	-9	-25	-35
Moose Factory	40	-56	-48	-48	-29	0	23	30	29	20	-3	-27	-41	-56
Nitchequon	9	-56	-55	-49	-31	-8	23	33	33	25	1	-25	-32	-56
Port Harrison	29	-51	-27	-49	-26	-5	15	20	28	12	-9	-29	-46	-57
Quebec	31	-34	-28	-20	0	19	34	43	43	30	18	-6	-32	-34
Sioux Lookout	15	-16	-50	-36	-22	14	27	37	32	23	12	-22	-39	-50
Aklavik (Territories)	25	-56	-62	-50	-44	-14	20	30	25	12	-22	-50	-54	-62
Aleric	6	-54	-53	-53	-50	-16	12	22	24	5	-12	-32	-39	-51
Arctic Bay	8-13	-52	-57	-49	-37	-14	11	22	24	9	-12	-42	-41	-57
Baker Lake	6	-57	-58	-53	-33	-9	14	27	28	18	-14	-34	-47	-58
Coppermine	30	-54	-58	-56	-45	-24	5	31	27	7	-28	-42	-49	-58
Dawson	53	-68	-73	-50	-41	9	25	30	17	9	-23	-50	-63	-73
Fort Simpson	52	-66	-69	-52	-37	-9	25	31	21	7	-18	-51	-64	-69
Mould Bay	8	-55	-58	-56	-38	-18	11	25	13	-13	-26	-46	-63	-63
Resolution Island	19	-16	-32	-22	-7	2	16	26	26	14	6	-10	-32	-36
Jakobhavn	58	-45	-46	-42	-35	-10	21	28	24	8	-6	-18	-34	-46
Narsartalik	32	-6	-5	-5	8	20	28	31	28	22	16	8	-1	-6
Scorebysund	12	-37	-32	-12	-8	20	22	28	26	10	-13	-27	-38	-42
Thule	6	-39	-47	-43	-32	-9	18	27	20	4	-20	-28	-36	-47
Reykjavik	11	6	8	8	13	24	33	39	34	27	18	10	13	6
Devils Lake B. S. A.	47	-44	-46	-32	-4	6	29	37	31	15	-1	-21	-37	-46
Eslena	73	-42	-41	-26	-10	22	31	36	29	6	-8	-22	-40	-42
Sault Ste. Marie	64	-32	-37	-27	-13	21	31	36	32	25	15	-12	-24	-37

Lowest value for period of record are underlined.

TABLE VI. ABSOLUTE MAXIMUM TEMPERATURE (°F)*

Station	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Barrow Alaska	33	31	30	42	45	70	78	83	71	43	39	34	78
Bethel	32	49	51	53	59	76	80	73	61	65	60	49	80
Dutch Harbor	22-26	56	60	69	61	66	73	79	80	65	58	56	80
PORT CRELLY	9	46	51	51	62	80	85	82	73	62	50	48	90
Junes	43-47	54	57	61	69	80	85	87	77	66	64	59	89
Kodiak	34-40	53	60	65	62	74	82	85	77	66	60	61	85
Nome	39	46	47	60	66	80	84	80	65	59	50	40	84
London British Isles	65	57	62	68	80	87	88	84	92	83	63	59	94
Stornoway	52	55	55	59	67	75	78	75	77	70	58	56	77
Calgary Canada (West)	59	61	76	75	85	90	95	96	90	85	71	67	97
PORT MURCHILL	61	39	31	41	62	87	96	90	84	65	45	34	96
Port Vermilion	30	52	55	62	76	92	90	94	89	81	61	50	101
Prince George	30	54	58	68	86	95	93	96	92	84	62	50	102
The Pas	39	44	56	59	87	93	93	96	90	80	58	48	100
Vancouver	40	59	61	68	79	83	92	91	86	77	63	60	92
Charlottetown (East)	30	53	53	60	80	83	88	92	88	82	57	51	98
Geander	14	50	53	53	71	78	91	91	84	76	67	53	91
Goose Bay	9	42	46	51	62	89	89	91	84	73	58	53	100
Moose Factory	33	43	44	61	75	95	96	96	91	84	64	50	97
Mitchequon	9	33	34	44	52	79	83	84	79	66	45	41	90
Port Harrison	25	33	41	38	44	74	78	80	73	62	43	35	80
Quebec	30	52	47	64	80	89	94	95	87	82	71	55	96
Sicux Lookout	15	44	48	66	87	95	94	96	89	78	58	48	96
Aklavik (Territories)	25	44	49	45	56	77	85	88	76	55	44	46	93
Alert	6	10	24	9	30	46	57	64	59	33	31	13	64
Arctic Bay	8-12	28	36	27	34	51	63	64	56	44	23	34	75
Baker Lake	6	10	6	25	31	42	63	62	66	43	36	24	82
Coppermine	30	21	34	29	46	78	82	83	72	57	36	37	87
Darscu	53	36	48	52	69	85	95	88	79	68	52	55	98
Fort Simpson	52	53	60	54	71	95	92	93	86	87	54	45	95
Mould Bay	8	10	-5	11	29	35	52	57	38	24	19	9	59
Resolution Island	19	34	34	37	55	67	71	66	58	43	38	35	61
Jakobshavn Greenland	36	59	49	51	55	62	67	67	58	54	54	52	71
Nanortalik	31	52	53	48	59	62	68	66	64	59	57	54	68
Scoresbysund	12	48	41	47	45	49	60	55	62	47	52	40	63
Thule	6	36	36	34	41	46	59	57	49	40	38	34	62
Reykjavik Iceland	11	49	48	51	58	60	64	64	61	56	52	49	69
Devils Lake U. S. A.	47	51	56	85	87	106	103	112	103	89	69	64	112
Salena	73	63	69	73	86	95	102	103	96	84	71	64	103
Sault Ste. Marie	64	48	50	75	83	91	93	98	92	82	75	54	98

*Highest values for period of record are underlined.

TABLE VII. MEAN SURFACE WIND SPEED (mph)*

Station	Yrs	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Barrow	28	10.8	11.4	10.8	11.4	11.6	11.3	11.8	12.6	13.3	14.0	12.5	11.0	11.9
Bethel	12	11.0	12.4	12.4	10.6	9.4	9.6	9.1	9.9	9.7	10.2	10.0	10.8	10.3
Dutch Harbor	21	5.4	8.5	13.0	8.9	10.1	10.3	11.2	10.1	11.6	10.5	8.5	8.1	9.7
FORT GREENLY	5	17.0	17.0	10.0	9.0	10.0	9.0	8.0	10.0	10.0	13.0	15.0	14.0	11.8
Juneau	12	8.7	8.4	9.0	8.7	7.6	7.4	7.1	7.1	7.6	9.3	7.9	8.8	8.1
Kodiak	7	8.7	9.5	9.5	9.4	8.4	6.1	5.9	6.0	7.9	8.6	11.0	9.9	8.1
Kome	25	10.7	10.7	10.3	10.2	8.9	8.9	9.4	10.2	10.7	10.6	11.0	9.9	10.1
London	35	8.2	8.7	8.9	8.6	7.8	7.1	6.6	6.7	6.2	6.7	7.6	8.4	7.6
Stornoway	15	10.4	6.9	5.8	5.2	5.8	5.8	4.6	4.6	6.9	6.9	6.9	6.9	6.5
Calgary	18	6.8	6.8	7.6	3.2	8.4	7.8	6.5	5.9	7.1	7.1	6.9	7.3	7.3
FORT CHURCHILL	19	14.9	14.6	14.2	14.3	13.4	12.2	11.5	13.0	16.3	17.1	16.9	14.9	14.4
Keg River	11	5.7	5.5	6.6	7.3	7.6	6.3	6.2	6.1	6.2	6.7	5.4	4.9	6.2
Prince George	8	4.5	3.8	5.0	4.4	4.3	3.6	3.3	3.2	3.3	4.5	4.7	4.3	4.1
The Pas	25	1.7	6.4	6.9	7.6	8.1	7.7	7.1	6.7	7.1	7.3	6.7	6.4	7.1
Vancouver	21	3.7	4.0	4.4	4.5	4.3	4.0	3.8	3.4	3.4	3.3	3.4	4.1	3.9
Charlottetown (East)	24	11.4	10.8	10.8	10.8	9.6	8.9	8.5	8.6	9.4	10.8	10.9	11.6	10.2
Gander	9	16.9	17.7	17.2	15.5	13.9	14.1	12.7	12.2	13.9	15.1	15.5	16.9	15.1
Goose Bay	4	10.7	9.3	10.2	9.7	9.2	8.8	8.9	8.7	9.6	10.7	10.3	10.9	9.8
Moosee	4-5	7.9	8.6	9.0	9.6	8.9	8.2	7.4	8.2	8.3	8.0	7.9	7.4	8.3
Nitchequon	3	12.5	7.8	7.7	7.5	7.9	6.9	8.6	7.9	9.9	10.4	8.6	9.1	8.4
Port Harrison	8	1.2	12.7	13.8	14.3	14.7	13.2	14.2	13.8	14.9	15.2	17.3	14.4	14.3
Quebec	25	15.0	16.1	15.3	14.4	14.4	13.2	11.6	10.7	11.5	12.4	14.0	13.9	13.5
St. Louis Lookout	10	8.1	8.4	9.2	9.5	9.5	9.2	8.2	8.7	9.7	10.1	10.2	9.1	9.2
Atlatvik (Territories)	6-8	5.6	6.2	7.9	7.4	7.8	7.7	6.6	6.9	6.5	5.3	4.1	3.4	6.3
Alert	1-2	1.2	1.8	2.3	2.5	2.9	4.9	9.0	6.5	7.4	7.2	3.3	4.2	4.7
Arctic Bay	7	3.4	2.8	3.5	3.5	6.0	8.2	8.8	7.6	8.1	6.2	6.1	3.4	5.6
Saier Lake	1-2	15.3	9.3	9.3	12.9	9.1	9.5	11.8	12.6	12.4	14.1	12.7	14.9	11.9
Coppermine	9	8.3	8.1	6.9	6.7	7.2	5.6	8.2	9.3	9.8	10.3	10.0	8.0	8.3
Dawson	4-6	3.9	2.2	3.8	3.1	3.8	3.6	2.9	3.1	3.4	3.8	4.5	3.6	3.5
Fort Simpson	16	5.8	7.2	8.7	6.7	6.0	6.6	7.2	7.7	6.4	7.2	6.9	6.9	6.9
Mould Bay	3	6.6	10.0	9.3	---	---	12.0	11.0	11.0	13.4	9.6	9.1	5.1	---
Resolution Island	16	18.4	18.1	16.8	14.4	14.4	13.0	12.6	12.9	14.8	17.7	18.9	18.2	16.1
Jakobshavn	30	8.1	6.8	6.4	5.4	6.4	6.8	4.7	5.5	6.8	8.6	9.6	9.1	7.2
Nanortalik	41	11.8	11.4	10.9	10.0	7.0	9.5	7.0	6.0	8.5	9.0	10.4	10.4	9.5
Scoresbysund	12	4.7	5.4	4.5	3.9	3.2	2.8	2.8	2.6	3.2	3.9	3.9	4.0	3.8
Thule	6	9.4	10.8	8.8	7.7	8.2	8.1	7.3	8.9	9.6	10.4	9.7	8.9	9.0
Reykjavik	19	14.2	14.2	13.2	11.6	10.6	10.6	9.1	9.1	10.1	11.6	11.6	12.6	11.5
Devils Lake	47	10.0	10.2	10.9	11.4	11.0	9.8	8.6	8.7	9.5	10.1	10.2	9.5	10.0
Belena	73	7.2	7.5	8.3	8.7	8.7	8.4	8.0	7.8	7.6	7.6	7.3	7.2	7.9
Sault Ste Marie	64	9.0	9.1	9.9	9.7	9.0	7.6	7.2	7.1	7.9	8.9	9.9	9.5	8.7

*Values for coldest month are underlined.

TABLE VIII: MEAN WINDCHILL (Kcal/m²/hr.)*

Station	Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Barrow	1945-46	1645	1705	1645	1455	1130	860	590	345	970	1210	1140	1570	1280
Bethel	"	1305	1335	1230	985	760	570	330	575	690	900	1125	1300	930
Dutch Harbor	"	770	835	915	810	740	685	630	575	655	750	805	840	745
Fort Greely	"	1630	1530	1260	930	630	315	455	560	710	1030	1430	1275	1005
Juneau	"	925	890	840	730	610	520	435	500	580	715	790	885	705
Kodiak	"	895	885	870	785	695	550	450	480	585	720	805	905	720
Koon	"	1360	1310	1255	1080	840	695	610	635	755	955	1165	1255	990
London	"	720	730	710	645	510	430	355	400	445	530	625	725	580
Brilliant Isles	"	760	750	665	690	570	520	455	455	540	610	665	710	615
Stornoway	"	1085	1025	925	735	590	450	330	560	685	885	875	1015	745
Calgary Canada (West)	"	1850	1790	1595	1270	985	750	575	625	825	1090	1450	1695	1200
PORT GEORGE	"	1085	1230	1020	765	610	475	415	455	570	735	1000	1170	815
Kag River	"	1085	1230	1020	765	610	475	415	455	570	735	1000	1170	815
Prince George	"	1255	1285	1185	1040	820	640	520	370	430	630	790	935	655
The Pas	"	1430	1275	1115	825	620	455	355	410	585	785	1025	1235	845
Vancouver	"	1665	1640	1600	1460	1260	1060	860	720	580	480	565	640	510
Charlottetown (East)	"	1125	1185	1005	850	645	435	360	480	590	660	850	1035	745
Quebec	"	1225	1255	1150	970	765	635	460	475	600	780	955	1125	865
Alert	"	1405	1300	1145	950	760	595	440	480	615	815	1025	1270	890
Goose Bay	"	1375	1270	1200	975	765	545	435	460	590	745	1005	1230	885
Moosehide	"	1495	1390	1215	995	790	580	510	515	695	870	1085	1370	960
Nitchequon	"	1715	1720	1585	1320	1040	835	720	715	825	1000	1270	1515	1185
Port Harrison	"	1345	1335	1135	885	655	465	370	410	545	750	995	1225	845
Quebec	"	1245	1300	1135	970	645	475	370	420	585	805	1065	1310	862
Sloux Lookout	"	1460	1400	1135	870	655	600	480	570	725	930	1165	1385	1025
Alert	"	1470	1470	1335	1160	855	600	480	570	725	930	1165	1385	1025
Arctic Bay	"	1140	1240	1290	1100	1075	795	770	795	1060	1325	1255	1420	1175
Baker Lake	"	1130	1335	1290	1135	970	795	705	710	895	1045	1285	1270	1100
Copernicus	"	2030	1835	1585	1480	1070	835	675	670	845	1165	1490	1760	1295
Dawson	"	1600	1625	1465	1250	970	730	605	655	830	1100	1445	1545	1190
Fort Simpson	"	1345	1105	1045	720	545	405	350	415	565	790	1140	1230	810
Mould Bay	"	1460	1440	1325	900	515	400	405	435	615	870	1210	1435	930
Resolution Island	"	1675	1690	1735	---	---	---	---	---	---	---	---	---	---
Jakobshavn	"	1580	1595	1430	1295	1055	915	865	890	925	1065	1225	1435	1180
Greenland	"	1320	1305	1205	1040	860	705	470	620	775	970	1110	1220	970
Nenortalik	"	1070	1065	1005	910	750	750	670	660	760	825	950	1010	870
Scorebysund	"	1110	1145	1085	940	745	615	555	575	665	865	940	1010	850
Thule	"	1515	1630	1595	1360	1060	810	715	785	960	1195	1315	1485	1185
Reykjavik	"	1000	1000	965	860	735	645	575	605	685	815	895	975	815
Devils Lake	"	1340	1240	1065	810	600	435	335	365	520	730	1005	1220	805
Helena	"	1090	990	855	690	570	450	325	355	495	640	815	990	670
Sault Ste. Marie	"	1125	1150	1030	795	620	450	375	390	500	680	890	1090	745

*Values for coldest month are underlined.

**Data from Tables II and VI: were combined without regard to period of record.

TABLE IX MEAN CLOUDINESS (tenths of sky cover)*

Station	Yrs	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
<u>Barrow</u>	5-14	4.2	5.2	5.0	5.9	8.5	7.9	9.0	9.0	9.2	8.8	4.7	4.8	6.8
<u>Bethel</u>	12	5.5	6.1	6.5	6.9	7.9	8.0	8.5	9.0	8.4	7.8	7.2	7.1	7.5
<u>Dutch Harbor</u>	10	7.6	7.6	7.4	7.5	8.0	8.0	7.7	7.5	8.0	7.9	7.5	7.2	7.7
<u>FORT GREENLY</u>	9	5.5	5.0	5.2	5.5	6.3	5.7	6.4	7.2	6.6	6.7	5.7	5.5	6.0
<u>Juneau</u>	12	7.6	8.1	7.9	8.2	7.8	7.8	8.5	7.9	8.4	8.7	8.3	8.4	8.1
<u>Kodiak</u>	10	7.6	6.4	6.4	6.7	7.7	7.1	7.1	6.9	6.3	6.9	6.7	7.0	6.8
<u>Ketchikan</u>	25	7.9	5.8	5.8	6.5	5.7	6.1	7.9	6.1	7.5	7.1	6.5	6.6	6.8
<u>London</u>	30	7.5	7.2	6.8	6.5	6.1	6.4	6.6	6.6	6.5	6.8	6.5	7.0	7.3
<u>Sitka</u>	17	7.7	7.9	7.5	7.4	7.5	7.8	8.3	8.1	8.0	8.0	7.6	7.6	7.8
<u>Calgary, Canada (West)</u>	10	6.4	6.4	6.5	6.8	6.9	6.9	4.2	5.3	5.3	5.6	6.3	6.2	5.9
<u>FORT CHURCHILL</u>	8-10	4.7	4.5	4.8	5.9	7.7	7.2	6.3	6.8	8.0	8.3	7.5	5.4	6.4
<u>Edmonton</u>	9	5.8	6.1	5.8	6.2	6.3	6.6	6.5	5.7	6.5	6.8	7.0	6.4	6.3
<u>Prince George</u>	10	7.3	6.9	6.3	7.1	6.2	7.1	6.6	6.1	5.7	6.9	7.5	7.5	6.8
<u>The Pas</u>	10	5.3	5.6	5.3	5.4	5.9	6.0	5.1	5.7	5.9	6.2	7.3	5.6	5.8
<u>Vancouver</u>	17	5.2	6.9	6.3	6.0	6.1	5.6	5.2	4.3	5.2	6.6	7.7	7.7	6.2
<u>Charlottetown (East)</u>	10	7.1	6.1	6.2	6.8	6.5	6.3	5.2	5.4	5.4	6.1	7.1	7.5	6.4
<u>Gander</u>	10	7.5	7.6	7.2	7.8	7.8	7.8	7.0	7.0	6.8	7.0	8.2	7.1	7.4
<u>Goose Bay</u>	10	6.1	6.3	6.2	7.2	7.6	7.9	7.7	7.2	7.2	7.5	7.5	6.5	7.1
<u>Moose Factory</u>	13	5.2	5.0	4.9	5.6	6.3	5.8	5.3	5.9	6.6	7.0	7.5	6.3	6.0
<u>Nitchequon</u>	10	5.5	5.8	6.1	6.6	7.4	7.4	7.5	7.4	7.8	8.4	8.0	7.3	7.1
<u>Port Harrison</u>	10	4.9	4.4	5.2	6.6	8.1	7.5	7.2	7.2	8.0	8.3	8.0	7.2	6.9
<u>Quebec</u>	51	6.5	6.1	6.0	5.8	6.2	6.0	5.9	5.6	5.7	6.4	7.4	6.8	6.2
<u>Sioux Lookout</u>	10	6.5	5.9	5.8	6.3	6.6	6.9	6.1	6.1	6.6	6.5	8.3	6.9	6.5
<u>Alert</u>	8-11	4.9	4.7	4.0	5.0	5.7	5.0	6.0	7.2	7.5	7.2	6.0	5.2	5.7
<u>Arctic Bay</u>	2-3	4.8	4.9	3.8	4.2	6.1	6.4	5.8	7.7	6.8	6.8	5.4	4.6	5.6
<u>Baker Lake</u>	13	3.8	3.2	4.5	4.1	6.1	6.9	6.6	7.5	8.1	7.0	5.2	3.6	5.5
<u>Coppermine</u>	6-7	3.7	3.7	4.2	4.9	6.9	6.9	6.3	6.7	8.2	8.2	6.7	5.5	5.9
<u>Dawson</u>	10	5.4	5.0	5.4	4.9	6.9	6.5	6.7	7.7	8.2	8.2	6.7	5.5	6.5
<u>Fort Simpson</u>	15	2.7	4.9	4.4	4.8	4.9	5.3	2.6	2.8	5.9	6.5	6.2	5.3	5.5
<u>Mould Bay</u>	10	5.4	6.0	5.0	6.0	6.8	6.8	6.6	6.3	6.2	6.9	6.5	6.3	6.2
<u>Resolute Island</u>	3	3.2	4.5	3.8	3.8	5.1	7.8	8.2	8.0	8.6	6.4	4.2	3.2	5.6
<u>Resolution Island</u>	11	7.6	7.3	7.5	7.9	8.2	8.1	7.8	8.2	8.1	8.6	8.6	8.6	8.5
<u>St. John's</u>	30	5.0	4.4	4.4	4.8	5.1	5.2	4.8	5.0	5.0	5.3	5.3	5.3	5.0
<u>St. John's</u>	41	6.7	6.5	6.5	6.4	6.6	6.8	6.5	6.6	6.7	6.8	6.3	6.3	6.5
<u>Nanortalik</u>	11	5.0	5.9	5.3	5.2	6.2	5.6	5.5	5.8	5.5	5.6	5.9	6.0	5.5
<u>Scoresbyund</u>	3	4.7	4.5	4.0	5.6	6.0	6.2	7.3	6.3	6.3	6.3	4.4	4.4	5.6
<u>Thule</u>	25	7.1	7.2	7.2	7.1	6.9	7.1	7.3	7.3	7.3	6.8	7.0	7.0	7.1
<u>Reykjavik</u>	47	6.0	5.9	6.0	5.7	5.8	5.6	4.5	4.6	5.3	5.7	6.5	6.2	5.7
<u>De lla Lake</u>	64	5.7	6.5	6.5	6.4	6.5	6.0	4.0	4.1	5.1	5.7	6.5	6.8	5.9
<u>Br. ena</u>	64	7.9	6.8	6.3	5.9	6.0	5.5	5.2	5.6	6.3	7.2	3.4	8.3	6.6
<u>Sault Ste. Marie</u>														

*Values for coldest and warmest months are underlined.

TABLE X: MEAN SNOW DEPTH (inches)*

Station	Yrs	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Barrow	4-6	12.2	14.5	15.6	15.9	12.6	5.3	T	0	0.4	3.2	7.4	9.4
Bethel	6-8	10.1	11.6	11.5	7.0	1.3	0.0	0.0	0.0	0.0	0.0	1.2	5.4
Fort Harbord	8	2.7	8.2	6.2	0.7	0.0	0.0	0.0	0.0	0.0	T	0.1	0.2
FORT GREENLY	5	10.7	17.0	14.9	11.6	9.6	2.0	0.0	0.0	T	4.2	7.0	10.5
Juneau	19	4.4	3.8	2.3	0.4	0.0	0.0	0.0	0.0	0.0	0.1	0.9	3.3
Kodiak	10	1.6	1.3	0.8	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.3	1.2
Nome	11	8.0	13.0	16.0	11.0	4.0	0.0	0.0	0.0	T	T	2.0	6.0
British Isles													
No Data													
Calgary Canada (West)	14-21	6.4	3.4	2.3	0.8	0.0	0.0	0.0	0.0	0.0	0.3	1.3	5.5
FORT CHURCHILL	7-10	16.0	20.9	25.1	20.4	10.6	4.0	T	0.0	0.0	0.7	4.4	10.2
Fort Vermilion	10	17.6	19.3	17.7	8.4	0.0	0.0	0.0	0.0	0.0	1.7	8.1	14.0
Prince George	15-22	9.4	9.9	7.0	2.7	0.0	0.0	0.0	0.0	0.0	0.1	2.9	6.7
The Pas	16-25	12.8	15.6	11.1	3.1	T	0.0	0.0	0.0	0.0	1.0	4.9	9.0
Vancouver	7-13	1.6	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.1	0.4
Charlottetown (East)	17-33	12.2	13.2	8.1	1.8	0.2	0.0	0.0	0.0	0.0	0.3	1.2	6.6
Gander	5-7	10.8	12.8	12.1	4.9	0.1	0.0	0.0	0.0	0.3	0.5	0.4	5.7
Goose Bay	8-10	28.5	36.5	36.7	25.1	8.8	0.1	0.0	0.0	0.0	1.5	7.8	18.6
Moosonee	8-12	15.0	20.3	20.0	11.1	1.9	0.0	0.0	0.0	0.0	0.1	3.9	9.4
Mitchouon	3-5	24.4	34.6	46.9	43.4	17.7	0.9	0.0	0.0	0.0	4.8	12.8	18.0
Port Harrison	7-11	13.2	15.7	18.1	21.7	13.4	2.7	0.0	0.0	0.0	2.4	7.2	10.3
Quebec	20-34	18.5	28.6	23.7	7.5	0.1	0.0	0.0	0.0	0.0	T	2.0	8.2
Sioux Lookout	13-25	1.5	22.5	18.7	6.7	0.4	0.0	0.0	0.0	0.0	0.5	4.1	10.9
Altiavik (Territories)													
Alert	5-9	14.6	20.9	21.7	16.7	5.6	0.0	0.0	0.0	0.0	2.5	7.5	9.1
Arctic Bay	5	14.1	15.2	14.7	12.8	10.7	4.7	0.4	0.9	5.6	11.2	12.5	13.0
Baker Lake	4-7	9.1	10.5	11.7	11.0	6.9	1.8	0.0	0.0	1.2	4.5	7.3	8.3
Coppermine	4-6	21.9	29.7	30.1	16.1	14.4	6.7	1.2	0.0	0.0	2.3	6.4	9.5
Dawson	17-33	20.6	24.2	23.2	12.1	1.4	0.0	0.0	0.0	0.0	5.3	11.1	16.1
Fort Simpson	4-10	19.9	21.5	18.6	8.1	1.3	0.0	0.0	0.0	0.0	0.0	7.8	9.3
Mould Bay	7	6.1	6.7	7.6	8.7	8.9	4.8	0.0	0.0	0.0	0.8	4.9	14.9
Resolution Island	4-7	26.1	26.9	35.2	27.1	6.5	0.0	0.0	0.2	1.6	3.5	4.6	5.4
Jakobshavn	5	17.2	20.8	22.1	21.6	11.6	0.4	0.0	0.0	0.0	1.3	6.2	18.7
Narsarsuaq	7	4.2	7.3	3.9	2.2	0.6	0.0	0.0	0.0	0.0	0.3	4.4	12.3
Scoresbysund	5	24.0	25.0	28.0	40.0	39.0	8.0	T	0.0	T	0.6	1.6	2.9
Thule	4-5	2.5	2.0	2.0	2.3	1.1	0.0	0.0	0.0	0.0	0.3	1.1	2.0
Reykjavik	15	3.0	3.4	1.7	0.8	0.2	0.0	0.0	0.0	T	0.5	1.4	2.1
Devils Lake	20	5.7	6.1	3.5	0.7	0.0	0.0	0.0	0.0	0.0	0.1	1.1	3.5
Elzevir	40	2.5	2.3	1.0	0.1	0.0	0.0	0.0	0.0	0.0	0.2	0.7	1.5
Sault Ste. Marie	20-40	12.1	15.0	8.5	1.2	0.0	0.0	0.0	0.0	0.0	0.2	1.9	6.2

*Values for month of greatest depth are underlined.

TABLE XI: MEAN SNOWFALL (inches)*

Station	Yrs Rec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
<u>Alaska</u>														
Barrow	21-33	2.6	2.7	2.0	2.7	1.8	0.4	1.0	0.7	3.0	7.2	4.3	3.9	33.0
Bethel	16-18	7.2	4.9	8.0	3.3	0.5	0.0	0.0	0.0	0.2	3.2	5.2	2.3	41.9
Dutch Harbor	9-23	15.6	13.7	10.5	6.7	0.1	T	0.0	T	0.4	0.5	5.8	10.3	69.7
<u>PORT GREENLY</u>														
Fort Churchill	9	7.0	2.9	5.6	1.3	0.3	T	0.0	T	0.4	6.4	5.3	4.6	33.8
Juneau	33-44	28.1	23.8	14.9	4.6	0.1	0.0	0.0	0.0	0.1	1.5	9.1	25.0	107.2
Kodiak	33-38	9.5	10.6	8.2	5.4	0.3	T	0.0	0.0	T	1.1	3.4	8.4	46.9
Nome	29-34	10.8	8.9	10.2	5.1	1.3	0.2	0.0	T	0.4	3.6	8.5	10.8	59.8
<u>British Isles</u>														
Londor	60	0.1	0.1	0.3	T	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.7
<u>Canada (West)</u>														
Calgary	67	5.0	5.8	8.1	6.4	4.2	0.3	T	0.1	2.6	4.2	7.1	5.4	49.2
Fort Churchill	30	4.8	6.1	8.5	7.7	1.8	1.4	T	0.0	1.7	8.0	10.3	6.6	56.9
Fort Vermilion	30	7.0	6.6	7.3	3.2	0.7	T	0.0	0.0	0.4	2.8	6.7	8.6	43.3
Prince George	27	16.2	10.1	8.1	0.6	T	T	0.0	0.0	0.2	1.7	10.0	15.8	62.7
The Pas	30	7.5	7.8	7.9	5.3	0.8	T	0.0	0.0	0.3	3.6	10.4	9.6	53.2
Vancouver	41	11.5	6.4	2.7	0.3	0.0	0.0	0.0	0.0	0.0	0.1	2.1	5.7	28.8
<u>Charlottetown (East)</u>														
Charlottetown	65	27.1	25.4	20.3	7.3	0.8	0.0	0.0	0.0	0.0	0.3	7.0	24.8	113.0
Gander	14	20.2	24.6	18.0	13.0	4.2	1.3	0.0	0.0	0.2	3.5	9.8	24.4	119.2
Goose Bay	10	22.1	21.8	23.0	15.2	7.7	T	0.0	0.0	0.8	10.5	18.6	24.4	144.1
Moose Factory	40	14.2	10.7	11.3	6.3	3.1	0.4	0.0	0.0	T	2.5	8.2	12.9	69.6
Niobeau	10	13.2	14.1	15.2	12.2	10.4	0.9	T	0.0	1.0	12.6	20.7	16.0	116.3
Port Harrison	20	6.6	3.1	8.2	7.0	5.3	1.7	T	0.0	1.8	10.5	19.3	9.6	73.3
Quebec	72	29.3	23.1	20.8	8.7	0.5	0.0	0.0	0.0	T	1.8	14.4	25.1	123.7
Stout Lookout	23	10.2	8.6	11.6	6.0	2.8	0.0	0.0	0.0	0.8	4.7	10.3	10.5	65.5
<u>Aklavik (Territories)</u>														
Aklavik	22	5.5	4.9	3.8	5.0	3.4	1.7	T	1.0	3.1	7.9	7.6	4.2	47.1
Alert	5	1.7	2.5	2.5	2.3	3.1	3.6	3.8	7.8	12.2	7.7	2.1	2.9	52.2
Arctic Bay	13	3.1	2.5	2.9	2.5	2.7	2.3	0.1	0.5	6.2	6.6	3.8	2.7	35.9
Baker Lake	6	2.2	2.0	3.1	2.7	0.8	1.0	T	0.1	0.4	3.1	3.4	2.9	21.9
Coppermine	30	6.0	4.0	7.1	4.8	4.7	1.4	T	0.1	3.8	10.2	7.3	5.6	55.5
Dawson	41	8.7	7.0	5.1	3.5	0.6	0.1	0.0	T	1.8	7.9	10.4	10.2	55.3
Fort Simpson	42	7.2	7.0	4.8	5.8	4.1	T	0.0	0.1	0.7	8.1	8.5	8.3	54.6
Mould Bay	7	0.8	0.6	1.0	1.4	2.8	1.7	1.1	2.0	2.8	1.2	1.2	1.2	21.3
Resolution Island	19	10.9	10.1	9.6	7.9	8.0	3.2	T	0.2	2.4	8.0	11.8	15.0	86.4
<u>Greenland</u>														
Narsarsuaq	7	17.1	18.0	8.7	7.8	3.2	T	0.0	0.0	1.9	3.9	10.9	11.2	74.7
Sandvig	4	3.5	1.1	2.1	2.0	1.2	0.0	0.0	0.0	0.0	2.0	5.8	6.1	31.0
Stordarfjord	6	0.6	0.5	1.5	3.3	1.1	0.8	0.0	0.0	0.2	1.5	2.5	1.9	14.9
Thule	M	20.8	14.9	8.4	8.6	0.4	T	0.0	0.0	T	1.2	3.3	10.2	67.9
<u>Iceland</u>														
Reykjavik	45	5.3	5.2	5.4	3.0	0.7	T	0.0	0.0	T	1.7	5.5	6.5	33.3
Devils Lake U. S. A.	73	9.9	8.1	9.2	5.1	1.9	0.1	T	1.0	3.7	6.8	8.8	8.8	54.6
Helena	50	18.7	13.4	10.2	3.5	0.6	0.0	0.0	0.0	T	2.3	12.6	18.1	79.3
Sault Ste. Marie														

* Values for month of greatest snowfall are underlined.

** % of Reykjavik, newly uncovered data, not shown on map (Fig 9) (period of record missing - M)

TABLE III: MEAN PRECIPITATION (inches)*

Station	Yrs Rec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Barrow	25-27	0.15	0.20	0.13	0.12	0.14	0.26	0.59	0.73	0.49	0.36	0.30	0.26	4.23
Betoul	18-19	0.96	0.61	0.77	0.53	0.74	1.13	2.37	3.89	3.31	1.60	0.98	0.79	17.68
Dutch Harbor	19-23	6.32	6.12	4.92	4.19	4.33	4.82	1.89	2.45	5.45	7.12	5.81	6.98	58.61
PORT GRIZLY	9	0.40	0.20	0.30	0.30	0.20	2.10	3.20	2.10	1.20	0.40	0.30	0.30	11.40
Juneau	46-51	7.16	5.53	5.51	5.39	5.22	4.04	3.14	7.38	10.17	11.41	9.60	7.58	83.72
Kodiak	46-52	4.82	4.68	3.94	4.20	5.94	4.86	3.56	5.04	5.48	7.15	5.69	6.17	61.94
Kodiak	35-37	1.09	0.81	0.84	0.67	0.64	1.12	2.50	1.82	2.71	1.58	0.97	1.11	17.26
LONDON BRITISH ISLES	64	2.00	1.67	1.83	1.84	1.76	2.03	2.24	2.23	1.95	2.61	2.36	2.49	28.71
Stornoway	58	5.62	4.18	3.79	2.96	2.59	2.46	2.97	3.99	4.20	5.43	5.48	5.84	49.51
Calgary	55	0.51	0.55	0.84	0.99	2.34	3.14	2.51	2.29	1.50	0.69	0.72	0.97	15.63
Canada (West)														
PORT CHURCHILL	30	0.41	0.36	0.63	0.69	0.65	1.51	2.61	2.94	1.97	1.50	0.76	0.56	14.41
Port Vermilion	30	0.71	0.66	0.74	0.61	1.36	1.67	1.67	1.66	1.27	0.61	0.74	0.86	12.76
Prince George	27	1.81	1.21	1.44	0.84	1.34	2.06	1.63	1.94	2.00	1.99	1.87	1.85	19.98
The Pas	30	0.77	0.78	0.83	0.93	1.38	2.30	2.42	2.30	2.06	1.11	1.13	0.97	16.98
Vancouver	41	8.57	8.72	5.03	3.34	2.84	2.45	1.22	1.69	3.63	5.78	8.28	8.76	57.38
Charlottetown (East)														
Gander	65	3.76	3.01	3.15	2.78	2.66	2.58	2.98	3.35	3.45	5.07	3.75	3.98	39.47
Goose Bay	14	2.63	3.28	2.29	2.57	2.56	2.79	3.61	3.59	3.66	4.09	4.21	3.72	39.50
Moose Factory	10	2.27	2.29	2.36	1.91	2.12	2.39	3.24	2.79	2.25	2.44	2.50	2.49	29.05
Hitchman	40	1.42	1.07	1.30	1.15	1.59	1.97	2.29	2.03	2.44	1.78	1.05	1.38	20.47
Port Harrison	10	1.34	1.24	1.65	1.55	2.65	3.75	4.07	4.12	3.12	3.11	2.44	1.67	30.88
Quebec	20	0.66	0.31	0.82	0.74	0.80	0.88	1.54	1.61	2.21	2.05	2.06	0.96	14.64
Sioux Lookout	72	3.45	2.74	3.02	2.35	3.15	3.68	4.02	3.98	3.60	3.41	3.23	3.22	39.85
Aklavik (Territories)	23	1.03	0.88	1.30	1.74	1.86	3.46	1.17	2.76	3.46	1.78	1.72	1.06	24.52
Alert	22	0.55	0.49	0.78	0.50	0.49	0.80	1.39	1.42	0.91	0.66	0.76	0.42	8.97
Arctic Bay	5	0.17	0.25	0.25	0.23	0.31	0.47	0.65	1.28	1.23	0.77	0.21	0.29	6.11
Baker Lake	13	0.31	0.25	0.29	0.25	0.29	0.44	0.68	1.17	0.81	0.66	0.36	0.27	5.80
Coppermine	6	0.22	0.20	0.31	0.27	0.21	0.70	0.86	1.09	0.83	0.50	0.34	0.29	5.82
Dawson	30	0.60	0.40	0.71	0.67	0.55	0.91	1.52	1.69	1.25	1.23	0.78	0.56	10.87
Port Simpson	41	0.87	0.70	0.52	0.51	1.01	1.18	1.53	1.57	1.41	1.17	1.13	1.03	12.61
Mould Bay	42	0.72	0.70	0.48	0.70	1.37	1.46	1.92	1.47	1.31	1.07	0.85	0.84	12.96
Resolution Island	7	0.08	0.06	0.16	0.14	0.28	0.27	0.63	0.63	0.48	0.28	0.12	0.12	3.25
Jakobshavn Greenland	19	1.02	1.02	0.97	0.83	1.14	1.30	1.83	1.59	2.02	1.27	1.22	1.21	15.71
Narsartalik	30	0.33	0.30	0.50	0.69	0.76	0.76	1.26	1.50	1.29	1.31	0.74	0.63	10.07
Scoresbysund	4	2.78	4.11	1.49	2.45	1.54	4.56	1.20	3.46	2.04	4.73	3.76	1.98	34.64
Thule	10	1.60	1.70	1.61	1.49	0.30	1.00	0.60	0.60	0.40	1.15	1.05	1.06	14.50
Reykjavik	6	0.11	0.08	0.10	0.13	0.10	0.15	0.41	0.60	0.37	0.13	0.19	0.18	2.52
Devils Lake U. S. A.	50	3.02	3.31	2.72	2.14	1.89	1.93	1.89	2.01	3.54	3.43	3.74	3.50	34.26
Salina	45	0.42	0.43	0.67	1.34	2.09	1.22	2.68	2.26	1.90	1.11	0.69	0.50	17.31
Sault Ste. Marie	73	0.76	0.57	0.76	0.99	1.93	2.26	1.10	0.80	1.20	0.82	0.67	0.72	12.58
	64	2.00	1.44	1.78	2.07	2.68	2.91	2.61	2.84	3.49	3.00	3.05	2.23	30.12

*Values for month of greatest precipitation are underlined.

TABLE XIII: MEAN RELATIVE HUMIDITY (%)

Station	Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Year
Barrow	11	64	62	65	74	87	93	92	93	93	92	86	76	80
Bethel	7	80	79	81	82	79	77	83	89	88	88	86	80	83
Fairbanks (for FORT GREENLY)	15	69	68	66	61	58	62	70	76	76	79	73	71	69
Juneau	12	80	79	78	75	76	74	78	81	82	82	84	81	80
Kona	15	78	79	79	80	80	81	86	86	83	81	81	81	81
London	30	84	82	79	75	73	73	73	76	79	85	86	86	79
Stornoway	18	87	86	84	82	80	80	85	85	85	86	86	87	84
Calgary Canada (West)	7	72	76	71	64	63	70	66	65	69	70	73	72	69
FORT CHERCHILL	10	91	90	90	89	85	76	72	72	85	86	93	94	85
Embaras	4-5	83	81	78	68	62	62	66	70	77	79	87	89	75
Prince George	6	88	80	72	67	62	72	73	73	77	81	89	92	78
The Pas	10	93	90	81	63	53	60	61	64	65	71	86	94	73
Vancouver	7	89	85	80	78	76	76	76	77	83	86	88	90	82
Charlottetown (East)	5	77	77	71	70	71	73	73	74	75	75	77	75	74
Gender	7	88	85	84	80	79	79	76	80	83	84	86	87	83
Goose Bay	5	84	83	74	71	68	67	63	69	70	73	80	87	74
Kapuskasing	7	91	90	84	72	70	71	73	72	83	86	90	90	82
Lake Dore	5-6	85	88	86	79	77	74	77	80	84	85	90	89	83
Port Harrison	6-7	--	--	--	93	93	89	87	90	91	93	94	94	--
Quebec	5	78	78	82	76	74	77	80	81	83	83	84	82	80
Sioux Lookout	6	90	87	75	65	68	73	73	77	81	79	89	91	79
Aklavik (Territories)	10	96	97	92	84	77	68	68	79	88	95	97	94	87
Arctic Bay	5	--	--	--	83	87	77	79	83	83	82	--	--	--
Baker Lake	6	70	78	83	87	87	73	81	82	87	92	92	82	83
Coppermine	9	--	--	--	--	--	57	60	66	71	--	--	--	--
Dawson	9	--	--	--	56	46	44	45	54	59	74	89	89	76
Fort Simpson	8	82	89	69	56	56	54	53	91	94	98	--	--	--
Resolution Island	7-9	97	--	--	--	97	77	73	71	73	78	78	77	76
Jakobhavn Greenland	30	76	74	79	83	77	73	73	83	75	72	76	75	78
Myggbukta	6	76	75	76	74	80	79	83	83	75	71	76	75	75
Nenortalik	23	75	76	74	72	72	73	76	78	76	75	74	75	77
Thule	6	76	76	74	75	76	80	82	74	76	79	80	76	77
Reykjavik Iceland	3	81	80	80	74	70	75	75	74	74	78	81	81	77
Lavila Lake U.S.A.	12	85	85	82	71	67	71	70	69	69	68	81	82	75
Belena	13-65	67	66	62	56	56	54	48	47	55	61	66	62	59
Sault Ste. Marie	9-61	84	83	78	76	74	76	76	77	81	80	82	84	79

Values for months of highest known relative humidity are underlined.

8. Bibliography

1. Anderson, Lois C., Variability of Monthly Mean Temperature, Tech Rpt EP-16, QM R&D Command, Natick, Mass., July 1955
2. Anstey, Robert L., Handbook of Thule, Greenland, Environment, Tech Rpt EP-34, QM R&D Command, Natick, Mass., Aug 1956
3. Boughner, C. C. and J. G. Potter, Snow Cover in Canada, Weatherwise, Dec 1953
4. Boughner, C. C. and M. K. Thomas, Climatic Summaries for Selected Meteorological Stations in Canada, Newfoundland, and Labrador, Vol. II (Wind Speed and Direction), Canadian Department of Transport, Meteorological Division, Toronto, 1948
5. Brand, W., Stündliche Werte des Luftdrucks und der Temperatur an DanmarksHavn (Hourly Values of Atmospheric Pressure and Temperature at Denmarkshavn), Danmark-Ekspeditionen til Grønlands Nordøstkyst 1906-08, Særtryk af Meddelelser om Grønland (Vol. 2.), Copenhagen, 1912
6. Canadian Department of Transport, Meteorological Division, Addendum to Volume I, Climatic Summaries for Selected Meteorological Stations in Canada (Average Values of Temperature and Precipitation), Toronto, 1954
7. ———, Average Wind Speed in Miles per Hour for Canadian Stations Equipped with Anemometers, Toronto, unpublished (no date)
8. ———, Climatic Summaries for Selected Meteorological Stations in the Dominion of Canada, Vol. I (Average Values of Mean and Extreme Temperature, Mean and Extreme Humidity, Sunshine, and Precipitation), Toronto, 1947
9. ———, Joint Arctic Weather Stations, Five Year Report 1944-51. A Review of the Establishment and Operation and Summary of Scientific Activities, compiled jointly with the U. S. Weather Bureau, Toronto, 1954
10. ———, Metstat, Telegraphic Meteorological Stations in Canada, Toronto, 1954
11. ———, Monthly and Annual Extreme Highest and Lowest Temperatures for Selected Canadian Stations, 1921-1950, Toronto, 1957(unpublished)

12. ———, Monthly Record of Meteorological Observations in Canada and Newfoundland, Toronto, 1920-1955

13. ———, Monthly Weather Map, Canada (mean and extreme temperatures, precipitation and snow depth data), Toronto, 1954-56

14. ———, Snow Cover Data, Eastern Canada, Winter 1954-55, Toronto, 1956 /First year of publication/

15. ———, Tabulations of Climatic Data for the Arctic Archipelago Joint U. S.-Canadian Weather Stations: Alert, Eureka, Isachsen, Mould Bay, and Resolute, Toronto 1956 (unpublished)

16. Chapman, L. J., The Climate of Northern Ontario, Canadian Journal of Agricultural Science, Ottawa, Jan-Feb 1953

17. Church, F. E. and W. Shallert, Tables for Alaska Climatic Atlas of the Aleutian Islands, DA Contract No. W-44-109-QM-1882, University of Washington, Dept. of Meteorology and Climatology, Seattle, 1950 (unpublished)

18. Connor, A. J., The Climate of Manitoba, Rpt. No. 16, Manitoba Economic Survey Board, Winnipeg, 1939

19. ———, The Temperature and Precipitation of Northern Canada, The Canadian Yearbook, Part VII, Sec. 1 (Climate and Meteorology), Dept. of Marine, Meteorological Service, Toronto, 1930

20. Court, Arnold, Windchill, Bull. Amer. Met. Soc., Vol. 29, No. 10, Dec 1948

21. dePercin, F. and S. J. Falkowski, Frequency and Duration of Low Temperatures at Fort Churchill, Manitoba, Canada, Tech. Rpt. KP-2, QM R&D Command, Natick, Mass., July 1954

22. ———, Frequencies of Selected Low Temperatures in Alaska, Tech. Rpt. KP-6, QM R&D Command, Natick, Mass., Feb 1955

23. dePercin, F. and S. J. Falkowski and R. C. Miller, Handbook of Big Delta, Alaska, Environment, Tech. Rpt. KP-5, QM R&D Command, Natick, Mass., Apr 1955

24. dePercin, F. and E. Bingham, Handbook of Devils Lake, North Dakota, Environment, Tech Rpt. KP-8, QM R&D Command, Natick, Mass., Apr 1955

25. dePercin, F., and L. W. White, Handbook of Fort Churchill, Manitoba, Environment, Tech. Rpt. EP-4, QM R&D Command, Natick, Mass., Aug 1954

26. Diamond Marvin, Precipitation Trends in Greenland During the Past 30 Years, SIFRE Rpt. No. 22, Corps of Engineers, U. S. Army, Wilmette, Illinois, Dec 1956

27. Dunbar, Moira and K. R. Greeraway Arctic Canada from the Air, Canadian Defense Research Board, Queen's Printer, Ottawa, 1956

28. Falkowski, S. J., Climatic Analogs of Fort Greely, Alaska and Fort Churchill, Canada in Eurasia, Tech. Rpt. EP-79, QM R&E Command, Natick, Mass., Dec 1957

29. Falkowski, S. J. and A. D. Hastings, Windchill in the Northern Hemisphere, Tech. Rpt. EP-82, QM R&E Command, Natick, Mass., Feb 1958

30. Georgi, J., Die Eismitte Station: Wissenschaftliche Ergebnisse Deutsche Grönland Expeditionen - Alfred Wegener 1929 und 1930-31 (The "Eismitte" Station: Scientific Results of the German Greenland Expeditions led by Alfred Wegener 1929 and 1930-31) (Vol. 4, Part I), Leipzig, 1935

31. Great Britain Air Ministry, Meteorological Office, Climatological Atlas of the British Isles, M. O. 488, HM Stationery Office, London, 1952

32. ———, Tables of Temperature and Relative Humidity, Oceans and Islands, M. O. 18, mimeographed, London, Sep 1948

33. Haro, F.R., The Climate of the American Northlands, OPNAV P03-32 U. S. Navy Dept., Office of Tech. Assist. to Chief of Naval Operations for Polar Projects, Washington, 1956

34. ———, The Climate of the Eastern Canadian Arctic and Subarctic and its Influence on Accessibility (PhD dissertation) 2 Vols., L'Université de Montreal, 1950

35. Hastings, A. D., Jr., Climatic Information Sources for Greenland, Research Study Report RER-11, QM R&D Command, Natick, Mass., Jan 1957

36. Henry, T. J. A. and G. R. Armstrong, Aerological Data for Northern Canada, Canadian Dept. of Transport, Meteorological Division, Toronto, 1949

37. Hogue, D. W., Temperatures of Northern North America, Research Study Report RER-9 QM RAD Command, Natick, Mass. (revised ed.), Oct 1957
38. Kols, R. S., Pole Vaulting for Weather, U. S. Air Force, 375th Recon. Sqdn-(VLR) Weather, Eielson AFB, Alaska, 1950
39. Kendrew, W. G., The Climates of the Continents, 4th ed., Oxford at the Clarendon Press, Amen House, London, 1951
40. Kendrew, W. G. and B. W. Currie, The Climate of Central Canada, Queen's Printer, Ottawa, 1955
41. Kendrew, W. G. and D. P. Kerr, The Climate of British Columbia and the Yukon Territory, Queen's Printer, Ottawa, 1955
42. Kimble, G. H. T. and Dorothy Good, Geography of the Northlands, Amer. Geog. Soc. and John Wiley & Sons, Inc., New York, 1955
43. Koeppe, C., The Canadian Climate, McKnight & McKnight, Bloomington, Illinois, 1931
44. Koepen, W., G. Geiger, and R. Geiger (Munich), Klima des Kanadischen Archipels und Grönlands (Climate of the Canadian Archipelago and Greenland), Handbuch der Klimatologie [Vol. II, Part K] verlag von Gebrüder Borntraeger, [coastal section by Helge Petersen, Icecap section by Fritz Loeve], Berlin, 1945
45. Lotz, J. R., Temperature Summaries at Knob Lake, Quebec, 1948-56, McGill University Subarctic Research Laboratory (extracted from manuscript records at Knob Lake) Montreal, unpublished, Jan 1957
46. Middleton, W. E. K., The Climate of the Gulf of St. Lawrence and Surrounding Regions in Canada and Newfoundland, as it Affects Aviation, Canadian Meteorological Memoirs No. 1, Dept. of Marine, Meteorological Service, Ottawa, 1935
47. Ministère des Travaux Publics, des Transports, et du Tourisme; Secrétariat Général à l'Aviation Civile et Commerciale, Expeditions Polaires Françaises, Recueil des Observations Météorologiques effectuées par la Station Française du Groenland entre le 20 Juillet 1949 et le 14 Juillet 1950, (Collection of the Meteorological Observations made at the French Greenland Station between 20 July 1949 and 14 July 1950), Météorologie Nationale, Paris, Sep 1950
48. Oliver, V. J. and M. B. Oliver, Ice Fogs in the Interior of Alaska, Bull. Amer. Met. Soc., Vol. 30, No. 1, Jan 1949

49. Petterssen, S., W. C. Jacobs, and B. C. Haynes, Meteorology of the Arctic, ONVAP P03-3, U. S. Navy Dept., Office of the Tech. Assist. to the Chief of Naval Operations for Polar Projects, Washington, Mar 1956
50. Putnam, D. F. (ed), B. Brouillette, D. P. Kerr, and J. L. Robinson, Canadian Regions: A Geography of Canada, Thomas Y. Crowell Co., New York, 1952
51. Rae, R. W., Climate of the Canadian Arctic Archipelago, Canadian Dept. of Transport, Meteorological Division, Toronto, 1951
52. Reed, W. W., Temperatures in Europe, U. S. Weather Bureau, prepared under the direction of the Chief, Division of Climate and Crop Weather, Washington, Nov 1932
53. Siple, Paul A. and C. F. Passel, Measurements of Dry Atmosphere Cooling in Subfreezing Temperatures, Proc. Amer. Phil. Soc., Vol. 89, 1945
54. Stone, Robert A., Fog in the United States and Adjacent Regions, The Geographical Review, Vol. 26, No. 1, Jan 1936
55. Stupart, Sir Frederick, The Climate of Canada, The Canada Yearbook, Part VII, Sec. 1 (Climate and Meteorology), Dept. of Marine, Meteorological Service, Toronto, 1930
56. Thomas, M. K., Climatological Atlas of Canada, Canadian Dept. of Transport, Meteorological Division, Toronto, 1947
57. Thompson, W. F., Environmental Handbook for Whittier, Alaska, EPD Report 226, QM R&D Command, Natick, Mass., Apr 1954
58. U. S. Air Force, Air Weather Service, Atlas of Temperature and Precipitation Data for Army Weather Stations in the Aleutian Islands, Washington, Sep 1947
59. ———, Climate of the Northeast Air Command Region, 5th Weather Group, 8th Weather Squadron, Tech Memo No. 6, Washington, Oct 1953
60. ———, Climatology of Thule, Greenland, Oct 1946-July 1952, Washington, 1953
61. ———, General Aspects of Fog and Stratus Forecasting, AWS Manual No. 105-44, Military Air Transport Service, Washington, 1954
62. ———, Climate, Weather, and Flying Conditions of Alaska and Eastern Siberia, Arctic Weather Central, 11th Weather Squadron, Fort Richardson, Alaska, May 1949

63. ———, Frequency of Daily Temperatures, Thule, Greenland, Oct 1946 - Dec 1954, National Weather Records Center, Data Control Division, Asheville, N. C., 1955 (unpublished)

64. ———, Local Forecasting Studies, Marsarssuak Area, Greenland, Aug. 1941 - Dec. 1953, National Weather Records Center, Data Control Division, Washington, 1953

65. ———, Summaries No. 6 and 7, Nord, Greenland, May 1952-July 1955, National Weather Records Center, Data Control Division, Asheville, N. C., 1956 (unpublished)

66. ———, Thule Site II, Greenland, July 1953-Dec. 1954, National Weather Records Center, Data Control Division, Asheville, N.C., 1955 (unpublished)

67. U. S. Air Force, Mean Monthly Temperature and Average Monthly Wind; Canadian Archipelago, Northern Quebec, Northern Manitoba, and Labrador, Regional Control Office, 8th Weather Region, Grenier AFB, Manchester, N. H., no date (unpublished)

68. U. S. Army Air Force, Air Weather Service, Studies on Local Forecasting, Greenland Weather Stations, Report No. 600-51, Washington, Aug 1944

69. ———, Climatology of the Arctic Regions, Parts I, II, and III, Special Study No. 58, National Weather Records Center, Data Control Division, prepared by Statistics Division, US Weather Bureau, Asheville, N. C., Dec 1946

70. ———, Frequency and Mean Temperature Data, Attahbury Pass, Greenland, Oct. 1943-Sept. 1944, National Weather Records Center, Data Control Division, Asheville, N. C., no date (unpublished)

71. ———, Summary of Temperatures, Precipitation, and Surface Winds, Nome, Alaska AAR Part of Evaporation, New Orleans 1945

72. ———, Climatic Data for Alaska, Report No. 444, Climatological Division, prepared by Climate and Crop Weather Division, US Weather Bureau, Washington, Sep 1943

73. ———, Climatic Data for Kaffin Island, AFSTW No. 211, Climatological Division, prepared with Statistics Division US Weather Bureau, WBSR No. 221, Washington, no date

74. ———, Climatic Statistics for Selected Stations in Canada, AFSTW No. 322, Climatological Division, prepared with Statistics Division, US Weather Bureau, WBSR No. 187, Washington, no date

75. ———, Climatology of Alaska, Supplement to Rpt. No. 444 (entry 72 above), Climatological Division, Washington, 1944

76. U. S. Army Air Force, Weather Information Branch, Preliminary Climatic Atlas of the World, Nov. 1942 and Supplement, July 1943, Special Series No. 1, prepared by the Institute of Meteorology, University of Chicago, Washington, 1942, 1943

77. U. S. Army Air Force, Weather Research Center, Climatic Statistics for Selected Stations in Alaska; Seattle, Wash; and Fort Snelling, Minn., Report No. 101, prepared by Statistics Division, US Weather Bureau, Washington, 1942

78. ———, Flying Conditions in the Southampton Island and Port Harrison Areas, Canada, Report No. 181, prepared by Statistics Division, US Weather Bureau, Washington, Sep 1942

79. ———, Preliminary Report on Climate and Weather of Northwestern Europe, Vol. III, No. 1, Washington, Jun 1942

80. ———, Preliminary Report on Topography and Climate of British Columbia and Northwestern Alberta, including Temperature, Precipitation, Rainfall and an Index of Meteorological Stations, Vol. II, No. 2, prepared by Statistics Division, US Weather Bureau, Washington, May 1942

81. U. S. Department of Agriculture, Climate and Man, Yearbook of Agriculture, House Document No. 27, 77th Congress, 1st Session, Washington, 1941

82. U. S. Department of Agriculture, Weather Bureau, Climatic Summary of the United States, Bulletin "W" (1st order station summaries), Washington, 1931

83. U. S. Department of Army, Corps of Engineers, Office of the Chief of Engineers, Depth of Snow Cover in the Northern Hemisphere, Investigation of Construction and Maintenance of Airdromes on Ice - FY 54, prepared by Arctic Construction and Frost Effects Laboratory, N. E. Division, Boston, Jun 1954

84. ———, Report of Investigations, Investigation of Construction and Maintenance of Airdromes on Ice - FY 50, prepared by 1st, Foundation, and Frost Effects Laboratory, N. E. Division, Boston, Jun 1950

85. U. S. Department of Army, GQG, Geography and Climatology of the Arctic Operations Area, Special Report No. 20, Military Planning Division, R & D Branch, prepared by the staff, Washington, May 1949

86. U. S. Department of Commerce, Weather Bureau, Climatic Summary of the United States, Supplement to Bulletin "W" (entry 82 above), Washington, [published for states at various times since 1952]
87. ——, Climatological Data, Alaska, Washington, annual summaries 1915-1954 [published by Dept. of Agriculture prior to 1940]
88. ——, Climatological Summary, Thule, Greenland, Oct. 1946-Dec. 1949, Washington, 1950
89. ——, Index of Climatic and Weather Data, Greenland (WB photostat summaries for 17 stations up to 1936), Washington, no dates
90. ——, Index of Climatic Data for Alaska, Great Britain, Iceland, and Scotland (WB photostat summaries for 52, 40, 14, and 7 stations respectively), Washington, no dates
91. ——, Local Climatological Data, Alaska, Washington, annually
92. ——, Local Climatological Data, United States, Washington, annually
93. ——, Maps of Climatic Means for Alaska and Western Canada (temperature extremes, precipitation, and snowfall), Washington, no date
94. ——, Quebec and Labrador, Report No. 51 (Preliminary Report No. 11), Washington, 1942
95. ——, Weekly Weather and Crop Bulletin, Washington, 1948 - 1956 [snow depth data for the United States]
96. U. S. Department of Interior, Bureau of Reclamation, Alaska Reconnaissance Report of the Potential Development of Water Resources in the Territory of Alaska, "Our Rivers" Series, House Document 197, 80th Congress, 1st Session, Washington, 1952
97. U. S. Navy Department, Bureau of Aeronautics, Project ARONA, Arctic Weather Maps (Berry, Owens and Wilson), Norfolk, Va., 1952
98. U. S. Navy Department, Hydrographic Office, Ice Atlas of the Northern Hemisphere, H. O. No. 550, 1st ed, Washington, 1946
99. ——, Naval Air Pilot, Greenland - Iceland Weather Summary, Supplement B to H. O. No. 252, GPO, prepared by the US Weather Bureau, Washington, 1943

100. ———, Naval Air Pilot, Nova Scotia, New Brunswick, Southeastern Quebec Weather Summary, H. O. No. 250, prepared by the US Weather Bureau, Washington, 1943

101. ———, Weather Summary, Alaska Area, H. O. No. 526, prepared by the US Weather Bureau, Washington, 1944

102. U. S. Navy Department, Office of the Technical Assistant to the Chief of Naval Operations for Polar Projects, Canadian North, OPNAV P03-4, Washington, 1956

103. Villeneuve, G.O., Climatic Conditions of the Province of Quebec and their Relationship to the Forests, Bulletin No. 6, Quebec Bureau of Meteorology, Montreal, 1946

PAGE 42 BLANK

9. Maps

- Figure 1. Station Location
- Figure 2. Major Physical Features
- Figure 3. Mean Temperature, Coldest Month
- Figure 4. Mean Daily Minimum Temperature, Coldest Month
- Figure 5. Absolute Minimum Temperature
- Figure 6. Mean Windchill, Coldest Month
- Figure 7. Mean Cloudiness, Coldest Month
- Figure 8. Mean Snow Depth, Month of Greatest Depth
- Figure 9. Mean Snowfall, Month of Greatest Snowfall
- Figure 10. Mean Temperature, Warmest Month
- Figure 11. Mean Daily Maximum Temperature, Warmest Month
- Figure 12. Mean Cloudiness, Warmest Month
- Figure 13. Mean Annual Precipitation
- Figure 13A. Mean Number of Days with Fog
- Figure 14. Composite Analogous Areas, Coldest Month
- Figure 15. Composite Analogous Areas, Fort Greely
- Figure 16. Composite Analogous Areas, Fort Churchill

Map Errata

Fig. 1 Bell Island (off Newfoundland) should read: Belle Isle
Skventna (southern Alaska) should read: Skventna

Fig. 2 Koskokvum Upland (central Alaska) should read: Kuskokwim Upland

Fig. 3 Tree Point (Alaskan Panhandle) - missing value is: 35J
Barter Island (Alaskan north coast) - value should be: -27F

Fig. 4 Lake Louise (Alberta) - value should be: -8

Fig. 5 In area of Amundsen Gulf (southeast of the Beaufort Sea) coastal area north of the -52° isotherm should be devoid of blue tint.

Fig. 10 In Siberia the blue tint should extend southward to include entire land area south of 64° N. latitude.

Fig. 13 Brynlandsfjord (Greenland north coast) - station missing and value is 2.3

Fig. 13A Quesnel (central British Columbia within island of dark blue tint) should be connected by a line to the following data printed on the map northeast of that station $\frac{5JA}{6J}$ $\frac{89}{158}$ and the station dot accompanying data should be removed

Jan Mayan (island in Greenland Sea) - station missing and data are $\frac{3MA}{10A}$ $\frac{58}{13J}$

Wiseman (central Alaska) - data should read $\frac{1JA}{1J}$ $\frac{16}{38}$

Eagle (eastern Alaskan boundary) - mean annual days should be 6
Yakutat (gulf coast of Alaska) - coldest month days should be 1JA

Taku Pass (near Juneau) - coldest month days should be <1JA

Dawson (Yukon) - coldest month days should be 3JA

Teslin (southern Yukon) - warmest month days should be 1J

Fort Norman (Mackenzie District, Canada) - coldest month should be January (JA)

Eureka (Ellesmere Island, Arctic Archipelago) - warmest month days should be <1J


Nord (north coast Greenland) - coldest month should be March (MA)

Prins Christians Sund (south coast Greenland) - coldest month days should be <1F


Godhavn (Disko Island, west Greenland) - warmest month days should be 17J

Tynemouth (east coast, England) - foggiest month should be January (JA)


CLIMATIC ANALYSIS OF FORT GREENE AND FORT CHICKADEE, AUGUST 1964



CLIMATIC ANALOGS OF FORT GREENE AND FORT CHARLES, NORTHERN CANADA



CLIMATIC ANALOGS OF FORT GREENE AND FORT CHURCHILL - NORTH AMERICA



ARCTIC OCEAN

BAFFIN BAY

MEAN DAILY MINIMUM TEMPERATURE IN FAHRENHEIT FOR THE MONTH OF JANUARY

FORT GREELY: 13 F FORT CHURCHILL: 27 F

Legend:

- 10 F and below
- 10 F to 15 F
- 15 F to 20 F
- 20 F to 25 F
- 25 F to 30 F
- 30 F and above

Source: Canadian Meteorological Service

Reliability: Fair to good

MEAN DAILY MINIMUM
TEMPERATURE
COLDEST MONTH

STANLEY WASSER

Method

County Auditor

Longstanding Goals
Are Achieved

THE

1

1

Abstract

1-800-828-6887

100%

21

1. *Staphylococcus aureus*

0 100 200 300 400 500 600 700 800 900 1000

CLIMATIC ANALOGS OF FORT GREELY AND FORT CHURCHILL - NOBIS AMERICA



MEAN JANUARY ISOTHERMS
PORT GREELY ETAH CHURCHILL


ISOTHERM	PORT GREELY	ETAH	CHURCHILL
50° F.	10.0	10.0	10.0
40° F.	10.0	10.0	10.0
30° F.	10.0	10.0	10.0
20° F.	10.0	10.0	10.0
10° F.	10.0	10.0	10.0
0° F.	10.0	10.0	10.0
-10° F.	10.0	10.0	10.0
-20° F.	10.0	10.0	10.0
-30° F.	10.0	10.0	10.0
-40° F.	10.0	10.0	10.0
-50° F.	10.0	10.0	10.0

Legend:
 - **PORT GREELY**: Indicated by a circle with a dot.
 - **ETAH**: Indicated by a circle with a cross.
 - **CHURCHILL**: Indicated by a circle with a plus sign.
 - **ISOTHERMS**: Lines showing mean January temperature in degrees Fahrenheit.

MEAN CLOUDINESS
CHILDREN MONTH
PORT GREELEY 55 TENTHS 
PORT CHURCHALL 47 TENTHS 

Cloud category (Tenths)	Strong winds	Cloud category	Sea swells
65 to 75	0	0	50 to 60
45 to 65	0	0	20 to 50
25 to 45	0	0	10 to 20
5 to 25	0	0	0 to 10

CLIMATIC ANALOGS OF FORT GREELY AND FORT CHURCHILL - NORTH AMERICA




CLIMATIC ANALOGS OF FORT GREELY AND FORT CHURCHILL - NORTH AMERICA




[illegible]

MEAN DAILY MAXIMUM
TEMPERATURE
WARMEST MONTH
FORT GREELY-69 F
CITY-64 F

CLIMATIC ANALOGS OF FORT GREELY AND FORT CHURCHILL - NORTH AMERICA



CLIMATIC ANALYSIS OF FORT GREELY AND FORT CHURCHILL



MEASUREMENT OF

THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

OF THE

100

CLIMATIC ANALOGS OF FORT GREEN AND FORT GREELY

ARCTIC OCEAN



CLIMATE
ANALOGS
FORT GREEN
FORT GREELY
FORT CHURCHILL
CLIMATE
ANALOGS
FORT GREEN
FORT GREELY
FORT CHURCHILL



CLIMATIC ANALYSIS OF FORT GREENE AND FORT GREENE

1901-1910



COMPOSITE
ANALYSIS OF
FORT GREENE

1. Average temperature
2. Average precipitation
3. Average humidity
4. Average wind speed
5. Average cloud cover
6. Average visibility
7. Average atmospheric pressure
8. Average relative humidity
9. Average dew point
10. Average frost depth

CLIMATIC ANALOGS OF FORT GREELY AND FORT CURCHILL - NORTH AMERICA

