

UNCLASSIFIED

AD NUMBER: AD0236174

LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited.

FROM:

Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; 1 Apr 1960. Other requests shall be referred to QUARTERMASTER RESEARCH AND ENGINEERING COMMAND NATICK MA

AUTHORITY

USAGREC ltr 18 May 1977 c/2 to a/1

UNCLASSIFIED

AD

236 174

Reproduced

Armed Services Technical Information Agency

ARLINGTON HALL STATION; ARLINGTON 12 VIRGINIA

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED

HEADQUARTERS

QUARTERMASTER RESEARCH & ENGINEERING COMMAND

U S ARMY

CATALOG OF NOTIA
AS AD No. 23617

XEROX

TECHNICAL REPORT

EP-132



CHANGES IN FOOT VOLUME AND CAPILLARY
FRAGILITY DURING CONTINUOUS COLD
EXPOSURE AND STARVATION



QUARTERMASTER RESEARCH & ENGINEERING CENTER
ENVIRONMENTAL PROTECTION RESEARCH DIVISION

APRIL 1960

NATICK, MASSACHUSETTS

HEADQUARTERS
QUARTERMASTER RESEARCH & ENGINEERING COMMAND, US ARMY
Quartermaster Research & Engineering Center
Natick, Massachusetts

ENVIRONMENTAL PROTECTION RESEARCH DIVISION

Technical Report
EP-132

CHANGES IN FOOT VOLUME AND CAPILLARY FRAGILITY
DURING CONTINUOUS COLD EXPOSURE AND STARVATION

Paul S. Norman, M. D.

Marlin B. Kreider, Ph. D.

P. F. Iampletro, Ph. D.

PHYSIOLOGY BRANCH

Project Reference:
7-83-01-006

April 1960

FOREWORD

PROTECTION OF THE FOOT AGAINST FROSTBITE IS IMPLICIT IN THE DESIGN AND DEVELOPMENT OF COLD-WEATHER FOOTGEAR. HOWEVER, IT IS KNOWN THAT FOOT INJURY MAY OCCUR UNDER CONDITIONS WHICH DO NOT PRODUCE FROSTBITE. A CONDITION CALLED "TRENCH FOOT" MAY OCCUR WHEN THE FEET ARE MAINTAINED AT COLD BUT STILL ABOVE-FREEZING TEMPERATURES FOR PROLONGED PERIODS OF TIME. PROPER EVALUATION OF THE PROTECTION AFFORDED BY ARCTIC FOOTGEAR REQUIRES THAT CONSIDERATION BE GIVEN TO TYPES OF FOOT INJURY OTHER THAN FROSTBITE. THIS REPORT DESCRIBES SOME OF THE CHANGES WHICH OCCUR IN FEET DURING PROLONGED EXPOSURE TO A COOL ENVIRONMENT IN WHICH FROSTBITE WOULD NOT BE A PROBLEM.

AUSTIN HENSCHEL, PH.D.
CHIEF
ENVIRONMENTAL PROTECTION RESEARCH
DIVISION

APPROVED:

PHILIP J. RORK, Lt COL, QMC
COMMANDING OFFICER
QM R&E CENTER LABORATORIES

DALE H. SIELING, PH.D.
SCIENTIFIC DIRECTOR
QM RESEARCH & ENGINEERING COMMAND

CONTENTS

	<u>PAGE</u>
ABSTRACT	iv
1. INTRODUCTION	1
2. PROCEDURE AND METHODS	1
3. RESULTS	3
A. FOOT VOLUME	3
B. CAPILLARY FRAGILITY	3
4. DISCUSSION	5
5. ACKNOWLEDGEMENT	7
6. REFERENCES	7

ABSTRACT

THE EFFECTS OF CONTINUOUS COLD EXPOSURE (2 WEEKS AT 60°F) AND STARVATION ON FOOT VOLUME AND CAPILLARY FRAGILITY WERE STUDIED IN 6 MEN. THE EXPERIMENTAL PERIOD WAS PRECEDED AND FOLLOWED BY 2-WEEK PERIODS (CONTROL AND RECOVERY) AT 80°F AND A CALORIC INTAKE OF 2,400 CALORIES. FOOT VOLUMES DECREASED 6.1 PERCENT IN THE FIRST 5 HOURS OF COLD EXPOSURE AND REMAINED CONSTANT FOR THE REMAINDER OF THE EXPERIMENTAL PERIOD. FOOT VOLUME WAS ALSO MEASURED AFTER 3 MINUTES OF ELEVATION OF THE FOOT AND OCCLUSION OF THE BLOOD SUPPLY. THIS VOLUME DECREASED LESS DURING THE EXPERIMENTAL THAN THE PRECEDING OR FOLLOWING PERIODS. CAPILLARY FRAGILITY DECREASED SLOWLY DURING THE EXPERIMENTAL PERIOD AND INCREASED IN THE RECOVERY PERIOD. ON THE THIRD DAY OF RECOVERY, FOOT VOLUMES WERE LARGER THAN AT ANY OTHER TIME. FOR THE REMAINDER OF RECOVERY, FOOT VOLUME WAS THE SAME AS THE CONTROL VOLUME, BUT THERE WAS A GREATER DECREASE IN VOLUME WITH FOOT ELEVATION. THE RESULTS SUGGEST THAT THERE WAS A CONSTRICTION OF ALL TYPES OF BLOOD VESSELS DURING THE EXPERIMENTAL PERIOD, WHILE IN THE RECOVERY PERIOD THERE WAS APPARENTLY A SIMULTANEOUS DILATATION OF LARGE BLOOD VESSELS AND A CONSTRICTION OF SMALL ONES. ONE SUBJECT HAD A GREATLY INCREASED VOLUME OF ONE FOOT ASSOCIATED WITH PAIN AND REDNESS DURING EARLY RECOVERY DAYS. THESE CHANGES MAY BE INDICATIVE OF EARLY TRENCH FOOT OR CAPILLARY DAMAGE. STARVATION PER SE APPEARED TO HAVE NO EFFECT ON FOOT VOLUME.

CHANGES IN FOOT VOLUME AND CAPILLARY FRAGILITY DURING CONTINUOUS COLD EXPOSURE AND STARVATION

1. INTRODUCTION

SUBJECTS EXPOSED TO A COLD ENVIRONMENT (NUDE AT 60°F) FOR PROLONGED PERIODS OF TIME OFTEN COMPLAINED OF TRANSIENT FOOT PAIN AND TENDERNESS DURING REWARMING (1, 2). THE POSSIBILITY OF MILD TRENCH FOOT WAS CONSIDERED BUT WAS NOT INVESTIGATED. TRENCH FOOT HAS BEEN A COMMON OCCURRENCE IN CHRONIC NON-FREEZING COLD ENVIRONMENTS (3, 4, 5). EDEMA FORMATION AND INCREASED CAPILLARY PERMEABILITY ARE BOTH FOUND IN MILD TRENCH FOOT (6). CAPILLARY CHANGES DUE TO LOCAL HYPOTHERMIA HAVE BEEN DEMONSTRATED BY LEWIS AND LOVE (7) AND WILLMAN AND HANLON (8).

RECENTLY, THE OPPORTUNITY WAS AFFORDED TO STUDY 6 MEN¹ LIVING NUDE FOR 2 WEEKS AT 60°F. DURING THIS PERIOD NO FOOD WAS CONSUMED. FOOT VOLUMES AND CAPILLARY FRAGILITY OF THE DORSUM OF THE FOOT WERE MEASURED TO ASCERTAIN IF A PATTERN SIMULATING MILD TRENCH FOOT WAS PRESENT.

2. PROCEDURE AND METHODS

SIX HEALTHY MEN, AGED 18 TO 27 YEARS, WEARING ONLY COTTON SHORTS, LIVED IN A TEMPERATURE-CONTROLLED ROOM FOR 6 WEEKS. THEIR ACTIVITIES WERE SEDENTARY, E.G., CARD PLAYING, READING, WRITING, WATCHING MOVIES AND TELEVISION. THE FIRST 2 WEEKS SERVED AS A CONTROL PERIOD WITH ROOM TEMPERATURE AT 80°F, RELATIVE HUMIDITY 40-50%, WIND LESS THAN 1 MPH. THEIR DIET (2400 CALORIES PER DAY) CONSISTED OF AN ENRICHED MILK DRINK, BUTTERED TOAST, AND COFFEE. THE SECOND 2-WEEK PERIOD WAS THE EXPERIMENTAL PERIOD WITH TEMPERATURE AT 60°F, RELATIVE HUMIDITY 40-50%, AND WIND LESS THAN 1 MPH. DIET DURING THIS PERIOD WAS NON-CALORIC (WATER AD LIBITUM AND BLACK COFFEE). THE CONDITIONS DURING THE THIRD 2-WEEK PERIOD (RECOVERY PERIOD) WERE THE SAME AS THOSE OF THE CONTROL PERIOD. IN ALL PERIODS, EACH SUBJECT WAS ALLOWED 1 WOOLEN ARMY BLANKET DURING THE SLEEPING HOURS.

MEASUREMENTS OF FOOT VOLUME WERE MADE 16 TIMES DURING THE 6 WEEKS (4 TIMES IN CONTROL, 6 TIMES IN EXPERIMENTAL, AND 6 TIMES IN RECOVERY). VOLUME OF THE FOOT WAS ESTIMATED BY MEASURING THE WATER DISPLACED BY THE FOOT IN A METAL BOOT. MEASUREMENTS WERE MADE WITH THE SUBJECT ERECT AND WITH HIS WEIGHT PLACED EQUALLY ON BOTH FEET. THE AMOUNT OF WATER IN THE BOOT BEFORE IMMERSION OF THE FOOT WAS KEPT CONSTANT. THE WATER WAS ALWAYS AT ROOM TEMPERATURE (80°F CONTROL AND RECOVERY, 60°F EXPERIMENTAL PERIOD). THE LOWER EXTREMITIES WERE KEPT IN A HORIZONTAL POSITION FOR 15 MINUTES PRIOR TO MEASUREMENT. THE VOLUME OF THE RIGHT FOOT WAS MEASURED. THE RIGHT FOOT WAS THEN SUPPORTED AT AN ELEVATION OF 18 INCHES FOR 3 MINUTES WITH THE SUBJECT SUPINE. WITH THE FOOT STILL ELEVATED, A PRESSURE OF




FIGURE 1. PERCENT CHANGE OF FOOT VOLUME OF 6 MEN DURING COLD-STARVATION AND RECOVERY PERIODS FROM PRECEDING CONTROL PERIOD.




FIGURE 2. CAPILLARY FRAGILITY OF 6 MEN DURING CONTROL, COLD-STARVATION, AND RECOVERY PERIODS. THE ORDINATE IS TIME IN MINUTES REQUIRED TO PRODUCE 12 PETECHIAE ON THE DORSUM OF THE RIGHT FOOT.

200 MM HG WAS APPLIED WITH A TOURNIQUET 3-4 INCHES BELOW THE KNEE TO CAUSE ARTERIAL OCCLUSION. THE FOOT VOLUME WAS AGAIN MEASURED WITH THE SUBJECT IN THE ERECT POSITION AND THE TOURNIQUET IN PLACE.

CAPILLARY FRAGILITY WAS MEASURED BY MEANS OF A SUCTION PETECHIOMETER SIMILAR TO THE APPARATUS OF KRAMAR (9). THE MAXIMAL NEGATIVE PRESSURE OF OUR APPARATUS WAS 60 CM HG. THE TIME REQUIRED TO PRODUCE 12 PETECHIAE UNDER THE SUCTION CUP (INSIDE DIAMETER OF SUCTION CUP 15 MM) WAS TAKEN AS THE END POINT. THIS VARIED FROM 0.5 TO 8.0 MINUTES WITH THE SUCTION CONSTANT AT 60 CM HG. MEASUREMENTS WERE TAKEN AFTER THE MEN HAD BEEN SUPINE FOR AT LEAST 15 MINUTES. FOUR AREAS ON THE MEDIAL SIDE OF THE DORSUM OF THE RIGHT FOOT, OVER THE METATARSAL BONES, WERE ROTATED AS THE TESTING SITE, WITH NO SITE BEING USED MORE THAN ONCE IN 48 HOURS.

3. RESULTS

A. FOOT VOLUME

AVERAGE FOOT VOLUME DURING THE CONTROL PERIOD WAS 1,279 CC. AFTER 5 HOURS IN THE COLD, FOOT VOLUME WAS 1,200 CC. THIS REPRESENTS A DECREASE IN THE COLD (FIG. 1) OF 6.1%. ALL THE MEN SHOWED THIS PATTERN (RANGE OF DECREASE 3.9% TO 7.3%). FOOT VOLUME DURING THE REMAINDER OF THE COLD PERIOD REMAINED ESSENTIALLY UNCHANGED AFTER THE FIRST 5 HOURS.

THE MEAN DECREASE IN FOOT VOLUME WITH ELEVATION AND ARTERIAL OCCLUSION WAS 29 CC (CONTROL), 14 CC (EXPERIMENTAL), AND 41 CC (RECOVERY); THE 3 PERIODS DIFFERED SIGNIFICANTLY FROM EACH OTHER. THE VOLUME OF THE FOOT AFTER ELEVATION AND ARTERIAL OCCLUSION WAS LESS DURING THE EXPERIMENTAL PERIOD THAN DURING CONTROL AND RECOVERY PERIODS (TABLE 1). THE VOLUME DURING RECOVERY AVERAGED 15 CC LOWER THAN THE CONTROL VOLUME (CONTROL, 1,250 CC; EXPERIMENTAL, 1,186 CC; AND RECOVERY, 1,235 CC).

FOOT VOLUME DURING RECOVERY DID NOT RETURN TO CONTROL VALUES UNTIL 3 DAYS HAD PASSED. FIVE HOURS AFTER LEAVING THE COLD ROOM, 5 OF THE 6 SUBJECTS' FOOT VOLUMES WERE MIDWAY BETWEEN CONTROL AND EXPERIMENTAL VALUES, BUT THE SIXTH SUBJECT HAD AN INCREASE OF 44 CC ABOVE HIS HIGHEST CONTROL VALUE. THE RIGHT FOOT OF THIS SUBJECT WAS PAINFUL, RED, AND SUBJECTIVELY WARM DURING THE FIRST 3 DAYS OF RECOVERY. ON THE THIRD DAY OF RECOVERY, THE AVERAGE FOOT VOLUME FOR 6 SUBJECTS WAS 24 CC ABOVE THE AVERAGE FOR THE CONTROL DAYS AND 29 CC ABOVE THE AVERAGE FOR THE REMAINING RECOVERY VALUES. FOOT VOLUME AFTER ELEVATION AND OCCLUSION ON THE THIRD DAY OF RECOVERY WAS 19 CC GREATER THAN ON ANY OTHER RECOVERY DAY. FROM THE FOURTH RECOVERY DAY UNTIL TERMINATION OF THE STUDY, FOOT VOLUME OF ALL THE MEN REMAINED UNCHANGED AT APPROXIMATELY CONTROL VALUES.

B. CAPILLARY FRAGILITY

CAPILLARY FRAGILITY ON THE DORSUM OF THE RIGHT FOOT DECREASED DURING THE EXPERIMENTAL PERIOD AND ROSE AGAIN DURING RECOVERY (FIG. 2).

TABLE 1: MEAN FOOT VOLUME OF 6 MEN FOLLOWING 15 MINUTES OF RECLINING (WITHOUT OCCLUSION) AND 3 MINUTES OF FOOT ELEVATION (WITH OCCLUSION)

<u>DAY OF PERIOD</u>	<u>VOLUME WITHOUT OCCLUSION</u> cc	<u>VOLUME WITH OCCLUSION</u> cc
CONTROL		
3	1274	1250
8	1272	1238
10	1281	1253
12	1288	1257
MEAN	1279	1250
COLD		
1	1203	1198
3	1197	1183
5	1209	1196
8	1198	1182
10	1202	1188
12	1191	1177
MEAN	1200	1186
RECOVERY		
1	1256	1218
3	1303	1257
4	1272	1230
5	1278	1239
8	1275	1235
10	1272	1228
MEAN	1276	1235

THE MEAN TIME TO REACH THE END POINT (FORMATION OF 12 PETECHIAE) WAS 2.3 MINUTES IN THE CONTROL, 5.2 MINUTES IN THE EXPERIMENTAL, AND 4.2 MINUTES IN THE RECOVERY PERIODS. THERE WAS MARKED INDIVIDUAL VARIATION IN THE RATE OF CHANGE OF CAPILLARY FRAGILITY IN BOTH THE EXPERIMENTAL AND RECOVERY PERIODS. THE END POINT DURING RECOVERY NEVER RETURNED TO CONTROL VALUES. SUBJECT D, WHO HAD A SWOLLEN RIGHT FOOT ON THE FIRST RECOVERY DAY, DEMONSTRATED MARKED CAPILLARY FRAGILITY (12 PETECHIAE PRESENT IN 1/2 MINUTE WITH 60 CM Hg) ON THIS DAY OF RECOVERY. THE MEAN TIME FOR THE REMAINDER OF THE GROUP (I.E., EXCLUDING SUBJECT D) TO REACH THE END POINT ON THE FIRST RECOVERY DAY WAS 5.15 MINUTES. SUBJECT D HAD A POSITIVE END POINT IN LESS THAN 1 MINUTE THROUGHOUT THE RECOVERY PERIOD, WHEREAS THE MEAN END POINT FOR THE OTHER MEN WAS 4.8 MINUTES.

4. DISCUSSION

NO PATTERN SIMULATING TRENCH FOOT WAS FOUND IN 5 OF THE 6 SUBJECTS, ALTHOUGH 2 SUBJECTS COMPLAINED OF FOOT PAIN AND TENDERNESS DURING THE FIRST 2 DAYS OF RECOVERY. THE 6TH SUBJECT, SUBJECT D, HOWEVER, HAD THE FINDINGS OF EARLY, MILD TRENCH FOOT.

FOOT VOLUME DECREASED WITHIN 5 HOURS AFTER ENTERING THE COLD CONDITIONS AND REMAINED UNCHANGED DURING THE REMAINDER OF THE COLD PERIOD. DURING THE EARLY RECOVERY PERIOD, FOOT VOLUME EXCEEDED THE CONTROL VOLUME AND LATER RETURNED TO CONTROL VALUES. THE SMALLER, CONSTANT FOOT VOLUME IN THE EXPERIMENTAL PERIOD CAN BE EXPLAINED BY A GENERAL VASCULAR CONSTRICTION. THREE MINUTES OF ELEVATION CAUSED A FURTHER REDUCTION IN FOOT VOLUME DUE TO A DECREASED AMOUNT OF INTRAVASCULAR FLUID. A FOOT ELEVATED FOR THIS TIME SHOULD LOSE POOLED VENOUS BLOOD AND POSSIBLY HAVE A DECREASED ARTERIAL INFLOW, BUT SHOULD NOT LOSE ANY EXTRAVASCULAR FLUID. SINCE THERE WAS A GREATER CHANGE IN FOOT VOLUME WITH ELEVATION DURING CONTROL AND RECOVERY PERIODS THAN DURING THE EXPERIMENTAL PERIOD, A GREATER AMOUNT OF BLOOD WAS PROBABLY PRESENT IN THE LARGE VESSELS DURING THE WARM CONDITION. ONE OF THE FIRST REACTIONS TO COLD IS PERIPHERAL VASOCONSTRICTION (5), WHICH WOULD REDUCE ARTERIAL AND ARTERIOLAR FLOW TO THE FEET. WOOD, BASS, AND LAMPIETRO (10) FOUND DECREASED BLOOD FLOW TO THE ARM, AS WELL AS VENOUS CONSTRICTION, WITH EXPOSURE AT 60°F AND NO FURTHER SIGNIFICANT CHANGES AS THE COLD PERIOD CONTINUED. IT IS LIKELY THAT SIMILAR CHANGES WOULD OCCUR IN THE FEET, THUS REDUCING VENOUS POOLING IN THE FEET AND PRODUCING SMALLER FOOT VOLUMES IN THE COLD.

CAPILLARY FRAGILITY MEASURES THE CAPACITY OF THE CAPILLARY WALL TO RESIST INCREASED INTRACAPILLARY PRESSURE OR EXTRACAPILLARY SUCTION (11). THE CAPILLARY WALLS CONSIST OF A SINGLE LAYER OF ENDOTHELIAL CELLS HELD TOGETHER BY CEMENT SUBSTANCE. SUBSTANCES PASSING THROUGH THE CAPILLARY WALL MUST PASS THROUGH EITHER THE ENDOTHELIAL CELL OR THE CEMENT SUBSTANCE (12). WHEN CAPILLARIES CONSTRICT, THEIR FLAT CELLS MIGHT BECOME MORE RECTANGULAR AND THE CEMENT SUBSTANCE, THEREFORE, MIGHT HAVE A

GREATER DEPTH FROM INTRACAPILLARY LUMEN TO CAPILLARY EXTERIOR. SANDERS, EBERT, AND FLORY (13) OBSERVED CAPILLARY CONTRACTION AS A RESULT OF SYMPATHETIC STIMULATION WITH SWELLING IN THE REGION OF THE ENDOTHELIAL CELL NUCLEI WHICH OCCLUDED THE LUMEN OF THE VESSEL. INCREASED CEMENT SUBSTANCE DEPTH OR INCREASED ENDOTHELIAL CELL WIDTH CAUSED BY SWELLING OF THE ENDOTHELIAL CELL COULD ACCOUNT FOR DECREASED CAPILLARY FRAGILITY DURING THE COLD PERIOD.

FOOT VOLUME ON THE THIRD RECOVERY DAY WAS GREATER THAN AT ANY OTHER TIME DURING RECOVERY, BUT THE DECREASE IN VOLUME AFTER ELEVATION AND OCCLUSION WAS ESSENTIALLY THE SAME AS ON ANY OTHER RECOVERY DAY. THIS INDICATED THAT THE INCREASED VOLUME OF THE FOOT ON THE THIRD DAY WAS PROBABLY DUE TO AN INCREASE IN EXTRAVASCULAR FLUID (EDEMA) RATHER THAN INCREASED VOLUME OF BLOOD IN THE VESSELS OF THE FOOT. THIS EXTRAVASCULAR FLUID HAD DISAPPEARED BY THE FOURTH RECOVERY DAY. SINCE THE FOOT AFTER ELEVATION AND ARTERIAL OCCLUSION REMAINED SMALLER DURING RECOVERY (AFTER THE THIRD DAY) THAN DURING CONTROL, AND SINCE CAPILLARY FRAGILITY INCREASED ONLY SLOWLY IN RECOVERY, IT IS POSSIBLE THAT CAPILLARY AND OTHER SMALL VESSEL CONSTRICTION CONTINUED INTO THE WARM CONDITION. THUS, IT IS POSSIBLE THAT A PARADOX OF DILATATION OF THE LARGE VESSELS AND CONSTRICTION OF THE CAPILLARIES MAY EXIST DURING RECOVERY.

THREE SUBJECTS COMPLAINED OF FOOT PAIN AND TENDERNESS ON THE FIRST RECOVERY DAYS, BUT 2 OF THESE SUBJECTS EXHIBITED NO OBJECTIVE FINDINGS. ANOTHER SUBJECT, ALONG WITH THE FINDINGS OF REDNESS, WARMTH, AND GREATLY INCREASED RIGHT FOOT VOLUME, ALSO HAD MARKEDLY INCREASED CAPILLARY FRAGILITY. THIS COULD INDICATE THAT IN ADDITION TO THE INCREASE IN LARGE-VESSEL BLOOD VOLUME, THE CAPILLARIES HAD AN INCREASED VOLUME AND WERE THINNER-WALLED OR THAT CAPILLARY DAMAGE OCCURRED DURING THE EXPERIMENTAL PERIOD. THE SIGNS OF EARLY TRENCH FOOT, INCLUDING INCREASED CAPILLARY PERMEABILITY WITH EXTRAVASATION OF EDEMA FLUID (6), WERE PRESENT IN THIS SUBJECT (SUBJECT D).

THE CHANGES IN FOOT VOLUME DURING THE EXPERIMENTAL PERIOD WERE PROBABLY A RESULT OF EXPOSURE TO COLD AND WERE NOT INFLUENCED BY STARVATION. THUS, THE FOOT ATTAINED ITS SMALLEST VOLUME WITHIN 5 HOURS IN THE COLD AND MAINTAINED THIS VOLUME FOR THE 2-WEEK PERIOD. AT THE TIME OF THE FIRST MEASUREMENT IN THE EXPERIMENTAL PERIOD ONLY ONE MEAL, BREAKFAST, HAD BEEN OMITTED. AFTER 14 DAYS OF A NON-VITAMIN, NON-CALORIC DIET, NO FURTHER CHANGES IN FOOT VOLUME WERE NOTED.

THE RELATIVE EFFECTS OF COLD AND STARVATION ON CAPILLARY FRAGILITY ARE MORE DIFFICULT TO EVALUATE. CAPILLARY FRAGILITY HAD DECREASED MARKEDLY BY THE SECOND DAY OF THE EXPERIMENTAL PERIOD (FIG. 2) AND THEN DECREASED MORE SLOWLY DURING THE REMAINDER OF THE PERIOD. IT WOULD APPEAR THAT THE EARLY RAPID CHANGE IN CAPILLARY FRAGILITY WAS THE RESULT OF COLD EXPOSURE AND THAT THE LATER, SLOWER CHANGE WAS THE RESULT OF STARVATION OR STARVATION AND COLD.

5. ACKNOWLEDGEMENT

THE AUTHORS THANK MR. EDWARD B. GREEN FOR SUGGESTING THAT CAPILLARY FRAGILITY BE STUDIED.

6. REFERENCES

1. IAMPINETRO, P.F., D.E. BASS AND E.R. BUSKIRK. J. APPL. PHYSIOL. 10:398 (1957).
2. KREIDER, M.B., P.F. IAMPINETRO, E.R. BUSKIRK AND D.E. BASS. J. APPL. PHYSIOL. 14:43 (1959).
3. BLAIR, J.R. TRANSACTIONS OF THE FOURTH JOSIAH MACY, JR. FOUNDATION, 1955, P. 9.
4. BURCH, G.E., H.L. MEYERS, R.R. PORTER AND N. SCHAFFER. JOHNS HOPKINS HOSPITAL BULLETIN 80:1 (1947).
5. ORR, K.D. AND D.C. FAINER. MEDICINE 31:177 (1952).
6. WHITE, J.C. AND W.B. SCOVILLE. NEW ENGLAND J. MED. 232:415 (1945).
7. LEWIS, T. AND W.S. LOVE. HEART 13:27 (1926).
8. WILLMAN, V.L. AND C.R. HANLON. ANN. SURG. 143:660 (1956).
9. KRAMÁR, J., V.W. MEYERS AND M. SIMAY KRAMÁR. J. LAB. & CLIN. MED. 47:423 (1956).
10. WOOD, J.E., D.E. BASS AND P.F. IAMPINETRO. J. APPL. PHYSIOL. 12:357 (1958).
11. KRAMÁR, J. AM. J. PHYSIOL. 175:69 (1953).
12. HAM, A.W. HISTOLOGY (2ND IMPRESSION). PHILADELPHIA, LIPPINCOTT, 1950, PP 89-90.
13. SANDERS, A.G., R.H. EBERT AND H.W. FLOREY. QUART. J. EXPER. PHYSIOL. 30:281 (1940).

DISTRIBUTION LIST

<u>ARMY (Cont)</u>	<u>ARMY (Cont)</u>	<u>ARMY (Cont)</u>	<u>NAVY (Cont)</u>
<p>1. Deputy Chief of Staff for Logistics Department of the Army Washington 25, D. C.</p> <p>1. Deputy Chief of Staff for Personnel Department of the Army Washington 25, D. C.</p> <p>1. Deputy Chief of Staff for Military Operations, Department of the Army Washington 25, D. C.</p> <p>1. Chief of Research & Development Department of the Army Washington 25, D. C.</p>	<p>1. Commanding Officer Signal Corps Engr. Lab. Ft. Monmouth, N. J.</p> <p>1. Office of Chief of Engineers Department of the Army Temp. Bldg. T-7, Gravelly Point Washington 25, D. C.</p> <p>4. CO, Chemical Warfare Laboratories Army Chemical Center, Maryland AHC Terminal (AR 19) Library</p> <p>1. Chief Chemical Officer Department of the Army Bldg. T-7, Gravelly Point Washington 25, D. C.</p> <p>1. CO, Hq., Medical Nutrition Lab. Florence Army Hospital Denver, Colorado (Dr. Friedman)</p> <p>1. Armed Forces Institute of Pathology Washington 25, D. C.</p> <p>1. CO, U. S. Armed Services Medical Development Agency 1101 E. St., Brooklyn 1, N. Y. Attn: Property Officer March 1. Post. DUES 811</p> <p>1. Chief of Transportation Department of the Army Temp. Bldg. T-7, Gravelly Point Washington 25, D. C.</p> <p>1. Commanding Officer Transportation Res. & Eng. Command U. S. Army Ft. Belvoir, Virginia Attn: Tech Services Div.</p> <p>1. The Army Library Fenton Bldg. Washington 25, D. C.</p> <p>1. Commanding, Chemical & General Staff College Ft. Leavenworth, Kansas</p> <p>1. Commanding, U. S. Military Academy West Point, New York</p> <p>1. Commanding Officer, Detroit Arsenal Shell Van Dyke Bld., Centerline, Mich. Attn: Res. & Engr. Div.</p> <p>1. Commanding General Hq., U. S. Army Medical R&D Command Main Navy Bldg. Washington 25, D. C. Attn: NRDPP Research Branch</p> <p>2. Commander QM Intelligence Agency, U. S. Army Washington 25, D. C.</p> <p>2. Executive Director Military Clothing and Textile Supply Agency 1990 S. 10th St., Philadelphia, Pa.</p> <p>1. Commanding Officer QM H&E Field Evaluation Agency, U. S. Army Aircraft Systems Test Div. Yuma Test Station Yuma, Arizona</p>	<p>1. Commanding Officer Cold Weather & Mountain Indoctrination School Fort Greeley, Alaska</p> <p>1. Commanding Officer Fort Greeley, Alaska 6th Post Library</p> <p>1. Department of Air Force Hq., USAF, Wash 25, D. C. (1 DC/S Material, 1 DC/S Dev.)</p> <p>1. Director Air University Library, Attn: 7875 Maxwell AFB, Alabama</p> <p>1. Commandant USAF School of Aviation Medicine Randolph AFB Base Randolph Field, Texas</p> <p>1. Commander, Arctic Ascomedical Lab APO 77, Seattle, Washington</p> <p>1. Commander Air Res. & Dev. Command Attn: RCDPTL Hq., Tech Lib. Br.) Address AF Base, Washington 25, D. C.</p> <p>1. Commander Wright Air Development Center Wright Patterson AF Base, Ohio Attn: Tech Library</p> <p>1. Commander Strategic Air Command Offutt AF Base, Nebraska</p> <p>1. Chief, Nutrition Div. Air Development Center Aero-Medical Lab. Wright Patterson AFB, Ohio Attn: Dr. Harry C. Lynn</p> <p>1. Commander AF Cambridge Research Center Air Research & Development Ctr. Lawrence G. Hanscom Field Bedford, Mass. Attn: DRGOTT-8</p>	<p>1. Commander, U. S. Naval Ord. Test Station, Chris Lake, Calif. Attn: Code 453</p> <p>1. Chief, Bureau of Aeronautics Dept. of the Navy, Wash 25, D. C. Attn: Code AF 12</p> <p>1. Chief, Bureau of Supplies & Accounts Department of the Navy Washington 25, D. C.</p>
<p><u>ARMY</u></p> <p>1. The Quartermaster General Department of the Army Washington 25, D. C.</p> <p>1. Commanding General Florence Army Hospital 1000 West 10th Street Florence, Colorado</p> <p>1. Commanding QM Property Control Institute for the Armed Forces, U. S. Army 100 W. Pennsylvania Washington 25, D. C.</p> <p>1. Commanding Officer QM Intelligence Agency, U. S. Army Washington 25, D. C.</p> <p>1. Commanding Officer, ACCG Wright Patterson Development Center Wright Patterson AF Base Dayton, Ohio</p> <p>1. Commanding Ft. Belvoir, Virginia Attn: Library</p> <p>1. Commanding General Bradford Arsenal, Phils 37, Pa. Attn: Engr. Technology Div. (LL)</p> <p>1. Hq., Army Electronic Proving Ground Ft. Huachuca, Arizona Attn: Aviation & Meteorological Depts. Tech. Information Br. Deputy Chief for Meteorology</p> <p>1. Commanding General The Engineer Center Ft. Belvoir, Va.</p> <p>1. Commanding Officer Personnel Ordnance Proc. Lab. Washington 25, D. C. Attn: Tech Reference Section (ORVT-204)</p> <p>1. Commanding General Aberdeen Proving Ground Aberdeen, Maryland</p> <p>1. Chief Signal Officer Department of the Army Washington 25, D. C. Attn: Res. & Dev. Div.</p>	<p><u>NAVY</u></p> <p>1. Director Hull Pattern Laboratory Hull Ordnance Bldg. 20, S. W. Washington 25, D. C.</p> <p>1. Chief, Bureau of Ordnance Department of the Navy Washington 25, D. C. Attn: R.D. Div.</p> <p>1. Naval Medical Research Institute National Naval Medical Center Bethesda, Md.</p> <p>1. Chief of Naval Research Washington 25, D. C. Attn: Code 400</p> <p>1. Chief, Bureau of Ships Department of the Navy Washington 25, D. C. Attn: Code 311</p> <p>1. Chief, Bureau of Med. & Surgery Dept. of the Navy, Wash 25, D. C. Attn: Code 30</p>	<p><u>CONARC</u></p> <p>1. C.O., U.S. Continental Army Ctr. H. H. Hays, Va.</p> <p>1. President U. S. Army Artillery Bn. Ft. Belvoir, Va. Attn: ATFA</p> <p>1. President US Army Armor Post Ft. Knox, Ky. Attn: ATFB</p> <p>1. President U. S. Army Infantry Bn. Ft. Belvoir, Va. Attn: ATFC</p> <p>1. President U. S. Army Air Defense Bn. Ft. Belvoir, Va. Attn: ATFD</p> <p>1. President U. S. Army Artillery and Mortar Bn. Ft. Belvoir, Va. Attn: ATFA</p> <p>1. President U. S. Army Cavalry Bn. Ft. Belvoir, Va. Attn: ATFC</p> <p>1. Commanding Officer U. S. Army Artillery Test Board H. H. Hays, Va. Attn: ATFA</p>	
<p><u>MICROFILM NEG.</u></p> <p>1. National Research Council 11 Constitution Ave., Washington, D.C. Attn: Advisory Bd. on QM 154</p> <p>1. Armed Services Technical Information Agency Arlington Hall Station Arlington, Va. Attn: THDR</p> <p>1. Gift and Exchange Division Library of Congress Washington 25, D. C.</p>	<p>1. U. S. Department of Commerce Weather Bureau Library, Washington, D. C.</p> <p>1. Central Intelligence Agency Collection & Dissemination Washington 25, D. C.</p> <p>1. National Library of Medicine Washington 25, D. C.</p> <p>1. General Reference State Historical Society Fenton, Mo. Ohio, N. Y.</p>	<p>1. Mater. Equip. Division 1 Mater. Development Center Main Corps Library Quantity 7.</p> <p>1. Office of Technical Services U. S. Department of Commerce Washington 25, D. C. Attn: Tech Equip. Sec. (THRU O&M)</p>	<p>1. U. S. Department of Agriculture Library Washington 25, D. C.</p> <p>1. Commanding Industrial College of the Armed Forces Ft. McNair, Washington 25, D. C.</p> <p>1. QM Reference Army Command and Staff College Department of the Army, Wash 25, D. C. Ft. Leavenworth, Kansas</p>

UNCLASSIFIED

AD

236 174

Reproduced

Armed Services Technical Information Agency

ARLINGTON HALL STATION; ARLINGTON 12 VIRGINIA

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OF OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED