

UNCLASSIFIED

AD 243 118

*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGED BY ASTIA

SMITHSONIAN CONTRIBUTIONS TO ASTROPHYSICS



Smithsonian Institution
Astrophysical Observatory

Volume 2, Number 7

**Distribution of
Meteoritic Debris
about the Arizona
Meteorite Crater**

ASTIA
SEP 27 1960

XEROX
Ln 60-4-4

88 000

Washington, D. C.

Smithsonian
Contributions to Astrophysics

VOLUME 2, NUMBER 7

DISTRIBUTION OF METEORITIC DEBRIS
ABOUT THE
ARIZONA METEORITE CRATER

by JOHN S. RINEHART



SMITHSONIAN INSTITUTION

Washington, D. C.

1958

Publications of the Astrophysical Observatory

This series, *Smithsonian Contributions to Astrophysics*, was inaugurated in 1956 to provide a proper communication for the results of research conducted at the Astrophysical Observatory of the Smithsonian Institution. Its purpose is the "increase and diffusion of knowledge" in the field of astrophysics, with particular emphasis on problems of the sun, the earth, and the solar system. Its pages are open to a limited number of papers by other investigators with whom we have common interests.

Another series is *Annals of the Astrophysical Observatory*. It was started in 1900 by the Observatory's first director, Samuel P. Langley, and has been published about every 10 years since that date. These quarto volumes, some of which are still available, record the history of the Observatory's researches and activities.

Many technical papers and volumes emanating from the Astrophysical Observatory have appeared in the *Smithsonian Miscellaneous Collections*. Among these are *Smithsonian Physical Tables*, *Smithsonian Meteorological Tables*, and *World Weather Records*.

Additional information concerning these publications may be obtained from the Editorial and Publications Division, Smithsonian Institution, Washington 25, D. C.

FRED L. WHIPPLE, *Director,*
Astrophysical Observatory,
Smithsonian Institution.

Cambridge, Mass.

Distribution of Meteoritic Debris About the Arizona Meteorite Crater¹

By John S. Rinehart²

The meteorite crater in northern Arizona, an outstanding topographic feature of the earth's surface, is the earth's largest authenticated meteorite crater. It is a large bowl-like depression lying in a sandy semiarid region of northern Arizona which is readily accessible by car. In outline, the crater is a rough square, about 4,100 feet across and 600 feet deep, with an elevated rim rising 160 feet above the surrounding plain.

The crater has been well known since 1870. By 1909 it had been exhaustively described (Gilbert, 1896; Barringer, 1905, 1909; Tilghman, 1905; Merrill, 1908) and most serious investigators agreed with the view that it had been blasted out by the impact of a large meteorite. Numerous surveys have been made since then with various objectives. The chief surveys were those of Barringer (1914) and Barringer, Jr. (1927), who were intent upon locating and, if possible, recovering the large meteorite that made the crater; and of Nininger (1951; 1956).

In a recent book Nininger (1956) has given an excellent description of the researches and surveys made since the discovery of the crater, and has critically reviewed all of the findings to date. He also speculates on the nature of the event that took place at the time of the earth's encounter with the meteorite. When did it strike? How fast was it moving? From what direction did it come? How large was it? Was it a single large meteorite or a swarm of meteorites? Speculations of Barringer (1914),

¹ This work was partially supported by Contract No. AF19(604)-1901 with the Geophysics Research Directorate, Air Force Cambridge Research Center, Air Research and Development Command.

² Smithsonian Astrophysical Observatory and Harvard College Observatory, Cambridge, Mass.

Nininger (1956), and others are inconclusive in many instances because of the great paucity of data.

The object of the survey reported here was to make a careful, systematic investigation of the distribution of the minuscule bits and pieces of meteoritic material that are scattered through the mantle of soil surrounding the crater, with a view to fixing more closely the mass of the meteorite that made the crater and its direction of flight. Our study arose from a suggestion by Nininger (1951), whose exploratory survey of the distribution of this material indicated that the amount might total several thousand tons, a weight many times greater than the 20 to 30 tons of meteorites that have been picked up in the surrounding territory thus far.

To carry out this survey, the Smithsonian Astrophysical Observatory sent an expedition into the field during the summer of 1956. This paper describes the results obtained.

Processing the soil sample

The expedition collected and processed some 700 soil samples. Developing techniques to extract the meteoritic material from the samples was a major problem. Nininger's method had employed a magnet, dragged through the soil. The adhering magnetic material was then collected from the magnet and separated by hand, or by the use of an inclined surface which divided spherical particles from those of more irregular shape. A disadvantage of this technique is that it collects both weakly and strongly magnetic material, although none of the weakly magnetic material has been found

to be of meteoritic origin. Such material, for example, never gives a positive test for nickel.

Since it seemed unwise to collect such material even in a preliminary extraction, and since we were interested only in the total amount of meteoritic material present whether its shape was flaky, chunky, or spheroidal, we did not use Nininger's method, but developed a special magnetic separator which is shown schematically in figure 1.

samples sometimes varied by several hundred grams. This volume of dirt can be put conveniently into two paper quart containers, and was found to contain amounts of meteoritic material entirely adequate for the determinations. In taking a sample, we first scraped away about an inch of surface dirt, vegetable matter, and rocks, and then with a shovel scooped up enough soil to fill our containers. Each sample was labeled at the site and brought




FIGURE 1.—Magnetic separator, shown schematically.

The separator consists basically of a vibrating hopper and a slanted trough down which the soil moves under the combined action of gravity and vibration. The trough is about one inch wide and two feet long. Three strong alnico magnets are suspended above the trough. As the soil moves downward, the magnetic material flies up and adheres to the suspended magnets. This method readily differentiates between weakly and strongly magnetic material, for the separation between magnet and trough is adjusted so that only the strongly magnetic material flies up. This technique worked especially well in the present instance because the unwanted constituents were, for the most part, only weakly magnetic and the line of demarkation between the two was very pronounced.

Various criteria for choosing the amount of soil to be processed were considered, and we fixed on a volume weighing about 2,000 grams on the average, as best, although individual

back to the laboratory where we separated it into four groups according to size by shaking it down in a series of U. S. Standard Sieves, Numbers 10, 40, and 100, which yielded four components. These sieves, which are commonly used for soil analysis, are made from woven wire mesh; the No. 10 sieve is woven with ten fairly coarse wires per inch; the No. 100 with 100 fine wires per inch.

All material larger than 2 mm in diameter was caught on the No. 10 sieve; this fraction contained stones, clods of dirt, vegetable matter, etc., and was always discarded. Some meteoritic material was lost in this way; its amount is roughly estimated in a later section. The material that passed the No. 10 sieve and was caught on the No. 40, referred to here as size 40 material, was between 2 mm and 0.42 mm in diameter, about the size of coarse sand. To be caught on the No. 100 sieve the particles (referred to henceforth as size 100) must be as large as 0.149 mm in diameter. The residue,

or size "pan" particles, were then all less than 0.149 mm in diameter. The No. 10 fraction averaged 17 percent by weight of the sample. The remainder of the material was usually divided approximately equally among the other three fractions. During the early stages of the program, each of these was run through the magnetic separator individually.

The sieving accomplished two things: it insured that large particles of soil or sand did not seriously interfere with the extraction of small magnetic particles while being processed in the separator; and it enabled us to study the relative abundances of nonmeteoritic magnetic materials, which, from our point of view, were contaminants.

The magnets collected three types of strongly magnetic material: a meteoritic particle that consisted mostly of nickel-iron; a meteoritic iron-oxide particle; and a black, shiny particle. In addition, many particles were a cross between the first two types: patches or veins of iron-oxide would contain bits of unoxidized iron. Some dirt and a thin layer of yellow limonite adhered to the meteoritic particles. The black, shiny particles were probably bits of magnetite. The three types are most easily identified by mounting them in plastic and then grinding the plastic so that the particles are seen in cross-section. The copper from a copper sulfate solution plates out immediately on the iron particles so that these can easily be distinguished from iron-oxide particles although both have a metallic lustre. The relative percentages of each type of material varied among the size groups.

The meteoritic particles had various shapes: flakes, angular chunks, and ball-like masses. No very serious attempt was made to classify them because Nininger (1956) has already done so well in this regard. A portion of each sample was mounted in plastic, and many of the specimens were polished and inspected to make sure that the contents of each sample were principally meteoritic.

About one particle in every ten of the size 40 group was wholly nickel-iron in samples whose concentration of meteoritic material was high. This ratio varied a great deal however from sample to sample. Only a few particles were of the black, shiny type; the amount of this contaminant was much greater in the smaller

sizes, with the No. 10 component containing about 15 percent. The "pan" material was found to be so highly contaminated that we could not determine the exact amount of meteoritic material present, but examination under a microscope showed that meteoritic fragments formed an exceedingly small percentage of the total magnetic component.

The correlation between the percentage of meteoritic debris in the finer material (size 100) and that in the coarser material (size 40) was found to be quite good; values for a number of representative samples are plotted in figure 2.




FIGURE 2.—Correlation between percents of size 100 and size 40 materials. Each point represents an individual sample. Ordinate is the percentage of size 100 material in the sample; abscissa is the percentage of size 40 material.

The abscissa is the percent by weight of meteoritic material in the size 40 component of soil; and the ordinate, the percentage in the size 100 component of the same sample. On the average, the concentration in the size 40 fraction is seen to be roughly two and one-half times that in the size 100 fraction. A similar plot of size "pan" versus size 40 showed only a very slight correlation between the two, indicating that a high percentage of the pan magnetic material was nonmeteoritic. Usually, although not always, the nickel test was negative with the pan material.

Sampling procedure

The objective of the expedition was to sample the entire area surrounding the crater and thus determine as accurately as feasible the distribution of meteoritic debris. Many factors influenced the manner of sampling and the

number of samples made. Because of limited time and funds, we decided to sample sparsely over a wide area rather than intensively over a small region, and to take mainly surface samples, relying on a few representative holes for an indication of the vertical distribution of material. We were anxious that our sampling area be large enough so that at its periphery the concentration of meteoritic material would be negligible. Our procedure was to begin in areas where we knew the concentration to be high and work out from these as far as we needed to.

Approximately 700 samples were taken from the locations shown in figure 3. The pattern is roughly a grid of 80 square miles with the crater near its center. The locations were separated by a distance of one-half mile. In the early phases of our work we occasionally took several samples at a single location, to check reproducibility of results or for other purposes; in general, however, we took two samples, about 30 feet apart. Each of these two samples was processed individually in order to obtain a rough indication of the local variation or irregularity in concentration. The two values were averaged for most purposes.

With the first 60 or 70 samples, we processed all fractions, except the size 10. It soon became clear, however, that processing the "pan" gave no reliable information because it was highly contaminated with nonmeteoritic material so this was abandoned. For most samples,




FIGURE 3.—Locations of sampling points. The crater is indicated at center. Open circles indicate surface points; solid circles, holes dug.




FIGURE 4.—Locations and sample numbers of holes dug.

therefore, meteoritic material was recovered only from the size 40 and size 100 fractions.

The first step in processing was the careful weighing of each soil fraction. The magnetic component extracted from each fraction was weighed on an analytical balance and the percentage of meteoritic material computed. These percentages are listed in table 1 for all size 40 and size 100 samples; figure 9 shows their locations.

To study the vertical distribution of material, we dug to bedrock at 25 locations, shown in figure 4, and sampled the soil every few inches along the wall of the hole. Great care was taken not to contaminate a sample with dirt from some other part of the hole. The two lines of holes were chosen so as to traverse areas that could be considered representative of the whole region over which surface samples were taken. Bedrock was usually only one to three feet below the surface. In most holes the amount of meteoritic material decreased rapidly with depth in an approximately exponential fashion. In a few holes the concentration remained about constant from top to bottom. The distribution within each hole is shown in figure 5. Detailed data are listed in table 2.

Estimate of total mass of meteoritic material

The data obtained provide the basis for estimating the total quantity of meteoritic material around the crater. Such an estimate suggests a minimum value for the mass of the meteorite that produced the crater. The percentages of size 40 magnetic material are the




FIGURE 5.—Vertical distribution of meteoritic debris in each hole; sample numbers are given at the top.

most reliable, and we have therefore used them throughout as primary data.

To estimate the total weight, W , we evaluate the integral

$$W = \int \rho_m dV \quad (1)$$

where ρ_m is the density (gm/cc or lb/ft³) of meteoritic material. This density of course varies from point to point. The volume over which the integration is carried out is the mantle of soil around the crater.

Certain assumptions and a number of approximations, some very good and others quite rough, must be made in order to carry out the integration.

First, we establish a standard soil sample. By weight this sample consists of 17 percent size 10 fraction, 25 percent size 40 fraction, 30 percent size 100, and 28 percent size "pan." These percentages were obtained by averaging values from the 700 samples taken.

Second, we assume that *in situ* the soil has a void ratio of 1/3 so that the volume, V , occupied

by soil of mass S is given by

$$V = \frac{3S}{2\rho} \quad (2)$$

where ρ is the average density of the soil particles themselves. Since the soil is mostly quartz sand we take ρ equal to 2.65 gm/cc or 2.65×62.4 lb/ft.³

Third, since the size 10 magnetic component was not measured and the size 100 and size "pan" were unduly contaminated with non-meteoritic material, we must infer the amounts of size 10, size 100, and size "pan," from determination of the amount of size 40 meteoritic material. The amount of size "pan" was taken as zero since there was so little correlation between size 40 and size "pan." The amount of size 100 component, as indicated in figure 2, averaged about 0.4 that of size 40. The size 10 fraction of soil usually consisted of small clods of dirt which broke up into sizes 40, 100, and "pan" when crushed. Hence we assumed that the amount of meteoritic material could best be approximated by taking it as equal to

the average of the other three components. Summarizing, we have

$$m_p = 0$$

$$m_{100} = 0.4 m_{40}$$

$$m_{10} = \frac{m_p + m_{40} + m_{100}}{3} = \frac{(0 + 1 + 0.4)}{3} m_{40} = 0.5 m_{40}$$

where m_p is the mass of size "pan" meteoritic material in the sample of total mass S ; m_{40} , the mass of size 40; m_{100} , the mass of size 100; and m_{10} , the mass of size 10. Our final estimate for the total mass, m , of meteoritic material in a soil sample of mass S is the sum of the masses of the individual fractions or

$$m = 1.9 m_{40}$$

Now, as mentioned above

$$S_{40} = 0.25 S$$

where S_{40} is the mass of the size 40 fraction of the soil so that

$$m = 0.5 \frac{m_{40}}{S_{40}} S. \quad (3)$$

Combining equations (1), (2), and (3) we have

$$W = \int 0.5 \left(\frac{m_{40}}{S_{40}} \right) \left(\frac{2}{3} \rho \right) dV \quad (4)$$

which can be integrated if we know only the ratio, m_{40}/S_{40} , for each point.

Fourth, while this ratio is well established at the surface of the soil mantle, we need to make some assumptions regarding the variation with depth. The vertical distribution of meteoritic material within the holes shows that the density decays roughly exponentially with depth and is reasonably well represented by the equation

$$\rho_m = \rho_{m_0}^{-1.44y} \quad (5)$$

where ρ_{m_0} is the density of meteoritic material at the surface and y is distance in feet below the surface. The constant in the exponent was chosen so that ρ_m would be equal to $\frac{1}{2}\rho_{m_0}$ for $y = \frac{1}{2}$ ft. The choice is in good accord with the observations made in the 25 holes.

Fifth, we assume that the average depth of

the mantle is two feet. The exact depth is not critical because the density of meteoritic material falls off so rapidly with depth; at two feet it is only 0.06 the surface density. The average density, $\rho_{m_{av}}$, in a mantle of this depth is

$$\rho_{m_{av}} = \frac{1}{2} \int_0^{\infty} \rho_{m_0} \exp^{-1.44y} dy = \frac{1}{2.88} \rho_{m_0}. \quad (6)$$

With these assumptions we can now perform the integration. We have according to equation (4)

$$W = 0.5 \left(\frac{2}{3} \rho \right) \sum \left(\frac{m_{40}}{S_{40}} \right)_i V_i \quad (7)$$

where $\left(\frac{m_{40}}{S_{40}} \right)_i$ is the average value of the ratio for the volume element V_i of the mantle. Now V_i can be replaced by $2A_i$ where A_i is a surface element of the mantle expressed in square feet and $\left(\frac{m_{40}}{S_{40}} \right)_i$ can be replaced by $\frac{1}{2.88} \left(\frac{m_{40}}{S_{40}} \right)_{s_i}$ where the subscript s denotes the surface value of the ratio. Substituting these values into equation (7) we have the weight in pounds,

$$\begin{aligned} W &= 0.5 \times 2 \times \frac{1}{2.88} \times \frac{2}{3} \times 2.65 \times 62.4 \sum \left(\frac{m_{40}}{S_{40}} \right)_{s_i} A_i \\ &= 38.2 \sum \left(\frac{m_{40}}{S_{40}} \right)_{s_i} A_i \end{aligned}$$

The value of the summation is most easily computed from the chart in figure 6 which gives average values of the ratio for segments of area. We find that

$$\sum \left(\frac{m_{40}}{S_{40}} \right)_{s_i} A_i = 6.46 \times 10^5 \text{ ft}^2.$$

Substituting, we have

$$\begin{aligned} W &= 38.2 \times 6.46 \times 10^5 \\ &= 24.7 \times 10^6 \text{ lb,} \\ &\text{or} \\ &= 12,000 \text{ tons.} \end{aligned}$$

Thus the total amount of finely divided meteoritic material present in the soil is about 12,000 tons. However, this figure is subject to a still further uncertainty. We noted that about 90

percent of the meteoritic material that we collected is iron-oxide, Fe_2O_3 , and contained 27.6 percent oxygen by weight. Thus, about one-quarter of the above estimated mass is very likely terrestrial oxygen that combined with the meteoric iron after its encounter with the earth.

Further, there seems to be no way to allow precisely for the amount of contaminating material present. Increasing the area of the soil mantle over which the integration was taken enlarges the error caused by contamination since in the outlying regions the ratio between contaminant and meteoritic material becomes ever larger. We have tried to choose an area that minimizes the effects of contaminants but at the same time includes almost all of the meteoritic material. It will be noted, how-

ever, that even for the area chosen, the density of magnetic material falls off less rapidly than $1/r^2$ (r equals distance from center of crater) and the integral does not converge. At distances of over about two miles from the crater the percentages given on the chart are little if any higher than the background contamination which on the average ranged from 0.002 to 0.005 percent. To obtain a true measure of the meteoritic material, the percentage in each segment should be reduced by this amount. When this is done our computed mass of 12,000 tons is reduced by about 10 to 20 percent.

Distribution of meteoritic debris

The distribution pattern of the meteoritic debris around the crater is shown in figure 7, based on the data for the size 40 material




FIGURE 6.—Average percentages of size 40 meteoritic material in the soil around the crater.

(given in table 1). The contour lines show areas having equal densities of debris; the solid lines represent 0.1 percent changes. The broken lines indicate the 0.05 percent level.

The chart indicates several things. First, the debris itself lies in a perfectly definite pattern. Second, the crater itself does not lie at the center of the pattern. Third, the debris is symmetrically distributed about a line that runs somewhat north of east. Its exact position is hard to fix more specifically from our data, but the outside limits are due east and 45° north of east. Fourth, the distribution, while symmetrical, is not smooth, but contains several local areas in which the density of meteoritic material is high. And, fifth, there is a concentration of material to the east of the crater.

Although the definite pattern of debris was not surprising, it was encouraging, for it represents evidence that our data were not seriously distorted by sample contamination with terrestrial magnetic material. The symmetry of the pattern, its relationship to the crater, and

the localization of material are our most significant results, and will be discussed later.

It is interesting to speculate on how the meteoritic debris reached the points where it is found today. The two chief hypotheses are: the material fell to its present location shortly before, just at, or shortly after, the instant of the meteor's impact; or the material was deposited, after the impact, by the action of wind and water or other carriers. The very close similarity between the distribution of ponderable chunks of meteoritic material, as shown in Barringer's (1909) chart, and the distribution of minute pieces as determined by our survey, constitutes strong evidence pointing to the first as the most probable hypothesis.

Nininger (1956) has suggested that a strong wind was blowing at the time of the meteor's impact, and that the meteoritic debris represents condensation products from a vast metallic cloud rising vertically from the crater and distributed asymmetrically about the crater by the wind. Our survey does not support this theory,




FIGURE 7.—Distribution pattern of meteoritic debris around the crater. Contour lines indicate areas of equal density, given in percents; solid lines represent 0.1 percent changes; broken lines indicate the 0.05 percent level.

however; it is not easy to imagine a wind strong enough to propel large chunks and minute pieces of the same material in exactly the same direction and for the same distance. The present juxtaposition of the two classes of material while not irrefutable evidence, certainly strongly suggests that both were deposited at the same time, and that any subsequent shifting by the wind has been negligible. Many of the bits are probably remnants of ponderable chunks.

Shifting of the material by the wind over the centuries is a reasonable thing to postulate, and, in fact, the direction of the observed asymmetry is that of the prevailing wind. Another possibility which must be considered is that the vaporized metal condensed in droplets which fell with radial symmetry about the crater and were subsequently blown northeasterly by the wind. The asymmetrical distribution of the large chunks, however, cannot be explained on this basis. If wind were a factor we might expect that the bits would be sorted by weight; no evidence of such a sorting was found.

A highly reasonable hypothesis is that the meteorite approached the earth from a south-westerly direction and, when it struck, pitched forward large quantities of meteoritic material to the position where it now rests. (Such a shoveling action occurs frequently with obliquely impacting high speed missiles.) Orbital arguments do not favor any particular direction of encounter except possibly east-west (or west-east) slightly over north-south (or south-north). When the meteorite struck it must have been almost completely shattered, into pieces that weighed from a thousand pounds or so downward. No piece larger than 2000 pounds has ever been found (Barringer, 1914) while thousands of smaller pieces weighing a few ounces or less have been recovered. Much melting and considerable vaporization would have accompanied this shattering. It seems unlikely, however, that much of the vaporized material would have condensed into metallic droplets. More probably, it would have oxidized and drifted off as a fine powder. On the other hand, the sprays of molten material would probably have moved along with the solid fragments so that

deposition of molten (which of course would quickly solidify) and of solid fragments would have occurred at the same time.³

Eventually, of course, both forms of material would gradually have disintegrated, by a process of oxidation and subsequent exfoliation. The taenite, being more resistant to oxidation than the kamacite, would be the last to go. It is significant that Nininger finds that the metallic bits contain about 17 percent nickel, a composition compatible with that of taenite. Thus, one would expect to find in the soil minute pieces of both oxide and metallic matter, and even to find them intimately mixed. The observed variety in particle shape is likewise consistent with the view that these bits are, in the main, remnants of larger chunks of meteoritic material. The exact nature of the bits is still being intensively studied and they will be described in a later report.

The several local areas of high concentration indicate that the rain of meteoritic debris was spotty. Each area would then represent a region in which a large swarm of fragments fell after being ejected from the crater. This theory receives strong support from the data yielded by our pairs of samples. Recall that two samples, about 30 feet apart, were taken from each location; and the percentage of meteoritic material in the two often differed by a factor of two or three, and had no relation to local irregularities of terrain. Local wind eddies may have affected the distribution slightly. Such highly localized variation in the distribution indicates a second order patchiness of the ejecta that is entirely consistent with our knowledge of the phenomena of high speed impact.

³ E. P. Henderson, of the U. S. National Museum, after reading this paper in manuscript, made the following comment:

"Possibly the mass that made the crater had some satellite masses. These, being smaller, were retarded during the fall through our atmosphere, so possibly they fell outside the area of the crater. Being small, they did not make a crater. The crater-forming mass retained its velocity and was ablating until the moment of impact. Quantities of material were removed and carried into the turbulent wake of the meteorite. Since these pieces were small, their velocity was quickly lost. They should settle to the ground at a considerable distance short of the end point of the large crater-producing mass. It would seem that such material should be tracked almost up to the rim of the crater. On the opposite side and in line with the trajectory of the falling meteorite, one should get pieces that broke from the large meteorite and rocks that were tossed out of the crater. This point was stressed earlier in the paper."

The direction of impact

The fact that the meteoritic debris is distributed symmetrically about a line gives us the direction but not necessarily the sense of the trajectory of the meteorite. Experience with the effects of high speed impact shows that one can unambiguously associate symmetry of the ejected missile and impact debris with direction of impact. The axis of symmetry and the trajectory of the missile always lie in the same plane. The same situation must obtain here. Thus, if we assume that the debris now lies where it originally fell, then there is little doubt that the meteorite approached the earth along the axis of symmetry of our pattern: roughly, north of east or south of west.

Whether the approach was from the east or from the west is debatable. If the meteorite approached its point of impact at a steep angle from the east, shedding material as it neared the ground, deposited a substantial amount of debris to the east of the crater and then buried itself beneath the floor of the crater, it could now lie under the southwest corner of the crater. The assumption of a steep approach would be required to account for the fact that very little meteoritic material is found to the west of the crater although a large field of ejected boulders lies on this side. On the other hand, the meteorite could well have approached from the west and thrown debris forward, to the east, where we find an even greater accumulation of large fragmented limestone boulders and other ejecta than on the west. The boulders east of the crater have been thrown farther than on the west, in some cases two or three miles (Barringer, 1909, 1914). A western approach would not require us to assume so steep an angle of impact since we are now permitted to assume also that meteoritic material was thrown forward by the force of the impact. The evidence for a western approach is therefore the stronger.

Impactite and rock flour are other forms of ejecta (Merrill, 1908; Nininger, 1954). Impactite is sandstone metamorphosed by the intense heat of the impact, and impregnated with fine bits of meteoritic material and, occasionally, bits of limestone. Two large patches have been found, southeast and northwest of the crater. The sandstone from which this material

was made lies deep in the crater. Rock flour is similar sandstone that was ground to fine powder at the time of impact. Large quantities lie within and piled more or less uniformly around the rim of the crater (Barringer, 1909). Large masses of meteoritic oxide and meteorites are occasionally found embedded in it. It is not at all obvious how the impactite and rock flour got to their present position. They probably were thrown out immediately after the limestone boulders and meteoritic debris were ejected, although it is difficult to demonstrate this conclusively.

The structures of recovered meteorites also suggest an approach from the southwest or west. Meteorites found to the east and northeast of the crater have been severely altered by heat and deformation. Those found far out on the plain to the southwest and west, however, are in their virgin state (Nininger, 1956), and may well be pieces that broke from the meteorite as it approached from the west. Usually these specimens are fairly large and are sparsely distributed.

Heretofore, it has been postulated that the meteorite approached from a north-northwesterly direction (Barringer, 1909, 1914; Nininger, 1956). This conclusion was based, at first, on a reconnaissance examination of the tilting of the rock strata; later, on extensive drillings made in the crater and on its south rim; and finally, on magnetic, electrical, and gravimetric surveys. These data are difficult to evaluate. Many investigators have attempted to do so and much of what has been written is naturally conjectural. The magnetic, electrical, and gravimetric surveys seem, in spite of some contradictory evidence, to favor a southwest to northeast direction. The observed anomalies, shown in figure 8 (from Nininger, 1956), are not pronounced but they all seem to lie either in or beyond the southwest part of the crater.

Barringer made extensive drillings over a long period of years, and thus established the fact that large masses of fragmented rock lie buried beneath the floor and south rim of the crater. Since drillings have not been made at other locations we have no way of knowing how much fragmented rock lies beneath the east, north, and west rims, and we know very little




FIGURE 8.—Results of various geophysical surveys of crater.

about the shape of the subsurface region that was shattered by the impact. Further data on this point could be of very great help in establishing the direction of impact.

As well as we can determine, the only study of the tilting of strata was a reconnaissance surface examination made by Barringer (1909) whose findings have been summarized by Watson (1936):

The least amount of tilt is found in the northern wall. The slope increases along both the eastern and western walls, until in the southeast and southwest the strata are practically vertical. In contrast to the strata so tilted, and separated from them by an abrupt discontinuity, there occurs, along the rim slightly east of south, a broad arch, 2,700 feet long, the almost flat strata of which have been raised vertically about 100 feet. The arch appears to be a localized uplift possibly due to steam explosion subsequent to the complete deceleration of the penetrating body. The peculiarity of the tilting combined with the rock structure beneath the crater, suggests that the impact was not vertical, as previously supposed, but at a considerable slant from the north.

To test this theory of the direction of impact, a churn-drill was put down through the center of the arch in the south rim of the crater wall. Below a depth of 1,200 feet the drill passed through a region increasingly rich in loose meteoritic material and finally stuck at 1,340 feet, a region very rich in nickel-iron and exceedingly resistant to boring. The drill was slowly forced through 30 feet of this material but it stuck and remains completely immovable at 1,376 feet. The behavior of the drill shows that the meteoritic penetration continued to a depth of over 1,300 feet and also that very probably a portion of the original mass lies buried beneath the southern rim of the crater. From the position of this mass, the various depths of the undisturbed strata under the crater, and the symmetrical tilting of the crater walls around the north-south axis, it was concluded by the Barringers and their associates that the body struck from the north and at an angle of approximately 45° .

It is hard to appreciate, however, what these tilt measurements do in fact signify. They show us only the condition at the surface and leave us completely in the dark as to how much tilting and faulting has occurred below the surface. Our best evidence on this point is the

might therefore bear little relationship to the direction of the impact.

Our argument here leaves much to be desired in the way of completeness. It is indeed unfortunate that we have so little experimental data on these matters.

Evidence for large buried mass

The data presented here give no direct indication as to whether a large mass of meteoritic material lies buried in the crater. If one accepts the low estimates, 10,000 to 15,000 tons (Rinehart, 1950; Wylie, 1943) of the mass needed to form the crater then no large mass could possibly remain. If, on the other hand, one accepts the 5,000,000 ton estimate of Öpik (1936) and of Rostoker (1953) as more realistic, then the amount of meteoritic debris so far recovered is insignificant and a large mass could well lie buried. According to F. L. Whipple (personal communication), a most reasonable mass is that given recently by Hill and Gilvarry (1956), between 80,000 and 400,000 tons.

TABLE 1.—Horizontal surface distribution of meteoritic material—Continued

Location of sample (see fig. 9)	Percent by weight of magnetic material					
	Size 40			Size 100		
	In samples	Average		In samples	Average	
48	.0277	.0366	.0322	.0251	.0128	.0190
49	.0115	.0195	.0155	.0181	.0246	.0214
50	.0022	.0008	.0015	.0027	.0059	.0043
51	.0611	.0123	.0367	.1190	.0062	.0626
52	.0056	.0082	.0044	.0294	.0039	.0152
53	.0236	.0218	.0227	.0845	.0622	.0774
54	.0700	.0037	.0854	.0207	.0588	.0423
55	.0069	.0035	.0052	.0145	.0130	.0138
56	.0054	.0065	.0060	.0058	.0130	.0094
57	.0949	.0789	.0869	.0624	.0054	.0814
58	.0023	.0055	.0089	.0032	.0023	.0028
59	.0254	.0883	.0284	.0214	.0084	.0139
60	.1124	.0850	.0987	.0204	.0042	.0123
61	.6280	.4150	.6215	.0217	.1395	.0606
62	.1348	.1922	.1635	.0197	.0530	.0364
63	.1655	.1438	.1546	—	—	—
64	.0607	.1290	.0648	.0542	.1200	.0871
65	.0456	.0361	.0408	.0230	.0309	.0270
66	.0372	.0073	.0072	—	.0106	—
67	.0004	.0002	.0003	.0047	.0010	.0028
68	.0020	.0013	.0016	.0017	.0017	.0017
69	.0014	.0015	.0014	.0294	.0183	.0208
70	.0206	.0184	.0180	.0483	.0157	.0320
71	.0160	.0175	.0166	.0146	.0380	.0268
72	.0064	.0062	.0063	.0112	.0214	.0163
73	.0147	.0146	.0146	.0203	.0228	.0216
74	.0832	.0138	.0235	.0155	.0347	.0251
75	.0276	.0311	.0294	.0507	.0257	.0382
76	.0089	.0005	.0047	.0124	.0037	.0080
77	.0046	.0027	.0037	.0148	.0202	.0176
78	—	—	—	—	—	—
79	.0674	.0120	.0487	.0418	.0140	.0279
80	.0100	.0063	.0082	.0126	.0264	.0195
81	.0136	.0110	.0123	.0118	.0208	.0163
82	.0012	.0014	.0013	.0085	.0078	.0082
83	.0062	.0094	.0078	.0459	.0332	.0398
84	.0078	.0007	.0042	.0107	.0107	.0107
85	.0126	.0178	.0153	.0234	.0314	.0274
86	.0124	.0131	.0128	.0132	.0262	.0197
87	.0501	.0897	.0729	.0226	.0627	.0426
88	—	—	—	—	—	—
89	.0765	.0588	.0682	.0462	.0402	.0432
90	.0169	.0849	.0509	.0143	.0067	.0120
91	.0453	.0143	.0298	.0464	.0206	.0335
92	.0124	.0103	.0114	.0138	.0303	.0220
93	.0053	.0050	.0052	.0060	.0130	.0110
94	.0005	.0003	.0004	.0033	.0007	.0020
95	.0144	.0241	.0142	.0239	.0404	.0322
96	.0063	.0062	.0062	.0062	.0082	.0186
97	.0108	.0077	.0092	.0158	.0248	.0203
98	.0080	.0094	.0062	.0111	.0066	.0098
99	.0191	.0128	.0160	.0621	.0126	.0374
100	.0411	.0846	.0628	.0434	.0995	.0714
101	.3850	.0611	.2230	.0289	.0443	.0371
102	.0063	.0177	.0220	.0274	.0128	.0201
103	.0433	.0322	.0378	.0214	.0138	.0176
104	.0005	.0000	.0002	.0002	—	—
105	.0008	.0004	.0006	.0004	.0020	.0012
106	.0013	.0026	.0020	.0000	.0066	.0033
107	.0008	.0002	.0005	—	.0004	—
108	.0001	.0004	.0002	.0005	.0003	.0004
109	.0010	.0029	.0020	.0011	.0002	.0006
110	.0005	.0084	.0044	.0005	—	—
111	—	.0004	—	.0005	.0001	.0003
112	.0004	.0008	.0006	—	.0030	—
113	.0001	.0006	.0004	.0008	.0014	.0011
114	.0007	.0003	.0005	.0004	.0005	.0004
115	.0590	—	—	.0009	.0009	.0009
116	.0009	.0005	.0007	.0012	.0008	.0005
117	.0000	.0005	.0002	.0001	.0014	.0007
118	.0009	.0003	.0006	.0005	.0017	.0011
119	.0014	.0006	.0010	.0106	.0014	.0080
120	.0253	.0313	.0283	.0066	.0433	.0264
121	.0062	.0051	.0056	.0018	.0019	.0018
122	.0601	.0019	.0010	.0024	.0022	.0023
123	.0085	.0095	.0090	.0200	.0191	.0200
124	.0025	.0098	.0062	.0028	.0126	.0077
125	.0012	.0012	.0012	.0023	.0067	.0045
126	.0048	.0019	.0034	.0338	.0053	.0066
127	.0000	.0035	.0018	.0011	.0127	.0069
128	.0044	.0008	.0026	.0029	.0018	.0024
129	.0103	.0281	.0162	.0391	.0118	.0254
130	.0000	.0002	.0001	.0013	.0017	.0015
131	.0070	.0023	.0046	.0054	.0072	.0063
132	.0007	.0003	.0005	.0016	.0319	.0018
133	.0011	.0005	.0008	.0046	.0053	.0050
134	.0004	.0007	.0006	.0039	.0139	.0365
135	.0159	.0047	.0103	.0154	.0037	.0096

TABLE 1.—Horizontal surface distribution of meteoritic material

Location of sample (see fig. 9)	Percent by weight of magnetic material					
	Size 40		Size 100			
	In samples	Average	In samples	Average		
1	.078	.076	.077	.025	.026	.026
3	.206	.211	.208	.077	.096	.086
5	.328	.218	.273	.136	.104	.120
7	.303	.888	.570	.089	.112	.100
9	.084	.084	.084	.018	.014	.016
11	.012	.032	.022	.011	.019	.015
13	.022	.009	.016	.019	.011	.015
15	.230	.112	.171	.097	.057	.077
17	.0119	.0166	.0142	.0198	.0262	.0230
18	.0331	.0199	.0265	.0334	.0245	.0290
19	.0532	.2640	.1666	.0779	.1434	.1106
20	.0680	.0833	.0506	.0332	.0351	.0342
21	.1090	.0478	.0779	.0865	.0606	.0736
22	.0129	.0063	.0096	.0268	.0119	.0194
23	.2960	.0796	.1878	.0675	.0566	.0570
24	.1565	.3600	.2612	.0456	.0591	.0524
25	.0947	.0029	.0038	.0111	.0107	.0109
26	.0023	.0020	.0022	.0008	.0126	.0112
27	.0055	.0024	.0040	.0244	.0150	.0197
28	.0075	.0054	.0064	.0209	.0136	.0172
29	.0332	.0753	.0542	.0522	.0637	.0580
30	.1876	.6300	.4088	.2050	.5400	.3770
31	.2630	.1456	.2138	.1296	.1020	.1159
32	.0412	.0169	.0290	.0162	.0063	.0128
33	.1105	.0833	.0969	.0651	.0541	.0596
34	.0502	.0447	.0474	.0463	.0569	.0516
35	.0011	.0033	.0022	.0175	.0269	.0222
36	.0900	.0914	.0857	.0570	.0604	.0587
37	.0400	.0488	.0444	.0723	.0606	.0664
38	.0013	.0016	.0014	.0007	.0059	.0078
39	.0008	.0006	.0007	.0018	.0067	.0042
40	.0030	.0026	.0023	.0292	.0063	.0178
41	.0022	.0021	.0022	.0088	.0072	.0080
42	.0066	.0031	.0045	.0113	.0261	.0187
43	.0032	.0023	.0011	.0008	.0019	.0014
44	.0561	.1568	.1064	.0125	.0107	.0116
46	.0741	.0707	.0724	.0124	.0417	.0270
46	.1295	.0053	.0674	.0184	.0094	.0134
47	.0099	.0296	.0148	.0061	.0212	.0136

TABLE 1.—Horizontal surface distribution of meteoritic material—Continued

Location of sample (see fig. 9)	Percent by weight of magnetic material					
	Size 40		Size 100			
	In samples	Average	In samples	Average		
136	.0015	.0114	.0064	.0221	.0249	.0285
137	.0008	.0004	.0004	.0025	.0050	.0038
138	.0022	.0018	.0010	.0089	.0100	.0094
139	.0065	.0070	.0068	.0318	.0014	.0166
140	.0010	.0027	.0042	.0138	.0038	.0021
141	.0016	.0067	.0042	.0138	.0028	.0093
142	.0002	.0007	.0004	.0037	—	—
143	.0000	.0002	.0001	.0009	.0002	.0006
144	.0017	.0074	.0046	.0037	.0079	.0058
145	.0012	.0007	.0010	.0022	.0006	.0014
146	.0049	.0016	.0032	.0003	.0005	.0004
147	.0010	.0087	.0038	.0172	.0047	.0110
148	.0012	.0045	.0028	.0006	.0021	.0014
149	.0045	.0003	.0024	.0028	.0028	.0028
150	.1760	.0048	.0904	.0269	—	—
151	.0244	.0230	.0237	.0292	.0300	.0296
152	.0013	.0004	.0011	.0005	.0002	.0004
153	.0067	.0021	.0044	.0047	.0041	.0044
154	.0933	.0512	.0722	.0121	.0125	.0123
155	.0061	.0007	.0034	.0049	.0001	.0025
156	.0001	.0007	.0004	.0059	.0033	.0043
157	.0023	.0020	.0022	.0088	.0026	.0037
158	.0030	.0035	.0022	.0012	.0077	.0044
159	.0039	.0012	.0026	.0159	.0030	.0094
160	.0013	.0018	.0016	.0007	.0073	.0040
161	.0088	.0202	.0145	.0096	.0064	.0080
162	.0003	.0000	.0002	.0022	.0004	.0013
163	.0003	.0002	.0002	.0013	.0028	.0020
164	.0009	.0036	.0022	.0030	.0037	.0033
173	.0125	.0157	.0141	.0068	.0064	.0051
179	.0113	.0159	.0136	.0038	.0010	.0024
180	.0118	.0035	.0076	.0072	.0102	.0087
181	.0157	.0080	.0118	.0235	.0084	.0150
182	.0191	.0136	.0164	.0030	.0126	.0078
183	.0004	.0004	.0004	.0024	.0023	.0023
184	—	—	—	—	—	—
185	.0032	.0035	.0034	.0011	.0015	.0013
186	.0056	.0034	.0045	.0097	.0066	.0092
187	.0017	.0032	.0024	.0015	.0061	.0038
188	.0043	.0108	.0076	.0048	.0128	.0088
189	.0231	.0786	.0408	.0127	.0188	.0148
190	.0017	.0110	.0064	.0020	.0105	.0062
191	.4830	.2490	.3660	.0814	.0120	.0217
192	.2160	.1169	.1664	.0285	.0423	.0354
193	.0138	.0657	.0358	.0072	.0176	.0124
194	.0342	.0167	.0164	.0083	.0028	.0064
195	.0376	.0285	.0320	.0028	.0046	.0037
206	.0083	.0103	.0093	.0171	.0151	.0181
209	.1595	.0659	.1127	.0241	.0256	.0248
210	.0087	.0083	.0085	.0023	.0047	.0035
211	.0523	.0020	.0672	.0213	.0275	.0244
212	.0307	.0568	.0438	.0071	.0115	.0093
213	.0057	.0034	.0046	.0311	.0073	.0192
214	.0035	.0063	.0049	.0052	.0060	.0071
215	.0045	.0054	.0050	.0050	.0125	.0088
216	.0189	.0062	.0126	.0046	.0564	.0305
217	.0009	.0020	.0014	.0038	.0018	.0028
218	.0008	.0007	.0006	.0006	.0008	.0006
219	.0017	.0029	.0023	.0003	.0032	.0052
220	.0002	.0007	.0004	.0011	.0040	.0026
221	.0180	.0147	.0164	.0187	.0082	.0110
222	.0085	.0013	.0039	.0053	.0100	.0076
223	.0024	.0033	.0011	.0016	.0014	.0032
224	.0005	.0017	.0011	.0016	.0188	.0038
225	.0000	.0000	.0000	.0003	.0014	.0008
226	.0008	.0005	.0006	.0005	.0015	.0010
227	.0125	.0002	.0064	.0001	.0007	.0004
228	.0042	.0005	.0024	.0054	.0042	.0048
229	.0010	.0039	.0025	.0061	.0035	.0048
230	.0020	.0025	.0022	.0036	.0044	.0040
231	.0009	.0002	.0006	.0011	.0007	.0009
232	.0005	.0003	.0004	.0012	.0016	.0014
233	.0283	.0299	.0278	.0053	.0185	.0119
234	.0001	.0017	.0009	.0003	.0005	.0004
235	.0154	.0172	.0168	.0084	.0115	.0100
236	.0003	.0000	.0002	.0007	.0009	.0008
237	.0203	.0129	.0166	.0248	.0040	.0078
238	.0117	.0056	.0086	.0107	.0048	.0078
239	.0090	.0110	.0100	.0022	.0178	.0100
240	.0032	.0023	.0028	.0022	.0034	.0028
241	.0099	.0773	.0086	.0061	.0062	.0062
242	.0023	.0040	.0032	.0001	.0012	.0012
243	.0008	.0066	.0067	.0066	.0066	.0082

TABLE 1.—Horizontal surface distribution of meteoritic material—Continued

Location of sample (see fig. 9)	Percent by weight of magnetic material					
	Size 40		Size 100			
	In samples	Average	In samples	Average		
244	.0013	.0015	.0014	.0119	.0067	.0088
245	.0019	.0016	.0018	.0132	.0196	.0164
246	.0065	.0081	.0073	.0462	.0262	.0362
247	.0006	.0006	.0006	.0104	.0026	.0035
248	.0001	.0002	.0002	.0003	.0007	.0005
249	.0001	.0007	.0004	.0058	.0030	.0044
250	.0005	.0023	.0014	.0017	.0013	.0015
251	.0026	.0066	.0056	.0065	.0140	.0102
252	.0109	.0086	.0098	.0094	.0148	.0121
253	.0188	.0065	.0122	.0349	.0409	.0379
254	.0142	.0083	.0112	.1361	.0728	.0994
255	.0039	.0053	.0050	.0925	.0494	.0694
256	.0006	.0007	.0006	.0015	.0014	.0014
257	.0013	.0036	.0025	.0029	.0016	.0022
258	.0225	.0090	.0308	.0257	.0089	.0173
259	.0065	.0008	.0036	.0088	.0008	.0048
260	.0002	.0011	.0006	.0086	.0027	.0082
261	.0005	.0027	.0016	.0059	.0166	.0098
262	.0112	.0080	.0086	.0088	.0226	.0182
263	.0068	.0042	.0055	.0215	.0059	.0137
264	.0114	.0106	.0060	.0041	.0041	.0041
265	.0370	.0329	.0350	.0046	.0131	.0088
266	.0028	.0027	.0025	.0039	.0042	.0040
267	.0018	.0042	.0030	.0023	.0038	.0030
268	.0114	.0025	.0020	.0079	.0047	.0034
269	.0032	.0014	.0023	.0010	.0026	.0018
270	.0035	.0022	.0028	.0040	.0033	.0036
271	.0008	.0014	.0011	.0007	.0014	.0010
272	.0065	.0196	.0080	.0062	.0062	.0062
273	.0182	.0028	.0192	.0123	.0073	.0096
274	.0099	.0098	.0086	.0084	.0180	.0112
275	.0098	.0116	.0107	.0113	.0262	.0188
276	.0080	.0044	.0062	.0050	.0080	.0065
277	.0171	.0113	.0142	.0123	.0242	.0182
278	.0264	.0162	.0213	.0212	.0171	.0192
279	.0068	.0063	.0068	.0062	.0066	.0064
280	.0180	.0118	.0149	.0083	.0060	.0072
281	.0024	.0172	.0068	.0474	.0196	.0385
282	.0114	.0043	.0078	.0088	.0089	.0294
283	.0214	.0185	.0190	.0196	.0320	.0258
284	.0049	.0137	.0118	.0182	.0072	.0117
285	.0108	.0042	.0106	.0302	.0229	.0296
286	.0064	.0174	.0119	.0188	.0089	.0114
287	.0098	.0100	.0099	.0162	.0276	.0219
288	.0066	.0154	.0110	.0224	.0321	.0272
289	.0133	.0115	.0124	.0193	.0140	.0166
290	.0057	.0043	.0050	.0092	.0042	.0045
291	.0074	.0109	.0062	.0079	.0042	.0060
292	.0050	.0060	.0050	.0077	.0074	.0076
293	.0062	.0085	.0084	.0090	.0080	.0107
294	.0009	.0024	.0016	.0099	.0132	.0116
295	.0092	.0069	.0080	.0050	.0063	.0056
296	—	—	—	—	—	—
297	.0623	.0197	.0410	.0003	.0284	.0188
298	.0117	.0340	.0223	.0181	.0052	.0116
299	.0032	.0094	.0063	.0377	.0334	.0356
300	.0060	.0060	.0060	.0306	.0231	.0268
301	.0058	.0272	.0165	.0396	.0545	.0470
302	.0049	.0085	.0067	.0432	.0206	.0319
303	.0065	.0080	.0073	.0335	.0383	.0059
304	.0031	.0049	.0040	.0083	.0022	.0052
305	.0035	.0017	.0028	.0039	.0047	.0043
306	.0196	.0095	.0146	.0068	.0294	.0181
307	.0037	.0121	.0170	.0074	.0174	.0124
308	.0146	.0241	.0190	.0040	.0080	.0035
309	.0048	.0088	.0043	.0021	.0020	.0020
310	.0152	.0190	.0171	.0027	.0113	.0070
311	.0844	.0146	.0245	.0047	.0000	.0024
312	.0005	.0009	.0007	.0007	.0007	.0007
313	.0100	.0037	.0085	.0053	.0128	.0088
314	.0075	.0100	.0088	.0076	.0088	.0067
315	.0021	.0072	.0048	.0068	.0079	.0088
316	.0013	.0000	.0006	.0057	.0016	.0036
317	.0026	.0104	.0065	.0038	.0013	.0026
318	.0016	.0081	.0048	.0039	.0067	.0053
319	.0012	.0022	.0017	.0076	.0061	.0064
320	.0129	.0034	.0062	.0151	.0069	.0125
321	.0029	.0052	.0040	.0117	.0021	.0024
322	.0045	.0002	.0024	.0022	.0011	.0016
323	.0005	.0002	.0004	.0008	.0008	.0008
324	.0040	.0006	.0023	.0060	.0024	.0037
325	—	—	—	—	—	—

TABLE 2.—Distribution of meteoritic material in vertical shafts

Location of sample	Depth (inches)	Percent by weight	
		Size 40	Size 100
164	0	.5600	.0302
	2	.3500	.1080
	12	.0890	.0408
165	24	.0084	.0087
	0	.0035	.0190
	2	.0033	.0021
166	12	.0011	.0188
	30	.0017	.0087
	0	.0748	.0889
167	18	.1900	.0771
	33	.0118	.0484
	0	.0084	.0183
168	1	.8980	.0207
	6	.2080	.0274
	15	.1400	.0288
169	12	.0485	.0483
	0	.2220	.0128
	1	.0788	.0489
170	6	.1860	.0180
	0	.0834	.0337
	1	.0680	.0224
171	12	.0386	.0386
	21	.0084	.0048
	0	.0320	.0431
172	1	.0485	.0189
	25	.0033	.0168
	54	.0002	.0042
173	0	.0082	.0284
	1	.0803	.0286
	12	.0274	.0066
174	27	.0013	.0008
	0	.3084	.0086
	1	.0086	.0080
175	15	.0082	.0014
	33	.0001	.0009
	0	.1285	.0411
176	1	.0810	.0241
	13	.0786	.0180
	27	.0685	.0187
177	0	.1884	.0685
	1	.0924	.0488
	18	.0430	.0088
178	30	.0280	.0084
	0	.0179	.0141
	1	.0103	.0143
179	12	.0086	.0018
	30	.0018	.0008
	0	.0000	.0000
180	1	.0870	.0144
	9	.0878	.0104
	0	.0781	.0280
181	0	.0880	.0184
	15	.1680	.0195
	30	.1425	.0174
182	0	.1748	.0688
	1	.0903	.0104
	9	.0171	.0384
183	18	.0426	.0023
	0	.1218	.0307
	1	.1031	.0180
184	21	.0089	.0013
	42	.0082	.0048
	0	.0297	.0080
185	1	.0081	.0049
	7	.0080	.0117
	0	.0072	.0017
186	1	.0101	.0027
	9	.0089	.0385
	0	.0082	.0124
187	1	.0082	.0075
	6	.0103	.0144
	15	.0164	.0586
188	0	.0212	.0117
	1	.0122	.0389
	18	.0010	.0007
189	36	.0008	.0080
	0	.0105	.0307
	1	.0221	.0083
190	13	.0223	.0388
	0	.0287	.0209
	1	.0060	.0141
191	18	.0006	.0005
	45	.0066	.0021

TABLE 2.—Distribution of meteoritic material in vertical shafts—Continued

Location of sample	Depth (inches)	Percent by weight	
		Size 40	Size 100
205	0	.0064	.0013
	1	.0012	.0031
	9	.0080	.0008
206	0	.0007	.0011
	1	.0030	.0005
	11	.0000	.0000
207	0	.0012	.0016
	1	.0001	.0008
	5	.0000	.0000

Acknowledgments

In conclusion I wish gratefully to acknowledge the assistance of several persons. The results could not have been obtained without the whole hearted enthusiasm and hard work, often under trying conditions, of members of the expedition. These were Mr. Nicholas Matalas, Mr. Robert O'Neil, and Mr. Robert Olson. The helpful cooperation of Mr. Ernest Chilson, vice president of the Bar-T-Bar Ranch, Mr. George Foster, curator of the crater Museum, Mr. D. Moreau Barringer of the Barringer Crater Company and Mr. Sid Griffin of Rimmy Jim's contributed greatly to the success of the program. Dr. F. Behn Riggs, Jr., designed the magnetic separator and helped in many other ways. Mr. Paul Hodge analyzed much of the data. Dr. Agnes M. Allen was good enough to let us use her facilities at the Northern Arizona State College, Flagstaff, for weighing our samples. And finally, Dr. Fred L. Whipple's constant enthusiasm for the project and his many stimulating suggestions and criticisms during all phases of it were invaluable.

References

- BARRINGER, E. M.
 1905. Proc. Acad. Nat. Sci. Philadelphia, vol. 57, p. 861.
 1909. Paper read before the National Academy of Sciences, at Princeton University Nov. 16, 1909. Privately printed.
 1914. Proc. Acad. Nat. Sci. Philadelphia, vol. 66, p. 556.
 BARRINGER, D. M., JR.
 1927. Scientific American, vol. 137, p. 52.

- GILBERT, G. K.
1896. The Geological Society of Washington; also, Science, new ser., vol. 3, p. 1.
- HAGER, D.
1953. Bull. Amer. Assoc. Petroleum Geologists, vol. 37, p. 821.
- HILL, J. E. AND GILVARRY, J. J.
1956. Journ. Geophys. Res., vol. 61, p. 501.
- MERRILL, G. P.
1908. Smithsonian Misc. Coll., vol. 50, p. 461.
- NININGER, H. H.
1951. Scientific Monthly, vol. 72, p. 75.
1954. Amer. Journ. Science, vol. 252, p. 277.
1956. Arizona's Meteorite Crater, American Meteorite Museum, Sedona, Ariz.
- ÕPIK, E.
1936. Publ. de L'Observatoire Astronomique de L'Université de Tartu, vol. 28, No. 6, 27 pages.
- RINEHART, J. S.
1950. Pop. Astron., vol. 58, p. 458.
- ROSTOKER, N.
1953. Meteoritics, vol. 1, p. 11.
- TILGHMAN, B. C.
1905. Proc. Acad. Nat. Sci. Philadelphia, vol. 57, p. 887.
- WATSON, F., JR.
1936. Pop. Astron., vol. 44, p. 2.
- WYLIE, C. C.
1943. Pop. Astron., vol. 51, p. 97.

UNCLASSIFIED

UNCLASSIFIED