

**UNCLASSIFIED**

---

---

**AD 259 075**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



---

---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S.

Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

FOR OFFICIAL USE ONLY.

MAY BE RELEASED ON

9 June 1961

# U. S. NAVAL AIR DEVELOPMENT CENTER

JOHNSVILLE, PENNSYLVANIA

259 075

CATALOGED BY ASTIA  
AS AD NO.

Aviation Medical Acceleration Laboratory

NADC-MA-6116

9 June 1961

The Effect of Ageing on the G-Tolerance of Rats

Bureau of Medicine and Surgery

Sub Task MR 005.15-0002.3

Report No. 5



**U. S. NAVAL AIR DEVELOPMENT CENTER  
JOHNSVILLE, PENNSYLVANIA**

**Aviation Medical Acceleration Laboratory**

**NADC-MA-6116**

**9 June 1961**

**The Effect of Ageing on the G-Tolerance of Rats**


**Bureau of Medicine and Surgery  
Subtask MR005.15-0002.3      Report No. 5**


**Prepared by:**

  
Elizabeth Reeves, LCDR, MSC, USN

**J. W. Frazer**

**Approved by:**

  
James D. Hardy, Ph.D.  
Research Director, Aviation Medical  
Acceleration Laboratory, and Professor  
of Physiology, University of Pennsylvania

  
F. K. Smith, Captain, MC, USN  
Director  
Aviation Medical Acceleration Laboratory

## SUMMARY

The effect of age on the G tolerance of rats was studied in five different age groups — one, three, four, six, nine and twelve months. Each group consisted of 24 male Sprague-Dawley rats. All of the rats were subjected to 20 positive G and their survival times were measured using an EKG end point. The results show that there was a gradual decline in acceleration tolerance with increase of age in this group of 98 rats.

## TABLE OF CONTENTS

	Page
SUMMARY.....	ii
INTRODUCTION.....	1
METHODS.....	1
RESULTS.....	5
DISCUSSION.....	5
BIBLIOGRAPHY.....	9

## LIST OF TABLES

Table		Page
I	Experimental conditions and comparison of survival time at 20 positive G.....	4
II	Significant difference between age groups or rats exposed to 20 positive G.....	6

## LIST OF FIGURES

Figure		Page
1	Probit plot of percent survival vs. log time of age.....	2
2	Probit plot of percent survival vs. log time of 4 month group vs. 3 and 6 month group.....	3
3	Median survival time vs. age.....	7

## INTRODUCTION

The question of the age of an animal in relation to that animal's ability to withstand positive acceleration has been raised during several experiments in this laboratory. The present study was designed, therefore, to determine if any relationship exists between age and tolerance to positive acceleration. If such a relationship were to be found, it was felt that a best range of age limits would be established for experimentation for this type of stress.

## METHODS

Animals used in this experiment were male Sprague-Dawley rats from Hormone Laboratories, Chicago, Ill. All the animals were born within a few days of each other. They were divided into groups of 24 each and held in the animal colony at the Aviation Medical Acceleration Laboratory (AMAL) for varying lengths of time—1, 3, 4, 6, 9, and 12 months. They were maintained on Purina Chow with water ad libitum. Temperature in the animal house was held at 23 to 24°C. Handling was limited to weighing the animals at weekly intervals. At the end of each ageing period, the animals from one group were loosely restrained in individual cages and placed horizontally on the eight-foot radius animal centrifuge for an acceleration of 20 positive G. Heart rate was monitored by means of a transistor amplifier method (1) developed at AMAL as a means for determining the physiological end point of an animal's tolerance to acceleration. Acceleration was stopped when the animal's heart rate decreased to 2 beats per second for ten seconds, at which time it was considered dead (2). The average heart beat under resting conditions before acceleration was 6 to 8 beats per second.

Since the survival time under an acceleration of 20 positive G does not follow a normal distribution, data were analyzed by the method advanced by Litchfield, et al (3, 4) for time-percent effect curves. This method gives a close approximation to the procedure developed by Bliss (5) of using logarithmic probability transformation for biological studies except that data in their original form may be plotted directly onto logarithmic-probability paper. By means of nomographs, the parameters and confidence limits of the time-percent effect curves may be made directly from original data. As is shown in Figures 1 and 2, such transformations effect a normal distribution which becomes a straight line by plotting on probit paper.




Fig. 1. Probit plot of percent survival vs. log time of age.




Fig. 2. Probit plot of percent survival vs. log time of 4 month group vs. 3 and 6 month group.

TABLE I

EXPERIMENTAL CONDITIONS AND COMPARISON OF  
SURVIVAL TIME AT 20 POSITIVE G

Age (months)	Number of Animals		$\bar{X}$ Med.	$\sigma$	$\sigma\bar{X}$ Med.
	Start	End			
1	24	24	18.2	2.01	0.58
3	24	24	15.0	1.99	0.57
4	48	47	13.0	1.91	0.39
6	26	23	9.5	1.89	0.59
9	24	17	4.8	1.95	0.68
12	24	10	2.4	1.91	0.86

## RESULTS

Figures 1 and 2 illustrate graphically the decline of tolerance when animals were tested at an acceleration of 20 positive G at 1, 3, 4, 6, 9, and 12 months of age. As is shown by Table I, the median survival time decreased as the age of the animal increased during this experiment (Figure 3). Rats in close age groups, i. e., 1 and 3, 1 and 4, 3 and 4, and 4 and 6, are not significantly different from each other; all other group differences are statistically significant from each other at the 0.5 level of confidence (Table II). A small aberrant group of animals with great tolerance to acceleration showed up in the 1 month group and will be further investigated.

## DISCUSSION

There have been observations that tolerance to different types of stress may vary with age and/or weight (6, 7, 8, 9). McCay, et al (7, 8) found that rats on diets of restricted caloric intake lived longer than those with similar diets but ad libitum. Thus it was shown that an actual improvement in physical condition must have taken place for those on the restricted diet because they not only lived longer but proved to be more resistant to chronic diseases as well as to tumor formation, and appeared to be chronologically younger than their control group. The fact that there were deaths in the older age groups before the termination of this experiment is probably to be explained by a lessening of physical condition as well as the usual chronic rat diseases although the animals accelerated to 20 positive G had very little disease apparent at autopsy. Sellers (10) and others also have noted age or weight factors under different stresses. They found that older rats do not appear to acclimatize to cold as well as younger animals, not being able to maintain weight in the cold and probably not being able to make as complete a metabolic adjustment as younger animals do. Heroux and Hart (6) found that the survival time at  $-29^{\circ}\text{C}$  was significantly correlated with individual body weight for animals which had been acclimatized to  $30^{\circ}\text{C}$ . Britton, et al (11) observed that young and nearly full-grown rats did not withstand G forces as well as mature animals and stated that there was a definite ageing factor involved. This observation however is just the opposite from that found in the present investigation. In the present study a decline in tolerance was noted with each increase of age. In addition, a group of fifty animals, four months of age and from the same supplier, which were used as controls in other experiments during this same time period, had a curve lying between the 3 and 6 month age groups (Figure 2). In rats under 6 months of age, differences between close age groups,

TABLE II

SIGNIFICANT DIFFERENCE BETWEEN AGE GROUPS  
OF RATS EXPOSED TO 20 POSITIVE G.

Age (months)	1	3	4	6	9	12
1	-	-	-	+	+	+
3	-		-	+	+	+
4	-	-		-	+	+
6	+	+	-		+	+
9	+	+	+	+		+
12	+	+	+	+	+	

Probability < .05




Fig. 3. Median survival time vs. age.

i. e. , between 1 and 3, 1 and 4, 3 and 4, and 4 and 6 were not significant; however differences between all other age groups show significance (Table II).

This decline in tolerance to positive acceleration might be caused by any of a number of physical and/or physiological variables, such as a decrease of elasticity of tissues, decreased ability to adapt, an increased inclination to disease and subclinical infections, increase in weight and fat deposits, increase in age, length of time on diet, length of time of forced inactivity, response of the rat to different seasons of the year, actual physical pull of the visceral organs downward to restrict or inhibit respiratory movement, (12) etc. The changes in elasticity of the tissues of the body with age, particularly the cardiovascular system and the tissues of the lungs could be a deciding factor with this type of stress. On autopsy it has been found that animals which have survived long periods of exposure to 20 positive G had little congestion or hemorrhagic areas in the lungs or disturbances in their cardiovascular system. On the other hand those which succumbed had congestion, large hemorrhagic areas sometimes involving whole lobes of the lungs as well as a persistent finding of engorgement of the vena cava and right heart failure leading to eventual death, most probably due to anoxic anoxia.

## BIBLIOGRAPHY

1. Sipple, W. C., and B. D. Polis: A physiological end point for the study of the tolerance of small mammals to high acceleration stress. U. S. Naval Air Development Center, Aviation Medical Acceleration Laboratory, Johnsville, Pa. NADC-MA-5906, 1959.
2. Polis, B. D.: Hormonal determinants of mammalian tolerance to acceleration stress. U. S. Naval Air Development Center, Aviation Medical Acceleration Laboratory, Johnsville, Pa., NADC-MA-6025, 12 Aug. 1960.
3. Litchfield, J. T. Jr., and F. Wilcoxon: A simplified method of evaluating dose-effect experiments. *J. Pharm. & Exper. Therap.*, 96:99, 1949.
4. Litchfield, J. T. Jr.: A method for rapid graphic solution of time-percent effect curves. *J. Pharm. & Exper. Therap.*, 97: 399, 1949.
5. Bliss, L. I.: The calculation of the time-mortality curve. *Ann. Appl. Biol.*, 24:815, 1937.
6. Heroux, O., and J. S. Hart: Adrenal cortical hormone requirements of warm and cold acclimated rats after adrenalectomy. *Amer. J. Physiol.*, 178:449, 1954.
7. McCay, C. M., M. F. Crowell, and L. A. Maynard: The effect of retarded growth upon the length of life span and upon ultimate body size. *J. of Nut.* 10:63, 1935.
8. McCay, C. M., G. Sterling, and L. L. Barnes: Growth, ageing, chronic diseases and life span in rats. *Arch. Biochem.* 2:469, 1943.
9. Reeves, E.: The effect of acclimatization to cold on the G tolerance of rats. U. S. Naval Air Development Center, Aviation Medical Acceleration Laboratory, Johnsville, Pa., NADC-MA-6117, 1961.

BIBLIOGRAPHY con't.

10. Sellers, E. A.: Adaptive and related phenomena in rats exposed to cold. *Rev. Canad. de Biol.*, 16:175, 1957.
11. Britton, S. W., E. L. Corey, and G. A. Stewart: Effects of high acceleratory forces and their alleviation. *Amer. J. Physiol.* 146:33, 1946.
12. Cranmore, D.: Lethal levels and gross pathology of rats exposed to positive and negative forces of acceleration. U. S. Naval Air Development Center, Aviation Medical Acceleration Laboratory, Johnsville, Pa., NADC-MA-5410, 30 July 1954.
13. Selye, H.: *Stress*, Acta, Montreal, Canada, 1950.

U. S. NAVAL AIR DEVELOPMENT CENTER, JOHNSVILLE, PA.  
AVIATION MEDICAL ACCELERATION LABORATORY

The Effect of Ageing on the G Tolerance of Rats, by Elizabeth Reeves, LCDR, MSC, USN and J. W. Frazer, 10 pp., 9 June 1961.

The effect of age on the G tolerance of rats was studied in five different age groups — one, three, four, six, nine and twelve months. Each group consisted of 24 male Sprague-Dawley rats. All of the rats were subjected to 20 positive G and their survival times were measured using an EKG end point. The results show that there was a gradual decline in acceleration tolerance with increase of age in this group of 98 rats.

1. Report NADC-MA-6116
2. BuMed Subtask MR005.  
15-0002.3

U. S. NAVAL AIR DEVELOPMENT CENTER, JOHNSVILLE, PA.  
AVIATION MEDICAL ACCELERATION LABORATORY

The Effect of Ageing on the G Tolerance of Rats, by Elizabeth Reeves, LCDR, MSC, USN and J. W. Frazer, 10 pp., 9 June 1961.

The effect of age on the G tolerance of rats was studied in five different age groups — one, three, four, six, nine and twelve months. Each group consisted of 24 male Sprague-Dawley rats. All of the rats were subjected to 20 positive G and their survival times were measured using an EKG end point. The results show that there was a gradual decline in acceleration tolerance with increase of age in this group of 98 rats.

1. Report NADC-MA-6116
2. BuMed Subtask MR005.  
15-0002.3

U. S. NAVAL AIR DEVELOPMENT CENTER, JOHNSVILLE, PA.  
AVIATION MEDICAL ACCELERATION LABORATORY

The Effect of Ageing on the G Tolerance of Rats, by Elizabeth Reeves, LCDR, MSC, USN and J. W. Frazer, 10 pp., 9 June 1961.

The effect of age on the G tolerance of rats was studied in five different age groups — one, three, four, six, nine and twelve months. Each group consisted of 24 male Sprague-Dawley rats. All of the rats were subjected to 20 positive G and their survival times were measured using an EKG end point. The results show that there was a gradual decline in acceleration tolerance with increase of age in this group of 98 rats.

1. Report NADC-MA-6116
2. BuMed Subtask MR005.  
15-0002.3

U. S. NAVAL AIR DEVELOPMENT CENTER, JOHNSVILLE, PA.  
AVIATION MEDICAL ACCELERATION LABORATORY

The Effect of Ageing on the G Tolerance of Rats, by Elizabeth Reeves, LCDR, MSC, USN and J. W. Frazer, 10 pp., 9 June 1961.

The effect of age on the G tolerance of rats was studied in five different age groups — one, three, four, six, nine and twelve months. Each group consisted of 24 male Sprague-Dawley rats. All of the rats were subjected to 20 positive G and their survival times were measured using an EKG end point. The results show that there was a gradual decline in acceleration tolerance with increase of age in this group of 98 rats.

1. Report NADC-MA-6116
2. BuMed Subtask MR005.  
15-0002.3

**UNCLASSIFIED**

**UNCLASSIFIED**