



NEW YORK UNIVERSITY
INSTITUTE OF
MATHEMATICAL SCIENCES

DISCARD

On Families of Sets Represented in Theories

HILARY PUTNAM

PREPARED UNDER
CONTRACT NO. AF49(638)-777
MATHEMATICAL SCIENCES DIRECTORATE
AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

ARIZONA STATE UNIVERSITY LIBRARY

AD 260956

REPRODUCTION IN WHOLE OR IN PART
IS PERMITTED FOR ANY PURPOSE
OF THE UNITED STATES GOVERNMENT.

New York University
Institute of Mathematical Sciences

ON FAMILIES OF SETS REPRESENTED IN THEORIES

Hilary Putnam

ABSTRACT: A necessary and sufficient condition is given for a family of sets to be the family of all sets representable in a theory.

"Qualified requestors may obtain copies of this report from the ASTIA Document Service Center, Arlington Hall Station, Arlington 12, Virginia. Department of Defense contractors must be established for ASTIA services, or have their "need-to-know" certified by the cognizant military agency of their project or contract".

The research reported in this document has been sponsored by the Mathematical Sciences Directorate, Air Force Office of Scientific Research, Washington 25, D. C., under Contract No. AF 49(638)-777.

The purpose of this paper is to answer the following questions: (1) Let F be a family of sets¹. What are necessary and sufficient conditions that F be the family of all sets represented in some consistent standard theory²? (2) What are necessary and sufficient conditions that F be the family of all sets represented in some consistent axiomatizable standard theory?³

I shall prove:

THEOREM 1. F is the family of all sets represented in some consistent standard theory if and only if F is closed under intersection, finite addition and subtraction⁴, and contains the null set and the "universal" set (i.e. the set N_n of all non-negative integers).

THEOREM 2. F is the family of all sets represented in some consistent axiomatizable standard theory if and only if F is a recursively enumerable family of recursively enumerable sets⁵; F contains the null set and the "universal" set; and F is closed under intersection and finite addition and subtraction.

As an example of a consequence of THEOREM 1, we may cite the fact that, since the Π_1 sets (the sets whose complements are recursively enumerable) satisfy the closure conditions mentioned in the theorem, there exists a theory T with the property that all and only Π_1 sets are represented in T . Similarly, it follows from THEOREM 2 that, since the recursive sets satisfy the conditions given (that they form a recursively enumerable family in the sense on n.5 was first proved by Dekker), there exists an axiomatizable theory T with the property that all and only recursive sets are

represented in T . This result has been previously obtained by Shoenfield (in a stronger form); and the result about $\overline{\Pi}_1$ sets can likewise be obtained by a quite different construction than the one used here. However, it is of interest to see these results not as isolated curiosities, but as special cases of very general theorems.

1. General Remarks. $\{F_1, F_2, \dots\}$ will be a family of sets which contains the null set, the "universal" set N_n , and is closed under intersection and finite addition and subtraction; $P_1, P_2, \dots$ will be an infinite list of monadic predicate letters; $\bar{n}$ will be the n th formal integer; T will be the theory whose axioms are $\bar{n} \neq \bar{m}$ for each pair n, m such that $n \neq m$; $P_i(\bar{n})$ for each i, n such that $n \in F_i$; and $(x)(P_{i_1}(x) \wedge \dots \wedge P_{i_K}(x) \vee P_{j_1}(x) \wedge \dots \wedge P_{j_N}(x))$ for each pair $\{P_{i_1}, \dots, P_{i_K}\}, \{P_{j_1}, \dots, P_{j_N}\}$ of disjoint finite sets ($K \geq 1, N \geq 1$) of predicate letters from the list $P_1, P_2, \dots$; $A_1, A_2, \dots, A_n \vdash B$ will be used to mean (where $n \geq 0$) that there is a proof of B from assumptions $A_1, A_2, \dots, A_n$ in first order predicate calculus with identity; and $\vdash_T B$ will mean that B is a theorem (valid sentence) of T .

2. Proofs. To prove Theorems 1 and 2 we need the following lemmas:

LEMMA 1: Let $\{F_1, F_2, \dots\}$ be the family of all sets represented in some consistent standard theory S . Then $F_1, F_2, \dots$ is closed under intersection, finite addition, and finite subtraction, and contains the null set and the universal set.

Proof: Closure under intersection is obvious, since if the w.f.f.

(well formed formula) $A(x)$ represents F_i and $B(x)$ represents F_j , then $A(x) \neq B(x)$ represents $F_i \cap F_j$. The null set is represented by any self contradictory w.f.f. with one free variable; the universal set is represented by any valid w.f.f. with one free variable; and finally the sets $F_i \cup \{n_1, n_2, \dots, n_k\}$ and $F_i - \{n_1, n_2, \dots, n_k\}$ are represented by the formulas $A(x) \vee x = \bar{n}_1 \vee \dots \vee x = \bar{n}_k$ and $A(x) \neq x \neq \bar{n}_1 \neq \dots \neq x \neq \bar{n}_k$ respectively.

LEMMA 2: P_i represents F_i in T .

Proof: If $n \in F_i$, then $P_i(\bar{n})$ is an axiom of T , and hence $\vdash_T P_i(\bar{n})$. Now suppose $n \notin F_i$, and consider the following interpretation of T : for all m , $\bar{m}$ designates n ; P_j is assigned the universal set as extension for $j \neq i$, and P_i is assigned as its extension the set $Nn - \{n\}$. This interpretation is a true interpretation of T , and according to it the sentence $P_i(\bar{n})$ is false. Hence $P_i(\bar{n})$ is not a theorem of T .

LEMMA 3. If $\vdash_T P_{i_1}(x) \neq \dots \neq P_{i_M}(x) \supset A(x)$, where $M \geq 0^6$ and $A(x)$ is a w.f.f. with one free variable, then $A(x)$ represents one of the F_i in T .

Proof: (By course-of-values induction on M .) Suppose $M = 0$. Then $\vdash_T A(x)$; hence $A(x)$ represents Nn .

Suppose the lemma holds for $M < N$, and let $\vdash_T P_{i_1}(x) \neq \dots \neq P_{i_N}(x) \supset A(x)$. Let $\bar{s}_1, \bar{s}_2, \dots, \bar{s}_k$ be all of the formal integers that occur in $A(x)$. If $A(\bar{t})$ is never provable unless $\vdash_T P_{i_1}(\bar{t}) \neq \dots \neq P_{i_N}(\bar{t})$ or $t \in \{s_1, s_2, \dots, s_k\}$, then $A(x)$ represents a set that can be obtained from $F_{i_1} \cap F_{i_2} \cap \dots \cap F_{i_N}$ by finite addition, and hence

one of the F_i . Now suppose that $\vdash_T A(\bar{t})$, where it is not the case that $\vdash_T P_{i_1}(\bar{t}) \& \dots \& P_{i_N}(\bar{t})$, and $t \neq s_k$. Let the following be all of the axioms needed for some proof of $A(\bar{t})$ in T :

$P_{j_1}(\bar{t}), \dots, P_{j_U}(\bar{t}); \bar{t} \neq \bar{n}_1, \dots, \bar{t} \neq \bar{n}_j; A_1, \dots, A_S$; where the A_i are all of the axioms not containing $\bar{t}$ used in the proof. If $U = 0$, then, by the Deduction Theorem, $A_1, \dots, A_S \vdash \bar{t} \neq \bar{n}_1 \& \dots \& \bar{t} \neq \bar{n}_j \supset A(\bar{t})$; hence, since t does not occur in $A_1, A_2, \dots, A_S$, and t is not one of the s_i , $A_1, A_2, \dots, A_S \vdash (x)(x \neq \bar{n}_1 \& \dots \& x \neq \bar{n}_j \supset A(x))$. Then $\vdash_T (x)(x \neq \bar{n}_1 \& \dots \& x \neq \bar{n}_j \supset A(x))$, and $A(x)$ represents $Nn - W$, where W has to be a subset of $\{n_1, \dots, n_j\}$, and hence finite. On the other hand, if $U \neq 0$, then by a similar argument

$A_1, A_2, \dots, A_S \vdash (x)(P_{j_1}(x) \& \dots \& P_{j_U}(x) \supset (x \neq \bar{n}_1 \& \dots \& x \neq \bar{n}_j \supset A(x)))$, and so $\vdash_T P_{j_1}(x) \& \dots \& P_{j_U}(x) \supset (x \neq \bar{n}_1 \& \dots \& x \neq \bar{n}_j \supset A(x))$. But we assumed $\vdash_T P_{i_1}(x) \& \dots \& P_{i_N}(x) \supset A(x)$, so

$$(1) \quad \vdash_T (P_{i_1}(x) \& \dots \& P_{i_N}(x) \cdot v. P_{j_1}(x) \& \dots \& P_{j_U}(x)) \supset (x \neq \bar{n}_1 \& \dots \& x \neq \bar{n}_j \supset A(x)).$$

If the P_i 's and the P_j 's are all distinct, then

$\vdash_T (x)(P_{i_1}(x) \& \dots \& P_{i_N}(x) \cdot v. P_{j_1}(x) \& \dots \& P_{j_U}(x))$, and hence $\vdash_T (x \neq \bar{n}_1 \& \dots \& x \neq \bar{n}_j \supset A(x))$ and $A(x)$ represents $Nn - w$, where W is a finite set. And if the P_i 's and the P_j 's are not all distinct, then $P_{i_1}(x) \& \dots \& P_{i_N}(x) \cdot v. P_{j_1}(x) \& \dots \& P_{j_U}(x)$ is quantificationally equivalent (in fact, equivalent by propositional calculus) to $F_{k_1}(x) \& \dots \& P_{k_H}(x) \& (P_{i_r}(x) \& P_{i_r'}(x) \& \dots \& P_{i_r}(D)(x)) \cdot v. P_{j_s}(x) \& P_{j_s'}(x) \& \dots \& P_{j_s}(Q)(x)$, where $P_{k_1}, P_{k_2}, \dots, P_{k_H}$ are all

of the P 's that occur both among the P_i and among the P_j , while $P_{i_r}, P_{i_{r'}}, P_{i_{r''}}, \dots, P_{i_r}(D)$ are the P_i that do not also occur among the P_j , and similarly $P_{j_s}, P_{j_{s'}}, P_{j_{s''}}, \dots, P_{j_s}(Q)$ are the P_j that do not also occur among the P_i . Moreover, D cannot $\equiv 0$ (otherwise $\vdash_{T} P_{i_1}(\bar{t}) \&\dots \& P_{i_N}(\bar{t})$, contrary to the choice of t), and we may assume that $Q \neq 0$ (since otherwise we would have $U < N$, and the lemma would follow by the induction hypothesis⁷). Thus $(x)(P_{i_1}(x) \&\dots \& P_{i_r}(D)(x) \cdot v. P_{j_s}(x) \&\dots \& P_{j_s}(Q)(x))$ is an axiom of T , so that $P_{i_1}(x) \&\dots \& P_{i_N}(x) \cdot v. P_{j_1}(x) \&\dots \& P_{j_U}(x)$ is provably equivalent to $P_{k_1}(x) \&\dots \& P_{k_H}(x)$, where $H < N$. Hence the lemma follows by the induction hypothesis and the fact that since (1) is a theorem of T , $\vdash_{T} P_{k_1}(x) \&\dots \& P_{k_H}(x) \supset (x \neq \bar{n}_1 \&\dots \& x \neq \bar{n}_j \supset A(x))$.

LEMMA 4. The family of all sets represented in an axiomatizable theory is a recursively enumerable family of recursively enumerable sets.

Proof: Let the w.f.f.s of S (where S is any axiomatizable theory) with one free variable be effectively listed as $A_1(x), A_2(x), \dots$. The predicate $P(i, n) =_{df} \vdash_S A_i(\bar{n})$ is a recursively enumerable predicate (to verify this, assuming Church's Thesis, note that it can be written in the form $(\exists x) \text{Prf}(x, n, i)$, where $\text{Prf}(x, n, i)$ is the decidable, and hence recursive, predicate " x is the gödel number of a proof of the formula that results when $\bar{n}$ is put for all occurrences of ' x ' in $A_i(x)$ ". Moreover, $A_i(x)$ represents $\{n \mid P(i, n)\}$, or $\{f(i)\}$ (cf. n. 5), where⁸ $f(i) = S_1^1(e, i)$ and e is a gödel number of P .

Proof of THEOREM 1. By LEMMA 1, we have "only if". To prove "if" (i.e., to show that the conditions given in the theorem are sufficient) we shall show that if $\{F_1, F_2, \dots\}$ satisfies the conditions, then $\{F_1, F_2, \dots\}$ is the family of all sets represented in T (where T is the theory mentioned in §1).

By LEMMA 2, F_i is represented in T (for $i = 1, 2, \dots$). So it suffices to show that for every w.f.f. $A(x)$ of T , $A(x)$ represents one of the F_i . Accordingly, let $A(x)$ be a w.f.f. of T with x as its only free variable, and let $\bar{s}_1, \bar{s}_2, \dots, \bar{s}_k$ be all the formal integers that occur in $A(x)$. If $\vdash_T A(\bar{t})$ only when $t \in \{s_1, s_2, \dots, s_k\}$, then $A(x)$ represents a finite set, and hence one of the F_i (noting that all finite sets can be obtained from the null set by finite addition). Now suppose $\vdash_T A(\bar{t})$ where $t \notin \{s_1, s_2, \dots, s_k\}$. Let the following be all of the axioms needed for some proof of $A(\bar{t})$ in T : $P_{i_1}(\bar{t}), \dots, P_{i_M}(\bar{t})$; $\bar{t} \neq n_1, \dots, \bar{t} \neq n_j$; $A_1, \dots, A_S$; where the A_k are all of the axioms not containing $\bar{t}$ used in the proof. Then $A_1, A_2, \dots, A_S \vdash P_{i_1}(\bar{t}) \& \dots \& P_{i_M}(\bar{t}) \supset (\bar{t} \neq n_1 \& \dots \& \bar{t} \neq n_j \supset A(\bar{t}))$; hence $A_1, A_2, \dots, A_S \vdash P_{i_1}(x) \& \dots \& P_{i_M}(x) \supset (x \neq n_1 \& \dots \& x \neq n_j \supset A(x))$; and hence $\vdash_T P_{i_1}(x) \& \dots \& P_{i_M}(x) \supset (x \neq n_1 \& \dots \& x \neq n_j \supset A(x))$. Then by LEMMA 3, $x \neq n_1 \& \dots \& x \neq n_j \supset A(x)$ represents one of the F_i , and hence $A(x)$ represents one of the F_i (cf. n.7).

Proof of THEOREM 2. The proof is similar to the proof of THEOREM 1, except that LEMMA 4 must also be used for the "only if" part of the theorem, and we must note that what we have given for this case is a recursively enumerable set of axioms. The axiomatizability of T (in the sense of recursive axiomatizability) then follows by Craig's Theorem.

FOOTNOTES

- 1) Terminology: In this paper "set" means set of non-negative integers, except when there is indication to the contrary. A formula $P(x)$ (with one free variable x) is said to "represent" a set S in a theory T if for all integers n , $n \in S$ if and only if $P(\bar{n})$ is a theorem of T (N.B. it is not required that $P(\bar{n})$ should be refutable in T --- i.e., that $\sim P(\bar{n})$ should be provable in T --- when $n \notin S$). The term "represent" comes from Undecidable Theories. ($n \notin S$ is an abbreviation for $\sim n \in S$.)
- 2) By a "standard theory" I mean a "theory in standard formalization" in the sense in which that term is used in Undecidable Theories, in which there are terms (called formal integers in the sequel), say $\bar{0}, \bar{1}, \bar{2}, \dots$ (which may be interpreted as designating $0, 1, 2, \dots$) such that $\bar{n} \neq \bar{m}$ is provable for all n, m such that $n \neq m$.
- 3) A theory in standard formalization is called "axiomatizable" in Undecidable Theories if the set of valid sentences is identical with the set of first-order consequences of some recursive subset (called the set of "axioms"). (Instead of "recursive" it would be better to say "solvable", in the sense of Post, since strictly speaking the recursiveness of a set of formulas depends upon the godel numbering employed, whereas "solvability" is defined directly for sets of expressions in any finite alphabet.)
- 4) A set B will be said to come from a set A by finite addition (resp. finite subtraction) if $B = A \cup W$ (resp. $A - W$) where W is a finite set.

5) Following Kleene, let $\{n\}$ be the n th partial recursive function in the standard enumeration. (This notation is not to be confused with the notation $\{n \mid \dots\}$, for the set of all n satisfying the condition $\dots$, nor with the notation $\{A_1, A_2, A_3, \dots\}$, for the set consisting of $A_1, A_2, A_3, \dots$.) We shall identify each partial recursive function with its domain, for the purpose of enumerating the recursively enumerable sets: thus $\{n\}$ will alternatively be thought of, where convenient, as "the n th recursively enumerable set, in the standard enumeration." A family F is called a "recursively enumerable family of recursively enumerable sets" if the members of F are $\{t(0)\}, \{t(1)\}, \dots$, for some general recursive function t .

6) If $M = 0$, the $\supset$ is to be understood as deleted.

7) More precisely, it would follow from the induction hypothesis that $x \neq \bar{n}_1 \dot{\vee} \dots \vee x \neq \bar{n}_j \supset A(x)$ represents one of the F_i . But $x \neq n_1 \dot{\vee} \dots \vee x \neq n_j \supset A(x)$ represents a superset with at most finitely many more members than the set represented by $A(x)$ (as is clear from the fact that this formula can also be written

$x = n_1 \vee x = \bar{n}_2 \vee \dots \vee x = \bar{n}_j \vee A(x)$ can be obtained from this F_i by finite subtraction. Hence $A(x)$ also represents one of the F_i (since the F_i are closed under finite subtraction).

8) $S_1^1(e, i)$ is a primitive recursive function whose value for any e, i is a Gödel number of $\{x \mid P_e(i, x)\}$, where P_e is the e th 2-place recursively enumerable predicate in the standard enumeration. This function is constructed in Introduction to Metamathematics.

BIBLIOGRAPHY

1. A. Tarski, A. Mostowski, and R. M. Robinson, Undecidable Theories, Amsterdam, North-Holland Publishing Co., 1953.
2. S. C. Kleene, Introduction to Metamathematics, New York, Van Nostrand, 1952.

DISTRIBUTION LIST AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

MATHEMATICAL SCIENCES DIRECTORATE

(ONE COPY UNLESS OTHERWISE NOTED)

ALABAMA

Commander
Army Rocket & Guided Missile
Agency
ATTN: ORDXR-OTL
Redstone Arsenal, Alabama

BELGIUM

Commander (3)
European Office, ARDC
47 Rue Cantersteen
Brussels, Belgium

CALIFORNIA

Applied Mathematics & Statistics
Laboratory
Stanford University
Stanford, California

Department of Mathematics
University of California
Berkeley, California

Commander
Air Force Flight Test Center
ATTN: Technical Library
Edwards Air Force Base,
California

The Rand Corporation (2)
Technical Library
1700 Main Street
Santa Monica, California

Commander
1st Missile Division
ATTN: Operation Analysis
Office
Vandenburg Air Force Base,
California

CONNECTICUT

Department of Mathematics
Yale University
New Haven, Connecticut

DISTRICT OF COLUMBIA

Office of Naval Research (2)
Department of the Navy
ATTN: Code 432
Washington 25, D.C.

Director
Department of Commerce
Office of Technical Services
Washington 25, D.C.

Administrator (6)
National Aeronautics and
Space Administration
ATTN: Documents Library
1520 H Street, N. W.
Washington 25, D.C.

Library
National Bureau of Standards
Washington 25, D.C.

Data Processing Systems Division
National Bureau of Standards
ATTN: Mr. Russel A. Kirsch
Washington 25, D.C.

Applied Mathematics Division
National Bureau of Standards
Washington 25, D.C.

Headquarters, USAF
Assistant for operations
Analysis
Deputy Chief of Staff,
Operations, AF00A
Washington 25, D.C.

Commander (2)
Air Force Office of Scientific
Research
ATTN: SRM
Washington 25, D.C.

Director
U.S. Naval Research Laboratory
ATTN: Library
Washington 25, D.C.

National Science Foundation
Program Director for Math-
ematical Sciences
Washington 25, D.C.

Commander, AFRD (2)
ATTN: Technical Library
Washington 25, D. C.

Canadian Joint Staff
ATTN: DRB/DSIS
2450 Massachusetts Avenue, N.W.
Washington, D.C.

ILLINOIS

Department of Mathematics
Northwestern University
Evanston, Illinois

Laboratories for Applied
Sciences
University of Chicago
Museum of Science and Industry
ATTN: Library, W-305
Chicago 37, Illinois

Department of Mathematics
University of Chicago
Chicago 37, Illinois

Department of Mathematics
University of Illinois
Urbana, Illinois

INDIANA

Department of Mathematics
Purdue University
Lafayette, Indiana

MARYLAND

Institute for Fluid Dynamics
and Applied Mathematics
University of Maryland
College Park, Maryland

Mathematics and Physics Library
The Johns Hopkins University
Baltimore, Maryland

Director
National Security Agency
ATTN: Dr. H. H. Campaign
Fort George G. Meade,
Maryland

MASSACHUSETTS

Department of Mathematics
Harvard University
Cambridge 38, Massachusetts

Department of Mathematics
Massachusetts Institute
of Technology
Cambridge 38, Massachusetts

Commander
Detachment 2, AFRD
ATTN: Technical Library
L. G. Hanscom Field
Bedford, Massachusetts

MICHIGAN

Department of Mathematics
Wayne State University
Detroit 1, Michigan

MINNESOTA

Department of Mathematics
Folwell Hall
University of Minnesota
Minneapolis, Minnesota

Department of Mathematics
Institute of Technology
Engineering Building
University of Minnesota
Minneapolis, Minnesota

MISSOURI

Department of Mathematics
Washington University
St. Louis 8, Missouri

Department of Mathematics
University of Missouri
Columbia, Missouri

NEBRASKA

Commander
Strategic Air Command
ATTN: Operations Analysis
Offutt Air Force Base
Omaha, Nebraska

NEW JERSEY

The James Forrestal Research
Center Library
Princeton University
Princeton, New Jersey

Library
Institute for Advanced Study
Princeton, New Jersey

Department of Mathematics
Fine Hall
Princeton University
Princeton, New Jersey

Commanding General
Signal Corps Engineering
Laboratory
ATTN: SIGFM/EL-RPO
Ft. Monmouth, New Jersey

NEW MEXICO

Commander
Air Force Missile Development
Center
ATTN: Technical Library, HDOI
Holloman Air Force Base,
New Mexico

Commander
Air Force Special Weapons Center
ATTN: Technical Library, SWOI
Kirtland Air Force Base
Albuquerque, New Mexico

NEW YORK

Professor J. Wolfowitz
Mathematics Department
White Hall
Cornell University
Ithaca New York

Department of Mathematics
Syracuse University
Syracuse, New York

Institute for Mathematical
Sciences
New York University
ATTN: Professor M. Kline
25 Waverly Place
New York, New York

Institute for Aeronautical
Sciences
ATTN: Librarian
2 East 64th Street
New York 16, New York

NORTH CAROLINA

Department of Mathematics
University of North Carolina
Chapel Hill, North Carolina

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina

Office of Ordnance Research (2)
Box CM
Duke Station
Durham, North Carolina

Department of Mathematics
Duke University
Duke Station
Durham, North Carolina

OHIO

P.O. Box AA
Wright-Patterson Air Force
Base
Ohio

Commander
Wright Air Development Division
ATTN: WCOSI
Wright-Patterson Air Force
Base
Ohio

Commander
Aeronautical Research
Laboratories
ATTN: Technical Library
Wright-Patterson Air Force
Base
Ohio

USAF Institute of Technology
Library (2)
ATTN: MCLI-ITLIB
Building 125, Area B
Wright-Patterson Air Force
Base
Ohio

PENNSYLVANIA

Department of Mathematics
Carnegie Institute of Technology
Pittsburgh, Pennsylvania

Department of Mathematics
University of Pennsylvania
Philadelphia, Pennsylvania

TENNESSEE

AEDC Library
ARO, Inc.
Arnold AF Station, Tennessee

U.S. Atomic Energy Commission
Technical Information Service
Extension
P.O. Box 62
Oak Ridge, Tennessee

TEXAS

Applied Mechanics Reviews (2)
Southwest Research Institute
8500 Culebra Road
San Antonio 6, Texas

Department of Mathematics
Rice Institute
Houston, Texas

VIRGINIA

Armed Services Technical
Information Agency (10)
ATTN: TIPDR
Arlington Hall Station
Arlington 12, Virginia

WISCONSIN

Department of Mathematics
University of Wisconsin
Madison, Wisconsin

Mathematics Research Center,
U.S. Army
ATTN: R. E. Langer
University of Wisconsin
Madison, Wisconsin