

273 950

Unclassified

Technical Report

No. 254


G. F. Dresselhaus

Ferro- and Antiferromagnetism
in a Cubic Cluster of Spins

11 January 1962

Lincoln Laboratory

MASSACHUSETTS INSTITUTE OF TECHNOLOGY



Unclassified

Unclassified

187

AFESD - 1DR - 62-71

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
LINCOLN LABORATORY


FERRO- AND ANTIFERROMAGNETISM
IN A CUBIC CLUSTER OF SPINS

G. F. DRESSELHAUS

Group 81

TECHNICAL REPORT NO. 254

11 JANUARY 1962



ABSTRACT

The Heisenberg exchange Hamiltonian has been solved exactly for a cubic array of eight spins (each with spin $1/2$). Both energy eigenvalues and thermodynamic functions have been calculated. A Curie and a Néel temperature can be defined and their values determined as a function of the strength of first, second and third neighbor interactions. For some values of the exchange constants, "spiral" antiferromagnetic states exist.

LEXINGTON

MASSACHUSETTS

Unclassified

FERRO- AND ANTIFERROMAGNETISM IN A CUBIC CLUSTER OF SPINS

I. INTRODUCTION

The exact three-dimensional solution of the Heisenberg model for ferromagnetism is not tractable for an infinite lattice. There are various approximate solutions, some of which involve the exact solution for some small cluster of spins.^{1,2} In addition, a number of solutions exist in the literature for a small number of spins in various configurations.³ This report gives the exact solution to the Heisenberg Hamiltonian for eight spins, each with spin 1/2, located on the corners of a cube. Both the eigenvalues and the thermodynamic functions have been calculated.

This simple cluster shows many of the features of an infinite magnetic material. In particular, a ferromagnetic Curie temperature T_C and an antiferromagnetic Néel temperature T_N can be defined, and their values computed as a function of the strength of second and third neighbor interactions. It is shown that, for some values of the interactions, "spiral" antiferromagnetic states exist.⁴ It is also shown that there are regions in which, as the temperature is lowered, ferromagnetic ordering begins, but at still lower temperatures, the system drops into the antiferromagnetic singlet state.

The approximate solution to the infinite simple cubic lattice, using the results contained herein, will be published in a subsequent report.

II. CALCULATION

The Heisenberg Hamiltonian is written

$$\mathcal{H} = -2J[\epsilon + x\mu + y\nu] \quad (1)$$

in which J is the exchange constant, x and y are proportional to the strength of next nearest and third nearest neighbor interactions, and

$$\begin{aligned} \epsilon = & \vec{S}_1 \cdot \vec{S}_2 + \vec{S}_2 \cdot \vec{S}_3 + \vec{S}_3 \cdot \vec{S}_4 + \vec{S}_4 \cdot \vec{S}_1 + \vec{S}_5 \cdot \vec{S}_6 + \vec{S}_6 \cdot \vec{S}_7 + \vec{S}_7 \cdot \vec{S}_8 + \vec{S}_8 \cdot \vec{S}_5 \\ & + \vec{S}_1 \cdot \vec{S}_5 + \vec{S}_2 \cdot \vec{S}_6 + \vec{S}_3 \cdot \vec{S}_7 + \vec{S}_4 \cdot \vec{S}_8 \end{aligned}$$

$$\begin{aligned} \mu = & \vec{S}_1 \cdot \vec{S}_3 + \vec{S}_2 \cdot \vec{S}_4 + \vec{S}_5 \cdot \vec{S}_7 + \vec{S}_6 \cdot \vec{S}_8 + \vec{S}_1 \cdot \vec{S}_6 + \vec{S}_2 \cdot \vec{S}_7 + \vec{S}_3 \cdot \vec{S}_8 + \vec{S}_4 \cdot \vec{S}_5 \\ & + \vec{S}_1 \cdot \vec{S}_8 + \vec{S}_2 \cdot \vec{S}_5 + \vec{S}_3 \cdot \vec{S}_6 + \vec{S}_4 \cdot \vec{S}_7 \end{aligned}$$

$$\nu = \vec{S}_1 \cdot \vec{S}_7 + \vec{S}_2 \cdot \vec{S}_8 + \vec{S}_3 \cdot \vec{S}_5 + \vec{S}_4 \cdot \vec{S}_6 \quad (2)$$




Fig. 1. Cube showing the numbering of the spins.




Fig. 2. Energy spectrum assuming only nearest neighbor interactions, $x = y = 0$. The spectrum is arranged according to the value of the total spin. The Néel state energy is indicated even though it is not a proper eigenstate of the system.

The spin operator for the i^{th} site \vec{S}_i follows the numbering of the sites shown in Fig. 1.

The eigenfunctions of Eq. (1) can be classified according to the total spin S , the z -component of the total spin M_S , and the irreducible representation of the cubic group. Table I lists the symmetry types and energy eigenvalues for the cubic cluster.⁵ Appendix A tabulates the wave functions. Appendix B gives the matrix elements of $\epsilon + x\mu + y\nu$ for the wave functions of Appendix A. The notation for the representations is such that a vector transforms as Γ_4^- . The spectrum for $x = y = 0$ is shown in Fig. 2.

For a given value of the total spin S , the center of gravity of the energy levels E_{CG} is given by the sum rule

$$E_{CG} = -2 \left[\sum_{i < j} J_{ij} \right] [S(S+1) - (3/4)N] / [N(N-1)] \quad (3)$$

in which N is the number of spins (each with spin $1/2$) and J_{ij} is the exchange constant between sites i and j . Equation (3) represents a convenient check of the calculated energy eigenvalues. The Heisenberg approximation for ferromagnetism⁶ corresponds to calculating the partition function, using Eq. (3) for the energy eigenvalues.

The partition function for the cubic cluster is calculated by using the eigenfunctions listed in Table I. From the partition function, the susceptibility and heat capacity for the system are calculated. The susceptibility χ given by the Curie law is written as

$$\chi = NS(S+1) \frac{g^2 \mu_B^2}{3kT} \quad (4)$$

in which N is the number of spins, S is the spin quantum number $g = 2$ and μ_B is the Bohr magneton. The cubic cluster obeys the Curie law with suitable identification of N and S . For high temperatures, $2J/kT \ll 1$, the necessary interpretation of N and S is

$$NS(S+1) = 6 \quad ,$$

and at low temperatures, $2J/kT \gg 1$, the interpretation is changed to $N = 1$ with

$$NS(S+1) = 20 \quad J > 0$$

and

$$NS(S+1) = 0 \quad J < 0 \quad .$$

A plot of the susceptibility vs $1/T$ for $x = y = 0$ is shown in Fig. 3 for both the ferromagnetic and antiferromagnetic states. For convenience, the plot is presented in dimensionless form in terms of $\alpha\chi$ vs $2J/kT$, with $\alpha = 6J/g^2\mu_B^2$. For the ferromagnetic case ($J > 0$) the slope of the curve in Fig. 3(a) has the two asymptotic values indicated by the dashed lines. The ferromagnetic Curie temperature T_C is defined as the position of maximum slope of the χ vs $1/T$ curve. The antiferromagnetic Néel temperature T_N is defined as the position of the maximum in the susceptibility curve. With these definitions, the Curie and Néel temperatures have been determined as a function of x and y . These results are shown in Fig. 4.

The heat capacity is shown for the same parameters in Fig. 5. The positions of T_C and T_N , as determined from the susceptibility, are also indicated. The maximum of the heat capacity curves differs somewhat from T_C and T_N , as determined from the susceptibility.

TABLE I SPIN, SYMMETRY TYPE AND ENERGY EIGENVALUES FOR THE CUBIC CLUSTER OF EIGHT SPINS		
Spin	Representation	Eigenvalue ($\lambda = E/(-2J)$)
4	Γ_1^+	$\lambda = 3 + 3x + y$
3	Γ_4^-	$\lambda = 2 + x$
	Γ_5^+	$\lambda = 1 + x + y$
	Γ_2^-	$\lambda = 3x$
2	Γ_3^+	$\left\{ \lambda = \frac{1}{2}(1-x) \pm \frac{1}{2}[(1-x)^2 + 4(1-y)^2]^{1/2} \right.$
	Γ_3^+	
	Γ_5^+	$\left\{ \lambda = \pm[(1-x)^2 + (1-y)^2 - (1-x)(1-y)]^{1/2} \right.$
	Γ_5^+	
	Γ_1^+	$\left\{ \lambda = -(1-x) \pm [4(1-x)^2 + (1-y)^2 - 3(1-x)(1-y)]^{1/2} \right.$
	Γ_1^+	
	Γ_5^-	$\lambda = 1 - x$
	Γ_4^-	$\lambda = -1 + x$
Γ_3^-	$\lambda = 0$	

TABLE I (Continued)
SPIN, SYMMETRY TYPE AND ENERGY EIGENVALUES
FOR THE CUBIC CLUSTER OF EIGHT SPINS

Spin	Representation	Eigenvalue ($\lambda = E/(-2J)$)
1	Γ_4^- Γ_4^- Γ_4^- Γ_4^+ Γ_4^+ Γ_2^- Γ_2^- Γ_5^+ Γ_5^+ Γ_5^- Γ_3^-	$\left\{ \begin{aligned} &\lambda^3 + 2\lambda^2(1+x+y) + \lambda(-2+7x+5y+3xy-x^2) \\ &\quad + (-2+2y+4x^2+6xy-2x^3) = 0 \end{aligned} \right.$ $\left\{ \begin{aligned} &\lambda = -\frac{1}{2}(1+3x) \pm \frac{1}{2}[5-2x+x^2+4y^2-8y]^{1/2} \end{aligned} \right.$ $\left\{ \begin{aligned} &\lambda = -2+x-y \pm [4-7x-y+4x^2+y^2-xy]^{1/2} \end{aligned} \right.$ $\left\{ \begin{aligned} &\lambda = -\frac{1}{2}(3+x) \pm \frac{1}{2}[5-10x+9x^2-8xy+4y^2]^{1/2} \end{aligned} \right.$ $\lambda = -2x$ $\lambda = -1-x$
0	Γ_1^+ Γ_1^+ Γ_1^+ Γ_3^+ Γ_3^+ Γ_1^- Γ_5^+ Γ_5^-	$\left\{ \begin{aligned} &\lambda^3 + \lambda^2(5+x+3y) + \lambda(-1+18x+14y-9x^2+6xy \\ &\quad -y^2) + 3(-9+3x+5y+3x^2+6xy-y^2-3x^3 \\ &\quad -3x^2y+3xy^2-y^3) = 0 \end{aligned} \right.$ $\left\{ \begin{aligned} &\lambda = -(1+2x) \pm [2-2x+x^2-2y+y^2]^{1/2} \end{aligned} \right.$ $\lambda = -3x$ $\lambda = -(1+x+y)$ $\lambda = -2-x$




Fig. 3. Susceptibility vs $2J/kT$ for (a) $J > 0$ and (b) $J < 0$ and $x = y = 0$. The Curie and Néel temperatures are indicated. See text for constant of proportionality, a .

Fig. 4. T_C and T_N vs x with y as a parameter. The upper curves refer to $J > 0$ and the lower curves to $J < 0$.





Fig. 5. Heat capacity vs $2J/kT$ for $x = y = 0$. T_C and T_N as determined from the susceptibility are indicated.

III. DISCUSSION

The calculation of the magnetic properties for an 8-spin cubic cluster shows interesting features. The antiferromagnetic state gives a good approximation to the magnetic state of the infinite solid. The susceptibility and heat capacity approach the high-temperature limits as the infinite lattice. For $x = y = 0$, the Curie temperature determined from the susceptibility of the 8-spin cluster is given by the Bethe, Peierls, Weiss value of 2.01 (see Ref. 1). The heat capacity, on the other hand, is not well represented by the small cluster. The heat capacity approaches the high-temperature limits, but the susceptibility has the correct low-temperature case (paramagnetic region). In addition, the Curie temperature given by $kT_C/J = 0.90$ compared with the Bethe, Peierls, Weiss value of 2.01.

In order to investigate the number of spins required to represent the magnetic state of an infinite solid, the Heisenberg approximation (i.e., assume the magnetic state of the infinite solid is carried out for clusters of 8, 64 and 216 spins. The results for the heat capacity are shown in Fig. 6. The Curie temperature for these clusters was found from the results are shown in Table II. A rather slow convergence is observed in the antiferromagnetic case. To put it another way, 216 spins are too few to represent the magnetic state of the infinite lattice. The results for the cluster show that fewer spins are needed to represent the antiferromagnetic state than the ferromagnetic state. Comparison with the infinite lattice is not useful for the antiferromagnetic case. The Heisenberg approximation for the antiferromagnetic state which get worse as N increases; that is




Fig. 6. Heat capacity for Heisenberg approximation with 216 spins.

TABLE II VALUES OF T_C CALCULATED IN THE HEISENBERG APPROXIMATION FOR CUBIC CLUSTERS OF VARIOUS NUMBERS OF SPINS	
N	kT_C/J
8	1.3
64	1.8
216	2.2
.	.
.	.
.	.
∞	3.0

Another interesting feature of the 8-spin cubic cluster is the existence of an antiferromagnetic state for $J > 0$. For example, for $y = 0$ and $x < -1/3$, an antiferromagnetic state exists with $J > 0$, which corresponds to a "classical" spiral state;⁴ that is, nearest neighbors are mostly parallel to take advantage of positive J and next nearest neighbors are mostly antiparallel. An examination of the susceptibility curve in the region $y = 0$ and $x \lesssim -1/3$ shows that the system begins to order ferromagnetically because of the greater weighting factor for the $S = 4$ state, but at low temperatures, the system becomes antiferromagnetic because an $S = 0$, Γ_4^+ state is lower in energy.⁷ The condition on x and y for the accidental degeneracy of the $S = 4$, Γ_4^+ state and an $S = 0$, Γ_1^+ state is given by

$$3(1 + 4x_0 + 3x_0^2) + 7y_0 + 12x_0y_0 + 2y_0^2 + 3x_0^2y_0 + 2x_0y_0^2 = 0 \quad (5)$$

In Fig. 4, the dashed branch of the $y = 0$ curve indicates this possibility of a "high" temperature ferromagnetic state. The $y = \pm 0.3$ calculations also allow this possibility but, for simplicity, these branches are not plotted in Fig. 4.

It is also possible to select values of the parameters x and y so that an $S = 0$, Γ_3^+ state is the lowest antiferromagnetic state of the system for $J < 0$. The classical analog to this situation is discussed in Ref. 4.

ACKNOWLEDGMENT

The author has profited from discussions with Drs. G.W. Pratt, Jr., and T.A. Kaplan. He is deeply indebted to Dr. M.S. Dresselhaus for many suggestions and for programming the computation for the IBM 7090.

REFERENCES

1. H.A. Brown and J.M. Luttinger, *Phys. Rev.* **100**, 685 (1955). This article contains references to much of the literature on the subject.
2. J.S. Smart, *J. Phys. Chem. Solids* **20**, 41 (1961).
3. R. Orbach, *Phys. Rev.* **115**, 1181 (1959); L.F. Mattheiss, *Phys. Rev.* **123**, 1209 (1961).
4. T.A. Kaplan, *Phys. Rev.* **116**, 888 (1959); J. Villain, *J. Phys. Chem. Solids* **11**, 303 (1959); A. Yoshimori, *J. Phys. Soc. Japan* **14**, 807 (1959).
5. The notation for the irreducible representations is $\Gamma_1^+ \rightarrow 1$; $\Gamma_2^- \rightarrow xyz$; $\Gamma_3^+ \rightarrow x^2 + \omega y^2 + \omega^2 z^2$, $x^2 + \omega^2 y^2 + \omega z^2$; $\Gamma_4^- \rightarrow x, y, z$; and $\Gamma_5^+ \rightarrow yz, xz, xy$.
6. J.H. VanVleck, *Electric and Magnetic Susceptibilities* (Oxford University Press, London, England, 1932) p. 322.
7. This effect has many of the same features as that used by H. Sato and A. Arrott, *Phys. Rev.* **114**, 1427 (1959), to explain the magnetic behavior of some Fe-Al alloys which are ferromagnetic at high temperatures but antiferromagnetic at low temperatures.

APPENDIX A
EIGENFUNCTIONS OF THE 8-SPIN PROBLEM

The notation is as follows:

$$\text{spin configuration } (137) \equiv \beta(1) \alpha(2) \beta(3) \alpha(4) \alpha(5) \alpha(6) \beta(7) \alpha(8)$$

in which $\alpha(i) \rightarrow$ spin "i" up and $\beta(i) \rightarrow$ spin "i" down. Only the maximum M_S state is listed, since the lower M_S states can be obtained by application of a lowering operator. For the representations Γ_3^\pm , only one eigenfunction is given; the other can be obtained by taking the complex conjugate of the given function. For Γ_4^\pm and Γ_5^\pm , only one function is given; the other two can be obtained by performing the appropriate symmetry operations. When the same representation occurs several times, a lower case Greek letter is used to distinguish the representation. An "i" is used to label the state in the case of a multidimensional representation.

The states are eigenstates of S^2 and S_z , but not necessarily of \mathcal{K} . The wave function is a sum of the spin configurations. The coefficient of each spin configuration is given in Table A-I. The absolute value of the square of the wave function is the last entry in the table.

TABLE A-I NUMERICAL COEFFICIENTS OF THE VARIOUS SPIN CONFIGURATIONS			
S = 3			
Spin Configuration	Γ_2^-	Γ_4^-	Γ_5^+
(1)	+	-	-
(2)	-	+	-
(3)	+	+	+
(4)	-	-	+
(5)	-	-	+
(6)	+	+	+
(7)	-	+	-
(8)	+	-	-
$ \Gamma_i ^2$	8	8	8

TABLE A-1 (Continued)
 NUMERICAL COEFFICIENTS
 OF THE VARIOUS SPIN CONFIGURATIONS

S = 2									
Spin Configuration	$\Gamma_{1\alpha}^+$	$\Gamma_{1\beta}^+$	$\Gamma_{3\alpha,i}^+$	$\Gamma_{3\beta,i}^+$	Γ_{3i}^-	Γ_{4i}^-	$\Gamma_{5\alpha,i}^+$	$\Gamma_{5\beta,i}^+$	Γ_{5i}^-
(12)	+	+	1			+			+
(23)	+	+	ω			+			-
(34)	+	+	1			+			+
(14)	+	+	ω			+			-
(56)	+	+	1			-			-
(67)	+	+	ω			-			+
(78)	+	+	1			-			-
(58)	+	+	ω			-			+
(15)	+	+	ω^2				+	+	
(26)	+	+	ω^2				-	-	
(37)	+	+	ω^2				+	+	
(48)	+	+	ω^2				-	-	
(17)		-6						-2	
(28)		-6						+2	
(35)		-6						-2	
(46)		-6						+2	

TABLE A-I (Continued)
 NUMERICAL COEFFICIENTS
 OF THE VARIOUS SPIN CONFIGURATIONS

S = 2									
Spin Configuration	$\Gamma_{1\alpha}^+$	$\Gamma_{1\beta}^+$	$\Gamma_{3\alpha,i}^+$	$\Gamma_{3\beta,i}^+$	Γ_{3i}^-	Γ_{4i}^-	$\Gamma_{5\alpha,i}^+$	$\Gamma_{5\beta,i}^+$	Γ_{5i}^-
(13)	-	+		1	1	-2	-	+	
(24)	-	+		1	-1	-2	+	-	
(57)	-	+		1	-1	2	-	+	
(68)	-	+		1	1	2	+	-	
(27)	-	+		ω^2	$-\omega^2$				
(36)	-	+		ω^2	ω^2				
(18)	-	+		ω^2	ω^2				
(45)	-	+		ω^2	$-\omega^2$				
(38)	-	+		ω	ω				
(47)	-	+		ω	$-\omega$				
(25)	-	+		ω	$-\omega$				
(16)	-	+		ω	ω				
$ \Gamma_i ^2$	24	168	12	12	12	24	8	24	24
where $\omega^3 = 1$.									

TABLE A-1 (Continued)
 NUMERICAL COEFFICIENTS
 OF THE VARIOUS SPIN CONFIGURATIONS

S = 1											
Spin Configuration	$\Gamma_2^{-\alpha}$	$\Gamma_2^{-\beta}$	Γ_3^{-i}	$\Gamma_4^{+\alpha, i}$	$\Gamma_4^{+\beta, i}$	$\Gamma_4^{-\alpha, i}$	$\Gamma_4^{-\beta, i}$	$\Gamma_4^{-\gamma, i}$	$\Gamma_5^{+\alpha, i}$	$\Gamma_5^{+\beta, i}$	Γ_5^{-i}
(123)	+	-	1			-	-1/2	-1/3	-	-	
(134)	+	-	1			-	-1/2	-1/3	-	-	
(568)	+	-	1			+	1/2	1/3	+	+	
(678)	+	-	1			+	1/2	1/3	+	+	
(236)	+	-	-e	+			+	1/6		+	+
(126)	+	-	+e ²	-			+	1/6		+	-
(367)	+	-	-e	-			-	-1/6		-	-
(348)	+	-	e ²	-			+	1/6		+	-
(378)	+	-	e ²	+			-	-1/6		-	+
(158)	+	-	-e	-			-	-1/6		-	-
(156)	+	-	e ²	+			-	-1/6		-	+
(148)	+	-	-e	+			+	1/6		+	+
(234)	-	+	-1			-	-1/2	-1/3	+	+	
(124)	-	+	-1			-	-1/2	-1/3	+	+	
(267)	-	+	e	+			-	-1/6		+	-
(145)	-	+	e	-			+	1/6		-	+
(256)	-	+	-e ²	-			-	-1/6		+	+
(347)	-	+	-e ²	+			+	1/6		-	-
(578)	-	+	-1			+	1/2	1/3	-	-	
(567)	-	+	-1			+	1/2	1/3	-	-	
(125)	-	+	-e ²	+			+	1/6		-	-
(478)	-	+	-e ²	-			-	-1/6		+	+
(237)	-	+	e	-			+	1/6		-	+
(458)	-	+	e	+			-	-1/6		+	-
(136)		2						+	+	-	
(138)		2						+	+	-	
(168)		2						-	-	+	
(368)		2						-	-	+	
(247)		-2						+	-	+	
(257)		-2						-	+	-	
(245)		-2						+	-	+	
(457)		-2						-	+	-	

TABLE A-I (Continued)
 NUMERICAL COEFFICIENTS
 OF THE VARIOUS SPIN CONFIGURATIONS

S = 1											
Spin Configuration	$\Gamma_{2^a}^-$	$\Gamma_{2^b}^-$	$\Gamma_{3^i}^-$	$\Gamma_{4^a,i}^+$	$\Gamma_{4^b,i}^+$	$\Gamma_{4^a,i}^-$	$\Gamma_{4^b,i}^-$	$\Gamma_{4^{\gamma},i}^-$	$\Gamma_{5^a,i}^+$	$\Gamma_{5^b,i}^+$	$\Gamma_{5^i}^-$
(127)	+	+	- ω		+	+	-1/2	1/6		-	+
(345)	+	+	- ω		+	+	-1/2	1/6		-	+
(456)	+	+	- ω		-	-	1/2	-1/6		+	-
(278)	+	+	- ω		-	-	1/2	-1/6		+	-
(235)	+	+	ω^2		-	+	-1/2	1/6		-	-
(246)	+	+	1			+	1/2	-2/3		-2	
(467)	+	+	ω^2		+	-	1/2	-1/6		+	+
(248)	+	+	1			+	1/2	-2/3		-2	
(147)	+	+	ω^2		-	+	-1/2	1/6		-	-
(157)	+	+	1			-	-1/2	2/3		2	
(258)	+	+	ω^2		+	-	1/2	-1/6		+	+
(357)	+	+	1			-	-1/2	2/3		2	
(238)	-	-	- ω^2		+	+	-1/2	1/6		+	-
(146)	-	-	- ω^2		+	+	-1/2	1/6		+	-
(268)	-	-	-1			-	-1/2	2/3		-2	
(135)	-	-	-1			+	1/2	-2/3		2	
(356)	-	-	ω		+	-	1/2	-1/6		-	-
(346)	-	-	ω		-	+	-1/2	1/6		+	+
(128)	-	-	ω		-	+	-1/2	1/6		+	+
(468)	-	-	-1			-	-1/2	2/3		-2	
(358)	-	-	- ω^2		-	-	1/2	-1/6		-	+
(178)	-	-	ω		+	-	1/2	-1/6		-	-
(167)	-	-	- ω^2		-	-	1/2	-1/6		-	+
(137)	-	-	-1			+	1/2	-2/3		2	
$ \Gamma_i ^2$	48	80	48	16	16	32	24	16/3	16	80	32

TABLE A-I (Continued)
 NUMERICAL COEFFICIENTS
 OF THE VARIOUS SPIN CONFIGURATIONS

S = 0								
Spin Configuration	$\Gamma_{1^a}^+$	$\Gamma_{1^b}^+$	$\Gamma_{1^c}^+$	Γ_1^-	$\Gamma_{3^a,i}^+$	$\Gamma_{3^b,i}^+$	$\Gamma_{5^i}^+$	$\Gamma_{5^i}^-$
(1234)	+	-	3		$-2\omega^2$			
(5678)	+	-	3		$-2\omega^2$			
(1256)	+	-	3		-2ω			
(3478)	+	-	3		-2ω			
(2367)	+	-	3		-2			
(1456)	+	-	3		-2			
(2347)	-		3				-	
(1348)	-		3				-	
(1245)	-		3				+	
(1236)	-		3				+	
(3678)	-		3				+	
(4578)	-		3				+	
(1568)	-		3				-	
(2567)	-		3				-	
(1368)	+	3	3					
(2457)	+	3	3					
(1278)			8			2		
(3456)			8			2		
(1467)			8			2ω		
(2358)			8			2ω		
(1357)			8			$2\omega^2$		
(2468)			8			$2\omega^2$		

TABLE A-1 (Continued)
 NUMERICAL COEFFICIENTS
 OF THE VARIOUS SPIN CONFIGURATIONS

S = 0								
Spin Configuration	$\Gamma_{1\alpha}^+$	$\Gamma_{1\beta}^+$	$\Gamma_{1\gamma}^+$	Γ_1^-	$\Gamma_{3\alpha,i}^+$	$\Gamma_{3\beta,i}^+$	Γ_5^+	Γ_5^-
(2345)		-	-2		ω^2	ω^2		
(1346)		-	-2		ω^2	ω^2		
(1247)		-	-2		ω^2	ω^2		
(1238)		-	-2		ω^2	ω^2		
(1678)		-	-2		ω^2	ω^2		
(2578)		-	-2		ω^2	ω^2		
(3568)		-	-2		ω^2	ω^2		
(4567)		-	-2		ω^2	ω^2		
(1378)		-	-2		ω	ω		-
(2478)		-	-2		ω	ω		-
(3457)		-	-2		ω	ω		+
(3468)		-	-2		ω	ω		+
(1356)		-	-2		ω	ω		-
(2456)		-	-2		ω	ω		-
(1257)		-	-2		ω	ω		+
(1268)		-	-2		ω	ω		+
(1367)		-	-2		1	1	-	+
(2467)		-	-2		1	1	+	+
(2357)		-	-2		1	1	+	-
(2368)		-	-2		1	1	-	-
(2458)		-	-2		1	1	-	+
(1358)		-	-2		1	1	+	+
(1468)		-	-2		1	1	+	-
(1457)		-	-2		1	1	-	-

TABLE A-1 (Continued)
 NUMERICAL COEFFICIENTS
 OF THE VARIOUS SPIN CONFIGURATIONS

S = 0								
Spin Configuration	$\Gamma_{1\alpha}^+$	$\Gamma_{1\beta}^+$	$\Gamma_{1\gamma}^+$	Γ_1^-	$\Gamma_{3\alpha,i}^+$	$\Gamma_{3\beta,i}^+$	Γ_{5i}^+	Γ_{5i}^-
(1237)		+	-2	+	- ω	ω		
(2348)		+	-2	+	-1	1	+	
(1345)		+	-2	+	- ω	ω		
(1246)		+	-2	+	-1	1	-	
(2678)		+	-2	+	- ω	ω		
(4568)		+	-2	+	- ω	ω		
(3467)		+	-2	+	- ω^2	ω^2		-
(2356)		+	-2	+	- ω^2	ω^2		+
(3578)		+	-2	+	-1	1	-	
(1567)		+	-2	+	-1	1	+	
(1478)		+	-2	+	- ω^2	ω^2		+
(1258)		+	-2	+	- ω^2	ω^2		-
(1235)		+	-2	-	-1	1	-	
(2346)		+	-2	-	- ω	ω		
(1347)		+	-2	-	-1	1	+	
(1248)		+	-2	-	- ω	ω		
(1267)		+	-2	-	- ω^2	ω^2		-
(2568)		+	-2	-	-1	1	+	
(1456)		+	-2	-	- ω^2	ω^2		+
(1578)		+	-2	-	- ω	ω		
(3458)		+	-2	-	- ω^2	ω^2		-
(3567)		+	-2	-	- ω	ω		
(4678)		+	-2	-	-1	1	-	
(2378)		+	-2	-	- ω^2	ω^2		+
$ \Gamma_i ^2$	16	72	720	24	72	72	24	24

APPENDIX B
MATRIX ELEMENTS

In this appendix the matrix elements of the form $\langle \Gamma_i | \epsilon + x\mu + y\nu | \Gamma_j \rangle$ are given. The non-degenerate states can be obtained directly from Table I, so only the matrix elements for degenerate states are listed.

For $S = 2$

	$\Gamma_1^+\alpha$	$\Gamma_1^+\beta$
$\Gamma_1^+\alpha$	$-\frac{3}{2} + \frac{5}{2}x - y$	$\frac{\sqrt{7}}{2}(1-x)$
$\Gamma_1^+\beta$	$\frac{\sqrt{7}}{2}(1-x)$	$-\frac{1}{2}(1+x) + y$

	$\Gamma_3^-\alpha, i$	$\Gamma_3^-\alpha, ii$	$\Gamma_3^-\beta, i$	$\Gamma_3^-\beta, ii$
$\Gamma_3^-\alpha, i$	$(1-x)$	0	0	$-\omega(1-y)$
$\Gamma_3^-\alpha, ii$	0	$(1-x)$	$-\omega^2(1-y)$	0
$\Gamma_3^-\beta, i$	0	$-\omega(1-y)$	0	0
$\Gamma_3^-\beta, ii$	$-\omega^2(1-y)$	0	0	0

	$\Gamma_5^+\alpha, i$	$\Gamma_5^+\beta, i$
$\Gamma_5^+\alpha, i$	$\frac{1}{2}(1+x) - y$	$\frac{\sqrt{3}}{2}(1-y)$
$\Gamma_5^+\beta, i$	$\frac{\sqrt{3}}{2}(1-y)$	$-\frac{1}{2}(1+x) + y$

For $S = 1$

	$\Gamma_4^{-\alpha, i}$	$\Gamma_4^{-\beta, i}$	$\Gamma_4^{-\gamma, i}$
$\Gamma_4^{-\alpha, i}$	$-\frac{1}{8}(1 + 14x + y)$	$\frac{1}{8\sqrt{3}}(11 - 6x - 5y)$	$\frac{1}{4}\sqrt{\frac{5}{3}}(1 - y)$
$\Gamma_4^{-\beta, i}$	$\frac{1}{8\sqrt{3}}(11 - 6x - 5y)$	$\frac{1}{24}(7 - 30x - 25y)$	$\frac{5\sqrt{5}}{12}(1 - y)$
$\Gamma_4^{-\gamma, i}$	$\frac{1}{4}\sqrt{\frac{5}{3}}(1 - y)$	$\frac{5\sqrt{5}}{12}(1 - y)$	$\frac{1}{6}(-13 + 6x - 5y)$

	$\Gamma_4^{+\alpha, i}$	$\Gamma_4^{+\beta, i}$
$\Gamma_4^{+\alpha, i}$	$\frac{1}{2} - \frac{3}{2}x - y$	$-\frac{1}{2}(1 - x)$
$\Gamma_4^{+\beta, i}$	$-\frac{1}{2}(1 - x)$	$-\frac{3}{2}(1 + x) + y$

	$\Gamma_2^{-\alpha}$	$\Gamma_2^{-\beta}$
$\Gamma_2^{-\alpha}$	$-\frac{1}{4} - x - \frac{3}{4}y$	$-\frac{\sqrt{15}}{4}(1 - y)$
$\Gamma_2^{-\beta}$	$-\frac{\sqrt{15}}{4}(1 - y)$	$-\frac{15}{4} + 3x - \frac{5}{4}y$

	$\Gamma_5^{+\alpha, i}$	$\Gamma_5^{+\beta, i}$
$\Gamma_5^{+\alpha, i}$	$-\frac{3}{2} + \frac{1}{2}x - 1$	$\frac{\sqrt{5}}{2}(1 - x)$
$\Gamma_5^{+\beta, i}$	$\frac{\sqrt{5}}{2}(1 - x)$	$-\frac{3}{2}(1 + x) + y$

For $S = 0$

	$\Gamma_1^+\alpha$	$\Gamma_1^+\beta$	$\Gamma_1^+\gamma$
$\Gamma_1^+\alpha$	$-3y$	$-\frac{3}{\sqrt{2}}(1-x)$	0
$\Gamma_1^+\beta$	$-\frac{3}{\sqrt{2}}(1-x)$	$-3+x-y$	$-\sqrt{\frac{5}{2}}(1-x)$
$\Gamma_1^+\gamma$	0	$-\sqrt{\frac{5}{2}}(1-x)$	$-2-2x+y$

	$\Gamma_3^+\alpha, i$	$\Gamma_3^+\beta, i$
$\Gamma_3^+\alpha, i$	$-2x-y$	$-(1-x)$
$\Gamma_3^+\beta, i$	$-(1-x)$	$-2-2x+y$