

UNCLASSIFIED

AD 274 596

*Reproduced
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

274596

274 596

MEMORANDUM
RM-2993-PR
APRIL 1962

CATALOGED BY ASTIA
AS AD NO. 1

A LINEAR PROGRAM OF PRAGER'S
Notes on Linear Programming
and Extensions - Part 60

Oliver Gross

PREPARED FOR:
UNITED STATES AIR FORCE PROJECT RAND

739 200

ASTIA
APR 23 1962
62-3-2
TISIA

The RAND Corporation
SANTA MONICA • CALIFORNIA

MEMORANDUM

RM-2993-PR

APRIL 1962

A LINEAR PROGRAM OF PRAGER'S
Notes on Linear Programming
and Extensions - Part 60

Oliver Gross

This research is sponsored by the United States Air Force under Project RAND - Contract No. AF 49(638)-700 - monitored by the Directorate of Development Planning, Deputy Chief of Staff, Research and Technology, Hq USAF. Views or conclusions contained in this Memorandum should not be interpreted as representing the official opinion or policy of the United States Air Force. Permission to quote from or reproduce portions of this Memorandum must be obtained from The RAND Corporation.

The **RAND** *Corporation*
1700 MAIN ST • SANTA MONICA • CALIFORNIA

PREFACE

An infinite-dimensional linear-programming problem proposed by W. Prager is solved in this Memorandum, which involves basic mathematical research. The problem arose in an elastico-plastic structural-design context. It is hoped that the solution of this problem will provide insight into the solution of many others of the same general character.

SUMMARY

For the problem of minimizing the integral $\int_0^1 f(x)dx$
subject to the constraints

$$\begin{aligned} - f(x) &\leq xg(y) \leq f(x) && \text{if } 0 \leq x \leq y \leq 1, \\ - f(x) &\leq xg(y) - x + y \leq f(x) && \text{if } 0 \leq y \leq x \leq 1, \end{aligned}$$

the author exhibits solutions and proves both that they satisfy
the constraints and that they have the extremizing property.

The problem, proposed by W. Prager, arose in an elastico-
plastic, structural-design context.

CONTENTS

PREFACE	iii
SUMMARY	v
Section	
1. STATEMENT OF THE PROBLEM AND OF ITS SOLUTION . .	1
2. VERIFICATION OF FEASIBILITY OF SOLUTION	2
3. PROOF OF OPTIMALITY OF SOLUTION	10
LIST OF RAND NOTES ON LINEAR PROGRAMMING AND EXTENSIONS .	17

A LINEAR PROGRAM OF PRAGER'S

1. STATEMENT OF THE PROBLEM AND OF ITS SOLUTION

In a recent letter to R. Bellman, W. Prager proposed the following problem:

It is required to find real-valued, nonnegative integrable functions f, g , on $[0, 1]$ so as to minimize

$$\int_0^1 f(x) dx,$$

subject to the constraints

$$\begin{aligned} -f(x) &\leq xg(y) \leq f(x) && \text{if } 0 \leq x \leq y \leq 1, \\ -f(x) &\leq xg(y) - x + y \leq f(x) && \text{if } 0 \leq y \leq x \leq 1. \end{aligned}$$

The foregoing problem arose in a structural-design context of elastico-plastic type. Although the writer is not familiar with the details of the model, it is hoped that the solution of the present problem will provide insight toward the solution of an entire class of problems of the same general character.

The solution (f_0, g_0) given below is not the only solution to the problem, as the subsequent analysis indicates. It turns out that g_0 can be perturbed slightly in its tail (beyond $\sqrt{2}/2$) without altering feasibility. Moreover, since the other component (f_0) is defined only in terms of the values of g_0 on the interval $[0, \sqrt{2}/2]$, integrability of g_0 beyond $\sqrt{2}/2$ is inessential. It is thus seen that there are 2^c solutions of the problem, where c is the cardinality of the continuum.

Finally we remark that, corresponding to an optimal (f, g) pair, if $\bar{f} \geq f$, with equality holding almost everywhere, then $(\bar{f}, g)$ is an optimal feasible solution to the problem.

Having dispensed with this digression, we now contend that the following (f_0, g_0) is a solution to Prager's problem:

$$g_0(x) = \max \left[0, \lambda - \log \left(x + \sqrt{x^2 + \frac{1}{2}} \right) \right],$$

where

$$\lambda = \frac{1}{2} \left[1 + \log \left(\frac{\sqrt{2} + 1}{2} \right) \right],$$

and

$$f_0(x) = \begin{cases} xg_0(x) & \text{if } 0 \leq x < \frac{\sqrt{2}}{2}, \\ -xg_0 \left(\sqrt{x^2 - \frac{1}{2}} \right) + x - \sqrt{x^2 - \frac{1}{2}} & \text{if } \frac{\sqrt{2}}{2} \leq x \leq 1. \end{cases}$$

In Sec. 2 we shall show that (f_0, g_0) satisfies the constraints imposed. In the final Sec. 3 we shall then show that (f_0, g_0) minimizes the objective integral over all pairs (f, g) satisfying the constraints.

2. VERIFICATION OF FEASIBILITY OF SOLUTION

It is clear that the given constraints can be rewritten as follows:

$$g(x) \geq 0, \quad (A)$$

$$f(x) \geq 0, \quad (B)$$

$$f(x) \geq xg(y) \quad \text{if } 0 \leq x \leq y, \quad (C)$$

$$f(x) \geq xg(y) - x + y \quad \text{if } 0 \leq y \leq x, \quad (D)$$

$$f(x) \geq -xg(y) + x - y \quad \text{if } 0 \leq y \leq x. \quad (E)$$

We remark that the given constraint

$$-f(x) \leq xg(y) \quad \text{if } 0 \leq x \leq y$$

is a trivial consequence of the nonnegativity conditions, (A) and (B).

Before we show that the given pair (f_0, g_0) satisfies the constraints (A) ... (E), a few preliminary remarks are in order.

Remark 1. g_0 is decreasing and convex.

To see this it is sufficient to show that on the set where $g_0 > 0$, its derivative is negative and increasing. But on this set, we have

$$(1) \quad g_0'(x) = -\frac{1}{\sqrt{x^2 + \frac{1}{2}}}.$$

It is immediate that the right-hand member of (1) has the requisite properties.

Remark 2. $g_0(0) < 1$.

A numerical calculation gives

$$g_0(0) \approx .9407 < 1.$$

Remark 3. $g_0(x) < 1$.

This is an immediate consequence of the two preceding remarks.

Remark 4. $g_0\left(\frac{\sqrt{2}}{2}\right) > 0$.

A numerical calculation gives

$$g_0\left(\frac{\sqrt{2}}{2}\right) \approx .0593 > 0.$$

We now proceed with the proof of feasibility of (f_0, g_0) .

That g_0 satisfies the constraint (A) is obvious from the definition of g_0 .

Moreover, it follows from (A) and the definition of f_0 that (B) is satisfied by f_0 on the interval $[0, \sqrt{2}/2)$. To see that f_0 is nonnegative on the remaining interval, it suffices to show that

$$f_0(x) = -xg_0\left(\sqrt{x^2 - \frac{1}{2}}\right) + x - \sqrt{x^2 - \frac{1}{2}} \geq 0 \quad \text{if } x \geq \frac{\sqrt{2}}{2}.$$

But, at $x = \sqrt{2}/2$, the right-hand member of the above equation has the value

$$-\frac{\sqrt{2}}{2} g_0(0) + \frac{\sqrt{2}}{2} = \frac{\sqrt{2}}{2} (1 - g_0(0)) > 0,$$

by virtue of remark 2. Consequently (B) will be established if we can show that f is increasing on the interval $[\sqrt{2}/2, 1]$. But, since the function $\sqrt{x^2 - 1/2}$ maps the interval $(\sqrt{2}/2, 1)$ into the interval $(0, \sqrt{2}/2)$, and g_0 satisfies $g_0 > 0$ on this latter interval by virtue of remarks 1 through 4, we may differentiate $f_0(x)$ using

$$g_0(x) = \lambda - \log \left(x + \sqrt{x^2 + \frac{1}{2}} \right).$$

Thus, we have

$$\begin{aligned} f_0'(x) &= -xg_0'\left(\sqrt{x^2 - \frac{1}{2}}\right) \frac{d}{dx} \sqrt{x^2 - \frac{1}{2}} - g_0\left(\sqrt{x^2 - \frac{1}{2}}\right) \\ &\quad + 1 - \frac{d}{dx} \sqrt{x^2 - \frac{1}{2}}. \end{aligned}$$

But, from (1) we see that

$$g_0' \left(\sqrt{x^2 - \frac{1}{2}} \right) = -\frac{1}{x}.$$

Substituting this value in the above formula for f_0' gives

$$f_0'(x) = 1 - g_0 \left(\sqrt{x^2 - \frac{1}{2}} \right) > 0,$$

by virtue of remark 3. This establishes the desired monotonicity of f_0 . Hence the constraint (B) is satisfied by (f_0, g_0) .

To see that (C) is satisfied by the proposed solution, we must prove that

$$f_0(x) \geq xg_0(y) \quad \text{if } 0 \leq x \leq y.$$

But, by remark 1, g_0 is decreasing. Hence the right-hand member of the desired relation will be maximal at $y = x$, and thus it is sufficient to establish that

$$(2) \quad f_0(x) \geq xg_0(x) \quad \text{for all } x \in [0, 1].$$

But this last relation holds with equality on the interval $[0, \sqrt{2}/2)$, by the definition of f_0 on this interval. Consequently, we need to show only that the relation holds on the interval $[\sqrt{2}/2, 1]$, i.e., from the definition of f_0 there, that

$$f_0(x) = -xg_0 \left(\sqrt{x^2 - \frac{1}{2}} \right) + x - \sqrt{x^2 - \frac{1}{2}} \geq xg_0(x) \quad \text{if } \frac{\sqrt{2}}{2} \leq x \leq 1.$$

On the set on which $g_0(x) = 0$, this last relation holds automatically since by (B) we have $f_0(x) \geq 0$. Consequently, we need verify the above only for points for which $g_0(x) > 0$

and $\sqrt{2}/2 \leq x \leq 1$. But at $x = \sqrt{2}/2$, we have

$$\begin{aligned} f_0\left(\frac{\sqrt{2}}{2}\right) &= -\frac{\sqrt{2}}{2} g_0(0) + \frac{\sqrt{2}}{2} \\ &= -\frac{\sqrt{2}}{2} \left\{ \frac{1}{2} \left[1 + \log\left(\frac{\sqrt{2}+1}{2}\right) \right] - \log\left(\frac{\sqrt{2}}{2}\right) \right\} \\ &\quad + \frac{\sqrt{2}}{2} . \end{aligned}$$

Upon simplification, using well-known properties of the logarithm and a little algebra, we obtain

$$f_0\left(\frac{\sqrt{2}}{2}\right) = \frac{\sqrt{2}}{4} - \frac{\sqrt{2}}{4} \log(\sqrt{2}+1) .$$

On the other hand, the right-hand member of the desired inequality yields, at $x = \sqrt{2}/2$,

$$\begin{aligned} \frac{\sqrt{2}}{2} g_0\left(\frac{\sqrt{2}}{2}\right) &= \frac{\sqrt{2}}{2} \left\{ \frac{1}{2} \left[1 + \log\left(\frac{\sqrt{2}+1}{2}\right) \right] - \log\left(\frac{\sqrt{2}}{2} + 1\right) \right\} \\ &= \frac{\sqrt{2}}{4} - \frac{\sqrt{2}}{4} \log(\sqrt{2}+1) . \end{aligned}$$

Thus, the desired inequality is sharp at $x = \sqrt{2}/2$. To establish it beyond this point on the set on which $g_0 > 0$, it suffices to show that the function

$$f_0(x) - xg_0(x) = -xg_0\left(\sqrt{x^2 - \frac{1}{2}}\right) + x - \sqrt{x^2 - \frac{1}{2}} - xg_0(x)$$

is increasing on this set.

Upon differentiating, we obtain

$$\begin{aligned} [f_0(x) - xg_0(x)]' &= \\ &= -xg_0' \left(\sqrt{x^2 - \frac{1}{2}} \right) \frac{d}{dx} \sqrt{x^2 - \frac{1}{2}} + 1 - \frac{d}{dx} \sqrt{x^2 - \frac{1}{2}} \\ &= -xg_0'(x) - g_0(x) . \end{aligned}$$

By the previous relation,

$$g_0' \left(\sqrt{x^2 - 1/2} \right) = -1/x,$$

this reduces to

$$\left(f_0(x) - xg_0(x) \right)' = \left(1 - g_0(x) \right)' - xg_0'(x) .$$

But by remark 3, we have $1 - g_0(x) > 0$, and by remark 1, we have $-xg_0'(x) > 0$, whence

$$[f_0(x) - xg_0(x)]' > 0.$$

Thus (C) is established, and we have moreover also shown that

$$(2) \quad f_0(x) \geq xg_0(x) \quad \text{for all } x \in [0, 1] .$$

To establish (D), we are required to show that

$$f_0(x) \geq xg_0(y) - x + y \quad \text{if } 0 \leq y \leq x .$$

By remark 1, g_0 is convex. Thus, the right-hand member of the desired inequality is convex in y and hence is maximal at an end point $y = 0$ or $y = x$. Hence it suffices to establish

the inequality at each of these points. But at $y = 0$, the right-hand member becomes $x [g_0(0) - 1] \leq 0$, by remark 2. But since $f_0(x) \geq 0$ we see that the inequality is satisfied at the lower end point. At $y = x$ the inequality reduces to

$$f_0(x) \geq xg_0(x) .$$

But this last is the inequality (2) already established. Thus condition (D) is satisfied.

Finally, to establish (E) we are required to show that

$$f_0(x) \geq -xg_0(y) + x - y \quad \text{if } 0 \leq y \leq x.$$

By remark 1, g_0 is convex; hence the right-hand member of the desired inequality is concave in y . Consequently, if there exists an interior point $y_0 \in (0, x)$ at which the derivative of the right-hand member with respect to y vanishes, this member will be maximal at that point. Upon differentiating with respect to y and equating to zero, we obtain

$$-xg_0'(y) - 1 = 0,$$

or

$$g_0'(y) = -\frac{1}{x} .$$

But recall that this equation is satisfied by

$$y_0 = \sqrt{x^2 - \frac{1}{2}} < x,$$

provided of course that $\sqrt{2}/2 \leq x \leq 1$. Moreover, y_0 is in the

range of $g_0 > 0$. Thus for $x \in [\sqrt{2}/2, 1]$ it is enough to show that

$$f_0(x) \geq -xg_0\left(\sqrt{x^2 - \frac{1}{2}}\right) + x - \sqrt{x^2 - \frac{1}{2}}.$$

On this interval, however, equality holds by the definition of f_0 . Thus it remains to show that the desired inequality holds on the interval $0 \leq x < \sqrt{2}/2$; i.e., we are required to show that

$$xg_0(x) \geq -xg_0(y) + x - y \quad \text{if } 0 \leq y \leq x < \frac{\sqrt{2}}{2}.$$

Now since g_0 is strictly convex on the set on which $g_0 > 0$, and $\sqrt{x^2 - 1/2}$ is imaginary for $x < \sqrt{2}/2$, we see the derivative of the right-hand member (with respect to y) cannot vanish at a point y_0 interior to $[0, \sqrt{2}/2)$. Thus the right-hand member is maximal at an end point $y = 0$ or $y = x$. But at $y = 0$, this member reduces to $-xg_0(0) + x$, and we are required to show that

$$xg_0(x) \geq -xg_0(0) + x \quad \text{if } 0 \leq x < \frac{\sqrt{2}}{2}.$$

Since this inequality holds trivially at $x = 0$, we can assume that $x > 0$, and this last desired inequality becomes

$$g_0(x) \geq -g_0(0) + 1 \quad \text{if } 0 \leq x < \frac{\sqrt{2}}{2}.$$

But since g_0 is decreasing and continuous on this interval, it suffices to establish the inequality at $x = \sqrt{2}/2$.

From a previous calculation, however, we have

$$g_0\left(\frac{\sqrt{2}}{2}\right) = \frac{1}{2} - \frac{1}{2} \log(\sqrt{2} + 1),$$

$$-g_0(0) + 1 = \frac{1}{2} - \frac{1}{2} \log(\sqrt{2} + 1).$$

The foregoing inequality is thus sharp at $y = 0$. Finally, at the upper end point $y = x$, the inequality

$$xg_0(x) \geq -xg_0(y) + x - y$$

reduces to the nonnegativity condition on g_0 already established. Thus, (E) is satisfied and the verification that (f_0, g_0) is in the constraint set is complete.

3. PROOF OF OPTIMALITY OF SOLUTION

The proof of optimality is not nearly so tedious as the rather messy analysis of Sec. 2, which showed that the proposed solution (f_0, g_0) is in the constraint set. The reason for this unbalance of effort is due primarily to the 2-dimensional character of the constraining inequalities and the fact (which we do not prove here) that equality of the constraints can hold only on a set of (2-dimensional) measure zero. This last assertion is an aside, however, and we shall turn our attention now to establishing the minimal property of (f_0, g_0) .

Let

$$J = \int_{\frac{\sqrt{2}}{2}}^1 \left(x - \sqrt{x^2 - \frac{1}{2}} \right) dx.$$

By elementary calculus and a numerical computation, we obtain

$$J = \frac{1}{4} \left(1 + \log(\sqrt{2} + 1) - \sqrt{2} \right) \approx .1168.$$

Moreover, we have

$$f_0(x) = xg_0(x) \quad \text{if } 0 \leq x < \frac{\sqrt{2}}{2},$$

whence

$$(3) \quad \int_0^{\frac{\sqrt{2}}{2}} f_0(x) dx = \int_0^{\frac{\sqrt{2}}{2}} xg_0(x) dx.$$

Also, we have

$$f_0(x) = -xg_0\left(\sqrt{x^2 - \frac{1}{2}}\right) + x - \sqrt{x^2 - \frac{1}{2}} \quad \text{if } \frac{\sqrt{2}}{2} \leq x \leq 1,$$

whence

$$(4) \quad \int_{\frac{\sqrt{2}}{2}}^1 f_0(x) dx = - \int_{\frac{\sqrt{2}}{2}}^1 xg_0\left(\sqrt{x^2 - \frac{1}{2}}\right) dx + J.$$

Adding equations (3) and (4) together gives

$$\int_0^1 f_0(x) dx = \int_0^{\frac{\sqrt{2}}{2}} xg_0(x) dx - \int_{\frac{\sqrt{2}}{2}}^1 xg_0\left(\sqrt{x^2 - \frac{1}{2}}\right) dx + J.$$

Upon making the change of variable

$$x' = \sqrt{x^2 - \frac{1}{2}}$$

in the second integral of the right-hand member of the above equation, we obtain

$$\int_{\frac{\sqrt{2}}{2}}^1 x g_0 \left(x^2 - \frac{1}{2} \right) dx = \int_0^{\frac{\sqrt{2}}{2}} x' g_0(x') dx' .$$

Thus, the integrals involving g_0 cancel and we are left with

$$(5) \quad \int_0^1 f_0(x) dx = J .$$

Next, let (f, g) be any pair of functions in the constraint set. Then (C) must be satisfied in particular for $y = x$ and $0 \leq x \leq \sqrt{2}/2$. Thus, we have

$$(6) \quad f(x) \geq xg(x) \quad \text{if } 0 \leq x \leq \frac{\sqrt{2}}{2} ,$$

whence we obtain

$$(7) \quad \int_0^{\frac{\sqrt{2}}{2}} f(x) dx \geq \int_0^{\frac{\sqrt{2}}{2}} xg(x) dx .$$

Moreover, if $\sqrt{2}/2 \leq x \leq 1$, we have $0 \leq \sqrt{x^2 - 1/2} \leq x$, so that (E) must be satisfied if $y = \sqrt{x^2 - 1/2}$ and $\sqrt{2}/2 \leq x \leq 1$; that is,

$$(8) \quad f(x) \geq -xg \left(\sqrt{x^2 - \frac{1}{2}} \right) + x - \sqrt{x^2 - \frac{1}{2}} \quad \text{if } \frac{\sqrt{2}}{2} \leq x \leq 1 .$$

Integrating this last inequality over the interval on which it holds yields

$$(9) \quad \int_{\frac{\sqrt{2}}{2}}^1 f(x) dx \geq - \int_{\frac{\sqrt{2}}{2}}^1 xg \left(\sqrt{x^2 - \frac{1}{2}} \right) dx + J .$$

Finally, adding (7) and (9) gives

$$\int_0^1 f(x) dx \geq \int_0^{\frac{\sqrt{2}}{2}} xg(x) dx - \int_{\frac{\sqrt{2}}{2}}^1 xg \left(\sqrt{x^2 - \frac{1}{2}} \right) dx + J .$$

But upon making the same change of variable as before, we see that the integrals involving g cancel and we are left with

$$\int_0^1 f(x)dx \geq J = \int_0^1 f_0(x)dx \quad (\text{from (5)}) ,$$

whence the minimal property of (f_0, g_0) is established.

A final closing note is in order. In Prager's original statement of the problem it was tacitly assumed that f is integrable (via the objective function). No such restriction (as we have made) was asserted about g . Nevertheless, a proof of optimality with this restriction relaxed can be given as follows:

If we replace x in (8) by $\sqrt{x^2 + 1/2}$, then the new variable ranges over the interval $[0, \sqrt{2}/2]$ and we obtain in place of (8) the equivalent inequality

$$(10) \quad f\left(\sqrt{x^2 + \frac{1}{2}}\right) \geq -\sqrt{x^2 + \frac{1}{2}} g(x) + \sqrt{x^2 + \frac{1}{2}} - x \quad \text{if } 0 \leq x \leq \frac{\sqrt{2}}{2} .$$

Multiplying this last inequality by the nonnegative number $x/\sqrt{x^2 + 1/2}$ we obtain

$$(11) \quad \frac{x}{\sqrt{x^2 + \frac{1}{2}}} f\left(\sqrt{x^2 + \frac{1}{2}}\right) \geq -xg(x) + x - \frac{x^2}{\sqrt{x^2 + \frac{1}{2}}} \quad \text{if } 0 \leq x \leq \frac{\sqrt{2}}{2} .$$

Adding the inequalities (6) and (11), which now hold over the same range, yields

$$f(x) + \frac{x}{\sqrt{x^2 + \frac{1}{2}}} f\left(\sqrt{x^2 + \frac{1}{2}}\right) \geq x - \frac{x^2}{\sqrt{x^2 + \frac{1}{2}}} \quad \text{if } 0 \leq x \leq \frac{\sqrt{2}}{2} ,$$

whence

$$\int_0^{\frac{\sqrt{2}}{2}} f(x) dx + \int_0^{\frac{\sqrt{2}}{2}} \frac{x}{\sqrt{x^2 + \frac{1}{2}}} f\left(\sqrt{x^2 + \frac{1}{2}}\right) dx \\ \geq \int_0^{\frac{\sqrt{2}}{2}} \left(x - \frac{x^2}{\sqrt{x^2 + \frac{1}{2}}}\right) dx .$$

Upon making the appropriate change of variable in the second integral in the left-hand member above, and the same change in the right-hand member, we arrive at the desired inequality,



$$\int_0^1 f(x) dx \geq J .$$

In accordance with an earlier assertion about the multiplicity of solutions to the problem, we contend that the analysis indicates that if (f^*, g^*) is any pair of functions satisfying the following conditions, then (f^*, g^*) is a solution:

$$g_0(0) \leq g^*(0) \leq 1 + g_0\left(\frac{\sqrt{2}}{2}\right), \\ g^*(x) = g_0(x) \quad \text{if } 0 \leq x \leq \frac{\sqrt{2}}{2}, \\ \max\left(0, g_0\left(\frac{\sqrt{2}}{2}\right) - x + \frac{\sqrt{2}}{2}\right) \leq g^*(x) \leq g_0\left(\frac{\sqrt{2}}{2}\right) \quad \text{if } \frac{\sqrt{2}}{2} < x \leq 1, \\ f^*(x) \geq f_0(x) \quad \text{with equality holding for almost all} \\ x \in [0, 1] .$$

(It is conjectured that these are the only solutions to the problem. It has been the writer's experience in such matters, however, that a proof would perhaps be rather long and will not, therefore, be attempted in this paper.)

Graphs of these solutions showing the allowable variation of g^* are appended. Notice that there is a discontinuity in the derivative of f^* at $x = \frac{\sqrt{2}}{2}$.



LIST OF RAND NOTES ON LINEAR PROGRAMMING
AND EXTENSIONS

- RM-1264 Part 1: The Generalized Simplex Method for Minimizing a Linear Form under Linear Inequality Restraints, by G. B. Dantzig, A. Orden, and P. Wolfe, April 5, 1954. Published in the Pacific Journal of Mathematics, Vol. 5, No. 2, June, 1955, pp. 183-195. (ASTIA No. AD 114134)
- RM-1265 Part 2: Duality Theorems, by G. B. Dantzig and A. Orden, October 30, 1953. (ASTIA No. AD 114135)
- RM-1266 Part 3: Computational Algorithm of the Simplex Method by G. B. Dantzig, October 26, 1953. (ASTIA No. AD 114136)
- RM-1267-1 Part 4: Constructive Proof of the Min-Max Theorem, by G. B. Dantzig, September 8, 1954. Published in the Pacific Journal of Mathematics, Vol. 6, No. 1, Spring, 1956, pp. 25-33. (ASTIA No. AD 114137)
- RM-1268 Part 5: Alternate Algorithm for the Revised Simplex Method Using Product Form for the Inverse, by G. B. Dantzig and W. Orchard-Hays, November 19, 1953. (ASTIA No. AD 90500)
- RM-1440 Part 6: The RAND Code for the Simplex Method (SX4) (For the IBM 701 Electronic Computer), by William Orchard-Hays, February 7, 1955. (ASTIA No. AD 86718)
- RM-1270 Part 7: The Dual Simplex Algorithm, by G. B. Dantzig, July 3, 1954. (ASTIA No. AD 114139)
- RM-1367 Parts 8, 9, and 10: Upper Bounds, Secondary Constraints, and Block Triangularity in Linear Programming, by G. B. Dantzig, October 4, 1954. Published in Econometrica, Vol. 23, No. 2, April, 1955, pp. 174-183. (ASTIA No. AD 111054)
- RM-1274 Part 11: Composite Simplex-Dual Simplex Algorithm--I, by G. B. Dantzig, April 26, 1954. (ASTIA No. AD 114140)
- RM-1275 Part 12: A Composite Simplex Algorithm--II, by William Orchard-Hays, May 7, 1954. (ASTIA No. AD 114141)
- RM-1281 Part 13: Optimal Solution of a Dynamic Leontief Model with Substitution, by G. B. Dantzig, June 15, 1954. Published in Econometrica, Vol. 23, No. 3, July, 1955, pp. 295-302. (ASTIA No. AD 90501)

- RM-1290 Part 14: A Computational Procedure for a Scheduling Problem of Edie, by G. B. Dantzig, July 1, 1954. Published in Operations Research, Vol. 2, No. 3, August, 1954, pp. 339-341. (ASTIA No. AD 109960)
- RM-1328 Part 15: Minimizing the Number of Carriers to Meet a Fixed Schedule, by G. B. Dantzig and D. R. Fulkerson, August 24, 1954. Published in Naval Research Logistics Quarterly, Vol. 1, No. 3, September, 1954, pp. 217-222. (ASTIA No. AD 109960)
- RM-1369 Part 16: The Problem of Routing Aircraft--A Mathematical Solution, by A. R. Ferguson and G. B. Dantzig, September 1, 1954. Published in Aeronautical Engineering Review, Vol. 14, No. 4, April, 1955, pp. 51-55. (ASTIA No. AD 90504)
- RM-1374 Part 17: Linear Programming under Uncertainty, by G. B. Dantzig, November 16, 1954. Published in Management Science, Vol. 1, Nos. 3-4, April-July, 1955, pp. 197-206. (ASTIA No. AD 90495)
- RM-1375 Part 18: Status of Solution of Large-scale Linear Programming Problems, by G. B. Dantzig, November 30, 1954. (ASTIA No. AD 86396)
- RM-1383 Part 19: The Fixed-Charge Problem, by W. M. Hirsch and G. B. Dantzig, December 1, 1954. (ASTIA No. AD 90494)
- RM-1400 Part 20: Maximal Flow through a Network, by L. R. Ford and D. R. Fulkerson, November 19, 1954. Published in Canadian Journal of Mathematics, Vol. 8, No. 3, 1956, pp. 399-404. (ASTIA No. AD 90541)
- RM-1418-1 Part 21: On the Min Cut Max Flow Theorem of Networks, by G. B. Dantzig and D. R. Fulkerson, April 15, 1955. Published in Linear Inequalities and Related Systems, Annals of Mathematics Study No. 38, edited by H. W. Kuhn and A. W. Tucker, Princeton University Press, 1956, pp. 215-221. (ASTIA No. AD 86705)
- RM-1475 Part 22: Recent Advances in Linear Programming, by G. B. Dantzig, April 12, 1955. Published in Management Science, Vol. 2, No. 2, January, 1956, pp. 131-144. (ASTIA No. AD 111056)
- RM-1432 Part 23: A Production Smoothing Problem, by S. M. Johnson and G. B. Dantzig, January 6, 1955. Published in Proceedings of the Second Symposium in Linear Programming (Washington, D. C., January 27-29, 1955), Vol. 1, U. S. Department of Commerce, Washington, D. C. 1956, pp. 151-176. (ASTIA No. AD 90506)

- RM-1470 Part 24: The Modification of the Right-hand side of a Linear Programming Problem, by H. M. Markowitz, April 20, 1955. (ASTIA No. AD 90543)
- RM-1452 Part 25: The Elimination Form of the Inverse and Its Application to Linear Programming, by H. M. Markowitz, April 8, 1955. (ASTIA No. AD 86956)
- RM-1489 Part 26: Computation of Maximal Flows in Networks, by D. R. Fulkerson and G. B. Dantzig, April 1, 1955. Published in Naval Research Logistics Quarterly, Vol. 2, No. 4, December, 1955, pp. 277-283. (ASTIA No. AD 90548)
- RM-1553 Part 27: Dilworth's Theorem on Partially Ordered Sets, by A. J. Hoffman and G. B. Dantzig, August 26, 1955. Published in Linear Inequalities and Related Systems, Annals of Mathematics Study No. 38, edited by H. W. Kuhn and A. W. Tucker, Princeton University Press, 1956, pp. 207-214. (ASTIA No. AD 88670)
- RM-1560 Part 28: A Simple Linear Programming Problem Explicitly Solvable in Integers, by O. A. Gross, September 30, 1955. (ASTIA No. AD 90546)
- RM-1604 Part 29: A Simple Algorithm for Finding Maximal Network Flows and an Application to the Hitchcock Problem, by L. R. Ford and D. R. Fulkerson, December 29, 1955. Published in Canadian Journal of Mathematics, Vol. 9, 1957, pp. 210-218. (ASTIA No. AD 90545)
- RM-1644 Part 30: A Class of Discrete-type Minimization Problems, by O. A. Gross, February 24, 1956. (ASTIA No. AD 90560)
- RM-1709 Part 31: A Primal-Dual Algorithm, by G. B. Dantzig, L. R. Ford, and D. R. Fulkerson, May 9, 1956. Published in Linear Inequalities and Related Systems, Annals of Mathematics Study No. 38, edited by H. W. Kuhn and A. W. Tucker, Princeton University Press, 1956, pp. 171-181. (ASTIA No. AD 111635)
- RM-1736 Part 32: Solving the Transportation Problem, by L. R. Ford and D. R. Fulkerson, June 20, 1956. Published in Management Science, Vol. 3, No. 1, October, 1956, pp. 24-32. (ASTIA No. AD 111816)
- RM-1737 Part 33: A Theorem on Flows in Networks, by David Gale, June 22, 1956. Published in Pacific Journal of Mathematics, Vol. 7, No. 2, 1957, pp. 1073-1082. (ASTIA No. AD 112371)

- RM-1798 Part 34: A Primal-Dual Algorithm for the Capacitated Hitchcock Problem, by L. R. Ford and D. R. Fulkerson, September 25, 1956. Published in Naval Research Logistics Quarterly, Vol. 4, No. 1, March, 1957, pp. 47-54. (ASTIA No. AD 112373)
- RM-1832 Part 35: Discrete-variable Extremum Problems, by G. B. Dantzig, December 6, 1956. Published in Operations Research, April, 1957. (ASTIA No. AD 112411)
- RM-1833 Part 36: The Allocation of Aircraft to Routes--An Example of Linear Programming under Uncertain Demand, by A. R. Ferguson and G. B. Dantzig, December 7, 1956. (ASTIA No. AD 112418)
- RM-1799 Part 37: Concerning Multicommodity Networks, by J. T. Robacker, September 26, 1956. (ASTIA No. AD 112392)
- RM-1864 Part 38: Note on B. Klein's "Direct Use of Extremal Principles in Solving Certain Problems Involving Inequalities," by G. B. Dantzig, January 29, 1957. Published in Operations Research, April, 1956. (ASTIA No. AD 123515)
- RM-1859 Part 39: Slightly Intertwined Linear Programming Matrices, by Richard Bellman, January 23, 1957. Published in Management Science, July, 1957. (ASTIA No. AD 123533)
- RM-1977 Part 40: Network Flows and Systems of Representatives, by L. R. Ford and D. R. Fulkerson, September 12, 1957. Published in Canadian Journal of Mathematics, Vol. 10, No. 1, 1958, pp. 78-84. (ASTIA No. AD 144263)
- RM-1981 Part 41: Constructing Maximal Dynamic Flows from Static Flows, by L. R. Ford and D. R. Fulkerson, September 17, 1957. Published in Operations Research, Vol. 6, No. 3, May-June, 1958, pp. 419-433. (ASTIA No. AD 144279)
- RM-2021 Part 42: Linear Programming and Structural Design, by W. Prager, December 3, 1957. (ASTIA No. AD 150661)
- RM-1976 Part 43: A Feasibility Algorithm for One-way Substitution in Process Analysis, by K. J. Arrow and S. M. Johnson, September 12, 1957. Published in Studies in Linear and Non-Linear Programming, by Kenneth J. Arrow, Leonid Hurwicz, and Hirofumi Uzawa, Stanford University Press, 1958, pp. 198-202. (ASTIA No. AD 144278)

- RM-2158 Part 44: Transient Flows in Networks, by D. Gale, April 11, 1958. (ASTIA No. AD 150686)
- RM-2159 Part 45: A Network-Flow Feasibility Theorem and Combinatorial Applications, by D. R. Fulkerson, April 21, 1958. Published in Canadian Journal of Mathematics, Vol. XI, No. 3, 1959. (ASTIA No. AD 156011)
- RM-2178 Part 46: Bounds on the Primal-Dual Computation for Transportation Problems, by D. R. Fulkerson, May 21, 1958. (ASTIA No. AD 156001)
- RM-2209 Part 47: Solving Linear Programs in Integers, by G. B. Dantzig, July 11, 1958. Published in Naval Research Logistics Quarterly, Vol. 6, No. 1, March, 1959. (ASTIA No. AD 156047)
- RM-2287 Part 48: Inequalities for Stochastic Linear Programming Problems, by Albert Madansky, November 13, 1958. Published in Management Science, January, 1960. (ASTIA No. AD 208311)
- RM-2321 Part 49: On a Linear Programming-Combinatorial Approach to the Traveling Salesman Problem, by G. B. Dantzig, D. R. Fulkerson, and S. M. Johnson, January 26, 1959. Published in Operations Research, Vol. 7, No. 1, January-February, 1959. (ASTIA No. AD 211642)
- RM-2338 Part 50: On Network Flow Functions, by L. S. Shapley, March 16, 1959. (ASTIA No. AD 214635)
- RM-2388 Part 51: The Simplex Method for Quadratic Programming, by Philip Wolfe, June 5, 1959. Published in Econometrica, Vol. 27, No. 3, July, 1959, pp. 382-398. (ASTIA No. AD 225224)
- RM-2425 Part 52: Computing Tetraethyl-Lead Requirements in the Linear-Programming Format, by G. B. Dantzig, T. T. Kawaratani, and R. J. Ullman, April 1, 1960. Published in Operations Research, Vol. 8, No. 1, January-February, 1960. (ASTIA No. AD 237380)
- RM-2480 Part 53: On the Equivalence of the Capacity-Constrained Transshipment Problem and the Hitchcock Problem, by D. R. Fulkerson, January 13, 1960. (ASTIA No. AD 235811)

- RM-2597 Part 54: An Algorithm for the Mixed Integer Problem, by Ralph Gomory, July 7, 1960. (ASTIA No. AD 243212)
- RM-2751 Part 55: On the Solution of Two-Stage Linear Programs under Uncertainty, by George Dantzig and Albert Madansky, July 1961. (ASTIA No. AD 263219)
- RM-2752 Part 56: Methods of Solution of Linear Programs under Uncertainty, by Albert Madansky, April 1961. (ASTIA No. 257816)
- RM-2813 Part 57: The Decomposition Algorithm for Linear Programming, by George Dantzig and Philip Wolfe, August 1961. (ASTIA No. AD 263628)
- RM-2956 Part 58: An Algorithm for Scaling Matrices, by D. R. Fulkerson and Philip Wolfe, February 1962.
- RM-2957 Part 59: Linear Programming in a Markov Chain, by George Dantzig and Philip Wolfe, March 1962.