

UNCLASSIFIED

---

---

AD 282 237

*Reproduced  
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA




---

---

UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

# DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

282 237

# 282 237

# COATING and CHEMICAL LABORATORY



ENGINEERING STUDIES ON SOLUBILIZATION ADDITIVES  
FOR WATER IN MINERAL OIL

Report No. CCL # 118  
OMS Code No. 5010.11.8020.0  
D. A. Project No 8R-593-210-61  
Author Charles B. Jordan  
Date 20 April 1962

~~This information is furnished for information purposes only with the understanding that it will not be released to any other nation without specific approval of the United States of America Department of the Army, and that it will not be used for other than military purposes. It is understood that the furnishing of the attached plans, specifications, technical data, and other information to the recipient does not in any way constitute a license to make, use, or sell the subject matter of the inventions which may be embodied or described in the information so furnished, and any manufacture, use or sale which the recipient makes of any such inventions disclosed therein is at the risk of the recipient.~~

## ABERDEEN PROVING GROUND MARYLAND

ASTIA AVAILABILITY NOTICE

Qualified requesters may obtain copies of this report from the Armed Services Technical Information Agency, Arlington Hall Station, Arlington 12, Virginia

Copies available at Office of Technical Services, \$ 0.75

THE FINDINGS IN THIS REPORT ARE NOT TO BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION.

This information is furnished for information purposes only with the understanding that it will not be released to any other nation without specific approval of the United States of America-Department of the Army, and that it will not be used for other than Military purposes. It is understood that the furnishing of the attached plans, specifications, technical data, and other information to the recipient does not in any way constitute a license to make, use, or sell the subject matter of any inventions which may be embodied or described in the information so furnished, and any manufacture, use or sale which the recipient makes of any such inventions disclosed therein is at the risk of the recipient.

TABLE OF CONTENTS

	<u>Page No.</u>
TITLE PAGE.....	ii
ABSTRACT.....	iii
INTRODUCTION.....	1
DETAILS OF TEST.....	1 - 2
DISCUSSION.....	2
ACKNOWLEDGEMENT.....	2
REFERENCES.....	2
APPENDIX.....	3
Table 1 .....	4 - 13
Table 2 .....	13 - 16
DISTRIBUTION LIST.....	17 - 18

UNCLASSIFIED

Report No. CCL # 118

Copy Number                     

ENGINEERING STUDIES ON SOLUBILIZATION ADDITIVES FOR  
WATER IN MINERAL OIL

By

Charles B. Jordan

20 April 1962

OMS Code No. 5010.11.8020.0

Dept of the Army Project No.  
BR - 593-210-61

Coating and Chemical Laboratory  
Aberdeen Proving Ground  
Maryland

UNCLASSIFIED

AUTHOR: *Charles B. Jordan*  
CHARLES B. JORDAN, Chemist  
Automotive Chemical Branch

REVIEWED BY: *Virgil O. Hatch*  
VIRGIL O. HATCH, Chief  
Automotive Chemical Branch

APPROVED BY: *C. F. Pickett*  
C. F. PICKETT, Director  
Coating & Chemical Laboratory

## ABSTRACT

The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of  $-65^{\circ}\text{F}$  to  $+400^{\circ}\text{F}$ .

Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

## I. INTRODUCTION

Aberdeen Proving Ground, Maryland, was directed by OCO, reference OMS 5010.11.8020.0, dated 21 April 1961, to conduct research on hydraulic brake fluids with emphasis on non-polar fluids.

Previous studies have shown that petroleum base fluids operate satisfactorily in brake systems and are good preservatives for metals. However, in field tests conducted in 1956 (CCL Report No. 18) it was found that water, which entered the brake system by condensation, was free to corrode metal parts due to its incompatibility with the petroleum base fluid. It is conceivable that an additive or mixture of additives could be placed in the fluid so that small amounts of water could be solubilized.

Military requirements dictate the use of brake fluids throughout all climatic extremes. In order to perform this function satisfactorily, the mineral oil/additive mixture should be capable of retaining a minimum of 1 - 2% water throughout temperature ranges of  $-65^{\circ}\text{F}$ . to  $+400^{\circ}\text{F}$ .

Since large amounts of additive would change the properties of mineral oil, only concentrations of additive less than 10% were studied.

## II. DETAILS OF TEST

### A. Mineral Oil

A refined white mineral oil of medium viscosity range was used in this test. The oil remained clear and fluid in the range of  $-65^{\circ}\text{F}$ . to  $+400^{\circ}\text{F}$ .

### B. Additives

Chemically pure additives or compounds of highest purity obtainable were used.

### C. Test Procedure

One drop of distilled water was placed in 90 drops of mineral oil. Up to 10% of the additive or mixture of additives was introduced in small portions with shaking. If solubilization occurred the minimum amount of additive required for a clear solution was determined. Clear solutions were exposed to  $-40^{\circ}\text{F}$  and  $-65^{\circ}\text{F}$  for periods up to 14 days. If solubilization did not occur at room temperature, the mixture was heated to  $180^{\circ}\text{F}$ . to effect solution if possible.

## III. RESULTS OF TESTS (See Appendix)

A sodium hydrocarbon sulfonate was the only single additive capable of giving a clear solution at room temperature. A minimum of nine percent or more of sulfonate in mineral oil was required to dissolve 1% water. Even with this quantity of additive, the water separated at  $-40^{\circ}\text{F}$ .

A mixture of sodium hydrocarbon sulfonates in mineral oil sometimes gave clear solutions at  $-40^{\circ}\text{F}$ . with 1% water. Results were inconsistent at  $-65^{\circ}\text{F}$  indicating that true solutions may not have been formed or supercooling had

occurred.

Mineral oil with mixtures of glyceryl mono oleate and some alcohols (3 - 5 carbons) gave clear solutions with 1% water at room temperature, but precipitates formed in all samples at  $-40^{\circ}\text{F}$ . The addition of deiceing agents did not improve results.

One mixture was found which would allow the mineral oil plus 1% water to remain clear at  $-40^{\circ}\text{F}$  and  $-65^{\circ}\text{F}$  for periods of time up to one week. This mixture contained 4 - 6% glyceryl mono oleate, 3% N, N-dimethyl capryl-caproamide, and percentages of isobutyl alcohol above 5%. However, small crystals were formed when the solution was cooled at  $-65^{\circ}\text{F}$  for one week, indicating water separation.

All other single additives and mixtures were unsatisfactory at room temperature.

#### IV. DISCUSSION

Hydrophilic substances which were soluble in mineral oil were first investigated. When none proved satisfactory, organic substances which were very hydrophilic at one end and very hydrophobic at the other end were tried. Later more than 300 compounds containing different functional groups were studied, in the event that some of these materials would exhibit unpredicted solubilization properties.

Mixtures of compatible hydrophilic and hydrophobic compounds were tried. N,N-dimethyl amides were given much attention because their solubility is inversely proportional to rise in temperature, and most difficulty in solubilization seems to be at lower temperatures.

The trend of solvency in this study followed normal lines in that increasing carbon content of homologous compounds reduced their affinity for water.

This study did not reveal any success by this approach. A new approach should be taken in future studies. The approach would consist of selecting an additive which will chemically react with water to give an oil soluble compound.

#### V. ACKNOWLEDGEMENT

Experimental data was collected and observations were made by Sp. 4 Jimmie G. Tolar.

#### VI. REFERENCES

- A. Authority: OMS 5010.11.8020.0, dated 21 April 1961.
- B. CCL Report No. 18, Field Test of Nitrile Type Synthetic Rubber Brake Cups in Trucks at Yuma, Arizona, dated 27 February 1957.

APPENDIX

Tables

TABLE I  
SINGLE COMPOUNDS

1 - 10% Added to 1% Water In Mineral Oil

(No single compound was capable of solubilizing 1% water in mineral oil over the range of -65°F to 400°F.)

---

- I. Alcohols
  - A. Mono hydroxy
  - B. Poly hydroxy
  - C. Mono hydroxy ethers
  - D. Aromatic
- II. Phenols
  - A. Mono hydric
  - B. Poly hydric
- III. Amines
  - A. Mono amines
  - B. Poly amines
- IV. Esters
- V. Ketones
- VI. Acids and Acid Anhydrides
- VII. Hydrocarbons
- VIII. Halogenated Hydrocarbons
- IX. Ethers
- X. Salts of Organic Acids
- XI. Sulfonates
- XII. Inorganic Esters
- XIII. Amine Salts
- XIV. Metal Organics
- XV. Benzothiazyls
- XVI. Nitriles
- XVII. Hydrazines
- XVIII. Acid Halides
- XIX. Oximes

TABLE I CONTINUED

XX. Isocyanates

XXI. Sulfonates

XXII. Peroxides

XXIII. Amides

## I. Alcohols

## A. Mono-hydroxy compounds

<u>No. Carbons</u>	<u>Name</u>
1	Methanol
2	Ethanol
2	2-Chloroethanol
3	Propanol-1
3	Propanol-2
3	Propyne-1-ol-3
4	Butanol-1
4	Butanol-2
4	2-Methyl Propanol-1
4	2-Methyl Propanol-2
5	Pentanol-1
5	2-Methyl-Butanol-1
5	2-Methyl-Butanol-2
5	2-Methyl-Butanol-3
5	Cyclopentanol
6	Hexanol-1
6	2-Methyl Pentanol-1
6	4-Methyl Pentanol-2
6	2-Ethyl Butanol-1
6	Cyclohexanol
6	Diacetone Alcohol
7	Heptanol-1
7	Heptanol-3
7	Heptanol-4
7	2,4-Dimethyl Pentanol-3
8	Octanol-1
8	Octanol-2
8	2-Ethyl Hexanol-1
9	N-Nonyl Alcohol
10	N-Decyl Alcohol
11	Undecyl Alcohol
11	Undecylenyl Alcohol
12	Dodecyl Alcohol

TABLE I CONTINUED

## B. Polyhydroxy compounds

<u>No. Carbons</u>	<u>Name</u>
2	Ethylene Glycol
3	1,2-Propylene Glycol
3	Glycerol
3	1,3-Propylene Glycol
4	Diethylene Glycol
4	2-Butyne-1,4-Diol
4	1,3-Butanediol
4	2,3-Butanediol
5	1,5-Pentanediol
6	2-Methyl-2,4-Pentanediol
6	Hexylene Glycol
6	Triethylene Glycol
6	Mannitol
8	Octylene Glycol
8	2,5-Dimethyl-3-Hexyne-2,5-Diol
18	Tetra Hydroxyl Castor Oil
18	Ricinoleyl Alcohol
18	12-Hydroxy Stearyl Alcohol
Polymer	Polymerized Trimethyl Dihydroxy Quinoline

## C. Mono Hydroxy ethers

<u>No. Carbons</u>	<u>Name</u>
3	Ethylene Glycol Mono-Methyl Ether
4	Ethylene Glycol Mono Ethyl Ether
4	Propylene Glycol Mono Methyl Ether
4	2,2'-Dimethoxy Ethanol
4	Methoxy Propanol
5	4-Methoxy Butanol-1
5	Propylene Glycol Mono Ethyl Ether
5	Diethylene Glycol Mono Methyl Ether
5	3-Methoxy Butanol-1
6	Ethylene Glycol Mono-n-Butyl Ether
6	Diethylene Glycol Mono-Ethyl Ether
7	2-Methoxy-2-Methyl-Pentanol-4
8	Ethylene Glycol Mono-n-Hexyl Ether
8	Diethylene Glycol Mono-n-Butyl Ether
16	Diethylene Glycol Monolaurate

## D. Aromatic Alcohols

Benzyl Alcohol  
 N-Phenoxy Ethanol  
 P-Tert-Amyl-Phenoxy Ethanol  
 O-Sec-Amyl Phenoxy Ethanol

## II. Phenols

## A. Mono-Hydric Phenols

TABLE I CONTINUED

## A. Mono-Hydric Phenols

<u>No. Carbons</u>	<u>Name</u>
6	Phenol
6	O-Chlorophenol
6	P-Chlorophenol
6	P-Nitro Phenol
6	Picric Acid
7	O-Cresol
7	P-Cresol
7	M-Cresol
7	Hydroquinone Mono Methyl Ether
9	8-Hydroxy Quinoline
9	Alkyl Phenol
10	Carvacrol
12	P-Hydroxy Diphenyl
12	O-Hydroxy Diphenyl
14	Alizarin
15	2,6-Di-t-butyl-p-cresol
15	Octyl Cresol
15	Bisphenol A
19	Phenol Red

## B. Polyhydric Phenols

<u>No. Carbons</u>	<u>Name</u>
6	Hydroquinone
6	Pyrogallol
6	Resorcinol
7	Toluhydroquinone
10	P-tert-butyl Catechol
10	1,5-dihydroxy Naphthalene
10	1,6-dihydroxy Naphthalene
12	Phenyl Hydroquinone
12	4,4-dihydroxy Diphenyl
14	Octyl Resorcinol
14	2,5-di-t-butyl Hydroquinone
20	Phenolphthalein
23	Cresolphthalein

## III. Amines

## A. Mono-amines

	<u>No. Carbons</u>	<u>Name</u>
1. Primary	4	Tris (hydroxyl methyl) Amino Methane
	7	Amino Benzoic Acid
2. Secondary	4	Diethyl Amine
	4	2-Pyrrolidone
	5	Pyridine
	6	Cyclohexyl Amine
	8	Di-n-butyl Amine

TABLE I CONTINUED

	<u>No. Carbons</u>	<u>Name</u>
	10	N-butyl-p-amino-phenol
	12	Dicyclohexyl Amine
	16	Di-2-ethyl Hexyl Amine
3. Tertiary	5	N-methyl Morpholine
	6	Triethyl Amine
	9	N-vinyl-2-pyrrolidone
	10	N-phenyl Morpholine
	10	Potassium Ethylene Diamine Tetracetate
	10	Sodium Ferric Ethylene Diamine Tetracetate
	12	N,N-di-n-propyl Aniline
	15	Tri-N-amyI Amine

## B. Poly Amines

	<u>No. Carbons</u>	<u>Name</u>
1. Primary - Secondary	4	Aminoethyl Ethanol Amine
2. Secondary	14	Disalicylal Propylene Diamine
	14	Disalicylidene Propylene Diamine
3. Tertiary	7	3-N,N-(dimethylamine) Propyne-1
	8	1,4-Bis(dimethylamine) Butyne-2
	14	N,N-di-sec-butyl-p-phenylene Diamine

## IV. Esters

	<u>No. Carbons</u>	<u>Name</u>
	4	2-Chloroethyl Acetate
	4	Butyrolactone
	5	Isopropyl Acetate
	6	Butyl Acetate
	6	Ethylene Glycol Monoethyl Ether Acetate
	7	N-amyI Acetate
	7	Propylene Carbonate
	8	Methyl Salicylate
	8	Diethylene Glycol Diacetate
	10	Propyl Gallate
	10	Isopropyl Benzoate
	12	N-Butyl Phthalate
	13	Phenyl Salicylate
	15	Tri-n-butyriI
	20	Diphenyl Phthalate
	21	Propylene Ricinoleate
	21	Glyceryl Mono Oleate
	31 - 42	Lanolin
	57	Glyceryl Trioleate
	Poly	Polyethylene Glycol 200-mono Ricinoleate

TABLE I CONTINUED

<u>No. Carbons</u>	<u>Name</u>
Poly	Sorbitan Fatty Acid Ester
Poly	Polyoxyethylene Ester of Fatty and Resin Acids
Poly	Polyethylene Glycol 400-di-tri-Ricinoleate

## V. Ketones

<u>No. Carbons</u>	<u>Name</u>
3	Acetone
4	Methyl Ethyl Ketone
4	2-pyrrolidone
6	Methyl Isobutyl Ketone
6	Cyclohexanone
6	Diacetone Alcohol
6	P-quinone
7	Methyl N-amyl Ketone
8	Methyl N-hexyl Ketone
9	Diisobutyl Ketone
9	N-vinyl-2-pyrrolidone
14	Alizarin
15	Alizarin Cyanine Green

## VI. Acids &amp; Acid Anhydrides

<u>No. Carbons</u>	<u>Name</u>
1	Formic
2	Acetic, Glacial
2	Oxalic
3	Propionic
3	Propiolic
3	Lactic
4	N-butyric
4	Iso-butyric
4	Acetic Anhydride
6	N-caproic
6	Iso-caproic
7	Heptanoic
7	P-toluene Sulfonic
7	Benzoic
7	Aminobenzoic
7	P-hydroxy Benzoic
7	Salicylic
7	Pyrogallic
8	N-caprylic
8	Octanoic
9	Trimethyl-n-caproic
9	Cinnamic
11	Undecanoic
12	Lauric
14	Tannic
15	Methyl Red

TABLE I CONTINUED

<u>No. Carbons</u>	<u>Name</u>
18	Stearic
18	Ricinoleic
18	Oleic
Unknown	Sulfonated Castor Oil
Unknown	Sulfonic Acid of Mineral Oil

## VII. Hydrocarbons

## A. Aliphatic

<u>No. Carbons</u>	<u>Name</u>
5-6	Petroleum Ether
5-8	Mineral Spirits
6	N-hexane
6	Cyclohexane
6	Cyclohexene
6	2-methyl Pentane
6-7	Naphtha
7	N-heptane
8	N-octane
8	2,2,4-trimethyl Pentane
8	Octene ( 1 & 2 )

## B. Aromatic

<u>No. Carbons</u>	<u>Name</u>
6	Benzene
7	Toluene
8	Xylene

## VIII. Halogenated Hydrocarbons

<u>No. Carbons</u>	<u>Name</u>
1	Chloroform
1	Iodoform
1	Carbon Tetrachloride
2	2-Chloroethanol
2	Dibromoethane
2	Tetrachloroethylene
6	Dichlorobenzene
6	Chlorobenzene
6	Bromobenzene
19	Triphenyl Chloromethane

## IX. Ethers

<u>No. Carbons</u>	<u>Name</u>
3	Trioxymethylene
4	Diethyl Ether
4	P-dioxane
4	Propylene Glycol Formal
6	Diisopropyl Ether

TABLE I CONTINUED

<u>No. Carbons</u>	<u>Name</u>
8	Triethylene Glycol Dimethyl Ether
10	Tetraethylene Glycol Dimethyl Ether
18	Epoxidized Castor Oil
20	Epoxidized Acetylated Castor Oil
20	Ethoxylated Castor Oil

## X. Salts of Organic Acids

<u>No. Carbons</u>	<u>Name</u>
1	Sodium Carbonate
2	Lead Acetate
2	Guanidine Carbonate
2	Sodium Methyl Carbonate
6	Sodium Citrate
7	Sodium Benzoate
7	Sodium Mercaptobenzothiozole
8	Potassium Acid Phthalate
9	Cinnamic Acid Sodium Salt
10	Copper-8-quinolinolate
10	Sodium Ferric Ethylene Diamine Tetra Acetate
10	Potassium Ethylene Diamine Tetra Acetate
11	Calcium Undecanoate
18	Copper Stearate
18	Zinc Dibutyl Dithiocarbamate
22	Zinc Diamyl Dithio Carbamate

## XI. Sulfonate

<u>No. Carbons</u>	<u>Name</u>
8	Sodium Xylene Sulfonate
10	Napthalene Sulfonic Acid Sodium Salt
26	* Sodium Hydrocarbon Sulfonate, Mol. Wt 430
28	Barium Dinonyl Naphthalene Sulfonate, Mol Wt. 1055
• 30+	Sodium Hydrocarbon Sulfonate, Mol. Wt. 500+
Unknown	Calcium Sulfonate

\* Capable of clear soln at room temperature if 9% or more sulfonate is added to 1% water in mineral oil. At -40°F solutions are cloudy with crystalline separation.

## XII Inorganic Esters

<u>No. Carbons</u>	<u>Name</u>
17	2,6-di-t-butyl-4-methyl-di-n- butyl Borate
18	Trihexylene Glycol Borate

TABLE I CONTINUED

	<u>No. Carbons</u>	<u>Name</u>
	18	Trihexylene Glycol Biborate
	22	2,6-di-t-butyl Phenyl-di-n-butyl Borate
	24	Tri-2-ethyl Hexyl Phosphate
	Unknown	Alkyl Phosphate 50% in Ethylene Glycol
XIII. Amine Salts		
	<u>No. Carbons</u>	<u>Name</u>
	6	Di-isopropanol Amine Phosphate
	6	Triethanol Amine Phosphate
	9	Tri-isopropanol Amine Borate
XIV. Metal Organics		
	<u>No. Carbons</u>	<u>Name</u>
	8	Ethyl Tri-ethoxy Silane
	12	Di-butyl Tin Maleate
	32	Dibutyl Tin Dilaurate
XV. Benzothiazyls		
		Benzothiazyl Disulfide
		Mercaptobenzothiazole
		Phenyl Biguanide Mercaptobenzothiazole Salt
XVI. Nitriles		
	<u>No. Carbons</u>	<u>Name</u>
	3	B-hydroxy Propionitrile
	7	Benzonitrile
XVII. Hydrazines		
		2,4-dinitrophenyl Hydrazine
XVIII. Acid Halides		
		3,5-dinitro benzoyl Chloride
XIX. Oximes		
		Salicylaldoxime
XX. Isocyanates		
		P-nitro Phenyl Isocyanate
XXI Aromatic Sulfonamides		
		P-toluene Sulfonamide

TABLE I CONTINUED

## XXII. Peroxides

## Benzoyl Peroxide

## XXIII. Amides

<u>No. Carbons</u>	<u>Name</u>
1	Formamide
4	N,N-dimethyl Acetamide
5	N,N-dimethyl Propionamide
6	N,N-dimethyl Butyramide
16	N,N-dimethyl Capryl-caproamide
14	N,N-dimethyl Lauramide
20	N,N-dimethyl Oleamide

TABLE II

## MIXTURES

Added to 1% water in mineral oil.

- I. Sulfonates
- II. Glyceryl Mono Oleate and Mono Hydroxy Compounds
- III. Glyceryl Mono Oleate and Poly Hydroxy Compounds
- IV. Glyceryl Mono Oleate and Glycol Ether
- V. Glyceryl Mono Oleate, Isobutyl Alcohol and N,N-dimethyl Capryl-caproamide

- I. Sulfonates (Mixture of two sodium hydrocarbon sulfonates), added to 1% water in mineral oil.

#1 Mol. Wt. Approx. 500, #2 Mol. Wt. Approx. 430.

Percent Added		<u>Appearance of Mixture</u>
<u>#1</u>	<u>#2</u>	
1 - 10%	0%	Cloudy at room temperature.
8 - 9%	1%	Clear at room temperature, frozen at -40°F.
8%	2%	Clear at room temperature, varying from clear to slightly cloudy at -40°F. Crystalline separation at -65°F.
1 - 7%	3%	Clear at -40°F. except 7% (inconsistent at -65°)
1 - 6%	4%	Clear at room temperature, clear at -40°F but extremely viscous.

TABLE II CONTINUED

#1	#2	Appearance of Mixture
0%	9 & 10%	Clear at room temperature, crystalline separation at -40°F.
1%	5% and above	Clear at room temperature, crystalline separation at -40°F.

II. A. 1 - 10% Glyceryl Mono Oleate, 1 - 10% Mono Hydroxy Compounds, Added to 1% Water in Mineral Oil.

## Alcohols Added

No. Carbons	Name	Appearance of Mixture
1	Methyl	Cloudy in all proportions at room temp.
2	Ethyl	Cloudy in all proportions at room temp.
3	N-propyl	Cloudy with less than 6% ester at room temp.
3	Iso-propyl	Cloudy with less than 6% ester at room temp.
4	N-butyl	Clear with 4% ester & 4% alc at room temp., crystalline separation at -40°F.
4	Isobutyl	Clear with 4% ester & 4% alc at room temp., crystalline separation at -40°F.
4	Sec Butyl	Clear with 5% ester & 3% alc at room temp., crystalline separation at -40°F.
4	T-butyl	Clear with 6% ester & 3% alc., crystalline separation at -40°F.
5	N-amyl	Cloudy in all proportions.
5	3-methyl butyl	Clear with 4% ester & 6% alc. at room temperature, crystalline separation at -40°F.
6	N-hexyl	Cloudy in all proportions at room temp.
6	4-methyl Pentanol-2	Cloudy in all proportions at room temp.
8	2-ethyl Hexyl	Cloudy in all proportions at room temp.

B. 4 - 6% Glyceryl Mono Oleate, 2 - 4% Isobutyl Alcohol, and Deicing Agents, Added, as Indicated, to 1% Water in Mineral Oil.

Deicing Agent - Percent Added	Appearance of Mixture
$\frac{1}{2}$ - 4% Pyrrolidone	Cloudy at room temperature.
$\frac{1}{2}$ - 4% N-methyl Pyrrolidone	Cloudy at room temperature.
$\frac{1}{2}$ - 4% N-vinyl-2-pyrrolidone	Cloudy at room temperature.
$\frac{1}{2}$ - 4% 1,2,6-hexanetriol	Clear at room temperature, cloudily at -40°F.

III. Glyceryl Mono-Oleate and Poly Hydroxy Compounds

A. 4% Glyceryl Mono Oleate, 1 - 10% Polyol, Added to 1% Water in Mineral Oil.

TABLE II CONTINUED

## Polyhydroxy Compound Added

<u>No. Carbons</u>	<u>Name</u>	<u>Appearance of Mixture</u>
2	Ethylene Glycol	Cloudy in all proportions at room temp.
3	Propylene Glycol	Cloudy in all proportions at room temp.
3	Glycerol	Cloudy in all proportions at room temp.
4	2,3-butane Diol	Cloudy in all proportions at room temp.
4	1,3-butane Diol	Cloudy in all proportions at room temp.
4	Diethylene Glycol	Cloudy in all proportions at room temp.
4	2-butyne-1,4-diol	Cloudy in all proportions at room temp.
6	2,4-dihydroxy-2-methyl Pentane	Cloudy in all proportions at room temp.
6	1,2,6-hexane-triol	Cloudy in all proportions at room temp.
6	Dipropylene Glycol	Cloudy in all proportions at room temp.
8	2,5 Dimethyl-3- hexyne-2,5-diol	Cloudy in all proportions at room temp.
8	Octylene Glycol	Cloudy in all proportions at room temp.

B. 8% Glyceryl Mono Oleate, 1 - 2% Same Polyhydroxy Compounds, Added to 1% Water in Mineral Oil.

Results: All mixtures were cloudy in all proportions at room temperature.

## IV. 4% Glyceryl Mono Oleate and 1 - 6% Glycol Ethers, Added to 1% Water in Mineral Oil

## Glycol Ethers Added

<u>No. Carbons</u>	<u>Name</u>	<u>Appearance of Mixture</u>
3	Ethylene Glycol Mono Methyl Ether	Cloudy in all proportions at room temp.
4	Ethylene Glycol Mono Ethyl Ether	Cloudy in all proportions at room temp.
4	Propylene Glycol Mono Methyl Ether	Cloudy in all proportions at room temp.
4	2,2'-dimethoxy Ethanol	Cloudy in all proportions at room temp.
5	Propylene Glycol Mono Ethyl Ether	Cloudy in all proportions at room temp.
5	4-Methoxy Butanol-1	Cloudy in all proportions at room temp.
6	Ethylene Glycol Mono- n-butyl Ether	Cloudy in all proportions at room temp.
6	Diethylene Glycol Mono- n-ethyl Ether	Cloudy in all proportions at room temp.
8	Ethylene Glycol Mono- n-hexyl Ether	Cloudy in all proportions at room temp.
8	Diethylene Glycol Mono- n-butyl Ether	Cloudy in all proportions at room temp.
16	Diethylene Glycol Mono Laurate	Cloudy in all proportions at room temp.

TABLE II CONTINUED

## V. Glyceryl Mono Oleate, Isobutyl Alcohol and N,N-dimethyl Capryl-caproamide.

## A. Percent-Added to 1% Water in Mineral Oil.\*

<u>Ester</u>	<u>Amide</u>	<u>Alcohol</u>	<u>Appearance of Mixture</u>
4%	3%	1 - 2%	Cloudy at room temperature.
		3 - 5%	Clear at room temperature.
5%	3%	1 - 2%	Cloudy at room temperature
6%	3%	1%	Cloudy at room temperature

\*Mixtures containing less than 4% ester were always cloudy regardless of amount of alcohol added. Alcohol above 5% gave clear solutions at room temperature and  $-40^{\circ}\text{F}$ . All solutions had crystalline separation at  $-65^{\circ}\text{F}$ .

## B. 4% Ester, 5% Alcohol, Above 2% Amide, Added to 1% Water in Mineral Oil.

Results: 2 - 12% amide, mixtures cloudy at  $-40^{\circ}\text{F}$ . 14% amide and above - clear and fluid at  $-40^{\circ}\text{F}$ .

## C. 4% Ester, 6% Alcohol, 1 - 24% Amide, Added to 1% Water in Mineral Oil

Results: Room temperature all mixtures clear.  
 $-40^{\circ}\text{F}$ : 1 - 18% amide crystalline separation, 19% and above slightly cloudy.  
 $-65^{\circ}\text{F}$ : All mixtures, crystalline separation.

## D. 4.3% Ester, 4.3% Alcohol, 1 - 20% Amide, Added to 1% Water in Mineral Oil.

Results: Room temperature: All mixtures clear.  
 $-40^{\circ}\text{F}$ : All mixtures cloudy

DISTRIBUTION LIST FOR OMS CODE NO. 5010.11.8020.0

<u>Department of Defense</u>	<u>No. of Copies</u>
Armed Services Technical Information Agency Arlington Hall Station ATTN: TIPDR Arlington 12, Virginia	10
 <u>Department of the Army - Technical Service</u>	
Chief of Ordnance Dept. of the Army ATTN: ORDTB - Fuels & Lubes ORDTB - Materials Washington 25, D. C.	1 1
Commanding General Aberdeen Proving Ground, Maryland ATTN: Technical Library	2
Commanding Officer Anniston Ord Depot ATTN: ORDKB-QC Laboratory Anniston, Alabama	1
Commanding General Frankford Arsenal ATTN: Pitman-Dunn Laboratory Group Philadelphia, Pa.	1
Commander Detroit Arsenal ATTN: ORDMX-B Center Line, Michigan	1
Commanding General Rock Island Arsenal ATTN: ORDBC-9320 Rock Island, Ill.	1
Commanding General Ordnance Weapons Command ATTN: ORDOW-IX Rock Island, Ill.	1
Commander Watertown Arsenal ATTN: Tech. Group Watertown 72, Mass.	1
Commanding Officer Diamond Ordnance Fuze Laboratories ATTN: Technical Reference Section Washington 25, D. C.	1

DISTRIBUTION LIST CONTINUED

	<u>No. of Copies</u>
Commander Ordnance Materials Research Office Watertown Arsenal ATTN: PS&C Division Watertown 72, Mass.	1
Commanding Officer U.S. Army Ord Test Activity Yuma, Arizona	1
Commanding Officer U. S Army Ord. Test Activity ATTN: Automotive Test Division (ORDBG-DP-FY-AU) Yuma, Arizona	1
Commander Ordnance Tank Automotive Command ATTN: ORDMC-REM.3 ORDMC-RRS.3 Detroit Arsenal Center Line, Michigan	2 1
Commander Officer Raritan Arsenal ATTN: NICP(46) Metuchen, New Jersey	1
<u>Department of the Army - Other Army Agencies</u>	
Ordnance Office, Headquarters U.S. Army Caribbean, Corozal Fort Clayton, Canal Zone	1
<u>Other Government Agencies</u>	
Office of Technical Services Acquisitions Section Department of Commerce Washington 25, D. C.	200
<u>Foreign Address</u>	
Ministry of Supply Staff British Joint Services Mission 1800 K Street, N. W. Washington 6, D. C.	2
Canadian Army Staff ATTN: GS0-1 A&R Section 2450 Massachusetts Avenue, N. W. Washington 8, D. C.	2

<p>AD No _____</p> <p>Coating &amp; Chemical Laboratory, Aberdeen Proving Ground, Md., CCL # 118 - ENGINEERING STUDIES ON SOLUBILIZATION ADDITIVES FOR WATER IN MINERAL OIL - Charles B. Jordan, Rpt No. 118, 20 April 1962, pgs, OMS 5010.11.8020.0 DA Proj No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>Unclassified</p>	<p>AD No _____</p> <p>Coating &amp; Chemical Laboratory, Aberdeen Proving Ground, Md., CCL # 118 - ENGINEERING STUDIES ON SOLUBILIZATION ADDITIVES FOR WATER IN MINERAL OIL - Charles B. Jordan, Rpt No. 118, 20 April 1962, pgs, OMS 5010.11.8020.0 DA Proj No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>Unclassified</p>
<p>AD No _____</p> <p>Coating &amp; Chemical Laboratory, Aberdeen Proving Ground, Md., CCL # 118 - ENGINEERING STUDIES ON SOLUBILIZATION ADDITIVES FOR WATER IN MINERAL OIL - Charles B. Jordan, Rpt No. 118, 20 April 1962, pgs, OMS 5010.11.8020.0 DA Proj No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>Unclassified</p>	<p>AD No _____</p> <p>Coating &amp; Chemical Laboratory, Aberdeen Proving Ground, Md., CCL # 118 - ENGINEERING STUDIES ON SOLUBILIZATION ADDITIVES FOR WATER IN MINERAL OIL - Charles B. Jordan, Rpt No. 118, 20 April 1962, pgs, OMS 5010.11.8020.0 DA Proj No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>Unclassified</p>

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

<p>AD No. <u>                    </u>  Coating &amp; Chemical Laboratory, Aberdeen  Proving Ground, Md., CCL # 118 -  ENGINEERING STUDIES ON SOLUBILIZATION  ADDITIVES FOR WATER IN MINERAL OIL -  Charles B. Jordan, Rpt No. 118, 20 April  1962, pgs, OMS 5010.11.8020.0 DA Proj  No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>Accession No. <u>                    </u>  Coating &amp; Chemical Laboratory, Aberdeen  Proving Ground, Md., CCL # 118 -  ENGINEERING STUDIES ON SOLUBILIZATION  ADDITIVES FOR WATER IN MINERAL OIL -  Charles B. Jordan, Rpt No. 118, 20 April  1962, pgs, OMS 5010.11.8020.0 DA Proj  No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>AD No. <u>                    </u>  Coating &amp; Chemical Laboratory, Aberdeen  Proving Ground, Md., CCL # 118 -  ENGINEERING STUDIES ON SOLUBILIZATION  ADDITIVES FOR WATER IN MINERAL OIL -  Charles B. Jordan, Rpt No. 118, 20 April  1962, pgs, OMS 5010.11.8020.0 DA Proj  No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>Accession No. <u>                    </u>  Coating &amp; Chemical Laboratory, Aberdeen  Proving Ground, Md., CCL # 118 -  ENGINEERING STUDIES ON SOLUBILIZATION  ADDITIVES FOR WATER IN MINERAL OIL -  Charles B. Jordan, Rpt No. 118, 20 April  1962, pgs, OMS 5010.11.8020.0 DA Proj  No. BR 593-210-61</p> <p>The object of this study was to find an additive capable of solubilizing small amounts of water in mineral oil throughout the temperature range of -65°F to + 400°F.</p> <p>Hydrophilic substances which were soluble in mineral oil were tried. When none proved satisfactory, hundreds of</p>	<p>Unclassified</p>	<p>Unclassified</p>
--	---	--	---	---------------------	---------------------

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.

Unclassified

available organic chemicals containing different functional groups were tried, in the event that some of these materials would exhibit unpredicted favorable solubilization properties. The trend of solvency in relationship to carbon content within each individual homologous series was studied.

Although there were instances where solubilization occurred, no single additive or mixture was found which would give the desired results over the specified temperature range.