

UNCLASSIFIED

AD NUMBER
AD291734
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; 13 Sep 1962. Other requests shall be referred to Army Medical Research and Development Command, Fort Detrick, MD.
AUTHORITY
USAMRL ltr, 26 Feb 1970

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD 291 734


*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.

291734

US ARMY MEDICAL RESEARCH LABORATORY

FORT KNOX, KENTUCKY

REPORT NO. 551

EFFECT OF SALMONELLA TYPHOSA ENDOTOXIN ON
RENAL FUNCTION, AND THE RELATIONSHIP OF
FUNCTION TO RENAL HEMODYNAMICS

CATALOGED BY ASTIA
AS AD NO.

291734

Capt J. Y. Gillenwater, MC
E. S. Dooley, Ph. D.
Capt E. D. Frohlich, MC

ASTIA
RECEIVED
DEC 27 1962
ASTIA



UNITED STATES ARMY
MEDICAL RESEARCH AND DEVELOPMENT COMMAND

13 September 1962

NO OTS

Report Submitted 9 August 1962

Authors

Capt Jay Y. Gillenwater, MC (M. D.)	Biophysics Division
Elmo S. Dooley, Ph. D.	Biophysics Division
Capt Edward D. Frohlich, MC (M. D.)	Biophysics Division

Technical Assistants

Sp 6 Johnny D. Wills	Biophysics Division
Pfc Robert Haber	Biophysics Division
Pfc George Angeloff	Biophysics Division
Pfc Frank Rozecki	Biophysics Division

Qualified requestors may obtain copies of this report from ASTIA.

Foreign announcement and dissemination of this report by ASTIA is limited.

AD	Accession No.	UNCLASSIFIED	AD	Accession No.	UNCLASSIFIED
US Army Medical Research Lab, Ft. Knox, Ky. EFFECT OF SALMONELLA TYPHOSA ENDOTOXIN ON RENAL FUNCTION, AND THE RELATIONSHIP OF FUNCTION TO HEMODYNAMICS - J. Y. Gillemwater, E. S. Dooley, and E. D. Frohlich w/tech asst J. Wills, R. Haber, G. Angeloff, and F. Rozecki Report No. 551, 13 Sep 62, 20 pp & ii - 7 illus - 2 tables - Project No. 6X59-01-001, Unclassified Report Injection of endotoxin into the perfused left renal artery produced an immediate (18 sec) transient vasoconstriction followed by a severe renal vasoconstriction (5 to 10 min) and a decrease in all renal function. The delayed response was blocked by locally infused phenolamine. It is concluded that the delayed vasoconstriction is induced by a systemic release of catecholamine. In a group of animals the hemodynamic effect was eliminated in one kidney by infusing phenolamine locally and holding renal blood flow constant, enabling the comparison of this kidney with the contralateral undisturbed kidney. In the kidney in which vascular effects were eliminated there was no alteration in renal functions after systemic or local injection of endotoxin. These data show that the primary effect of endotoxin on the dog kidney is hemodynamic and not nephrotoxic.		UNCLASSIFIED 1. Renal Function 2. Renal Hemodynamics 3. Endotoxin 4. Salmonella typhosa	US Army Medical Research Lab, Ft. Knox, Ky. EFFECT OF SALMONELLA TYPHOSA ENDOTOXIN ON RENAL FUNCTION, AND THE RELATIONSHIP OF FUNCTION TO HEMODYNAMICS - J. Y. Gillemwater, E. S. Dooley, and E. D. Frohlich w/tech asst J. Wills, R. Haber, G. Angeloff, and F. Rozecki Report No. 551, 13 Sep 62, 20 pp & ii - 7 illus - 2 tables - Project No. 6X59-01-001, Unclassified Report Injection of endotoxin into the perfused left renal artery produced an immediate (18 sec) transient vasoconstriction followed by a severe renal vasoconstriction (5 to 10 min) and a decrease in all renal function. The delayed response was blocked by locally infused phenolamine. It is concluded that the delayed vasoconstriction is induced by a systemic release of catecholamine. In a group of animals the hemodynamic effect was eliminated in one kidney by infusing phenolamine locally and holding renal blood flow constant, enabling the comparison of this kidney with the contralateral undisturbed kidney. In the kidney in which vascular effects were eliminated there was no alteration in renal functions after systemic or local injection of endotoxin. These data show that the primary effect of endotoxin on the dog kidney is hemodynamic and not nephrotoxic.		UNCLASSIFIED 1. Renal Function 2. Renal Hemodynamics 3. Endotoxin 4. Salmonella typhosa
US Army Medical Research Lab, Ft. Knox, Ky. EFFECT OF SALMONELLA TYPHOSA ENDOTOXIN ON RENAL FUNCTION, AND THE RELATIONSHIP OF FUNCTION TO HEMODYNAMICS - J. Y. Gillemwater, E. S. Dooley, and E. D. Frohlich w/tech asst J. Wills, R. Haber, G. Angeloff, and F. Rozecki Report No. 551, 13 Sep 62, 20 pp & ii - 7 illus - 2 tables - Project No. 6X59-01-001, Unclassified Report Injection of endotoxin into the perfused left renal artery produced an immediate (18 sec) transient vasoconstriction followed by a severe renal vasoconstriction (5 to 10 min) and a decrease in all renal function. The delayed response was blocked by locally infused phenolamine. It is concluded that the delayed vasoconstriction is induced by a systemic release of catecholamine. In a group of animals the hemodynamic effect was eliminated in one kidney by infusing phenolamine locally and holding renal blood flow constant, enabling the comparison of this kidney with the contralateral undisturbed kidney. In the kidney in which vascular effects were eliminated there was no alteration in renal functions after systemic or local injection of endotoxin. These data show that the primary effect of endotoxin on the dog kidney is hemodynamic and not nephrotoxic.		UNCLASSIFIED 1. Renal Function 2. Renal Hemodynamics 3. Endotoxin 4. Salmonella typhosa	US Army Medical Research Lab, Ft. Knox, Ky. EFFECT OF SALMONELLA TYPHOSA ENDOTOXIN ON RENAL FUNCTION, AND THE RELATIONSHIP OF FUNCTION TO HEMODYNAMICS - J. Y. Gillemwater, E. S. Dooley, and E. D. Frohlich w/tech asst J. Wills, R. Haber, G. Angeloff, and F. Rozecki Report No. 551, 13 Sep 62, 20 pp & ii - 7 illus - 2 tables - Project No. 6X59-01-001, Unclassified Report Injection of endotoxin into the perfused left renal artery produced an immediate (18 sec) transient vasoconstriction followed by a severe renal vasoconstriction (5 to 10 min) and a decrease in all renal function. The delayed response was blocked by locally infused phenolamine. It is concluded that the delayed vasoconstriction is induced by a systemic release of catecholamine. In a group of animals the hemodynamic effect was eliminated in one kidney by infusing phenolamine locally and holding renal blood flow constant, enabling the comparison of this kidney with the contralateral undisturbed kidney. In the kidney in which vascular effects were eliminated there was no alteration in renal functions after systemic or local injection of endotoxin. These data show that the primary effect of endotoxin on the dog kidney is hemodynamic and not nephrotoxic.		UNCLASSIFIED 1. Renal Function 2. Renal Hemodynamics 3. Endotoxin 4. Salmonella typhosa

REPORT NO. 551

**EFFECT OF SALMONELLA TYPHOSA ENDOTOXIN ON
RENAL FUNCTION, AND THE RELATIONSHIP OF
FUNCTION TO RENAL HEMODYNAMICS**

Capt J. Y. Gillenwater, MC
E. S. Dooley, Ph.D.
Capt E. D. Frohlich, MC

Biophysics Division
US ARMY MEDICAL RESEARCH LABORATORY
Fort Knox, Kentucky

13 September 1962

Radiation and Thermal Burns
Task 03
Surgery
USAMRL Project No. 6X59-01-001

Report No. 551
USAMRL Project No. 6X59-01-001-03

ABSTRACT

EFFECT OF *SALMONELLA TYPHOSA* ENDOTOXIN ON RENAL FUNCTION, AND THE RELATIONSHIP OF FUNCTION TO RENAL HEMODYNAMICS

OBJECT

This study was designed to determine whether the decrease in renal function seen after endotoxin administration is hemodynamic or due to nephrotoxicity of endotoxin.

RESULTS

The injection of endotoxin locally or systemically caused intense renal vasoconstriction and decrease in all renal functions. Locally infused phentolamine blocked the vasoconstriction and the renal function remained normal after endotoxin injection.

CONCLUSIONS

These studies demonstrate that after injection of endotoxin there is intense renal vasoconstriction which is apparently due to systemic release of catecholamine. The alterations in renal function are due to changes in renal hemodynamics and not nephrotoxicity.


RECOMMENDATIONS

Since the preparation utilized was found to be stable and the vascular responses predictable, this may provide a reliable bio-assay for microwave-irradiated endotoxin.

ACKNOWLEDGEMENT


The authors wish to thank Drs. A. D. Keller, F. J. Haddy, and G. D. Webster for their helpful criticisms and review of this paper.

APPROVED:



SVEN A. BACH
Colonel, MC
Director, Biophysics Division

APPROVED:



FLOYD A. ODELL, Ph. D.
Technical Director of Research

APPROVED:



SVEN A. BACH
Colonel, MC
Commanding

EFFECT OF SALMONELLA TYPHOSA ENDOTOXIN ON RENAL FUNCTION, AND THE RELATIONSHIP OF FUNCTION TO RENAL HEMODYNAMICS

I. INTRODUCTION

Impaired renal function due to endotoxin has been reported in humans (1, 2) and experimental animals (3, 4). The altered renal function has been attributed to intense renal vasoconstriction and a fall in renal perfusion pressure. The vasoconstriction has been postulated to be due to both a direct local action of endotoxin on the renal vascular bed and to secondary systemic release of vasoconstrictor agents during the systemic hypotension (3). In an attempt to separate the hemodynamic from a possible nephrotoxic action of endotoxin, the local effects of Salmonella typhosa endotoxin were observed simultaneously in the dog kidney perfused at a constant blood flow and in the contralateral undisturbed kidney perfused naturally. The results of these studies are the subject of this report.

II. METHOD

Perfusion Procedures. Dogs, weighing 10 to 15 kg, were anesthetized with intravenous sodium pentobarbital (33 mg/kg). Heparin sodium (6 gm/kg) was administered intravenously. The kidney was perfused by the technique of Hardin et al (5). In brief, the abdominal aorta was exposed retroperitoneally through a left flank incision and the lumbar vessels tied. Blood was then withdrawn from a carotid artery by a precalibrated, pressure-independent Sigmamotor pump (Model T-6) and perfused into the abdominal aorta in a cephalad direction through a right angle glass cannula tied in position 3 cm below the left renal artery. Taking care not to disturb the left kidney, the aorta was occluded with a Potts clamp placed between the left and right renal arteries. This diverted the entire output of the pump through the left kidney. Renal blood flow was adjusted to an average value of 150 ml/min (range 95 to 220 ml/min) which produced a mean arterial perfusion pressure of approximately 100 mm Hg. This provided a blood flow of approximately 3.0 ml/gm of kidney tissue.

Renal arterial, renal venous, and systemic arterial pressures were measured by a needle inserted in the perfusion tubing just proximal to the glass cannula, by directly needling the renal vein, and by needling the proximal aorta, respectively. Pressures were recorded on a direct writing oscillograph recorder connected to Statham pressure transducers. Renal vascular resistances were calculated by dividing the pressure gradient from renal artery to renal vein by the blood flow and expressed as mm Hg/ml/min.

Endotoxin. Lethal doses (0.6 mg/kg) of Salmonella typhosa 0901 endotoxin (Bacto lipopolysaccharide, Difco Laboratories) were injected in one bolus directly into either the perfusion system just proximal to the renal artery (15 dogs) or systemically into the inferior vena cava (8 dogs). Renal vascular and systemic pressures were recorded for at least 10 minutes previous to and for 30 minutes following injection of endotoxin. Five additional animals were observed over a 30 minute period to obtain technique control values.

Contribution of Catecholamines. The contribution of catecholamines was delineated by the continuous infusion of phentolamine methanesulfonate (5-10 μ gm/min) directly into the renal perfusion circuit. This amount of phentolamine has been shown by others (6, 7) to be of sufficient concentration to block the local catecholamine effect without depressing systemic arterial pressure. After the phentolamine had been infused and renal and systemic vascular pressures remained stable, endotoxin was injected directly into the renal artery. Pressures were recorded over the following 30 minutes.

The tissue catecholamine levels were diminished in another group of dogs by pretreatment for three days with reserpine (8) (0.13 mg/kg intraperitoneally daily). On the fourth day endotoxin was administered locally into the renal artery and pressures recorded as described above.

Renal Function. Differential renal function was measured by comparing several renal functions of the undisturbed right kidney with the phentolaminized, perfused, left kidney. Urine was collected separately from each kidney by inserting polyethylene tubing retrograde into both ureters through a suprapubic incision. Three 10 minute control samples were obtained prior to the systemic administration (inferior vena cava) of endotoxin. Glomerular filtration rate (creatinine clearance), maximal tubular resorption of p-aminohippuric acid (TmPAH), sodium excretion ($U_{Na}V$), potassium excretion (U_KV), osmolal clearance (C_{OSM}), and free water clearance (CH_2O) were obtained from each kidney for six 10 minute collection periods following systemic injection of endotoxin. Blood levels of creatinine (16-20 mg%) and of PAH (45-50 mg%) were maintained in all experiments and adequate urine flow (greater than 2 ml/min) was insured by mannitol diuresis. PAH was determined by the method of Smith et al (9), and creatinine by the method of Bonsnes and Tausky (10). Na and K were determined on a Coleman flame photometer. Osmolalities were determined on a Fiske osmometer. Osmolal clearance (C_{OSM}) equals urine osmolality (U_{OSM}) times the urine volume (V) (ml/min) divided

by plasma osmolality (POSM). $T^C_{H_2O} = C_{OSM} - V$ and represents the quantity of water that must be added to a minute volume of urine to dilute it to the concentration of plasma.

The division of animals in the different experiments is outlined in Table I. To eliminate the possible effects of pentobarbital anesthesia, morphine sulfate (15 mg) and chloralose (75 mg/kg) anesthesia were administered in one half of the animals in the renal function group.

The responses of local and systemic administration were compared by testing them statistically according to the student "t" test (11).

III. RESULTS

Initial Transient Vasoconstriction. As shown by the recording of a representative experiment in Figure 1, subsequent to injection of endotoxin into the perfused left renal artery, there was an immediate (18 sec) transient vasoconstriction followed by a secondary prolonged renal vasoconstriction (5 to 10 min). Both local and systemic injection of endotoxin caused this immediate transient vasoconstriction (Fig. 2) which persisted for 30-40 seconds. This increase in renal vascular resistance peaked at 18 seconds (average) after local injection and 28 seconds after systemic injection. Control injections of isotonic saline and pyrogen-free distilled water were without effect. The reserpinized animals and the dogs receiving the local infusion of phentolamine responded with similar transient increases in vascular resistances (Fig. 2).

Renal Artery Administration. There was a marked secondary prolonged renal vasoconstriction in all the dogs (Fig. 3) after the local administration of endotoxin into the renal artery. Although there was some variation in the intensity of the response, the maximum renal vasoconstriction was found between 5 and 10 minutes after the endotoxin injection and seemed to coincide with the drop in systemic pressure (Fig. 1). At 5 minutes the average renal arterial pressure had doubled and was maintained at this level for another 5 minutes (Fig. 4A). At 30 minutes the pressure had returned half-way to pre-injection levels. The renal venous pressure was increased by 90 per cent at 5 minutes and declined to an average of 20 per cent above the pre-endotoxin pressures at 30 minutes. Systemic pressure dropped maximally at 10 minutes (20 per cent), with an average 10 per cent below pre-injection pressures at the end of 30 minutes.

(Fig. 4B). No significant changes were observed in renal artery, renal vein, or systemic arterial pressure of the five technique control dogs (Fig. 4).

Systemic Administration. The local administration caused an earlier and more intense secondary renal vasoconstriction than did the injection of endotoxin into the inferior vena cava. The comparison of the average renal and systemic arterial responses to endotoxin given by two different routes is shown in Figure 4. Local injection produced a 100 per cent increase in renal artery pressure at 5 minutes as compared to a 78 per cent increase if endotoxin was given systemically. The difference in renal arterial pressure at 5 minutes after local and systemic administration was statistically significant ($p < 0.01$). At 30 minutes there was a 40 per cent average increase in renal artery pressure after arterial injection, while intravenous endotoxin caused a 35 per cent increase. Systemic pressure declined in both experiments, but there was a greater decline in the systemic pressure after the local injection at the 5 and 10 minute interval which was barely statistically significant ($p = 0.05$).

Contribution of Catecholamines. In the five animals in which phentolamine was administered locally there was no secondary increase in the renal artery perfusion pressure after the local injection of endotoxin (Figs. 5 and 6). At 30 minutes there was a 30 per cent decrease in pressure. Systemic arterial pressure remained unchanged during the first 10 minutes and then gradually fell to 20 per cent of pre-injection levels at 30 minutes. In the five dogs pretreated with reserpine there was also an intense secondary renal vasoconstriction associated with the systemic arterial hypotension (Fig. 5).

Differential Renal Functions. Table II and Figure 7 show the results of the renal functions measured in the perfused phentolaminized left kidney as compared with the undisturbed right kidney for each of the eight animals. Following systemic administration of endotoxin there was a progressive decrease in function of the right kidney in all eight animals. The average urinary output decreased from 4.4 ml/min to 1.2 ml/min and glomerular filtration rate decreased from 20 ml/min to 7.6 ml/min. The TmPAH decreased from 7.3 mg/min to 1.3 mg/min; $U_{Na}V$ decreased from 156 μ Eq/min to 22 μ Eq/min, the $U_{K}V$ declined from 39 μ Eq/min to 6 μ Eq/min; C_{OSM} decreased from 5.3 ml/min to 1.9 mg/min; and the $T^{C}H_2O$ decreased from 0.9 ml/min to 0.2 ml/min.

The phentolaminized left kidney, perfused at a constant rate, did not demonstrate the decreased renal functions measured in the naturally

perfused right kidney. The average urine volume increased from 2.4 ml/min to 3.9 ml/min, GFR 11.6 ml/min to 12.4 ml/min, $^{22}\text{Na V}$ 54 $\mu\text{Eq/min}$ to 126 $\mu\text{Eq/min}$, and C_{OSM} 2.7 ml/min to 4.1 ml/min. There was a slight decline in TmPAH from 4.8 ml/min to 4.1 ml/min, U_{KV} 30 $\mu\text{Eq/min}$ to 28 $\mu\text{Eq/min}$, and $\text{T}^{\text{C}}_{\text{H}_2\text{O}}$ from 0.5 ml/min to 0.3 ml/min. The serum potassium levels decreased in all animals during the hour in which they were observed, while the plasma sodium remained unchanged.

IV. DISCUSSION

These data demonstrate that administration of lethal doses of Salmonella typhosa endotoxin directly into the renal artery produced intense renal vasoconstriction with consequent impairment of renal function.

Hinshaw and Bradley suggested that endotoxin had an initial direct effect on the renal vasculature followed by a secondary vasoconstriction at 10 minutes (3, 4), on the basis of a sudden fall in kidney weight, 15 to 35 seconds after endotoxin administration, followed by a minimum weight within 10 to 20 minutes. The pressure findings reported herein confirm and extend this impression of Hinshaw *et al* of the initial vasoconstriction followed by the secondary intense pressor phase at 10 minutes. The prompt vasoconstriction observed at 18 seconds during these experiments and in Hinshaw's experiments suggests that either the endotoxin molecule itself is a renal vasoconstrictor, or that it causes an immediate, transient release of pressor substances from the blood elements or renal tissue. Although there is a significant release of serotonin into the platelet-free plasma in the rabbit 15 seconds after endotoxin injection (12) this finding was not confirmed by the same workers in the dog (13). The initial pressor response could not be duplicated by large doses of serotonin injected locally into the renal artery. Pretreatment with reserpine and local infusion of phentolamine did not abolish this initial transitory pressor phase observed during the first minute (Fig. 2). It, therefore, seems that this initial, transient pressor phenomenon may be produced by the endotoxin molecule itself; however, the rapid local release of a pressor agent cannot be ruled out. It is interesting to note that this finding was not observed in other vascular beds studied in this laboratory (6, 14).

The injection of endotoxin directly into the renal artery caused a significantly more intense secondary renal vasoconstriction at the 5 minute interval than was observed when the endotoxin was administered systemically (Fig. 3A). This difference could be attributed to a

direct effect of endotoxin altering the reactivity of blood vessels to epinephrine (15-19). In addition, the concentration of endotoxin reaching the renal vascular bed could have been decreased by several factors, including systemic hemodilution as well as detoxification in the liver, lungs, and spleen after the systemic injection.

A detailed review of the hemodynamics of endotoxin shock found clinically and experimentally has been reported by Gilbert (20). Various studies have been reported on the effects of endotoxin on isolated, perfused vascular beds (6, 14, 21-23). Studies on the forelimb (24), lung (23), and gastric (6) circulations demonstrated vasoconstriction while the coronary vascular bed (14) responded by vasodilation. These responses have been and can be explained by a systemic release of pressor amines, since the effect of endotoxin on these vascular beds mimics the responses of epinephrine and norepinephrine (25-29). Evidence to support this concept is offered by the experiments with the phentolamine-treated kidneys. In these animals small doses of a phentolamine, administered locally into the perfused renal artery, prevented the renal vasoconstriction usually observed after endotoxin injection.

Further confirmation of this hypothesis may be offered by the data of the reserpine-treated dogs. Reserpine has been shown to diminish body stores of catecholamines (8). It has been demonstrated that blood vessels of reserpine-treated animals are hyper-responsive to these same catecholamines (30). These findings suggest that after the injection of the endotoxin, the remaining body catecholamines were liberated into the systemic circulation. The greater vasoconstriction observed during the first 5 minutes in reserpine-treated dogs could have been produced by an exaggerated vascular response of catecholamine-depleted renal vasculature. Then, as suggested by Burn and Rand (30) when norepinephrine was repleted in the renal vasculature, the constrictor response was similar to those animals not treated by reserpine.

There was no significant change in plasma sodium during the experiment. In contrast, serum potassium levels decreased markedly (Table II) after endotoxin administration. Since there was no increase in urinary excretion of potassium, it must have moved intracellularly. We have no explanation for this apparent shift of potassium intracellularly.

Hinshaw et al (4) showed that lethal doses of E. coli endotoxin produced a decreased RBF, GFR, PAH extraction, and urinary output.

These effects were attributed to the effects of systemic hypotension, renal vasoconstriction, and a possible nephrotoxic action. As they suggested, it was impossible from their study to measure changes in renal vascular pressures, and there was the further disadvantages of evaluating renal function in the presence of low urine flow rates. The present investigation was designed to measure changes in renal vascular pressures at a controlled, constant renal blood flow. Furthermore, in the renal function studies, adequate initial urine volumes were insured by osmotic diuresis, and the vascular effects were eliminated by local vascular blockage of catecholamine effects. In addition, this enabled comparison of the renal functions of one kidney not vasoconstricted with the other kidney exposed to all the local and systemic variables.

The undisturbed right kidney demonstrated similar decreases in kidney functions as were described by Hinshaw et al (4). There was a marked decrease in all of the measured functions. The average urine volume of all eight animals decreased by 75 per cent, TmPAH decreased 80 per cent, and GFR decreased by 60 per cent. The left kidney, which was maintained at a constant blood flow, and relatively unchanged pressure, demonstrated no significant decrease in any of the measured renal functions after the endotoxin injection. The observed increase in urine flow of the left kidney may be explained either by a possible diuretic action of phentolamine or by its action of vasodilation. This explanation is supported by the slight increase in GFR and the doubled sodium output of the left kidney during the phentolamine infusion.

In the renal function experiments, the only difference was that the vascular effect of catecholamines was eliminated in the left kidney; since endotoxin was administered systemically both kidneys should have received equal doses of endotoxin. In one animal (Dog No. 6), endotoxin was inadvertently given directly into the left renal artery. In this animal there was still no nephrotoxic effect demonstrated. The data, therefore, show that the primary effect of endotoxin on the kidney is hemodynamic and not nephrotoxic.

V. SUMMARY AND CONCLUSIONS

The injection of lethal doses of Salmonella typhosa endotoxin in the dog produced an initial (18 sec) transient vasoconstriction followed by a secondary intense vasoconstriction within 5 to 10 minutes after the injection. Renal arterial pressure remained elevated for 30 minutes. Since the local infusion of phentolamine blocked the secondary renal vasoconstriction it is suggested that the persistent vasoconstriction

is due to catecholamine release during the systemic hypotension. After endotoxin injection there is a marked decrease in all renal functions. In a group of animals the hemodynamic effect was eliminated in one kidney by infusing phentolamine locally and holding renal blood flow constant, enabling comparison of this kidney with the contralateral and undisturbed kidney. In the kidney in which vascular effects were eliminated there was no alteration in renal functions after systemic or local endotoxin injection. These data, therefore, show that the primary effect of endotoxin on the kidney is hemodynamic and not nephrotoxic.

VI. REFERENCES

1. Brande, A. J., J. Sieminski, D. Williams, and J. P. Sanford. Overwhelming bacteremic shock produced by gram-negative bacteria, a report of four cases with one recovery. *Univer. Michigan M. Bull.* 19: 23, 1953.
2. Brande, A. J., D. Williams, J. Sieminski, and R. Murphy. Shock-like state due to transfusion of blood contaminated with gram-negative Bacilli. *A. M. Arch. Int. Med.* 92: 75, 1953.
3. Hinshaw, L. B. and G. M. Bealey. Alterations in kidney weight produced by Escherichia coli endotoxin. *Am. J. Physiol.* 189: 329, 1957.
4. Hinshaw, L. B., W. W. Spink, J. A. Vick, E. Mallet, and J. Finstad. Effect of endotoxin on kidney function and renal hemodynamics in the dog. *Am. J. Physiol.* 201 (1): 144, 1961.
5. Hardin, R. A., J. B. Scott, and F. J. Haddy. Relationship of pressure to blood flow in the dog kidney. *Am. J. Physiol.* 199: 1192, 1960.
6. Dooley, E. S., J. B. Scott, E. D. Frohlich, and E. D. Jacobson. Hemodynamics of the stomach. III. Effects of Salmonella typhosa endotoxin on the resistance to blood flow. *USAMRL Report No. 509*, Ft Knox, Ky., 1961.
7. Haddy, F. J., J. B. Scott, M. Fleishman, and D. Emanuel. Effect of change in flow rate upon renal vascular resistance. *Am. J. Physiol.* 195: 111, 1958.

8. Waud, D. R., S. R. Kottegoda, and O. Krayner. Threshold dose and time course of norepinephrine depletion of the mammalian heart by reserpine. *J. Pharmacol. and Exper. Therap.* 124: 340, 1958.
9. Smith, H. W., N. Finkelstein, L. Aliminos, B. Crawford, and M. Graver. The renal clearance of substituted hippuric acid derivatives and other aromatic acids in dog and man. *J. Clin. Invest.* 24: 388, 1945.
10. Bosnes, R. W. and H. H. Tausky. On the colorimetric determination of creatinine by the Jaffe reaction. *J. Biol. Chem.* 158: 581, 1945.
11. "Student." The probable error of a mean. *Biometrika*, 6: 1, 1908.
12. Davies, R. B., W. R. Meeker, and W. L. Bailey. Serotonin release by bacterial endotoxin. *Proc. Soc. Exper. Biol. and Med.* 108: 774, 1961.
13. Davies, R. B., W. R. Meeker, and D. G. McQuarrie. Immediate effects of intravenous endotoxin on serotonin concentration and blood platelets. *Circulation Res.* 8: 234, 1960.
14. Frohlich, E. D., J. B. Scott, and E. S. Dooley. Hemodynamic alterations due to Salmonella typhosa endotoxin with special reference to the coronary vascular bed. *J. Clin. Invest.* 41: 147, 1962.
15. Zweifach, B. W., A. L. Nagler, and L. Thomas. The role of epinephrine in reaction produced by endotoxin of gram negative bacteria. *J. Exper. Med.* 104: 881, 1956.
16. Thomas, L., B. W. Zweifach, and B. Benacerraf. The role of epinephrine in reaction produced by endotoxin of gram negative bacteria. *J. Exper. Med.* 104: 865, 1956.
17. Landy, M. and M. J. Shear. Similarity of host response elicited by polysaccharides of animal and plant origin and by bacterial endotoxins. *J. Exper. Med.* 106: 77, 1957.

18. Zweifach, B. W. and L. Thomas. The relationship between the vascular manifestations of shock produced by endotoxin, trauma, and hemorrhage. *J. Exper. Med.* 106: 385, 1957.
19. Thomas, L., B. W. Zweifach, and B. Benacerraf. Mechanisms in the production of tissue damage and shock by endotoxin. *Tr. A. Am. Physicians*, 70: 84, 1957.
20. Gilbert, R. P. Mechanism of the hemodynamic effects of endotoxin. *Physiol. Rev.* 40: 245, 1960.
21. Gilbert, R. P. and P. J. Gordon. Peripheral vascular reactions in experimental shock due to endotoxin. *Clin. Res. Proc.* 4: 110, 1956.
22. MacLean, U. D., M. H. Weil, W. W. Spink, and M. B. Visscher. Canine intestinal and liver weight changes induced by *E. coli* endotoxin. *Proc. Soc. Exper. Biol. and Med.* 92: 602, 1956.
23. Kuida, H., L. B. Hinshaw, R. P. Gilbert, and M. B. Visscher. Effect of gram-negative endotoxin on pulmonary circulation. *Am. J. Physiol.* 192: 335, 1958.
24. Hinshaw, L. B., J. A. Vick, M. M. Jordan, and L. E. Wittmers. Vascular changes associated with development of irreversible endotoxin shock. *Am. J. Physiol.* 202: 103, 1962.
25. Haddy, F. J., M. Fleishman, and D. A. Emanuel. Effect of epinephrine, norepinephrine and serotonin upon systemic, small, and large vessel resistance. *Circulation Res.* 5: 247, 1957.
26. Aviado, D. M., Jr. and C. F. Schmidt. Effects of sympathomimetics drugs on pulmonary circulation; with special reference to a new pulmonary vasodilator. *J. Pharmacol. and Exper. Therap.* 120: 512, 1957.
27. Jacobson, E. D. Effect of histamine, norepinephrine, and acetylcholine on gastric vascular resistance. *The Physiologist* (in press).

28. Berne, R. M. Effects of epinephrine and norepinephrine on coronary circulation. *Circulation Res.* 6: 644, 1958.
29. Hardin, R. A., J. B. Scott, and F. J. Haddy. Effect of epinephrine and norepinephrine on coronary vascular resistance in dogs. *Am. J. Physiol.* 201: 276, 1961.
30. Burn, J. H. and M. J. Rand. Noradrenalin in artery walls and its dispersal by reserpine. *Brit. M. J.* 1: 903, 1958.

TABLE I

Experimental design

A.	Renal artery administration:	
1.	Effect of local injection.	15 dogs
2.	Local injection with phentolamine infusion.	5 dogs
3.	Local injection in reserpinized animals.	5 dogs
B.	Inferior vena cava administration:	
1.	Effect of systemic injection.	8 dogs
2.	Renal function studies.	8 dogs
	(a) Pentobarbital anesthesia	4 dogs
	(b) Morphine and chloralose anesthesia	4 dogs
C.	Technique controls.	5 dogs

TABLE II

Effect of *Salmonella pyramus* endotoxin on renal function

Dog no.	Time min.	Syst. BP	L. RA Press.	Urine vol.		CFR		TMPAH		COSM		T _C H ₂ O		UNAV		UKV		Blood Na K	
		mm/Hg	mm/Hg	L R	L R	L R	L R	L R	L R	L R	L R	L R	L R	L R	L R	L R	L R	mmEq/L	mmEq/L
1	Control*	160	80	1.7	4.6	9.2	23.0	4.0	6.2	2.1	5.6	0.4	1.0	6	65	21	10	140	3.6
	0-10	155	95	0.7	3.7		14.5		5.3		4.2		0.5	2	38	6	7	139	3.2
	10-20	135	95	1.7	3.1	6.4	12.2	3.6	5.0	1.8	3.5	0.1	0.4	20	21	14	7	139	2.6
	20-30	120	105	1.8	2.4	7.2	10.1	3.0	4.9	2.0		0.2		22	12	13	5	139	2.6
	30-40	110	110	2.1	1.6	9.1	8.5	3.3	3.9	2.3	2.7	0.2	1.1	39	10	13	2	142	2.6
2	Control*	165	140	1.4	3.8	8.9	21.6	7.1	11.9		4.9		1.1	140	122	17	22	141	2.8
	0-10	160	155	1.8	4.1	10.8	19.9				5.0		0.9	36	145	13	16	144	2.4
	10-20	135	110	1.9	4.0	9.4	16.8	5.7	8.9		4.8		0.8	40	132	10	13	139	2.2
	20-30	115	90	2.1	3.1	7.2	11.8	6.5	1.8		3.7		0.6	94	100	9	9	140	2.0
	30-40	100	80	2.8	2.2	12.3					3.6		1.4	96	43	10	8	138	1.9
3	Control*	125	70	2.0	5.7	11.1	21.9			2.3	6.2	0.3	0.5	26	230	17	42	141	3.8
	0-10	120	77	1.8	1.7	9.3	9.2			1.9	1.9	0.1	0.2	26	27	17	20	141	4.2
	10-20	110	80	2.2	2.2	12.2	13.7			2.3	2.5	0.1	0.3	24	20	19	30	143	3.5
	20-30	90	85	2.5	2.6	10.8	12.9			2.6	2.7	0.1	0.1	39	27	20	19	145	3.1
	30-40	80	85	3.1	2.9	11.3	13.4			3.1	3.1	0.0	0.2	73	40	21	20	140	3.0
4	Control*	150	82	2.3	2.8	14.0	19.7	4.2	3.8	3.1	3.9	0.6	1.1	19	51	35	42	142	3.4
	0-10	140	82	2.6	2.9	15.6	15.3	4.5	2.1	3.1	3.4	0.5	0.5	22	29	33	32	140	3.2
	10-20	127	82	2.4	2.4	12.8	10.5	2.9	2.7	2.7	2.8	0.3	0.4	17	24	30	30	140	3.3
	20-30	122	82	2.4	1.9	11.0	9.0	3.6	2.7	2.7	2.6	0.3	0.7	16	19	30	28	132	3.1
	30-40	95	87	2.6	1.5	11.8	6.7	3.0	2.8	2.9	2.3	0.3	0.8	19	15	28	23	139	2.9
5	Control*	137	90	3.0	5.7	14.7	19.4	3.8	7.0	3.9	7.1	0.9	1.4	111	423	28	55	140	3.8
	0-10	125	107	3.4	5.2	11.5	18.5	3.0	3.5	3.9	6.1	0.5	0.9	159	264	31	49	144	4.0
	10-20	105	105	4.0	4.9	12.9	15.6	3.3	3.6	4.4	5.5	0.4	0.6	189	230	28	45	143	3.5
	20-30	100	102	3.9	4.4	16.1	15.8	2.6	4.2	4.2	5.0	0.3	0.6	191	194	17	33	147	3.2
	30-40	87	102	4.0	4.0	14.0	16.5	2.2	2.1	4.3	4.6	0.3	0.6	205	165	17	17	147	2.7
6	Control*	112	95	3.7	3.8	12.9	19.6	7.0	9.1	3.2	4.3	0.5	0.7	41	43	36	79	136	5.0
	0-10	100	97	3.2	3.1	14.5	14.6	10.2	6.5	3.6	3.4	0.4	0.3	26	41	64	63	138	4.6
	10-20	77	112	3.0	3.2	18.0	11.3	10.9	7.3	3.5	2.3	0.5	0.1	9	44	59	33	138	3.2
	20-30	60	102	4.1	1.6	15.7	9.2	9.0	5.6	4.9	1.6	0.8	0.0	11	37	84	16	139	3.2
	30-40	50	107	4.5	1.0	18.2	6.2	11.6	4.9	5.6	1.0	1.1	0.0	14	25	88	7	141	2.9
7	Control*	175	112	2.8	6.1	12.9	21.9	3.7	5.1	3.2	6.5	0.4	0.4	67	271	39	54	132	3.2
	0-10	165	102	3.1	4.8	12.4	16.4	3.2	4.0	3.4	5.1	0.3	0.3	65	169	33	43	131	2.3
	10-20	150	90	3.3	5.0	12.9	18.3	3.5	6.2	3.4	5.2	0.1	0.2	86	173	25	39	130	2.4
	20-30	138	97	3.7	4.9	14.6	17.6	4.6	5.1	4.0	5.2	0.3	0.3	123	176	51	23	130	2.3
	30-40	125	100	3.7	3.9	12.1	14.0	3.8	3.8	3.8	4.1	0.1	0.2	139	120	17	17	129	2.2
8	Control*	160	135	1.7	2.9	8.8	14.4	4.1	4.6	1.9	3.6	0.2	0.7	18	45	29	12	148	4.0
	0-10	110	150	1.9	1.4	5.2	6.9	3.9	1.9	2.1	1.8	0.2	0.4	62	15	18	5	147	3.4
	10-20	122	127	2.5	0.5	9.9	2.5	5.8	1.4	2.8		0.3		72		28		147	2.7
	20-30	115	132	2.6	0.7	10.7	3.5	4.6	3.2	2.8		0.3		73		16		147	2.5
	30-40	87	130	2.9	0.4	10.2	2.0	4.1	1.5	3.1		0.3		87		16		150	2.3
	40-50	72	140	3.2	0.1	12.8		5.1		3.4		0.3		102		15		144	2.3
	50-60	67	132	3.6	0.1	9.3		5.1		3.7		0.1		121		17		145	2.3

*Average of three 10-minute samples.




Fig. 1. A representative experiment in which endotoxin was injected directly into the renal artery. Note the initial transient vasoconstriction was followed by an intense renal vasoconstriction at 5 to 10 minutes.




Fig. 2. Average increase in renal vascular resistance during the immediate transient vasoconstriction observed in the animals given endotoxin locally (15 dogs), systemically (8 dogs), pretreated with reserpine (5 dogs), and during the local infusion of phentelamine (5 dogs). ± 1 S. E. M. is indicated.




Fig. 3. The increase in renal arterial perfusion pressure of the 15 dogs given endotoxin locally into the renal artery.




Fig. 4. Comparison of change in average renal arterial and systemic pressure after the local (15 dogs) and systemic (8 dogs) administration of endotoxin and the technique controls (5 dogs). ± 1 S. E. M. is indicated.




Fig. 5. Comparison of change in average renal arterial pressure after local endotoxin injection (15 dogs), local injection during phentolamine infusion (4 dogs), local injection in reserpinized animals (5 dogs). ± 1 S.E.M. is indicated.




Fig. 6. A representative experiment in which endotoxin was injected into the renal artery during the local infusion of phentolamine. Note that there was still the initial transient vasoconstriction, but that the intense vasoconstriction usually seen at 5 to 10 minutes was blocked.




Fig. 7. Renal functions of a representative animal in which the hemodynamic effect was eliminated in the left kidney by infusing phentolamine locally and holding renal blood flow constant, enabling the comparison of this kidney with the contralateral undisturbed right kidney. In the left kidney in which vascular effects were eliminated there was no alteration in renal function.

DISTRIBUTION LIST OF USAMRL REPORTS
Project No. 6X59-01-001 Biophysics Division

No. of
Copies

AGENCY - DEFENSE

10 Armed Services Technical Information Agency, Arlington Hall Sta, Arlington Hall, Virginia

AGENCY - US ARMY

6 Army Attache, Box 79, Navy 100 Fleet Post Office, New York, New York, Attn: John C. Cressler, Lt Colonel, Asst. Army Attache (Medical)

6 Armed Forces Institute of Pathology, 6825 16th Street N. W., Washington 25, D. C.

1 Brooke Army Medical Center, US Army Surgical Research Unit, Library, Building 2653, Fort Sam Houston, Texas

1 Brooke Army Medical Center, Army Medical Service School, Fort Sam Houston, Texas

1 Brooke General Hospital, Medical Library, Box 151, Fort Sam Houston, Texas

5 Brooke General Hospital, Radioisotope Clinic, Brooke Army Medical Center, Fort Sam Houston, Texas

1 Brooke General Hospital, Radioisotope Section, Brooke Army Medical Center, Fort Sam Houston, Texas

1 Chief of Chemical Corps, Department of the Army, Washington 25, D.C., Attn: DDCm10/SA

1 Chief of Engineers, Department of the Army, Washington 25, D. C., Attn: ENGNB

1 Chief of Ordnance, CRDTB, Washington 25, D. C., Attn: Res & Spec, Projects Section

1 Chief Research and Development, Army Research Office, Department of the Army, Washington 25, D. C., Attn: F. E. Quimby, Chief of Life Sciences Division

1 Chief Research and Development, Department of the Army, Room 3D-442, Washington 25, D. C. Attn: Lt Colonel James B. Hartgering, Life Sciences Division, Army Research Office

1 Chief Research and Development, Department of the Army, Army Research Office, Washington 25, D. C., Attn: Scientific Information Branch

1 Fitzsimons General Hospital, US Army Medical Research and Nutrition Laboratory, Denver 30, Colorado

1 Fitzsimons General Hospital, Medical Technical Library, Denver 30, Colorado

1 Letterman General Hospital, Medical Library, San Francisco, California

1 Madigan General Hospital, Medical Library, Tacoma, Washington

1 Quartermaster Food and Container Institute, Library Branch, 1819 W. Pershing Road, Chicago 9, Illinois

1 Quartermaster Research and Engineering, Field Evaluation Agency, Fort Lee, Virginia, Attn: Technical Library

1 Quartermaster Board, US Army, Fort Lee, Virginia, Attn: President

2 Quartermaster Research and Engineering Center, Natick, Massachusetts, Attn: Technical Library

1 Redstone Arsenal, Redstone Arsenal, Alabama, Attn: CRDDW-SSI

2 US Army, Alaska, Office of the Surgeon, APO 949, Seattle, Washington

2 US Army Chemical Warfare Laboratories, Directorate of Medical Research, Army Chemical Center, Maryland

1 US Army Chemical Corps Research and Development Command, US Army Biological Warfare Laboratories, Fort Detrick, Maryland, Attn: Technical Library Branch, Documents Room

1 US Army Command and General Staff College, Fort Leavenworth, Kansas, Attn: Archives

1 US Army Dispensary, Post Surgeon, Springfield Armory, Springfield, Massachusetts

3 US Army Engineer Research and Development Laboratories, Fort Belvoir, Virginia, Attn: Technical Documents Center

1 US Army Environmental Health Laboratory, Army Chemical Center, Maryland

10 US Army, Europe, Medical Division, APO 403, New York, New York

1 US Army, Medical Command, Japan, APO 343, San Francisco, California

2 US Army Medical Liaison Branch, Office of the Chief Surgeon, Box 503, Balboa Heights, Canal Zone

No. of
Copies

US ARMY - CONTINUED

- 3 US Army Medical Research and Development Command, Commanding General, Main Navy Building, Washington 25, D. C.
- 1 US Army Medical Research Unit, Institute for Medical Research, Kuala Lumpur, Malaya
- 1 US Army Ordnance Arsenal Frankford, Philadelphia 37, Pennsylvania, Attn: Human Factors Section, 1734
- 1 US Army Ordnance Arsenal, Frankford, Philadelphia 37, Pennsylvania, Attn: A. Charles Karr, Chief Human Factors Section, CRDBA-1753
- 2 US Army Ordnance Tank Automotive Command, Detroit Arsenal, Center Line, Michigan, Attn: ORDMC-REO
- 1 US Army, Pacific, APO 958, San Francisco, California, Attn: Surgeon
- 1 US Army Signal Research and Development Laboratory, Fort Monmouth, New Jersey, Attn: SIGRM/ADT-E
- 1 US Army Standardization Group, Canada, Office of the Senior Standardization Representative, c/o Directorate of Equipment Policy, Canadian Army Headquarters, Ottawa, Canada, Attn: Colonel Joseph R. Blair, MC, Medical Liaison Officer
- 2 US Army Transportation Research and Engineering Command, Fort Eustis, Virginia, Attn: Chief, Technical Information Division
- 2 US Army Tropical Research Medical Laboratory, APO 851, New York, New York
- 1 US Army War College, Carlisle Barracks, Pennsylvania, Attn: Library
- 1 US Continental Army Command, Medical Section, Fort Monroe, Virginia
- 5 US Atomic Energy Commission, Technical Information Service Extension, P. O. Box 82, Oak Ridge, Tennessee
- 1 Valley Forge General Hospital, Phoenixville, Pennsylvania, Attn: Medical Library
- 1 Walter Reed Army Institute of Research, Division of Neuropsychiatry, Washington 12, D. C.
- 1 Walter Reed Army Institute of Research, Department of Atomic Casualties Studies, Walter Reed Army Medical Center, Washington 12, D. C.
- 5 Walter Reed Army Institute of Research, Office of the Director, Walter Reed Army Medical Center, Washington 12, D. C.
- 1 Walter Reed Army Institute of Research, Walter Reed Army Medical Center, Washington 12, D. C., Attn: Paul S. Marshall, Adjutant
- 1 Walter Reed Army Institute of Research, Walter Reed Army Medical Center, Washington 12, D. C., Attn: Stanley W. Weitzner, M. D.
- 2 Walter Reed General Hospital, Audiology and Speech Center, Walter Reed Army Medical Center, Washington 12, D. C., Attn: James P. Albrite, Major, MC
- 1 Walter Reed Army Medical Center, US Army Medical Service, Historical Unit, Washington 12, D. C., Attn: Lucy W. Lazarus, Chief, General Reference and Research Branch
- 1 Watervliet Arsenal CRDEF-R, Technical Library, Watervliet, New York

AGENCY - US NAVY

- 1 Bureau of Aeronautics (AE-14) Department of the Navy, Washington 25, D. C.
- 2 Bureau of Medicine and Surgery, Research Division (71) Department of the Navy, Washington 25, D. C.
- 1 Bureau of Naval Operations, (Op03EG) Operations Evaluation Group, Navy Department, Washington 25, D. C.
- 1 Bureau of Naval Weapons, (RAAE-14) Navy Department, Washington 25, D. C.
- 2 Bureau of Naval Weapons, Department of the Navy, Washington 25, D. C., Attn: Code RMO, Pyrotechnics Branch
- 2 Bureau of Ships, Physics and Oceanography Branch, Main Navy Building, 18th and Constitution Avenue, Washington 25, D. C.
- 1 Bureau of Yards and Docks, Department of the Navy, Washington 25, D. C., Attn: Code D-440
- 1 Marine Corps Educational Center, Archives and Historical Group, Marine Corps School, Quantico, Virginia
- 1 Naval Air Crew Equipment Laboratory, Naval Air Material Center, Philadelphia 12, Pennsylvania
- 1 Naval Air Engineering Facility, Naval Air Material Center, Philadelphia 12, Pennsylvania, Attn: Technical Library
- 1 Naval Medical Field Research Laboratory, Camp LeJeune, North Carolina, Attn: Library

No. of
Copies

US NAVY - CONTINUED

- 1 Naval Medical Research Institute, Technical Reference Library, National Naval Medical Center, Bethesda 14, Maryland
- 1 Naval Research (Code 408) Special Assistant for Medical and Allied Sciences, Department of the Navy, Washington 25, D. C.
- 10 Naval Research Branch Office, Navy 100, Box 39, Fleet Post Office, New York, New York
- 1 US Naval Air Development Center, Aviation Medical Acceleration Laboratory, Johnsville, Pennsylvania, Attn: Library
- 1 US Naval Air Station, Chief of Naval Air Technical Training, Memphis 75, Tennessee, Attn: Staff Medical Officer
- 1 US Naval Civil Engineering Laboratory, Commanding Officer and Director (Code 131) Port Hueneme, California
- 1 US Naval Medical Research Laboratory, Box 100, US Navy Submarine Base, New London, Connecticut, Attn: Technical Information Section Library
- 2 US Naval Medical School, National Navy Medical Center, Bethesda 14, Maryland
- 2 US Naval Missile Center, Point Mugu, California, Attn: Librarian Code 21211
- 1 US Naval Radiological Defense Laboratory, San Francisco 24, California, Attn: Director (222)
- 1 US Naval Ordnance Test Station, Medical Officer, Station Hospital, Code 88, China Lake, California
- 1 US Naval School of Aviation Medicine, US Naval Aviation Medical Center, Director of Research, Pensacola, Florida
- 1 US Naval School of Aviation Medicine, US Naval Aviation Medical Center-54, Pensacola, Florida, Attn: Commanding Officer
- 2 US Naval School of Aviation Medicine, US Naval Aviation Medical Center, Pensacola, Florida, Attn: Asst Director of Research
- 1 US Naval Supply Research and Development, Facility Technical Library, 29th Street and Third Avenue, Brooklyn 32, New York
- 1 US Naval Weapons Plant, Experimental Diving Unit, Washington 25, D. C.

AGENCY - AIR FORCE

- 1 Air Force Flight Test Center, (FTFTH) Edwards Air Force Base, California, Attn: Lt Colonel Burt Rowen, USAF (MC)

AIR RESEARCH AND DEVELOPMENT COMMAND

- 1 Air Research and Development Command, US Air Force (RDSFC-LS) Andrews Air Force Base, Washington 25, D. C.
- 2 Wright Air Development Center, Wright-Patterson Air Force Base, Ohio, Attn: (WCLDOR)
- 1 Wright Air Development Center, (WCLDAT-Library), Wright-Patterson Air Force Base, Ohio
- 1 Air Proving Ground Center, Air Research and Development Command, (PGTRI) US Air Force, Eglin Air Force Base, Florida
- 2 Air Training Command, (ATCSG) Randolph Air Force Base, Texas
- 1 Arctic Aeromedical Laboratory, APO 731, Seattle, Washington, Attn: Librarian
- 1 Assistant for Ground Safety, DIGS, US Air Force, Washington 25, D. C.
- 1 Human Factors Division, US Air Force, Director of Research and Development, Washington 25, D. C. Attn: Colonel Charles H. Roadman, USAF (MC)
- 2 Langley Research Center, National Aeronautics and Space Administration, Langley Field, Virginia, Attn: Mrs Elizabeth R. Gilman, Librarian
- 2 US Air Force Aerospace Medical Center (ATC) School of Aviation Medicine (SAM-41.2) Brooks Air Force Base, Texas
- 1 US Air Force Aerospace Medical Center (ATC) US Air Force Hospital, Lackland Air Force Base, Texas, Attn: Medical Library
- 1 US Air Force, Strategic Air Command, Office of the Surgeon, Offutt Air Force Base, Nebraska

GOVERNMENTAL AGENCIES

- 1 Argonne National Laboratory, P. O. Box 299, Lemont, Illinois, Attn: Heylande D. Young

No. of
Copies

GOVERNMENTAL AGENCIES - CONTINUED

- 1 Civil Aeromedical Research Center, Aeronautical Center, P. O. Box 1082, Oklahoma City, Oklahoma
- 1 Civil and Defense Mobilization, Technical Library, Room G-2-B, Battle Creek, Michigan
- 1 Chief Defense Atomic Support Agency, Washington 25, D. C., Attn: Document Library Branch
- 1 Library of Congress, Science and Technology Division, Washington 25, D. C., Attn: Dr. A. J. Jacobius
- 1 National Institutes of Health, Library, Bethesda 14, Maryland
- 1 National Institutes of Health, National Cancer Institute, Radiation Branch, Bethesda 14, Maryland
- 1 National Institutes of Health, Division of Research Grants, Bethesda 14, Maryland
- 1 National Library of Medicine, 7th and Independence Avenue, S. W., Washington 25, D. C.
- 2 National Research Council, Division of Medical Sciences, Medical Records, 2101 Constitution Avenue, Washington 25, D. C.
- 2 Research Library, Brookhaven National Laboratory, Upton, Long Island, New York
- 1 US Atomic Energy Commission, Division of Biology and Medicine, Washington 25, D. C., Attn: Mr Robert L. Crossie, Director, CETO

OTHER AGENCIES

- 1 Arctic Health Research Center, Library, P. O. Box 980, Anchorage, Alaska
- 1 Boeing Airplane Company, Central Medical Library, 11-40, P. O. Box 3707, Seattle 24, Washington
- 1 Boeing Airplane Company, Wichita Division, Library, Wichita 1, Kansas
- 2 Boeing Airplane Company, Aero-Space Division, Biosciences Section, Seattle 24, Washington, Attn: Dr. Rosney H. Lowry
- 2 Department of Health, Education and Welfare, Division of Radiological Health - BSS, Room 5830, South HEW Building, Washington 25, D. C., Attn: Information Officer
- 2 Department of Health, Education and Welfare, Division of Radiological Health - BSS, Penthouse, South Building, Washington 25, D. C. Attn: DRH Reference Room
- 1 Electric Boat, Division of General Dynamics Corporation, Special Projects Division, Groton, Connecticut, Attn: Mr William J. King, Human Factors Section
- 1 General Electric Company, Technical Military Planning Operation, 735 State Street, Santa Barbara, California
- 1 The John Crerar Library, 88 East Randolph Street, Chicago 1, Illinois
- 1 Kings County Hospital, Department of Anesthesiology, Brooklyn, New York Attn: Dr. Stanley W. Weitzner
- 1 Lankenau Hospital, Lancaster and City Line Avenues, Philadelphia 31, Pennsylvania, Attn: Dr. Steven M. Horvath, Division of Research
- 1 Linfield Research Institute, McMinnville, Oregon
- 1 Mayo Clinic, Section of Biophysics, Rochester, Minnesota, Attn: Dr. Kenneth N. Ogle
- 1 Mead Johnson and Company, Medical Department, 2404 W. Pennsylvania Street, Evansville 21, Indiana
- 7 National Aeronautics and Space Administration, 1520 H. Street, N. W., Washington 25, D. C., Attn: Mr. Bertram A. Mulcahy, Director of Technical Information
- 1 Sensory Biophysics Group, Pattern Recognition Department, IBM Research Center, Box 218, Yorktown Hts, New York
- 1 Space Technology Laboratories, Inc, Information Services, P. O. Box 59001, Document Acquisition Group, Los Angeles 45, California
- 1 The Rand Corporation, 1700 Main Street, Santa Monica, California, Attn: Library
- 1 Yerkes Laboratories of Primate Biology, Orange Park, Florida

MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS

- 1 Albany Medical College Library, New Scotland Avenue, Albany 8, New York
- 1 Bowman Gray School of Medicine, Winston-Salem, North Carolina, Attn: Library
- 1 College of Medical Evangelists, White Memorial Medical Library, 1720 Brooklyn Avenue, Los Angeles 33, California
- 1 College of Medical Evangelists, Vernier Radcliffe Memorial Library, Loma Linda, California

No. of
Copies MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS CONTINUED

- 1 College of Physicians of Philadelphia, 19 South 22nd Street, Philadelphia 3, Pennsylvania, Attn: Library, College of Physicians of Philadelphia
- 1 Columbia University, Medical Library, 630 West 168th Street, New York 32, New York
- 1 Cornell University, Medical College Library, 1300 York Avenue, New York 21, New York
- 1 Creighton Medical Library, 302 North 14th Street, Omaha 2, Nebraska
- 1 Dartmouth College Medical Library, Baker, Hanover, New Hampshire
- 1 Florida State University, Department of Psychology, Tallahassee, Florida, Attn: W. W. Dawson
- 2 George Washington University, Human Resources Research Office, P. O. Box 3596, Washington 7, D. C., Attn: Library
- 1 Harvard Medical Library, 25 Shattuck Street, Boston 15, Massachusetts
- 1 Indiana University Medical Center, School of Medicine Library, 1100 West Michigan Street Indianapolis 7, Indiana
- 1 Jefferson Medical College Library, 1025 Walnut Street, Philadelphia 7, Pennsylvania
- 1 Johns Hopkins University, Operations Research Office, 6935 Arlington Road, Bethesda, Maryland, Washington 14, D. C., Attn: Library
- 1 Johns Hopkins University, Welch Medical Library, 1900 East Monument Street, Baltimore 5, Maryland
- 1 Marquette University, Medical-Dental Library, 580 North 18th Street, Milwaukee 3, Wisconsin
- 1 Medical College of Virginia, Tompkins-McCaw Library, Richmond 19, Virginia
- 1 New York Academy of Medicine, 2 East 103rd Street, New York 29, New York, Attn: Library
- 1 New York University, Bellevue Medical Center Library, 550 First Avenue, New York 16, New York
- 1 Northwestern University Medical School, Archibald Church Library, 303 East Chicago Avenue, Chicago 11, Illinois
- 1 Ohio State University, Chemical Abstracts, Columbus 10, Ohio, Attn: Library
- 1 Ohio State University, Operations Research Group, Engineering Experiment Station, 156 West Tenth Avenue, Columbus 10, Ohio
- 1 Ohio State University, School of Optometry, Topaz Library of Visual Science, 338 West Tenth Avenue, Columbus 10, Ohio
- 1 Ohio State University, Topaz Library, School of Optometry, 338 West Tenth Avenue, Columbus 10, Ohio
- 1 Rush Medical College Library, 1758 West Harrison Street, Chicago 12, Illinois, Attn: Frida Pliefke, Librarian
- 1 St. Louis University, School of Medicine, 1422 South Grand Avenue, St. Louis 4, Missouri Attn: Library
- 1 Stanford University, Department of Physiology, Stanford, California, Attn: Dr. J. M. Crismon
- 1 Stanford University Medical Center, 300 Pasteur Drive, Stanford, California, Attn: Lane Medical Library
- 1 State University of Iowa, College of Medicine Library, Medical Laboratories Building, Iowa City, Iowa
- 1 State University of New York, Downstate Medical Center, Department of Anesthesiology, 450 Clarkson Avenue, Brooklyn, New York, Attn: Dr. Stanley W. Weitner
- 1 State University of New York, Downstate Medical Center, Medical Library, 450 Clarkson Avenue, Brooklyn 3, New York
- 1 Texas Medical Center Library, Jesse H. Jones Library Building, Texas Medical Center, Houston 25, Texas
- 2 Tufts University, Institute for Applied Experimental Psychology, Medford 55, Massachusetts, Attn: Paul G. Ronco
- 1 Vanderbilt University School of Medicine, Department of Medicine, Nashville 5, Tennessee, Attn: George R. Meneely, M.D. Asso. Prof. of Medicine, Director Radioisotope Center
- 1 University of Alabama, Medical Center Library, 1919 Seventh Avenue South, Birmingham 3, Alabama
- 1 University of Arkansas, Medical Center Library, 4301 West Markham, Little Rock, Arkansas

No. of
Copies

MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS CONTINUED

- 1 University of Buffalo, Medical-Dental Library, Buffalo 14, New York
- 1 University of Buffalo, Department of Psychology, Buffalo 14, New York
- 1 University of California, U. S. Naval Medical Research Unit No. 1, Berkeley 4, California
- 1 University of California Medical Center, Biomedical Library, Los Angeles 24, California
- 1 University of California, Civil Defense Research Project, 1301 South 46th Street,
Richmond 4, California
- 1 University of Chicago, US Air Force Radiation Laboratory, 930 E. 58th Street, Chicago 37,
Illinois, Attn: Dr. Kenneth P. DuBois
- 1 University of Chicago, Faculty Exchange Box 184, Chicago 37, Illinois, Attn: W. D. Neff,
Laboratory of Physiological Psychology
- 1 University of Cincinnati, Kettering Laboratory, Eden and Bethesda Avenue, Cincinnati 19,
Ohio
- 1 University of Illinois, Department of Psychology, Urbana, Illinois, Attn: Dr. Wm E.
Kappauf
- 1 University of Illinois, Aeromedical Laboratory, 840 South Wood Street, Chicago 12,
Illinois, Attn: Dr. John P. Marbarger
- 1 University of Kansas Medical Center, Clendenen Medical Library, Kansas City 12, Kansas
- 1 University of Louisville, School of Medicine, Louisville Medical Library, 101 West
Chestnut Street, Louisville 2, Kentucky
- 1 University of Maryland, Health Sciences Library, 111 South Greene Street, Baltimore 1,
Maryland
- 1 University of Miami, School of Medicine, Jackson Memorial Library, 1000 N. W. 17th Street,
Miami 36, Florida
- 1 University of Michigan, Serials and Documents Section, General Library, Ann Arbor, Michigan
- 1 University of Minnesota Library, Gift and Exchange Department, Minneapolis 14, Minnesota
- 1 University of Mississippi, Medical School, Rowland Medical Library, Jackson, Mississippi
- 1 University of Missouri, Medical Library, Columbia, Missouri
- 1 University of Nebraska, College of Medicine Library, 42nd and Dewey Avenue, Omaha 5,
Nebraska
- 1 University of North Carolina, Division of Health Affairs Library, North Carolina Memorial
Hospital, Chapel Hill, North Carolina
- 1 University of Oklahoma Medical Center, 801 N. E. 13th Street, Oklahoma City, Oklahoma,
Attn: Library
- 1 University of Oregon Medical School Library, 3181 S. W. Sam Jackson Park Road, Portland
1, Oregon
- 1 University of Pittsburgh, Falk Library of the Health Professions, Pittsburgh 13,
Pennsylvania
- 1 University of Rochester, Atomic Energy Project, Technical Report Control Unit, P. O.
Box 287, Station 3, Rochester 20, New York
- 1 University of Southern California, School of Medicine Library, University Park, Los
Angeles 7, California
- 1 University of South Dakota, Medical Library, Vermillion, South Dakota, Attn: Medical
Library
- 1 University of Tennessee, Department of Bacteriology, Knoxville, Tennessee, Attn: Dr. D.
Frank Holtman
- 1 University of Tennessee, Institute of Clinical Investigation, Clinical Physiological
Laboratories, Memphis 3, Tennessee, Attn: Dr. Gerald E. Spurr
- 1 University of Tennessee, Mooney Memorial Library, 62 S. Dunlay, Memphis 3, Tennessee
- 1 University of Texas, The Radiobiological Laboratory, RFD 4 Box 189, Austin, Texas
- 1 University of Texas, Medical Branch Library, Galveston, Texas
- 1 University of Utah, Medical Library, Salt Lake City 12, Utah
- 1 University of Vermont, College of Medicine Library, Burlington, Vermont
- 1 University of Virginia Hospital, Medical Library, Charlottesville, Virginia
- 1 University of Washington, Health Sciences Library, Seattle 5, Washington

No. of MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS - CONTINUED
Copies

- 1 University of Wisconsin Medical Library, S. M. I. Building, N. Charter Street, Madison 6, Wisconsin
- 1 University of West Virginia, Medical Center Library, Morgantown, West Virginia

FOREIGN

- 6 British Army and Navy Medical Liaison Officer, B. J. S. M. (Army Staff) 1800 K. Street, N. W., Washington 25, D. C.
- 8 Canadian Liaison Officer, Office of the Surgeon General, US Army, Room 2709, Main Navy Building, Canadian Joint Staff, (Washington) Washington 25, D. C.
- 2 Defence Research Member, Canadian Joint Staff, (Washington) Room 2709, Main Navy Building, Washington 25, D. C.
- 1 Laboratoire de Pathologie et Therapeutique Generales, 32 Boulevard de la Constitution, Liege, Belgique. Attn: Directeur Professeur Z. M. Bacq
- 1 Physics Research Laboratories, Upper Redlands Road, Reading, Berks, England. Attn: Prof. R. W. Ditchburn
- 1 Royal Society of Medicine Library, 1 Wimpole Street, London W. 1, England

UNCLASSIFIED

UNCLASSIFIED