

UNCLASSIFIED

AD 296 242

*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

Copy 112
19 November 1962

**ACCELERATED INDUCTION OF HEPATOMAS IN
FAST NEUTRON IRRADIATED MICE INJECTED WITH
CARBON TETRACHLORIDE**


by
L. J. Cole
P. C. Nowell*

*Dept of Pathology, School of Medicine, University of Pennsylvania, Philadelphia, Pa.

**U.S. NAVAL RADIOLOGICAL
DEFENSE LABORATORY**

SAN FRANCISCO 24, CALIFORNIA

12ND. P7463



296 242
AS AD NO. 296 242

296 242

EXPERIMENTAL PATHOLOGY BRANCH
L. J. Cole, Head

BIOLOGICAL AND MEDICAL SCIENCES DIVISION
E. L. Alpen, Head

ADMINISTRATIVE INFORMATION

This work was accomplished under the Bureau of Medicine and Surgery Task MR005.08-5200, Subtask 3, Technical Objective AW-6, as described in the U.S. Naval Radiological Defense Laboratory Annual Report to the Bureau of Medicine and Surgery (OPNAV Form 3910-1) of 31 December 1961, and is listed in the U.S. Naval Radiological Defense Laboratory Fiscal Year 1963 Technical Program under Program A3, Problem 2. This study was supported through funds provided by the Bureau of Medicine and Surgery, and the Defense Atomic Support Agency under DASA Subtask 2 of RD 42-52, Subproject 03.027.

Eugene P. Cooper

Eugene P. Cooper
Scientific Director

E. B. Roth

E. B. Roth, CAPT USN
Commanding Officer and Director

ABSTRACT

Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control non-irradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP X-rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 61%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

SUMMARY

The Problem:

Although it is well established that ionizing radiation is a carcinogenic agent in animals and man, there is still a lack of information as to the relative potency of high LET radiations (such as fast neutrons) versus low LET radiations (such as X-radiation) with respect to long-term carcinogenic hazard. Furthermore, in the light of the accumulating evidence that the development of tumors involves at least two separate phases, i.e., so-called "initiation," and "promotion," it is of some importance to gain knowledge on the role of promoting agents in radiation carcinogenesis. In the system investigated here, carbon tetrachloride (CCl_4) was used to induce mitotic activity in the liver after irradiation.⁴ The hypothesis tested was whether the application of this specific proliferative stimulus would act as a promoting agent for tumor formation in the target tissue, in this case liver, under these conditions.

The Findings:

Young adult (C57L x A) F_1 hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl_4 , given either at 2, 12, 15, or 18 months post-irradiation. Control non-irradiated mice received a single injection of CCl_4 . Other groups of mice were exposed to a single 500 rad dose of 250 KVP X-rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl_4 , the hepatoma incidence attained a value of 61%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron- CCl_4 mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl_4 but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl_4 seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on

cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

INTRODUCTION

It has been generally accepted that carcinogenesis is a multi-stage process, involving at least two separate phases, i.e., an "initiating action", and a "promoting phase" (1). With respect to carcinogenesis by ionizing radiations, it is becoming increasingly more evident that the physiological, hormonal, and local tissue events implicated in the final manifestation of many neoplasms are in fact specific proliferative stimuli or growth regulators for target tissues, and therefore can be considered as promoting factors acting on radiation-altered cells (2,3). The experimental studies of Furth et al (4) on radiation-induced mammatropic tumors, and those of Cronkite et al (5) on mammary tumors provide excellent examples of this concept.

However, our knowledge of the nature of the "initiating" action of ionizing radiation is very limited. The investigation of carcinogenesis following exposure to high LET versus low LET radiations appears to offer some possible new insights into this problem, particularly in view of recent developments in karyotype analysis of tumor and irradiated cells (6), and of the application of cloning techniques for investigating radiation effects at the cellular level(7).

In previously published studies (8,9,10) on fast neutron carcinogenesis in mice we observed a large increase in the incidence of gastrointestinal neoplasms in LAF₁ mice exposed to a single high sublethal dose of fast neutrons, whereas the occurrence of such neoplasms was much less frequent in mice exposed to 250 KVP X-rays. The incidence of neoplastic lesions in the kidneys of the neutron-irradiated mice was only slightly increased over that of non-irradiated controls, while hepatomas were not more frequent after irradiation. The question arises whether the difference in carcinogenic response of the gastrointestinal epithelium versus liver or kidney with respect to fast neutrons, may be related to the fact that the latter tissues exhibit a low mitotic rate and cell turnover, whereas gastrointestinal epithelium is a rapidly proliferating tissue. On the supposition that ionizing radiation initiates a latent pre-neoplastic, heritable change of a more or less permanent nature (2,11) in the cells of liver and kidney, as well as in the gastrointestinal epithelium, it was of interest to establish whether the tumorigenic effect of fast neutron radiation could be more readily evoked in the

liver or kidneys, by applying a specific mitotic stimulus to these otherwise non-proliferating tissues.

The persistence of latent radiation-induced mitotic aberrations in the resting liver for periods of several months after irradiation is well established. These cytological changes become overt when the liver is induced to proliferate by administration of CCl_4 (12) or by partial hepatectomy (13,14). Similar radiation-induced latent injury to proliferative capacity has been observed also in the kidneys (15).

MATERIALS AND METHODS

The mice employed were (C57L x A) F_1 hybrids (so-called LAF_1) of both sexes. At age 10-15 weeks old, the mice received a single whole body exposure to fast neutrons (with a spectrum similar to that of fission neutrons), produced in the 60-inch cyclotron at the Crocker Laboratory of the University of California, Berkeley by the $\text{Be}(p,n)\text{B}$ reaction. The irradiation procedure has been described previously (8); and the physical characteristics of these neutrons, as well as the details of dosimetry have been reported by Tochlin et al (16). The dose rate was approximately 30 rad per minute. The neutron doses employed were in the range from 165 rad to 306 rad. Other groups of LAF_1 mice 10-15 weeks old, were exposed to a single sublethal dose of 250 KVP X rays (500 rad) at dose rate of approximately 28 rad per minute. Subgroups of these irradiated mice then received a single subcutaneous injection of 0.15 ml of 40% CCl_4 in sesame oil, given either at 2, 12, 15 or 18 months post-irradiation. Control non-irradiated mice received a single injection of CCl_4 . Tissues were taken from mice which died or were sacrificed up to 28 months post-irradiation. The tissues were fixed in formalin, and stained with hematoxylin and eosin after routine processing.

RESULTS

Hepatomas: A summary of the findings is given in Table I. Of 58 neutron-irradiated mice examined to date, which did not receive CCl_4 , 11 (19%) showed hepatomas. By comparison, in 42 mice exposed to 500 rad of X-rays, only 1 hepatoma was found (2%). Thus, the hepatoma incidence in the neutron-irradiated mice is markedly increased over that of the X-irradiated group. In the neutron-irradiated mice injected with CCl_4 , the hepatoma incidence attained a level of 61% (11 out of 18), three times that found in the mice which were exposed to fast neutrons only.

TABLE I

INCREASED INCIDENCE OF HEPATOMAS IN IRRADIATED MICE
RECEIVING A SINGLE INJECTION OF CARBON TETRACHLORIDE

TREATMENT	AGE AT DEATH (mos)	NO. OF MICE	HEPATOMAS	
			No.	%
Neutrons (165-306 rad)	20-30	58	11	19
Neutrons + CCl ₄ (171-306 rad)	20-31	18	11	61
X-rays (500 rad)	18-33	41	1	2
CCl ₄ only (0 rad)	17-22	6	0	0

The tumors varied in size from 5 mm to several cm. diameter (Figure 1) the largest tumors being in the neutron-CCl₄ mice. Histologically, the tumors in all groups were similarly composed of cords and nests of relatively well-differentiated hepatic cells (Figure 2). They were generally well-circumscribed, and metastases were not observed. In addition, of the 7 mice in the neutron-CCl₄ group which did not have hepatomas, 5 showed marked pleomorphism and atypicality of liver cell nuclei (Figure 3). Similar, but generally less severe nuclear changes were observed in only 6 of the 47 mice, irradiated with neutrons only, which did not have hepatomas. Finally, of the small group of mice which received CCl₄ but no irradiation, and were sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities.

Other tumors: The incidence of tumors other than hepatomas was comparable, in the various groups, to that observed in previous reports by us (10), and by Upton et al (17).

DISCUSSION

Although the number of animals involved in these experiments is relatively small, certain conclusions appear to be warranted from the data. First, that fast neutron radiation may have a relatively greater potency than X-rays in eliciting hepatomas in these mice. This observation is at variance with earlier studies by ourselves (10) and by Upton et al (17) in which fast neutron irradiation and X-irradiation resulted in similar hepatoma incidence (2-3%). The marked increase in incidence in the present series may be due, in part, to the more careful examination of the livers. However, further data are obviously required. It is of interest that Rosen et al (18) have recently reported a relative potency between 2 and 3 for fission-spectrum neutrons versus X-rays for the production of renal neoplasms in rats.

Second, a single inoculation of CCl₄ to neutron-irradiated mice resulted in a three-fold increase in hepatoma incidence. Although the incidence of hepatomas in non-irradiated mice can be greatly increased by multiple feedings of CCl₄, as reported by Edwards (19), a single injection of CCl₄ did not result in hepatoma formation (19). The absence of liver neoplasms in our small non-irradiated group which received CCl₄ alone, is in accord with this observation.

It appears, therefore, that in this system CCl₄ acts as a promoting agent for liver carcinogenesis. The proliferative response of the liver to this hepatotoxic agent is well known. However, this proliferative effect may not be the sole basis for the promoting action of CCl₄. Ingle and Baker (20) carried out multiple (12 times) partial hepatectomies



Fig. 1 Gross view of hepatoma in neutron irradiated- CCl_4 mouse.
The tumor mass occupies most of the liver.




Fig. 2 Low power view of well-circumscribed hepatoma from 28 month old LAF₁ mouse irradiated with fast neutrons (300 rad) at age 3 months, followed by a single injection of CCl₄ 12 months later. (x 35)




Fig. 3 High power view showing nuclear variability in liver cells of another mouse similarly treated. This mouse did not have a hepatoma. (x 250)

in non-irradiated rats during a period of 1 year. Under these conditions of repetitive hyperplasia, no development of liver neoplasia was found. Since repeated feeding of CCl_4 does result in hepatomas in non-irradiated mice (19), it is possible that the promoting action of CCl_4 in the present experiments involves factors in addition to a proliferative stimulus. Pertinent to this is the recent finding of Smuckler et al (21) of an early intracellular defect in protein synthesis associated with alterations in the endoplasmic reticulum in the livers of rats, following a single intubation with CCl_4 .

There is considerable evidence that high LET radiation produces greater acute damage to cells, when compared with low LET radiations. In animal experiments, this is reflected in the greater histological damage, and the relatively slower regeneration of the intestinal epithelium (22) and of the bone marrow (23) after exposure of mice to fast neutrons. At the cellular level, Barendsen (24) has shown that the surviving fraction of cultured human kidney cells irradiated *in vitro* with α -radiation, decreases exponentially as dose increases. No shoulder was evident in the curve on a semi-logarithmic plot. The LD_{37} was only 65 rads. The α -irradiated cells exhibited no repair on dose fractionation. These data are interpreted by Barendsen as a "single event" type of action, by this high LET radiation. By contrast, the survival curves for such cells after exposure to low LET radiation (200 KV X-rays) exhibit a distinct shoulder, and show recovery when the radiation dose is fractionated (25); (cf. also 26). Such effects are characteristic of the "cumulative" type of action of low LET radiation, according to Barendsen. Similar results have been obtained by Andrews and Berry (27, 28) in studying the effect of fission-spectrum fast neutron radiation on reproductive capacity of mouse lymphocytic leukemia cells, measured by means of transplantability. The response of these cells to fast neutron irradiation, according to these authors, is a "no-threshold, no-recovery one".

These differences in the mode of action of high LET versus low LET radiations, reflected in the relatively more severe intracellular defect, probably at the chromosomal level (cf.7), in cells exposed to high LET radiations, may well be the basis for the marked difference in carcinogenic potency between fast neutrons and X-rays, as observed in the liver and in the gastrointestinal epithelium. According to this concept, carcinogenic "initiation" is correlated with the degree and persistence of a radiation-induced cellular defect or alteration within a population of cells. The fact that fast-neutron irradiated cells show the "single event", no-recovery type of response with respect to reproductive capacity, as discussed above implies that any radiation-induced genetic damage incurred by these cells would persist for more prolonged periods without repair, as compared with X-irradiated cells. Although the true nature

of this intracellular lesion is unknown, it would appear to reside at the chromosomal or subchromosomal level, and to be associated with an increased genetic instability or mutability in this population; as a consequence, the eventual appearance of a small number of the postulated "mutant" neoplastic cells becomes statistically more probable. The establishment of such "mutant" cell clones and their emergence as an actual tumor mass, presumably by selective proliferation, would depend, at least in part, upon the level of proliferative activity in the specific tissue involved. Under proper conditions (e.g. mitotic stimulation by CCl_4 damage) these aberrant cells and their progeny would be more and more likely to change in the direction of release from hormonal or other regulatory control (cf. 29). Similar accelerated induction of renal neoplasms has been shown to occur in X-irradiated mice following the proliferative stimulus of unilateral nephrectomy (30).

What is the promoting agent in the case of the neutron-irradiated mice which did not receive CCl_4 ? One would have to postulate here that the cell turnover in the liver although extremely slow, would, over a period of 2 years or more, permit the eventual emergence of the radiation-initiated neoplastic cell clones. Any endogenous factors (for example, hormonal) or extrinsic influences (for example, dietary or toxic) which increase cell turnover in the liver, would then be expected to hasten the carcinogenic process under these conditions.

REFERENCES

1. I. Berenblum. Carcinogenesis and Tumor Pathogenesis. Adv. in Cancer Res. 2: 129-175, 1954.
2. J. Furth and K. Yokoro. Components of Radiation Carcinogenesis in "Proc. 3rd Australasian Conference on Radiation Biology", Butterworths Scientific Publications, London, 1960.
3. J. Furth. Radiation Neoplasia and Endocrine Systems in "Radiation Biology and Cancer", University of Texas Press, Austin, 1959, pp. 7-25.
4. J. Furth. Vistas in the Etiology and Pathogenesis of Tumors. Federation Proc. 20(2): 865-873, 1961.
5. E. P. Cronkite, C. J. Shellabarger, V. P. Bond, and S. W. Lippincott. Studies on Radiation-Induced Mammary Gland Neoplasia in the Rat. I. The Role of the Ovary in the Neoplastic Response of Breast Tissue to Total or Partial X-irradiation. Radiation Res. 12: 81-93, 1960.
6. T. S. Hauschka. The Chromosomes in Ontogeny and Oncogeny. Cancer Res. 21: 957-974, 1961.
7. T. T. Puck and P. I. Marcus. Action of X-rays on Mammalian Cells. J. Exper. Med. 103: 653-666, 1956.
8. P. C. Nowell, L. J. Cole, and M. E. Ellis. Induction of Intestinal Carcinoma in the Mouse by Whole-Body Fast Neutron Irradiation. Cancer Research 16: 873-876, 1956.
9. P. C. Nowell, L. J. Cole, and M. E. Ellis. Neoplasms of the Glandular Stomach in Mice Irradiated With X rays or Fast Neutrons. Cancer Res. 18: 257-260, 1958.
10. P. C. Nowell, and L. J. Cole. Late Effects of Fast Neutrons Versus X-rays in Mice: Nephrosclerosis, Tumors, Longevity. Radiation Res. 11: 545-556, 1959.
11. F. M. Burnet. Cancer: Biological Approach. I. Processes of Control. Brit. Med. J. 1: 779-786, 1957.
12. M. D. Albert. X-irradiation Induced Mitotic Abnormalities in Mouse Liver Regenerating After Carbon Tetrachloride Injury. I. Total Body Irradiation. J. Nat. Cancer Inst. 20: 309-319, 1958.
13. M. D. Albert, and N. L. R. Bucher. Latent Injury and Repair in Rat Liver Induced to Regenerate at Intervals After X-radiation. Cancer Res. 20: 1514-1522, 1960.
14. G. F. Leong, R. L. Pessotti and J. S. Krebs. Liver Cell Proliferation in X-irradiated Rats After Single and Repetitive Partial Hepatectomy. J. Nat. Cancer Inst. 27: 131-143, 1961.
15. L. J. Cole and V. J. Rosen, Jr. Latent Radiation-Inhibition and Recovery of Mitotic Activity in Mouse Kidney. Exper. Cell Res. 23: 416-418, 1961.
16. E. Tochlin, B. Shumway, G. D. Kohler, and R. Golden. Cyclotron Neutron and X-ray Dosimetry for Animal Irradiation Studies. Radiation Res. 4: 158-173, 1956.

17. A. C. Upton, A. W. Kimball, J. Furth, K. W. Christenberry, and W. H. Benedict. Some Delayed Effects of Atom-Bomb Radiations in Mice. *Cancer Res.* 20: (Part 2): 1-62, 1960.
18. V. J. Rosen, Jr., T. J. Castanera, D. J. Kimeldorf, and D. C. Jones. Pancreatic Islet Cell Tumors and Renal Tumors in the Male Rat Following Neutron Exposure. *Lab. Invest.* 11: 199-203, 1962.
19. J. E. Edwards. Hepatomas in Mice Induced With Carbon Tetrachloride. *J. Nat. Cancer Inst.* 2: 197-199, 1941.
20. D. J. Ingle, and B. L. Baker. Histology and Regenerative Capacity of Liver Following Multiple Partial Hepatectomies. *Proc. Soc. Exper. Biol. Med.* 95: 813-815, 1957.
21. E. A. Smuckler, O. A. Iseri, and E. P. Benditt. An Intracellular Defect in Protein Synthesis Induced By Carbon Tetrachloride. *J. Exper. Med.* 116: 55-72, 1962.
22. S. Leshner, and H. H. Vogel, Jr. A Comparative Histological Study of Duodenal Damage Produced by Fission Neutrons and Co⁶⁰ Gamma-Rays. *Radiation Res.* 9: 560-571, 1958.
23. W. E. Davis, Jr., and L. J. Cole. Comparative Effects of Fast Neutrons and X-rays on Marrow Deoxyribonucleic Acid (DNA) Content in Mice. *Radiation Res.* 14: 104-116, 1961.
24. G. W. Barendsen. Damage to the Reproductive Capacity of Human Cells in Tissue Culture by Ionizing Radiations of Different Linear Energy Transfer in "The Initial Effects of Ionizing Radiations on Cells". ed. R. J. C. Harris. Academic Press, London, 1961.
25. G. W. Barendsen. Dose-Survival Curves of Human Cells in Tissue Culture Irradiated With Alpha-, Beta-, 20-K.V.X- and 200 K.V. X-Radiation. *Nature* 193: 1153-1155, 1962.
26. M. M. Elkind, and H. Sutton. Radiation Response of Mammalian Cells Grown in Culture. I. Repair of X-ray Damage in Surviving Hamster Cells. *Radiation Res.* 13: 556-593, 1960.
27. J. R. Andrews and R. J. Berry. Fast Neutron Irradiation and the Relationship of Radiation Dose and Mammalian Tumor Cell Reproductive Capacity. *Radiation Res.* 16: 76-81, 1962.
28. R. J. Berry and J. R. Andrews. Modification of Radiation Effects in Mammalian Tumor Cells by Pharmacological Agents. *Nature* 196: 185-186, 1962.
29. M. M. Maini, and N. F. Stich. Chromosomes of Tumor Cells. III. Unresponsiveness of Precancerous Hepatic Tissues and Hepatomas to a Mitotic Stimulus. *J. Nat. Cancer Inst.* 28: 753-762, 1962.
30. V. J. Rosen, Jr., and L. J. Cole. Accelerated Induction of Kidney Neoplasms in Mice After X-Radiation (690 rad) and Unilateral Nephrectomy. *J. Nat. Cancer Inst.* 28: 1031-1041, 1962.

DISTRIBUTION

Copies

NAVY

1-3	Chief, Bureau of Ships (Code 335)
4	Chief, Bureau of Ships (Code 320)
5-6	Chief, Bureau of Medicine and Surgery
7	Chief of Naval Operations (Op-07T)
8	Chief of Naval Research (Code 104)
9-11	Director, Naval Research Laboratory (Code 2021)
12	Office of Naval Research (Code 422)
13	Office of Naval Research (Code 441)
14-23	Office of Naval Research, FPO, New York
24-26	Naval Medical Research Institute
27	OIC, Radiation Exposure Evaluation Laboratory
28	Director, Aviation Medical Acceleration Laboratory
29	U.S. Naval Postgraduate School, Monterey
30	Commander, Naval Ordnance Laboratory, Silver Spring
31	Naval Missile Center (Code 5700)
32	U.S. Naval Hospital, San Diego
33	CO, Naval Medical Research Unit No. 2
34	CO, Naval Medical Field Research Laboratory, Camp Lejeune

ARMY

35	Chief of Research and Development (Atomic Division)
36	Chief of Research and Development (Life Science Division)
37	Deputy Chief of Staff for Military Operations (CBR)
38	Chief of Engineers (ENGMC-DE)
39	Chief of Engineers (ENG CW)
40	CG, Army Materiel Command (AMCRD-DE-NE)
41	CG, USA CBA Agency
42	President, Chemical Corps Board
43-45	CO, BW Laboratories
46	CO, Chemical Corps Training Command
47	Commandant, Chemical Corps Schools (Library)
48	CO, Chemical Research and Development Laboratories
49	Commander, Chemical Corps Nuclear Defense Laboratory
50	CO, Army Environmental Hygiene Agency
51	CG, Aberdeen Proving Ground
52	CO, Army Medical Research Laboratory

54 Army Medical Research and Nutrition Laboratory (MEDEN-AD)
 55 CO, Army Medical Service Combat Development Agency
 56-57 Medical Field Service School, Fort Sam Houston (Stimson Lib.)
 58 Brooke Army Medical Center (Dept. Prev. Med.)
 59 Director, Surgical Research Unit, Fort Sam Houston
 60 Director, Walter Reed Army Medical Center
 61 Hq., Army Nuclear Medicine Research Detach., Europe
 62 CG, Combat Developments Command (CDCMR-V)
 63 CG, Quartermaster Res. and Eng. Command
 64 Hq., Dugway Proving Ground
 65-67 The Surgeon General (MEDNE)
 68 Office of the Surgeon General (Combat Dev.)
 69 CG, Engineer Res. and Dev. Laboratory
 70 Director, Office of Special Weapons Development
 71 CG, Munitions Command
 72 CO, Frankford Arsenal
 73 CG, Army Missile Command

AIR FORCE

74 Assistant Chief of Staff, Intelligence (AFCIN-3B)
 75-80 CG, Aeronautical Systems Division (ASAPRD-NS)
 81 CO, Radiological Health Laboratory Division
 82 Director, USAF Project RAND
 83 Commandant, School of Aerospace Medicine, Brooks AFB
 84 CO, School of Aviation Medicine, Gunter AFB
 85 6571st Aeromedical Research Lab., Holloman AFB
 86 Radiobiological Laboratory
 87 Office of the Surgeon (SUP3.1), Strategic Air Command
 88 Office of the Surgeon General
 89 CG, Special Weapons Center, Kirtland AFB
 90 Director, Air University Library, Maxwell AFB
 91-92 Commander, Technical Training Wing, 3415th TTG
 93 Hq., Second Air Force, Barksdale AFB
 94 Commander, Electronic Systems Division (CRZT)

OTHER DOD ACTIVITIES

95 Chief, Defense Atomic Support Agency (Library)
 96 Commander, FC/DASA, Sandia Base (FCDV)
 97 Commander, FC/DASA, Sandia Base (FCTG5, Library)
 98 Commander, FC/DASA, Sandia Base (FCWT)
 99-100 Office of Civil Defense, Washington
 101-102 Civil Defense Unit, Army Library
 103 Armed Forces Institute of Pathology
 104-113 Armed Services Technical Information Agency
 114 Director, Armed Forces Radiobiology Research Institute

AEC ACTIVITIES AND OTHERS

115	Research Analysis Corporation
116	Life Science Officer, AEC, Washington
117	Director, Division of Biology and Medicine
118	NASA, Ames Research Center, Moffett Field
119	Naval Attache, Stockholm (for Commodore Troell)
120	Aerojet General, Azusa
121-125	Argonne Cancer Research Hospital
126-135	Argonne National Laboratory
136-137	Atomic Bomb Casualty Commission
138	AEC Scientific Representative, France
139	AEC Scientific Representative, Japan
140-142	Atomic Energy Commission, Washington
143-146	Atomic Energy of Canada, Limited
147-149	Atomics International
150-151	Battelle Memorial Institute
152	Borden Chemical Company
153-156	Brookhaven National Laboratory
157	Chicago Patent Group
158	Colorado State University
159	Columbia University (Rossi)
160	Committee on the Effects of Atomic Radiation
161-163	Defence Research Member
164-165	duPont Company, Aiken
166	duPont Company, Wilmington
167	Edgerton, Germeshausen and Grier, Inc., Goleta
168	Edgerton, Germeshausen and Grier, Inc., Las Vegas
169-170	General Dynamics, Fort Worth
171-172	General Electric Company, Cincinnati
173-180	General Electric Company, Richland
181	General Electric Company, St. Petersburg
182	General Scientific Corporation
183	Hughes Aircraft Company, Culver City
184	Iowa State University
185	Journal of Nuclear Medicine
186	Knolls Atomic Power Laboratory
187-188	Los Alamos Scientific Laboratory (Library)
189	Lovelace Foundation
190	Martin-Marietta Corporation
191	Massachusetts Institute of Technology
192	Mound Laboratory
193	National Academy of Sciences
194-195	NASA, Scientific and Technical Information Facility
196	National Bureau of Standards (Taylor)
197	National Cancer Institute
198	National Lead Company of Ohio
199	National Library of Medicine
200-201	Nevada Operations Office
202	New Jersey State Department of Health
203	New York Operations Office

204 New York University (Eisenbud)
 205 Oak Ridge Institute of Nuclear Studies
 206 Office of Assistant General Counsel for Patents
 207-208 Phillips Petroleum Company
 209-212 Pratt and Whitney Aircraft Division
 213-214 Public Health Service, Washington
 215 Public Health Service, Las Vegas
 216 Public Health Service, Montgomery
 217 Sandia Corporation, Albuquerque
 218 Union Carbide Nuclear Company (ORGP)
 219-223 Union Carbide Nuclear Company (ORNL)
 224 Union Carbide Nuclear Company (Paducah Plant)
 225 United Nuclear Corporation (NDA)
 226 U.S. Geological Survey, Denver
 227 U.S. Geological Survey, Menlo Park
 228 U.S. Geological Survey, Naval Weapons Plant
 229 U.S. Geological Survey, Washington
 230 U.S. Weather Bureau, Washington
 231 University of California, Davis
 232-234 University of California Lawrence Radiation Lab., Berkeley
 235-236 University of California Lawrence Radiation Lab., Livermore
 237 University of California, Los Angeles
 238 University of California, San Francisco
 239 University of Chicago Radiation Laboratory
 240 University of Hawaii
 241 University of Puerto Rico
 242 University of Rochester (Atomic Energy Project)
 243 University of Tennessee (UTA)
 244 University of Utah
 245 University of Washington (Donaldson)
 246 Wayne State University
 247 Westinghouse Electric Corporation (Mahilly)
 248 Westinghouse Electric Corporation (NASA)
 249 Western Reserve University (Friedell)
 250-274 Technical Information Extension, Oak Ridge

USNADL

275-310 USNADL, Technical Information Division

DISTRIBUTION DATE: 15 January 1962

Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. UNCLASSIFIED Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over) UNCLASSIFIED	1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200.
Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. UNCLASSIFIED Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over) UNCLASSIFIED	1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200.

X rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

X rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

UNCLASSIFIED

UNCLASSIFIED

Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over) UNCLASSIFIED	1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200. UNCLASSIFIED
Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over) UNCLASSIFIED	1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200. UNCLASSIFIED

X rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

X rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

UNCLASSIFIED

UNCLASSIFIED

Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. UNCLASSIFIED Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over)	1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200. UNCLASSIFIED
Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. UNCLASSIFIED Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over)	1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200. UNCLASSIFIED

X-rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

X-rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed.

UNCLASSIFIED

UNCLASSIFIED

Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. UNCLASSIFIED Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over) 1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200. UNCLASSIFIED	Naval Radiological Defense Laboratory USNRDL-TR-596 ACCELERATED INDUCTION OF HEPATOMAS IN FAST NEUTRON IRRADIATED MICE INJECTED WITH CARBON TETRACHLORIDE by L.J. Cole and P.C. Nowell 19 November 1962 19 p. table illus. 30 refs. UNCLASSIFIED Young adult (C57L x A)F₁ hybrid mice received a single whole body exposure to fission spectrum fast neutrons (165-306 rad). Subgroups of these mice then received a single subcutaneous injection of CCl₄, given either at 2, 12, 15, or 18 months post-irradiation. Control nonirradiated mice received a single injection of CCl₄. Other groups of mice were exposed to a single 500 rad dose of 250 KVP (Over) 1. Tumors - Production. 2. Fast neutrons - Biological effects. 3. Liver - Mitosis. I. Cole, L.J. II. Nowell, P.C. III. Title. IV. MR005.08-5200. UNCLASSIFIED
X rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed. UNCLASSIFIED	X rays. The incidence of hepatomas was markedly increased in the neutron-irradiated mice (19%) as compared with that in the X-irradiated mice (2%). In the neutron-irradiated mice injected with CCl₄, the hepatoma incidence attained a value of 51%, three times that in the mice irradiated with neutrons only. In addition, marked pleomorphism and atypicality of liver cell nuclei was evident in almost all of the neutron-CCl₄ mice, but was observed in only 6 of the 47 mice exposed to neutrons only. Of the small group of mice which received CCl₄ but no irradiation, and sacrificed up to 22 months later, none exhibited hepatomas or nuclear abnormalities. In this system therefore, CCl₄ seems to act as a promoting agent in liver carcinogenesis. These findings, taken together with other data in the literature, support the concept that hepatoma induction is accelerated as a consequence of the action of a specific proliferative stimulus on cells bearing a latent radiation-induced alteration. The general question of the role of proliferative stimuli in radiation carcinogenesis is discussed. UNCLASSIFIED