

UNCLASSIFIED

AD NUMBER	
AD379107	
CLASSIFICATION CHANGES	
TO:	UNCLASSIFIED
FROM:	CONFIDENTIAL
LIMITATION CHANGES	
TO: Approved for public release; distribution is unlimited.	
FROM: Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; 31 JAN 1967. Other requests shall be referred to Office of Naval Research, One Liberty Center, 875 North Randolph Street, Arlington, VA 22203-1995.	
AUTHORITY	
ONR ltr dtd 4 May 1977 ONR ltr dtd 4 May 1977	

THIS PAGE IS UNCLASSIFIED

SECURITY

MARKING

The classified or limited status of this report applies to each page, unless otherwise marked.

Separate page printouts MUST be marked accordingly.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE LAWS, TITLE 18, U.S.C., SECTIONS 793 AND 794. THE TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW.

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CONFIDENTIAL

201628

ADVANCED OXIDIZER RESEARCH (U)

SEMIANNUAL REPORT RMD-5043-SA-2

Report Period: 1 July 1966 to 31 December 1966

Office of Naval Research Contract No. NONr 4364(00)

January 31, 1967

GROUP 4
DOWNGRADED AT 3-YEAR INTERVALS;
DECLASSIFIED AFTER 12 YEARS

Thiokol
CHEMICAL CORPORATION
REACTION MOTORS DIVISION
DENVER, NEW JERSEY

CONFIDENTIAL

"This document contains information affecting the National defense of the United States within the meaning of the Espionage Laws, Title 18, U. S. C., Sections 793 and 794. Its transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law."

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

ADVANCED OXIDIZER RESEARCH (U)

SEMIANNUAL REPORT RMD-5043-SA-2

Report Period: 1 July 1966 to 31 December 1966

Office of Naval Research Contract No. NONr 4364(00)

Joseph Dvorak
Archie R. Young, II

Submitted by:

Joseph Green
JOSEPH GREEN, Supervisor
Non-metallic Materials Section

Approved by:

Stanley Tannenbaum
STANLEY TANNENBAUM
Manager, Propellants and
Materials Department

David J. Mann
DAVID J. MANN
Director of Research
and Technical Staff

— i —
CONFIDENTIAL

Report RMD-5043-SA-2

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

FOREWORD

This report was prepared by the Thiokol Chemical Corporation, Reaction Motors Division, Denville, New Jersey, and summarizes work in the area of oxidizer chemistry conducted over the period from July 1, 1966 to December 31, 1966. This research was sponsored by the Office of Naval Research with Mr. Richard L. Hanson serving as Scientific Officer.

Contributors to the work described in this report are as follows:

Dr. Archie R. Young, II, Project Supervisor

Mr. Joseph Dvorak, Principal Investigator

Dr. Josephine Readio, N.M.R. Analysis

Mr. Boyd Fagan, X-ray Analysis

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	1
INTRODUCTION	2
DISCUSSION	3
A. Reactions of Trifluorodiazyl (N_2F_3^+) Salts	3
B. Fluorination Studies	26
EXPERIMENTAL	29
REFERENCES	33

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

ABSTRACT

The existence of trifluorodiazyl ions in BrF_5 and in IF_5 was established by F^{19} n.m.r. and infrared analyses. Solutions of $\text{N}_2\text{F}_3\text{AsF}_6$ in IF_5 treated with perchlorates showed absorptions of N_2F_3^+ and ClO_4^- in the infrared. Attempts to isolate trifluorodiazyl perchlorate from solution were not successful.

An attempt to prepare N_3F_5 by reaction of $\text{N}_2\text{F}_3\text{AsF}_6$ with C_3CNF_2 at -70 resulted in the formation of N_2F_4 and trans- N_2F_2 .

Attempts to prepare new interhalogen oxides by the fluorination of $\text{Cl}_2 \cdot \text{O}_2\text{AsF}_6$ and $\text{NOCl} \cdot \text{O}_2\text{AsF}_6$ were unsuccessful.

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

INTRODUCTION

This research effort is directed toward the synthesis of new inorganic oxidizers with potential application as liquid or solid propellants. The scope of our synthesis research includes the chemistry of nitrogen fluorides, oxygen fluorides, and interhalogen compounds. During this report period emphasis has been placed on attempts to prepare trifluorodiazyl perchlorate ($N_2F_3ClO_4$). Evidence of the coexistence of trifluorodiazyl and perchlorate ions in iodine pentafluoride was established, however, efforts to isolate the salt were not successful.

CONFIDENTIAL


CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION



DISCUSSION

A. Reactions of Trifluorodiazyl ($N_2F_3^+$) Salts

A number of stable salts containing cationic N-F species have been reported in recent years. Salts of N_2F^+ (Refs 1, 2, 3), $N_2F_3^+$ (Refs 4, 5), and ONF_2^+ (Refs 6, 7) are prepared by reactions of covalent nitrogen fluorides with fluorides of Group V elements, as shown in equation 1. Salts of NF_4^+ (Refs 8,



9, 10) are obtained in reactions involving elemental fluorine, as shown in equations 2 and 3. Finally, NH_3F^+ (Ref 4) salts are obtained by reactions of N-fluorocarbamates with strong protonic acids (equation 4).



The fluorammonium (NH_3F^+) cation is the only N-F cation which has been demonstrated to form stable salts with oxygen-containing anions. Our

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

most recent work in this area of N-F chemistry has been directed toward the synthesis of a perchlorate of the trifluorodiazyl (N_2F_3^+) ion.

1. Reaction of $\text{N}_2\text{F}_3\text{AsF}_6$ with Perchlorates

We have previously reported attempts to react $\text{N}_2\text{F}_3\text{AsF}_6$ with perchlorates in hydrogen fluoride (Ref 12), and in sulfur dioxide at subambient temperatures (Ref 13). We found that the addition of soluble perchlorates to solutions of $\text{N}_2\text{F}_3\text{AsF}_6$ in hydrogen fluoride results in the evolution of tetrafluorohydrazine. Similarly, because of a slow reaction of the trifluorodiazyl ion with sulfur dioxide, we obtained NOClO_4 from mixtures of $\text{N}_2\text{F}_3\text{AsF}_6$ and $(\text{CH}_3)_4\text{ClO}_4$ in that solvent.

The behavior of $\text{N}_2\text{F}_3\text{AsF}_6$ in other liquids has now been briefly investigated in the hopes of uncovering a suitable medium for the desired metathesis. The results of these tests are presented in Table I.

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

Table I
N₂F₃AsF₆ Solubility Study

<u>Solvent</u>	<u>Temperature</u>	<u>Results</u>
HSO ₃ F	ambient	Soluble with rapid decomposition
CF ₃ OF	-95°	Insoluble
SF ₄	-78° → -40°	Insoluble
CF ₃ Cl	ambient	Insoluble, slight gassing
(CF ₃) ₂ CO	-28°	Insoluble
BrF ₅	-60° → ambient	Soluble
IF ₅	ambient	Soluble

Confirmation of the existence of trifluorodiazyl ions in BrF₅ and in IF₅ was obtained by F¹⁹ n.m.r. (Figure 1) and liquid phase infrared analyses (Figure 2).

Efforts to achieve an exchange reaction between N₂F₃AsF₆ and perchlorates were resumed using IF₅ as a solvent medium. Because of the low volatility of IF₅, it was convenient to carry out most of the operations involved in these experiments in a glove box.

Equimolar Mixtures of N₂F₃AsF₆ and CsClO₄ in IF₅

When IF₅ solutions containing equimolar quantities of N₂F₃AsF₆ and CsClO₄ were mixed in an open tube (under dry nitrogen in a

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION




Figure 1. A. F^{19} n.m.r. of Trifluorodiazyl Ion in BrF_3 at -60°

B. F^{19} n.m.r. of Trifluorodiazyl Ion in IF_3 at Room Temperature

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION




Figure 2. Infrared Spectrum of $N_2F_3AsF_6$ in IF_5
(NaCl Liquid Cell)

Report RMD 5043-SA-2

- 7 -

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

glove box) vigorous gas evolution was observed. In spite of this indication of decomposition of trifluorodiazyl ions, the infrared spectrum of the residual solution shows a small peak at 920 cm^{-1} , characteristic of N_2F_3^+ , and a very strong perchlorate absorption at $1150\text{--}1040\text{ cm}^{-1}$ (Figure 3). The higher frequency absorptions of N_2F_3^+ (Figure 2) are not apparent in Figure 3, but they may well be masked by the perchlorate and window absorptions. Other features of the spectrum (Figure 3) include an increase (see Figure 2) in the absorption at 2300 cm^{-1} , which is probably due to nitrosonium ion (NO^+), and a new absorption at 1240 cm^{-1} (8.1μ) in the nitrite ion region.

Although there was obviously some loss of N_2F_3^+ in this run, the results were encouraging. The strong concentration of perchlorate in the residual solution seemed to rule out any conclusion that the loss was due to instability of trifluorodiazyl perchlorate.

The equimolar reaction of $\text{N}_2\text{F}_3\text{AsF}_6$ with CsClO_4 in IF_5 was repeated in a closed Kel-F apparatus, in order to determine the composition

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION




Figure 3. Infrared Spectrum of IF₅ Solution Upon Mixing
Equimolar Amounts of N₂F₃AsF₆ and CsClO₄

(NaCl Liquid Cell)

Report RMD 5043-SA-2

- 2 -

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

and approximate quantity of gas liberated on mixing the reagents. In this experiment, one millimole each of the reagents were observed to produce approximately 0.3 millimole of gas. The gas was identified as N_2F_4 with a trace of ClO_3F (Figure 4). Since the N_2F_4 liberated on initial mixing of the reagents was significantly less than the $N_2F_3AsF_6$ charged, there was reason to assume that $N_2F_3^+$ and ClO_4^- ions were still present in solution. The solvent was removed by vacuum distillation at room temperature and the residual solid was examined by infrared analysis (Figure 5). It exhibited only ClO_4^- and AsF_6^- absorptions. One must conclude, therefore, that the residual $N_2F_3^+$ was discharged as N_2F_4 during the distillation of the IF_5 .

Reaction of $CsClO_4$ with Excess $N_2F_3AsF_6$

An IF_5 solution containing approximately 2 millimoles of $N_2F_3AsF_6$ was mixed with a solution of 1 millimole $CsClO_4$. The solutions were mixed in a closed Kel-F apparatus under an atmosphere of nitrogen. A rapid exothermic reaction ensued, and a suspension of needle-like white

CONFIDENTIAL

CONFIDENTIAL




Figure 4. Gaseous Products from Equimolar Reaction
of $\text{N}_2\text{F}_3\text{AsF}_6$ with CsClO_4 in IF_5

Thiokol
REACTION MOTORS DIVISION

Report RMD-5043-SA-2

- 11 -

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION




Figure 5. Solid Residue from IF_5 Solution
of $\text{N}_2\text{F}_3\text{AsF}_6$ and CsClO_4

Report RMD-5043-SA-2

- 12 -

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

crystals appeared. After 24 hours the crystals settled to the top of the solvent, because of the design of the apparatus however, it was not possible to separate the fine crystals from the liquid. A sample of the liquid (24 hours after mixing the reagents) was withdrawn for infrared analysis. As in the case of the equimolar reaction, it showed a strong perchlorate absorption and an N-F absorption at 920 cm^{-1} (Figure 6). After removal of the solvent, an infrared of the residual solid showed no N-F absorptions. Thus, even though a two-fold excess of N_2F_3^+ to perchlorate had been present initially, only perchlorate (and AsF_6^-) was recovered in the residual solid.

Reaction of $\text{N}_2\text{F}_3\text{AsF}_6$ with KClO_4

The reaction of $\text{N}_2\text{F}_3\text{AsF}_6$ with KClO_4 (equimolar) bears mentioning because KClO_4 was observed to be only sparingly soluble in IF_5 , thus the concentration of N_2F_3^+ in solution was initially far greater than that of perchlorate. Indeed, the infrared spectrum of the solution (Figure 7) obtained immediately after the addition of KClO_4 , shows no trace of perchlorate. After

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION




Figure 6. Residual IF₅ Solution. Reaction of CsClO₄
with Excess N₂F₃AsF₆

(NaCl Liquid Cell)

Report RMD-5043-SA-2

- 14 -

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION




Figure 7. Solution Phase. Equimolar Mixture
of KClO_4 and $\text{N}_2\text{F}_3\text{AsF}_6$ in IF_5

Report RMD-5043-SA-2

- 15 -

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

twelve hours, the solution still showed no significant absorption in the perchlorate region (9-10 μ), yet the N_2F_3^+ absorptions at 900 cm^{-1} , 1100 cm^{-1} , 1300 cm^{-1} , and 1500 cm^{-1} had decreased considerably (Figure 8). After two days, a sample of the gas phase was examined by infrared analysis. It contained approximately equivalent amounts of ClO_3F and NF_3 along with a smaller quantity of trans- N_2F_2 (Figure 9).

Although these results suggest the possibility of a reaction between N_2F_3^+ and perchlorate, the residual solid (Figure 10) showed a strong perchlorate absorption.

Summary of Results of IF_5 Studies

The results obtained in our studies of the behavior of IF_5 solutions of the trifluorodiazyl ion in the presence of perchlorate, do not lend themselves to simple interpretation. Infrared evidence was obtained that N_2F_3^+ and ClO_4^- may coexist in solution, yet no N_2F_3^+ salt could be recovered from such solutions. There was also evidence of decomposition of perchlorate

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION





Figure 8. Solution Phase. Equimolar Mixture of KClO_4 and $\text{N}_2\text{F}_3\text{AsF}_6$ in IF_5 After 12 Hours

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION



CONFIDENTIAL

Figure 9. Gaseous Products. Reaction Of
 $\text{N}_2\text{F}_3\text{AsF}_6$ with KClO_4

Report RMD-5043-SA-2

- 18 -

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION




Figure 10. Solid Residue. Equimolar
 $N_2F_3AsF_6-KClO_4$ Reaction

Report RMD-5043-SA-2

- 19 -

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

via the formation of ClO_3F , yet even when an excess of $\text{N}_2\text{F}_3\text{AsF}_6$ was used, most of the perchlorate was recovered. An additional confusing fact is that the trifluorodiazyl ion in some runs was converted to N_2F_4 and in others to a mixture of NF_3 and trans- N_2F_2 . We do not have the experimental data necessary to completely describe the processes that occur in $\text{N}_2\text{F}_3\text{AsF}_6\text{-MClO}_4\text{-IF}_5$ systems. We are fairly certain, however, that the IF_5 plays an active role via a reaction with perchlorate ions. For this reason we plan to investigate the behavior of tetrafluoroperchlorate (ClF_4^-) salts with solutions of $\text{N}_2\text{F}_3\text{AsF}_6$ in IF_5 .


CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

2. Reactions of $N_2F_3Sb_2F_{11}$ with Perchlorates

An alternate approach to the synthesis of trifluorodiazyl perchlorate involves the use of the low melting ($74-78^\circ$) tetrafluorohydrazine-antimony pentafluoride adduct (Ref 4) as a solvent and as a source of the trifluorodiazyl cation (equation 5).



Compatibility of $N_2F_3Sb_2F_{11}$ with $CsClO_4$

When approximately equimolar amounts of $N_2F_3Sb_2F_{11}$ and $CsClO_4$ were ground together in an agate mortar, no visual indication of reaction was observed. However, a more recently prepared sample of $N_2F_3Sb_2F_{11}$ did deflagrate upon mixing with $CsClO_4$. For reasons to be discussed later, this result is believed to be anomalous. Additional confirmation that the materials are compatible is being sought.

Compatibility of $N_2F_3Sb_2F_{11}$ with $CsClO_4$ at $90^\circ C$

When a mixture of $N_2F_3Sb_2F_{11}$ and $CsClO_4$ was heated to $90^\circ C$,

CONFIDENTIAL

CONFIDENTIAL**Thiokol**
REACTION MOTORS DIVISION

well above the melting point of the antimony complex, a small amount of gas was liberated which contained ClO_3F and NF_3 . However, the bulk of the solid material charged was recovered and the presence of ClO_4^- , SbF_6^- and N_2F_3^+ was established by infrared analysis. The results of X-ray analysis of the residual solid mixture cannot be interpreted unambiguously (Table II). The diffraction patterns of $\text{N}_2\text{F}_3\text{Sb}_2\text{F}_{11}$ and CsSbF_6 are quite similar and one might expect the same to hold true for $\text{N}_2\text{F}_3\text{ClO}_4$ and CsClO_4 .

Table II

Comparison of Diffraction Patterns (Principal Lines)

$\text{N}_2\text{F}_3\text{Sb}_2\text{F}_{11}$		$\text{CsSbF}_6 (\text{CsSb}_2\text{F}_{11}?)$		CsClO_4		Reaction Product	
d, Å	I/I ₀	d, Å	I/I ₀	d, Å	I/I ₀	d, Å	I/I ₀
4.65	60	----	----	----	----	----	----
3.89	100	3.95	100	----	----	3.90	100
----	----	----	----	3.81	75	3.76	15
3.56	100	3.53	70	----	----	3.49	100
----	----	----	----	3.42	100	3.38	20
----	----	----	----	2.32	85	2.30	30
2.18	1	2.19	35	2.14	2	2.17	100

However, extraction of the reaction product at low temperatures with SO_2

yielded an insoluble solid which showed only SbF_6^- and ClO_4^- in the infrared.

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

A small unidentified SO_2 soluble fraction was presumed to be unreacted

$\text{N}_2\text{F}_3\text{Sb}_2\text{F}_{11}$.

In a similar experiment CsClO_4 was added to molten $\text{N}_2\text{F}_3\text{Sb}_2\text{F}_{11}$ at 80°C . Upon addition of the CsClO_4 , a vigorous reaction ensued, accompanied by a pressure surge. Although the evolved gases were not completely trapped, a portion was identified by infrared analysis as NF_3 and ClO_3F . The solid residue has an infrared absorption attributable to SbF_6^- ($\text{Sb}_2\text{F}_{11}?$), and is apparently pure CsSbF_6 or $\text{CsSb}_2\text{F}_{11}$ (equation 6).

$$\text{N}_2\text{F}_3\text{Sb}_2\text{F}_{11} + \text{CsClO}_4 \xrightarrow{?} [\text{N}_2\text{F}_3\text{ClO}_4] \longrightarrow \text{NF}_3 + \text{ClO}_3\text{F} + \text{CsSb}_2\text{F}_{11} \quad (6)$$


Since $\text{N}_2\text{F}_3\text{Sb}_2\text{F}_{11}$ and CsClO_4 react vigorously at elevated temperatures, an attempt was made to prepare $\text{N}_2\text{F}_3\text{ClO}_4$ by suspending CsClO_4 in SbF_3 and subsequently adding N_2F_4 . In this manner a white amorphous solid (I) was obtained which contained N_2F_3^+ , SbF_6^- and ClO_4^- by infrared. Heating a portion of I to 80°C resulted in the liberation of NF_3 and ClO_3F , and left a residue of CsSbF_6 . Extraction of I with SO_2 at low temperatures resulted in the

CONFIDENTIAL

CONFIDENTIAL


Thiokol
REACTION MOTORS DIVISION

isolation of $N_2F_3Sb_2F_{11}$. The insoluble fraction contained ClO_4^- and SbF_6^- absorptions in the infrared. This indicates that the reaction product (I) is simply a mixture of $N_2F_3Sb_2F_{11}$ and $CsClO_4$ (equation 7). It is also additional evidence that the two salts are compatible at room temperature.



Compatibility of $N_2F_3Sb_2F_{11}$ with $CsClO_4$ by
Differential Thermal Analysis

Since $N_2F_3Sb_2F_{11}$ and $CsClO_4$ appear to be compatible at room temperature but extremely reactive at elevated temperatures, an attempt is currently being made to determine by differential thermal analysis whether the desired perchlorate is being formed at some intermediate temperature and subsequently decomposing (equation 8). It was during the initiation of these



studies that difficulty was encountered in preparing mixtures of $N_2F_3Sb_2F_{11}$ and $CsClO_4$. When freshly prepared $N_2F_3Sb_2F_{11}$ was mixed with $CsClO_4$ on an agate

CONFIDENTIAL


CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

mortar deflagration occurred. Pre-grinding the materials and then mixing them resulted in a delayed ignition. Although the infrared spectrum of the $N_2F_3Sb_2F_{11}$ is consistent with prior samples, the salt appeared to undergo several transformations near room temperature in the differential thermal analysis apparatus. Since these results appear to be anomalous we believe that the difficulty is associated with one particular sample of $N_2F_3Sb_2F_{11}$ and plan to pursue this investigation further, after preparing a new supply of $N_2F_3Sb_2F_{11}$.

3. Reaction of $N_2F_3AsF_6$ with ϕ_3CNF_2

An attempt was made to isolate N_3F_5 via the reaction of $N_2F_3AsF_6$ with ϕ_3CNF_2 (equation 9). Since the solid reactants deflagrate upon



contact, ϕ_3CNF_2 was added to a solution of $N_2F_3AsF_6$ in liquid SO_2 at $-70^\circ C$.


An immediate reaction took place and a gaseous product was obtained which contained N_2F_4 and possibly N_2F_2 . Removal of the solvent at $-50^\circ C$ gave a solid

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

residue which was shown to be Cl_3CAsF_6 . The reaction probably proceeds as shown in equation 10.



B. Fluorination Studies

In addition to pursuing the synthesis of new solid oxidizers from the Group V pentafluoride adducts of simple N-F compounds, we have initiated a study of the fluorination and oxyfluorination of chlorine-containing compounds with the view of obtaining new interhalogen compounds. Thus we have investigated the fluorination of Cl_2NF as a possible route to F_3ClNF and the fluorination of " $\text{Cl}_2 \cdot \text{OAsF}_6$ " as a route to interhalogen oxides (Ref 13). Additional work was done in this area during this report period.

1. Reaction of " $\text{Cl}_2 \cdot \text{O}_2\text{AsF}_6$ " with Fluorine

Chlorine forms an uncharacterized purple complex with O_2AsF_6 which is stable at -78°C (Ref 14). An investigation of the fluorination of this

CONFIDENTIAL


CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

complex had been initiated with the purpose of preparing new interhalogen oxides. In a previously reported experiment (Ref 13), the fluorination of $\text{Cl}_2 \cdot \text{O}_2\text{AsF}_6$ produced a mixture of gaseous products including Cl_2 , O_2 , ClO_2 , and an unidentified minor product. The reaction has since been repeated at a higher fluorine pressure in an attempt to increase the amount of the minor constituent for identification. However, the only gaseous products obtained were O_2 , Cl_2 , ClO_2 and ClO_3F .

2. Reaction of NOCl with O_2AsF_6

The possibility of preparing a complex between NOCl and O_2AsF_6 was investigated, also with the view of preparing new interhalogen compounds (equations 11, 12).



In an initial experiment NOCl was condensed on O_2AsF_6 and the reaction mixture was allowed to warm to room temperature. No complex

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

was formed and the reactants were substantially recovered with only a trace of NO_3F being formed. When the reaction was repeated and the reagents brought in contact for a longer period of time, only two thirds of the NOCl charged was recovered and a non-condensable gas, presumably oxygen, was liberated. Upon hydrolysis, the white solid product gave a positive test for NO_2^- indicating the presence of NO^+ . Since the overall reaction appears to be the formation of NOAsF_6 with the liberation of O_2 rather than the formation of the desired complex, we do not intend to pursue this reaction further.

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

EXPERIMENTAL

Reaction of $N_2F_3Sb_2F_{11}$ and $CsClO_4$

A Kel-F reactor was charged with a mixture of 0.296 g (0.4 mmole) of $N_2F_3Sb_2F_{11}$ and 0.696 g (0.3 mmole) of $CsClO_4$ and heated to 70°C when a yellow gas was evolved. The reactor was pumped on and heating continued to 90°C. The evolved gas was found to contain ClO_3F and NF_3 . The infrared spectrum of the residual solid (0.83 g) showed the presence of $N_2F_3^+$, ClO_4^- and SbF_6^- . Extraction of the reaction product at low temperatures with SO_2 yielded an insoluble solid which showed only SbF_6^- and ClO_4^- in the infrared. A small SO_2 soluble fraction was presumed to be unreacted $N_2F_3Sb_2F_{11}$.

In a similar experiment 0.7559 (1.4 mmole) of $N_2F_3Sb_2F_{11}$ was heated to 80°C and then 0.696 g (0.3 mmole) of $CsClO_4$ was added. A vigorous reaction ensued with the evolution of NF_3 and ClO_3F . The residual solid has an infrared absorption attributable to SbF_6^- and is apparently pure $CsSbF_6$.

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

Reaction of $N_2F_3Sb_2F_{11}$ with $CsClO_4$ in Excess SbF_5

To a suspension of 0.696 g (13 mmole) of $CsClO_4$ in 2 ml of SbF_5 was added N_2F_4 . The pressure of N_2F_4 in the system was kept below 100 mm. The reaction mixture turned yellow and finally orange. After standing overnight the reaction mixture was pumped on in vacuo at $40^\circ C$ for several hours to yield an amorphous solid (I) which contained $N_2F_3^+$, SbF_6^- and ClO_4^- by infrared analysis. Extraction of I with SO_2 resulted in the isolation of $N_2F_3Sb_2F_{11}$. The insoluble fraction contained SbF_6^- and ClO_4^- absorptions in the infrared.

Reaction of $N_2F_3AsF_6$ with ϕ_3CNF_2

To 0.274 g (1 mmole) of $N_2F_3AsF_6$ in approximately 10 ml of SO_2 at $-70^\circ C$ was added 0.295 g (1 mmole) of ϕ_3CNF_2 . The reaction mixture immediately turned brown and then yellow. The reaction mixture was pumped on at $-70^\circ C$ and N_2F_4 was collected in a $-196^\circ C$ trap. The reactor was then allowed to warm slowly to room temperature while being pumped on. The

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

materials collected in this way contained SO_2 and N_2F_4 . The residual solid was identified as O_3CAsF_6 .

Reaction of NOCl with O_2AsF_6

To 0.276 g (0.8 mmole) of O_2AsF_6 at -196°C was added 1.75 mmole of NOCl . The reaction mixture was allowed to warm slowly to room temperature with stirring. The NOCl was recondensed on the solid several times. Finally the reaction mixture was stirred at -6°C for an hour. No evidence for the formation of a complex was obtained. In addition to recovering two thirds of the NOCl , a non-condensable gas, presumably O_2 was liberated. The residual solid upon hydrolysis gave a positive test for NO_2^- indicating the presence of NO^+ . The overall reaction appears to be the formation of NOAsF_6 and oxygen.

IF_5 Studies

The IF_5 used in the attempts to achieve a metathesis between $\text{N}_2\text{F}_3\text{AsF}_6$ and perchlorates was obtained from Mathieson Co. The commercial material contains colored impurities (I_2 and lower fluorides) which were removed by

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

bubbling fluorine through the liquid at room temperature. The colorless IF_5 was then distilled in an all glass apparatus. The distillate was stored under dry nitrogen in a Kel-F tube having a tightly fitting Teflon cap. $\text{N}_2\text{F}_3\text{AsF}_6$ dissolves readily in IF_5 , treated as described above, with no evidence of gassing.

All of the operations performed in the studies of the behavior of IF_5 solutions of $\text{N}_2\text{F}_3\text{AsF}_6$ with perchlorates were carried out in a dry nitrogen glove box, with the exceptions of gas collection and solvent distillation procedures.

A Teflon-glass syringe was used to introduce samples of IF_5 into NaCl liquid cells for infrared analysis. The NaCl cell was attacked somewhat by the IF_5 solutions, giving rise to absorptions at 4.7μ , $5.8-6.5\mu$, and 8.4μ . In spite of this, it was possible to identify the presence of N_2F_3^+ in the infrared even after repeated use of the cell.

CONFIDENTIAL

CONFIDENTIAL

Thiokol
REACTION MOTORS DIVISION

REFERENCES

1. D. Moy and A. R. Young, II, J. Am. Chem. Soc., 87, 1889 (1965).
2. American Oil Company, Fifth Quarterly Report, Contract No. DA-31-124-ARO(D)-78, May 1964.
3. H. Roesky, O. Glemser, D. Bormann, Chem. Ber., 99, 1589 (1966).
4. J. Ruff, Inorg. Chem., 5, 1791 (1966).
5. A. R. Young, II and D. Moy, Inorg. Chem., 6, 178 (1967).
6. N. Bartlett and J. Passmore, Chem. Comm., 213 (1966).
7. W. Fox, et al., J. Am. Chem. Soc., 88, 2604 (1966).
8. J. Guertin, K. Christe, and A. Pavlath, Inorg. Chem., 5, 1921 (1966).
9. W. Tolberg, et al., Inorg. Nucl. Chem. Letters, 2, 79 (1966).
10. E. Lawless and G. Hennon, Midwest Research Institute, Report AFRPL-TR-66-227, October, 1966, Contract AF04(611)-10921.
11. Aerojet-General Corporation, Report 3132, October, 1965, Contract NOmr 2655(00).
12. Thiokol Chemical Corporation, Reaction Motors Division, Report RMD-5043-65F, January, 1966, Contract NOmr 4364(00).
13. Thiokol Chemical Corporation, Reaction Motors Division, Semiannual Report RMD-5043-66, June, 1966, Contract NOmr 4364(00).
14. Thiokol Chemical Corporation, Reaction Motors Division, Report RMD-5043-64F, January, 1965, Contract NOmr 4364(00).

CONFIDENTIAL

CONFIDENTIAL

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) THIokol CHEMICAL CORPORATION Reaction Motors Division Denville, New Jersey		2a. REPORT SECURITY CLASSIFICATION Confidential
		2b. GROUP IV
3. REPORT TITLE ADVANCED OXIDIZER RESEARCH (U)		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Semiannual Report - July 1, 1966 to December 31, 1966.		
5. AUTHOR(S) (Last name, first name, initial) Young, A. R., Dvorak, J.		
6. REPORT DATE 31 January 1967	7a. TOTAL NO. OF PAGES 33	7b. NO. OF REFS 14
8a. CONTRACT OR GRANT NO. NONr 4364(00)	9a. ORIGINATOR'S REPORT NUMBER(S) RMD-5043-SA-2	
8b. PROJECT NO. 5043 (0101000406)-022	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. AVAILABILITY/LIMITATION NOTICES		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Department of the Navy Office of Naval Research Washington, D.C.	
13. ABSTRACT The existence of trifluorodiazyl ions in BrF_5 and in IF_5 was established by F^{19} n.m.r. and infrared analyses. Solutions of $\text{N}_2\text{F}_3\text{AsF}_6$ in IF_5 treated with perchlorates showed absorptions of N_2F_3^+ and ClO_4^- in the infrared. Attempts to isolate trifluorodiazyl perchlorate from solution were not successful. An attempt to prepare N_3F_5 by reaction of $\text{N}_2\text{F}_3\text{AsF}_6$ with O_3CNF_2 at -70° resulted in the formation of N_2F_4 and <u>trans</u>-N_2F_2. Attempts to prepare new interhalogen oxides by the fluorination of $\text{Cl}_2\cdot\text{O}_2\text{AsF}_6$ and $\text{NOCl}\cdot\text{O}_2\text{AsF}_6$ were unsuccessful.		

DD FORM 1 JAN 64 1473

CONFIDENTIAL

Security Classification

CONFIDENTIAL

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Liquid Oxidizers						
Solid Oxidizers						
NF Chemistry						
Trifluorodiazyl Salts						
Trifluorodiazyl Perchlorate						
Fluorination						
Halogen Oxides						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task numbers, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

CONFIDENTIAL

Security Classification