

UNCLASSIFIED

AD NUMBER
AD401273
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Foreign Government Information; 04 SEP 1961. Other requests shall be referred to US Library of Congress, Attn: Aerospace Technology Division, Washington, DC.
AUTHORITY
ATD ltr, 2 Dec 1965

THIS PAGE IS UNCLASSIFIED

401 273

5 17511

P/045/62/021/005/001/009
I069/I269

ASTIA FILE COPY

STEP
AUTHOR:

8 Rubinowicz, A.

TITLE:

6 On a source-free tensor field, and its tensor potential, when associated with an electromagnetic field

PERIODICAL:

15 Acta physica polonica, v.21; no.5, 1962 pp. 451-468

TEXT:

For incident divergence of convergent spherical waves the diffraction wave is derivable from a source-free vector field \underline{V} . This is uniquely determined by the solution of the wave equation describing the incident wave. The vector potential \underline{V}' for the field \underline{V} can be established by a simple geometric method (A. Rubinowicz, Acta phys. Polon., 21, 61 (1962)). For electromagnetic fields \underline{V} and \underline{W} being themselves vector fields, have to be replaced by tensorial

Card 1/3

Ed

On a source-free tensor field...

fields and potentials. The Helmholtz-Huygens principle in Kirchhoff's electromagnetic theory was formulated by H.A. Lorentz and Larmor (A. Rubinowicz, Die Beugungswelle in der Kirchhoffschen Theorie der Beugung, Warsaw, 1957, p.238) and also differently by F. Kottler (Ann. Phys. (Leipzig), (4) 71, 457 (1923)). However, both formulations, if applied to the Maxwell equations, are equivalent and both can be expressed in terms of two tensor fields \hat{V}_e and \hat{V}_m (carets denote tensors of the second rank in a three-dimensional space), which determine the electric and magnetic fields. These, in turn, are uniquely determined by a given solution of the Maxwell equations. These vector fields having vanishing divergence are derivable from tensorial potentials thereby providing two separate formulations of the electromagnetic Helmholtz-Huygens principle. These are of importance for Kirchhoff's theory of diffraction when isolated for arbitrary incidence of the light waves. An integral theorem for tensorial fields is applied

Card 2/3

analogous to Stoke's theorem for vectorial fields. Utilising the Larmor and Kottler formulations of the Helmholtz-Huygens principle, the tensorial potentials for arbitrary electromagnetic fields is derived geometrically. The author similarly derived the Miyamoto-Wolf vector potentials associated with scalar wave equations.

ASSOCIATION: Polish Academy of Sciences (Polish Academy of Science, Warsaw)

SUBMITTED: September 4, 1961