

UNCLASSIFIED

AD **401 564**

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

ASTIA FILE COPY

401 564

5 17511

7

S/185/62/007/010/004/020
D234/D308

24,2600
STEP

AUTHORS: 8 Brodovyy, V. A. and Lyashenko, V. I.

TITLE: 6 Temperature dependence of the kinetics of photoconductivity of single crystals of Sb_2S_3

PERIODICAL: 15 Ukrayins'kyy fizychnyy zhurnal, v. 7, 10, 1962.
pp. 1062-1066

TEXT: The temperature range was -100° to $+100^{\circ}C$. The kinetics of photoconductivity were studied by analyzing the decrease of photocurrent after illuminating the samples by single rectangular pulses of white light. There are two groups of specimens with different behavior, which is illustrated by graphs taken from two specimens, the resistance of specimen 1 being about 10^9 ohm.cm and that of specimen 2 about 10^8 ohm.cm. Each component of the photocurrent is considered separately. Heating of specimen 1 leads to more intense poly-molecular recombination. At $68^{\circ}C$ the decrease of the first component of the photocurrent is exponential, and so is that of the second component at $44^{\circ}C$. Decrease of temperature only di-

Card 1/2

Temperature dependence f ...

S/185/62/007/010/004/020
D234/D308

minishes the intensity of the first component, which practically disappears at $-35^{\circ}C$. In specimen 2 there is no change of recombination mechanism with temperature increase up to $80^{\circ}C$. The stationary photoconductivity of the specimens has a maximum about $-10^{\circ}C$ and a minimum about $35^{\circ}C$. There are 4 figures.

ASSOCIATION: Kyivsk'yy universytet im. T. H. Shevchenka (Kiev University im. T. H. Shevchenko)

SUBMITTED: March 13, 1962

Card 2/2