

**UNCLASSIFIED**

**AD** **401 569**

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

401 569

(3) 17511

u

STEP  
3.2310

11492  
S/033/62/039/005/005/011  
E032/E514

ASTIA FILE COPY

AUTHOR: (8) Getmantsev, G.G.

TITLE: (6) On the heating of interstellar gas by cosmic rays

PERIODICAL: (15) Astronomicheskiy zhurnal, v.39, no.5, 1962, 856-861

TEXT: It is argued that the heating of interstellar gas due to ionization losses can only be effective at relatively high temperatures of the medium and can hardly be responsible for an increase in the temperature if the initial temperature is much less than  $10^4$  deg. This paper is concerned with a second possible mechanism which is effective provided the temperature is not too high. It is assumed that a cosmic-ray stream moves along randomly oriented magnetic lines of force. Owing to the high conductivity of the interstellar gas, the magnetic field due to the cosmic-ray stream cannot be produced in a short interval of time and the stream is almost completely compensated by electron currents in the interstellar plasma. The gas should then become heated owing to Joule losses. It is shown that these losses are given by  $W_2^* = 3 mc^2 N^2 / T^{3/2}$ , where  $N$  is the cosmic-ray density,  $T$  is the temperature and  $m$  is the mass of the cosmic-ray particle.  
Card 1/3

4

On the heating of interstellar ...

S/033/62/039/005/005/011  
E032/E514

ray particles. In deriving this expression it is assumed that the particle velocity is  $v_2 = c/3$  and the conductivity of the interstellar gas is  $\sigma \sim e^2 T^{3/2} / 27 m$ . These losses are then shown to be greater than the rate of emission of energy through recombinational processes so that the temperature of the gas should increase and this increase will continue until the two effects just balance each other. Detailed examination of this type of effect leads to the conclusion that cosmic rays cannot in fact ensure any appreciable heating of interstellar gas in those regions of the galaxy where the gas concentration is large ( $n \sim 0.1-1 \text{ cm}^{-3}$ ) unless it is assumed that there are cosmic rays with  $E/mc^2 \sim 0.1-0.01$  and  $N \sim 10^{-4} - 10^{-5} \text{ cm}^{-3}$ . If, on the other hand, the gas density in the halo is of the order of  $10^{-3} - 5 \times 10^{-4} \text{ cm}^{-3}$  or if there are regions in which the concentration is of this magnitude, then primary cosmic rays might be able to heat the gas. If these conditions are satisfied and if the sources of cosmic rays ensure that the concentration of cosmic electrons with energies of  $10^4 - 10^5 \text{ eV}$  is of the order of  $3 \times 10^{-8} \text{ cm}^{-3}$ , then an interstellar medium with  $n \sim 10^{-3} - 5 \times 10^{-4} \text{ cm}^{-3}$

4

Card 2/3

On the heating of interstellar ...

S/O33/62/039/005/005/011  
EO32/E514

may become heated to a limiting temperature of a few million deg. If in addition the initial temperature is low, then in the initial stage of the heating process the latter should be due to the "induction" mechanism mentioned above and the temperature should rise to about  $10^5$  deg in a time interval of the order of  $10^{12}$ - $10^{13}$  sec.

ASSOCIATION: Radiofizicheskiy institut Gor'kovskogo gos. universiteta imeni N. I. Lobachevskogo  
(Radiophysics Institute of the Gor'kiy State University imeni N. I. Lobachevskiy)

SUBMITTED: May 5, 1961 (initially)  
October 9, 1961 (after revision)

Card 3/3