

63-3-3

403011

403 011

**Photodissociation of Molecules
For Excitation of an
Optical Maser**

by

J. R. Singer

CATALOGED BY ASTIA
AS AD NO.

Series No. 60, Issue No. 429

Contract No. AF 49(638)-102

January 8, 1962

ELECTRONICS RESEARCH LABORATORY

UNIVERSITY OF CALIFORNIA

BERKELEY, CALIFORNIA

"Qualified requesters may obtain copies
of this report from ASTIA"

AFOSR 2251

**Electronics Research Laboratory
University of California
Berkeley, California**

**PHOTODISSOCIATION OF MOLECULES FOR EXCITATION
OF AN OPTICAL MASER**

by

J. R. Singer

**Institute of Engineering Research
Series No. 60, Issue No. 429**

**Physics Division
Air Force Office of Scientific Research
Contract No. AF 49(638)-102
Division File No. 13-25-J**

January 8, 1962



One of the most interesting aspects of optical masers is the fact that a very narrow bandwidth wave is obtainable in the infrared and visible wavelengths. The bandwidth of the output signal $\Delta \nu$ is related to the molecular bandwidth $\Delta \nu_a$ by the formula¹

$$\Delta \nu = \frac{4\pi kT(\Delta \nu_a)^2}{P_a} \quad (1)$$

where k is Boltzmann's constant, T is the absolute temperature, and P_a is the power emitted by the atomic transitions. It is advantageous to obtain a very narrow emission source for a number of reasons; perhaps the major reason is that the peak intensity of the source is enhanced, and the coherence is also usually increased. Intense coherent sources are of interest to provide a new means of studying molecular bonding, harmonic excitation of molecular vibrations, and they may be utilized as communication channel carriers in addition to a large number of other possibilities.

The best method of obtaining a very narrow optical emission source is to eliminate doppler broadening by observing atomic transitions perpendicular to the path of an atomic beam. This method can be adapted to optical masers and we will discuss our approach to this problem.

The system which we are presently investigating involves the photodissociation of molecules in which one dissociated atom appears in an excited state. The general process may be described by



where $h\nu_0$ is a high energy photon interacting with a molecular compound AB composed of atoms A and B . The dissociation of AB leaves A in an excited state and B in the ground state or in a metastable state. KE is the kinetic energy available to conserve energy in the reaction. Following the dissociation, A^* decays to the ground state with the emission of a photon $h\nu_1$ which is lower in energy than $h\nu_0$.

There is the advantage here that the excitation occurs by a band absorption, thus an incoherent radiation source may be used efficiently. On the other hand, the dissociative process has a threshold, and the

exciting light frequencies exceeding the threshold provide energy which reappears as kinetic energy of the dissociated atoms. This kinetic energy is not desirable because it increases doppler broadening of the emission.

If a static system of molecular vapor is dissociated by radiation, the process is necessarily pulsed with a very long recovery time since the dissociated atoms A and B will not, in general, recombine rapidly. The most appropriate system, therefore, is a molecular beam configuration.

If the lifetime of the excited atomic state occurring after dissociation is sufficiently long, the optical maser system configuration may consist of a molecular beam passing through an illuminated region where the molecule is dissociated; the excited atoms then pass into a region surrounded by a Fabry-Perot etalon (or a confocal spherical optical resonator) where coherent radiation could take place. This system required atomic lifetimes of at least 10^{-5} seconds since molecular beam velocities are of the order of 10^5 cm/sec. It is, however, more usual to observe lifetimes of 10^{-8} seconds.

With the shorter atomic lifetimes, the excitation by dissociation must take place within the optically resonant structure. Also, shorter lifetimes imply more difficulty in obtaining continuous operation since the ground state atoms may accumulate faster than they can be swept out of the active region of the maser. That is, the molecular beam velocity may be too slow relative to the radiative lifetime to permit the inverted population distribution to exist for an extended time. The factors leading to a restoration of the Boltzmann distribution are spontaneous emission, and the expectation that atoms attaining the ground state will not be re-excited by the UV pumping radiation, but rather will be acting as absorbers of the coherent output.

Some molecules which are candidates for a molecular beam optical maser are listed in Table 1.

While it would be desirable to select a transition with a long lifetime because then excited atoms could be accumulated, however, the transitions listed here are strong transitions with consequent short-lived states. The substance which we are presently investigating is RbI. The advantages of this material is that the excitation source is easily obtained in the form of the $2537 \text{ \AA}$ line of a mercury lamp, and that the emission is conveniently in the near infrared.

The proposed experimental setup is shown in Fig. 1. The general plan is to photodissociate the molecules within an optically resonant structure and have the excited atoms decay with the emission of coherent light because of the positive regeneration of the structure.

As mentioned previously, it is desirable to have a very narrow atomic linewidth. The major broadening mechanism is the doppler shift, hence one wants a molecular effusion process from the oven rather than turbulent flow. The requirement for a molecular effusion is from a thin walled aperture³

$$\omega \sim L$$

where ω is the slit width, and L is the mean free path within the oven. Assuming 10. mm of Hg pressure within the oven, (.03 cm for the oven mean free path), the number of molecules effusing per second is given by

$$I \approx 10^{22} p A_s / (MT)^{1/2}$$

where p is the pressure in mm of Hg, A_s is the area of the source, M is the molecular weight, and T is the absolute temperature (approximately 1000°K for RbI). Using a meter long oven, one obtains a beam intensity of approximately 10^{21} molecules/second. While this beam is not well collimated it enters the emissive region essentially without loss of molecules because of the closeness and breadth of the optically resonant region.

Assuming reflecting areas of 4 cm^2 separated by a meter for the optically resonant region, the number of molecules from the beam within the region is about 10^{16} molecules.

We now ask is this sufficient to satisfy the oscillation condition,⁴

$$N_{\min} > \frac{3hV \Delta \nu}{4\pi^2 \mu^2 Q}$$

and observe that $N_{\min}$ is about 10^{14} to 10^{15} molecules, hence the oscillation condition should be satisfied providing the cross-section for the photodissociation is sufficiently large. Sufficient leeway is available so that dissociation need not occur in all of the molecules, but we would hope to dissociate at least ten percent of the molecules within the resonant region. Previous work has demonstrated that photodissociation of RbI is easily observed,^{5,6} but little quantitative data on cross-sections is available.

TABLE I

<u>Molecule</u>	Metal atom emission		Metal Ion Emission
	<u>λ output</u>	Minimum <u>excitation λ</u>	<u>Transition</u>
LiI	6708A	2350A	$2^2P_{1/2} \rightarrow 2^2S_{1/2}$
NaI	5896	2450	$3^2P_{1/2} \rightarrow 3^2S_{1/2}$
NaI	3303	1854	$3^2P_{3/2} \rightarrow 3^2S_{1/2}$
RbI	7947	2650	$5^2P_{1/2} \rightarrow 5^2S_{1/2}$
RbI	7800	2650	$5^2P_{3/2} \rightarrow 5^2S_{1/2}$
RbI	4216	2080	$6^2P_{1/2} \rightarrow 5^2S_{1/2}$
CsI	4555	2085	$7^2P_{3/2} \rightarrow 6^2S_{1/2}$
LiBr	6708	2040	$2^2P_{1/2} \rightarrow 2^2S_{1/2}$
NaBr	5896	2144	$3^2P_{1/2} \rightarrow 3^2S_{1/2}$
TlI	5351	2080	$7^2S_{1/2} \rightarrow 6^2P_{3/2}$
TlBr	5351	1915	$7^2S_{1/2} \rightarrow 6^2P_{3/2}$
TlCl	5351	1850	$7^2S_{1/2} \rightarrow 6^2P_{3/2}$

(RbI oven temperature is approximately 700°C)




Figure 1. Schematic plan of molecular beam photodissociation type optical maser




Figure 2. Experimental arrangement for measurement of the RbI photo-dissociation cross-section

REFERENCES

1. Gordon, J. P. , Zeiger, H. J. and Townes, C. H. , Phys. Rev. , Vol. 99 (1955), p. 1264; Phys. Rev. , Vol. 95 (1954), p. 284.
2. Franken, P. , Hill, A. E. , Peters, C. W. and Weinreich, G. , Phys. Rev. Letters , Vol. 7 (1961), p. 118.
3. A general discussion of molecular effusion is given in Chapter 2 of Ramsey, N. F. , Molecular Beams, Oxford Univ. Press, 1956.
4. Schawlow, A. L. and Townes, C. H. , "Optical Masers," Phys. Rev. , Vol. 112 (1958), p. 1940.
5. Visser, G. H. and Van Heel, A. C. S. , Uber die Einwirkung des lichtetes auf Rubidiumjodiddampf, Z. fur Phys. , Vol. 70 (1931), p. 605; Vol. 86 (1933), p. 694.
6. Desai, M. S. , Uber die einwirkung des Lichtes auf Rubidiumjodiddampf, Z. fur Phys. , Vol. 85 (1933), p. 360.

**AIR FORCE OFFICE OF SCIENTIFIC RESEARCH
FLUOROTUBE RESEARCH
CONTRACT NO. AF 49(630)-102**

DISTRIBUTION LIST

ADDRESSEE	NO COPIES	ADDRESSEE	NO COPIES	ADDRESSEE	NO COPIES
AFOSR ATTN: Technical Library Washington 25, D. C.	4	High Speed Flight Station (NASA) ATTN: Technical Library Edwards AFB, California	1	The Ohio State University Antenna Laboratory 1024 Neil Avenue Columbus 10, Ohio	1
AFOSR (SRPP) Washington 25, D. C.	1	Langley Research Center (NASA) ATTN: Technical Library Langley AFB, California	1	Major General Casemiro Montenegro Filho Central Technico de Aeronautica (CTA) São José dos Campos São Paulo, Brazil	1
ASD ATTN: Technical Library Wright-Patterson AFB, Ohio	1	Lewis Research Center (NASA) ATTN: Technical Library 21000 Brookpark Road Cleveland 15, Ohio	1	Purdue University Lafayette, Indiana ATTN: Richard L. Funkhouser Engineering Librarian	1
AFRL ATTN: Technical Library 1. G. Hanscom Field Bedford, Massachusetts	1	Goddard Space Flight Center (NASA) ATTN: Technical Library Greenbelt, Maryland	1	Professor J. Van Bladel Electrical Engineering Department College of Engineering University of Wisconsin Madison 6, Wisconsin	1
EOAR The Shell Building 47 Rue Carterstein Brussels, Belgium	(Unclassified Reports)	George C. Marshall Space Flight Center (NASA) ATTN: Technical Library Redstone Arsenal, Alabama	1	Professor W. Low Department of Physics The Hebrew University of Jerusalem Jerusalem, Israel	1
ARL ATTN: Technical Library Wright-Patterson AFB, Ohio	1	Wallops Station (NASA) ATTN: Technical Library Wallops Island, Virginia	1	University of Illinois Department of Electrical Engineering Urbana, Illinois ATTN: H. Von Kester	1
ASTIA (ITPCR) Arlington Hall Station Arlington 12, Virginia	10	Institute of Aeronautical Sciences 2 East 64th Street New York 21, New York	(Unclassified Reports)	Ohio State University Department of Electrical Engineering Columbus, Ohio	1
Director of Research Headquarters, USAF ATTN: AFRDR Washington 25, D. C.	1	Applied Mechanics Reviews Southwest Research Institute 8000 Culebra Road San Antonio 6, Texas	(Unclassified Reports)	The University of Michigan Department of Electrical Engineering Flintston Physics Laboratory Ann Arbor, Michigan ATTN: Professor J. E. Rowe	1
Office of Naval Research Department of the Navy ATTN: Code 420 Washington 25, D. C.	1	Linda Hall Library ATTN: Document Division 605 Cherry Street Kansas City 10, Missouri	(Unclassified Reports)	Stanford University Electronics Research Laboratory Stanford, California	1
Naval Research Laboratory ATTN: Technical Library Washington 25, D. C.	1	AFOSR (SRAL) ATTN: Technical Library Holloman AFB, New Mexico	2	Massachusetts Institute of Technology Research Lab. of Electronics Room 20N-221, Document Office Cambridge 38, Massachusetts ATTN: J. H. Hewitt	1
Chief, Research and Development ATTN: Scientific Information Branch Department of the Army Washington 25, D. C.	1	AFSWC (SWOI) Kirtland AFB, New Mexico	1	Harvard University Cruft Laboratory Cambridge 38, Massachusetts ATTN: Technical Reports Collection	1
Chief, Physics Branch Division of Research, U. S. Atomic Energy Commission Washington 25, D. C.	1	Advanced Research Projects Agency Washington 25, D. C.	1	Technical Information Libraries Bell Telephone Laboratories, Inc. Whippany Laboratory Whippany, New Jersey ATTN: Technical Reports Librarian	1
U. S. Atomic Energy Commission Technical Information Extension P. O. Box 82 Oak Ridge, Tennessee	1	Rand Corporation 1700 Main Street Santa Monica, California	1	California Institute of Technology Pasadena 4, California ATTN: R. Gould	1
National Bureau of Standards ATTN: Technical Library Room 203, Northwest Building Washington 25, D. C.	1	Chairman Canadian Joint Staff (DRR/DRI) 2450 Massachusetts Avenue, N. W. Washington 25, D. C.	(Unclassified Reports)	Sylvania Electric Company Mountain View, California ATTN: D. H. Goodman	1
Physics Program National Science Foundation Washington 25, D. C.	1	Officer in Charge Office of Naval Research Navy No. 100 Fleet Post Office New York, New York	1	Professor William H. Surder Electrical Engineering Department Princeton University Princeton, New Jersey	1
Director, Army Research Office, Durham Box CM, Duke Station Durham, North Carolina	1	Office of Technical Services Department of Commerce Technical Reports Branch Washington 25, D. C.	1	General Electric Company Electron Tube Div. of the Research Lab The Knolls Schenectady, New York ATTN: E. D. McArthur	1
AEDC (AEOM) ATTN: Technical Library Arnold Air Force Station Tullahoma, Tennessee	1	M. D. Adcock, Head American Systems Incorporated 1625 E. 126 Street Hawthorne, California	1	General Electric Company Missile and Space Vehicle Department 3198 Chestnut Street Philadelphia 4, Pennsylvania ATTN: Aerodynamic Engineering Operation R. F. Perch, Manager	1
AFTTC (AFOTL) ATTN: Technical Library Edwards AFB, California	1	Dr. Harold Glaser Office of Naval Research Washington 25, D. C.	1	Hughes Aircraft Company Florence at Tule Street Culver City, California ATTN: Documents Group, Bldg 6, Rm. X2015	1
AFMDC (MDF) Holloman AFB, New Mexico	1	Professor Charles Townes Department of Physics Columbia University New York 27, New York	1	ACA Laboratories Princeton, New Jersey ATTN: Dr. W. M. Webster, Director Electronics Research Lab	1
AFMDC (NDO) Holloman AFB, New Mexico	1	Professor Harvey Brooks Department of Physics Harvard University Cambridge 38, Massachusetts	1	Varian Associates 611 Hansen Way Palo Alto, California ATTN: Technical Library	1
ARGMA (ORDXR-OTL) Redstone Arsenal, Alabama	1	Professor P. Kusch Department of Physics Columbia University New York 27, New York	1	Westinghouse Electric Corporation Electronic Tube Division P. O. Box 184 Elmira, New York ATTN: Mr. Sheldon S. King, Librarian	1
Institute of Technology (AU) Library MCL-LIB, Bldg. 125, Area B Wright-Patterson AFB, Ohio	1	Professor N. Bloembergen Department of Physics Harvard University Cambridge 38, Massachusetts	1	Professor Zohrab Kaprielian University of Southern California School of Engineering Department of Electrical Engineering University Park Los Angeles 7, California	1
AFSC (SCRE) Andrews AFB Washington 25, D. C.	1	Dr. Irving Rowe Office of Naval Research 346 Broadway New York, New York	1	B. J. Maxam Delmo-Victor Research Laboratories Belmont, California	1
Signal Corps Engineering Laboratory ATTN: (SCF)MEL-RPO Fort Monmouth, New Jersey	1	W. A. Kosamplich, Manager Lockheed Aircraft Corporation Missiles and Space Division Technical Information Center 3251 Manover Street Palo Alto, California	1	Headquarters Air Force Office of Scientific Research Office of Aerospace Research USAF Washington 25, D. C. ATTN: Marshall C. Harrington General Physics Division	1
Headquarters National Aeronautics and Space Administration ATTN: Technical Library Washington 25, D. C.	1	Mr. E. Ohress Sperry Gyroscope Company Electron Tube Division Mail Station 1, 240 Great Neck, New York	1		
Ames Research Center (NASA) ATTN: Technical Library Moffett Field, California	1	DOFL (ORDTL 012) Washington 25, D. C.	1		
Technical Information Division Bldg. 30, Room 101 Radiation Laboratory Berkeley, California	1	Prof. J. H. Mulligan, Jr. Chairman, Department of Electrical Engineering New York University 25 Waverly Place New York, New York	1		
Hans Motz Oxford University Oxford, England	1				
Miss May Dorin Department of Archives 303 Library University of California Berkeley 4, California	1				