

UNCLASSIFIED

AD

403 884

*Reproduced
by the*

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-34

CATALOGED
AS AD NO. 403884

OFFICE OF NAVAL RESEARCH

Contract Monr-1677(01)

Project No. NR-051-362

TECHNICAL REPORT NO. 7

Continuous Representations in the Statistical
Theory of Electronic Energies

by

Sidney Golden
Department of Chemistry
Brandeis University

and

George Handler
Department of Chemistry
Tufts University

MAY 17 1963

TJIA A

April 15, 1963

403884

**Continuous Representations in the Statistical
Theory of Electronic Energies**

by

**Sidney Golden
Department of Chemistry
Brandeis University**

and

**George Handler
Department of Chemistry
Tufts University**

Abstract

The use of continuous bases of representation other than a plane-wave basis is considered in the theory developed originally by Thomas and Fermi. The bases considered here are the sets of eigenfunctions of Hamiltonians corresponding to a particle subjected to a field of force which varies inversely as the cube of the distance to some fixed point. Variation of the strength of the interaction varies the basis continuously. Calculations of the energy of the hydrogen atom is carried out, with results that are appreciably closer to the quantum mechanical result than is obtained with the original Thomas-Fermi theory. Numerical results suggest the possible existence of a statistical analogue of the Rayleigh-Ritz variation principle.

1. Introduction

The statistical theory originally introduced by Thomas⁽¹⁾ and Fermi⁽²⁾ for describing the behavior of many-electron systems, had its origin in a quasi-classical statistical mechanical viewpoint which incorporated the Pauli exclusion principle. Subsequent investigations dealing with this theory have provided it with a better relation to quantum mechanical theory than it had originally. As a result, several modifications of the original Thomas-Fermi theory exist which incorporate features of a distinctly quantum mechanical character.⁽³⁾ Of all of these, the "inhomogeneity correction" of von Weizsacker⁽⁴⁾ has the most significant corrective influence upon the results obtained with the use of the statistical theory. Although this addition to the Thomas-Fermi theory was introduced originally in an ad hoc manner,⁽⁵⁾ its presence, with slight modification from the original, seems assured from generalized treatments of the statistical method.⁽⁶⁾

However, in spite of any improvements which have been made on the original Thomas-Fermi theory, there remains one glaring deficiency in the theory: the density of electrons is either singular in the vicinity of a nucleus⁽⁷⁾ or, on the basis of somewhat arbitrary auxiliary conditions, is set equal to zero.⁽⁸⁾ It is to this aspect of the theory that the present paper is directed. The use of the momentum basis, traditionally associated with Thomas-Fermi theory, is a consequence of particular partitioning of the Hamiltonian operator. Here we address ourselves to the problem of alternative partitionings, preserving the spirit of the original theory by limiting ourselves to those partitionings that lead to continuous bases.

2. Formal Theory

In terms of a general formulation of the statistical theory based upon the use of the density matrix of many-electron systems,⁽⁹⁾ the original Thomas-Fermi density may be derived entirely from the assumption that the kinetic energy and potential energy of a particle are commutable observables. This assumption, in fact, is basically the one made by both Thomas and Fermi. Such an assumption is illustrative of a certain type which generates a set of quasi-classical approximations.⁽¹⁰⁾ For these approximations the Hamiltonian operator of the system is divided into two parts, the choice of the parts being a matter of convenience. The assumption that the two parts commute immediately produces an approximation to the density matrix which is the generalized analogue of the Thomas-Fermi theory.

In the interest of brevity, the formalism of (I) will be employed here. The density matrix (representative) of a many-particle system then may be expressed as⁽¹⁰⁾

$$\rho_N(x_1, x_2) = \sum_{n=1}^{\infty} \left\{ \theta(\lambda_N - \underline{H}) \psi_n(x_1) \right\}^* \left\{ \theta(\lambda_N - \underline{H}) \psi_n(x_2) \right\}, \quad (2.1)$$

where $\underline{H}$ is the Hamiltonian operator of the system, $\{\psi_n(x)\}$ is any properly symmetrized (i.e., with respect to exchange of identical particles) complete orthonormal set of functions of the correct boundary conditions. x stands for the entire set of configurational and spin coordinates of the particles, and λ_N is a parameter determined by

$$N = \text{Tr } \rho_N = \int dx \rho_N(x, x) = \sum_{n=1}^{\infty} \langle \psi_n | \theta(\lambda_N - \underline{H}) | \psi_n \rangle. \quad (2.2)$$

The spectral operator $\theta(\lambda_N - \underline{H})$ can be expressed most simply in terms of its matrix elements which are evaluated in a basis of eigenfunctions of $\underline{H}$. Then, we have for such a set $\{\varphi_n(x)\}$ which satisfies

$$\langle \varphi_m | \underline{H} | \varphi_n \rangle = E_n \delta_{m,n} \quad (2.3)$$

the matrix elements

$$\langle \varphi_m | \theta(\lambda - \underline{H}) | \varphi_n \rangle = \theta(\lambda - E_n) \delta_{m,n} \quad (2.4)$$

with

$$\left. \begin{aligned} \theta(y) &= 1, y > 0, \\ \theta(y) &= 0, y < 0. \end{aligned} \right\} \quad (2.5)$$

In these terms the "density" of the many-particle system in its configuration and spin space is given by $\rho_M(x, x)$, but corresponds to the density associated with the M lowest energy eigenstates of the system.⁽¹¹⁾

The procedure to be exploited here involves making approximations to the Laplace transform of the density matrix defined in Eq. (2.1). For convenience, we may deal formally with the spectral operator and obtain⁽¹²⁾

$$\mathcal{L}\{\theta(\lambda - \underline{H})\} = \int_0^\infty d\lambda e^{-\underline{z}\lambda} \theta(\lambda - \underline{H}), \quad \text{Re}(\underline{z}) > 0 \quad (2.6)$$

$$= \frac{e^{-\underline{z}\underline{H}}}{\underline{z}}. \quad (2.7)$$

As a consequence, a representation for the spectral operator is obtainable in terms of the inverse Laplace transform, viz.,

$$\theta(\lambda - \underline{H}) = \frac{1}{2\pi i} \mathcal{P} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{d\underline{z}}{\underline{z}} e^{\underline{z}\lambda} e^{-\underline{z}\underline{H}}, \quad \text{Re}(\gamma) > 0. \quad (2.8)$$

Approximations which may be made for $\exp(-\underline{z}\underline{H})$ give rise to approximations to $\theta(\lambda - \underline{H})$.

In particular, the quasi-classical approximation to which reference has been made arises as follows. Let

$$\underline{H} = \underline{H}_0 + \underline{H}_1(x) \quad (2.9)$$

and set

$$\exp(-\beta \underline{H}) = \exp(-\beta \underline{H}_1) \exp(-\beta \underline{H}_0). \quad (2.10)$$

If we work directly in terms of a complete orthonormal basis of properly symmetrized eigenfunctions of $\underline{H}_0$, say $\{\varphi_n^0(x)\}$, which satisfy

$$\langle \varphi_m^0 | \underline{H}_0 | \varphi_n^0 \rangle = E_n^0 \delta_{m,n} \quad (2.11)$$

we obtain the quasi-classical approximation to the density matrix

$$\begin{aligned} [\rho_M(x_1, x_2)]_{q.c.} = & \sum_{n=1}^{\infty} \left\{ \theta(\lambda_M - E_n^0 - \underline{H}_1(x_1)) \varphi_n^0(x_1) \right\}^* \\ & \times \left\{ \theta(\lambda_M - E_n^0 - \underline{H}_1(x_2)) \varphi_n^0(x_2) \right\}. \end{aligned} \quad (2.12)$$

The preceding analysis is general and can be taken over in a single-particle approximation for a many-electron system. In such an approximation, $\underline{H}_0$ is an appropriate single-electron Hamiltonian and $\underline{H}_1$ is a suitable interaction potential. The quantity M is taken equal to the number of single-electron eigenstates which, on account of spin degeneracy, is usually equated to half the number of electrons of the system.⁽¹³⁾ In the Thomas-Fermi theory $\underline{H}_0$ is identified with the kinetic energy operator, $\{\varphi_n^0(x)\}$ is a free particle basis and $\underline{H}_1(x)$ is a self-consistent potential which is determined from Poisson's equation and certain physically plausible boundary conditions upon the potential. Regarding the latter, the condition that the potential has singularities corresponding to an attractive Coulomb field at any point immediately leads to a singularity there in the density. By contrast, it will be noted, quantum mechanical solutions yield no such behavior for the density. As a result, the choice of a basis will be examined with the object of alleviating this difficulty in a quasi-classical approximation.

3. The Basis Functions

Because the object of the present paper is mainly constructive rather than analytical, no extensive analysis will be made of the seat of the aforementioned disparity in the density behavior. However, we may take note of the fact that for the systems generally of interest the binding energy is a finite quantity. This may be stated in more quantum mechanical terms as corresponding to a lowest finite eigenvalue of the Hamiltonian of the system. When measured relative to its lowest value, the eigenvalues of the Hamiltonian are non-negative as is, thus, the (modified) Hamiltonian itself. However, in the quasi-classical approximation which leads to the Thomas-Fermi theory, the Hamiltonian of a single particle is represented as the sum of two operators: the first is the kinetic energy and is non-negative while the second is the potential energy of the electron and is non-positive and can become unbounded for systems of interest. As a consequence, while the Laplace transform of the spectral distribution operator, Eq. (2.7), is a bounded operator in general, the approximation employed in Eq. (2.10) does not conform to this behavior.

Now, regardless of the choice of a non-negative $\underline{H}_0$ in Eq. (2.9), it is apparent from Eq. (2.12) that any negative singularity in $\underline{H}_1$ at, say, $\vec{r}_K$,⁽¹⁴⁾ yields a divergent value for the density. To avoid this behavior, $\underline{H}_1$ must be chosen also to be bounded from below (i.e., $> -\infty$). For systems which involve only electrons and nuclei, the interaction between the particles involves only Coulomb terms which exhibit singularities that are simple poles. In any approximation of a self-consistent sort, we may suppose that no singularity of greater order need be introduced. Hence, specializing to a single-particle approximation, we wish to examine the possibility of finding $\underline{H}_0$ and $\underline{H}_1$ in Eq. (2.9) such that both are bounded from below. That is, letting

$$\underline{H} = \underline{T} + \underline{V} , \quad (3.1)$$

where $\underline{T}$ is the kinetic energy operator for an electron and $\underline{V}$ its potential energy function depending only upon its configurational coordinates, we seek a potential function such that

$$\underline{H}_0 = \underline{T} + \underline{V}_0 \quad (3.2)$$

$$\underline{H}_1 = \underline{V} - \underline{V}_0$$

are both bounded from below. Since $\underline{V}$ has no singularities other than simple poles, it is apparent that if, in the vicinity of the singularity $\vec{r}_K$,

$$V_0 \sim \frac{c}{|\vec{r} - \vec{r}_K|^s} , \quad s > 1 \quad (3.3)$$

with c a positive quantity, $\underline{H}_1$ will be bounded from below. Similarly, if $\underline{V}_0$ has the behavior exhibited in Eq. (3.3) with $s < 2$, $\underline{H}_0$ will also be bounded from below.⁽¹⁵⁾ It is evident that a range of choices is possible for both c and s for the purpose at hand. Hence, considerations of convenience may be included in any final choice of $\underline{V}_0$.

In order to maintain some similarity to the original Thomas-Fermi theory, it is appropriate to deal with an $\underline{H}_0$ which admits only a continuum of positive eigenvalues. Such eigenvalues correspond to states for which the particle is unbound and is reminiscent of the free-particle basis of the Thomas-Fermi theory. At the same time it is desirable to have a $\underline{V}_0$ for which the minimum value of $\underline{H}_1$ has as small a magnitude as possible. Because the potential energy function $\underline{V}$ is to be kept general and unspecified, except for its singularities, a condition leading to fulfillment of this desire is impossible to formulate explicitly. Nevertheless, a certain choice of c and s in Eq. (3.3) will adequately fulfill the requirements which have been mentioned.

Consider the central field Hamiltonian

$$\underline{H}_0 = -\frac{\hbar^2}{2m} \nabla^2 - \frac{\hbar^2}{8m} \frac{c}{r^2} \quad (3.4)$$

The value of c in Eq. (3.4) may range from zero to unity. The latter value is the maximum which an attractive inverse square potential may have if the ground state of the system is to be unbound. Any slight increase in the value of the parameters yields a ground state energy value which corresponds to a "fall" of the particle to the origin,⁽¹⁵⁾ i.e., a negative infinite value of the energy. Thus, the system corresponding to $c = 1$ in the Hamiltonian of Eq. (3.4) is the limit of unbounded systems having an attractive center varying inversely as some power of the distance from that center.⁽¹⁶⁾ With $c = 0$, the free-particle basis results.

Each Schrodinger equation corresponding to Eq. (3.4) generates a complete, orthogonal and continuous set of functions which normalize accordingly. Because of spherical symmetry of $\underline{H}_0$, the Schrodinger equation is separable in terms of spherical polar coordinates. Hence, if

$$\underline{H}_0 \psi_{K\ell m} = E_{K\ell m} \psi_{K\ell m} \quad (3.5)$$

and

$$\psi_{K\ell m} = f_{K\ell}(r) Y_{\ell m}(\theta, \varphi), \quad |m| \leq \ell = 0, 1, 2, \dots, \quad (3.6)$$

where $Y_{\ell m}(\theta, \varphi)$ is a normalized spherical harmonic of the Eulerian angles θ and φ ⁽¹⁷⁾, we obtain the radial equation

$$\frac{d^2 f_{K\ell}(r)}{dr^2} + \frac{2}{r} \frac{df_{K\ell}(r)}{dr} + \left(K^2 - \frac{\ell(\ell+1)}{r^2}\right) f_{K\ell}(r) = 0 \quad (3.7)$$

with

$$K^2 = \frac{2m}{\hbar^2} E_{K\ell m} \quad (3.8)$$

and

$$\gamma_\ell = \ell(\ell + 1) - c/4 \quad (3.9)$$

With the transformation

$$R = Kr \quad (3.10)$$

we obtain

$$\frac{d^2 [R f_\ell(R)]}{dR^2} + \left(1 - \frac{\gamma_\ell}{R^2}\right) [R f_\ell(R)] = 0 \quad (3.11)$$

which is Bessel's equations. Its solutions yield

$$f_\ell(R) = R^{-1/2} J_{u_\ell}(R), \quad (3.12)$$

$$\text{with} \quad u_\ell = \pm \sqrt{(\ell + \frac{1}{2})^2 - c/4} \quad (3.13)$$

and $J_u(x)$ is a Bessel function⁽¹⁸⁾ of the first kind of order u .

For each $\ell \neq 0$ there is one value of u_ℓ which is positive and one which is negative. The latter correspond to solutions which diverge at the origin, while the former do not. Moreover, they diverge so strongly that the integral of the square of the eigenfunction over any finite volume enclosing the origin also diverges. Hence, they comprise unacceptable eigenfunctions. Thus, we shall take the positive sign in Eq. (3.13). For $c \neq 0$ the eigenfunction corresponding to $\ell = 0$ also diverges at the origin. However, the integral of its square over any finite volume enclosing it does not, so we retain it.⁽¹⁹⁾

Since the Bessel functions in Eq. (3.12) are not quadratically integrable, we examine the integral⁽²⁰⁾

$$\begin{aligned} H_a(K_1, K_2) &= \int_0^\infty dr r^2 e^{-a^2 r^2} \frac{J_u(K_1 r) J_u(K_2 r)}{\sqrt{K_1 K_2} r} \\ &= \frac{1}{2a^2 \sqrt{K_1 K_2}} e^{-\frac{K_1^2 + K_2^2}{4a^2}} I_u\left(\frac{K_1 K_2}{2a^2}\right), \end{aligned}$$

where $I_u(x)$ is a Bessel function of the first kind of purely imaginary argument. From the asymptotic properties of the latter, ⁽²¹⁾ the dominant term is given by

$$I_u\left(\frac{K_1 K_2}{2a}\right) \sim \left(\frac{a^2}{\pi K_1 K_2}\right)^{\frac{1}{2}} e^{-\frac{K_1 K_2}{2a^2}}$$

so that

$$\lim_{a^2 \rightarrow 0} N_a(K_1, K_2) = \frac{1}{2K_1 K_2} \left[\lim_{a^2 \rightarrow 0} e^{-\frac{(K_1 - K_2)^2}{8a^2}} \frac{1}{(\pi a^2)^{\frac{1}{2}}} \right]. \quad (3.14)$$

The term in brackets is a well-known representation of a delta function. As a consequence,

$$\int_0^\infty dK_1 K_1^2 \lim_{a^2 \rightarrow 0} N_a(K_1, K_2) = 1, \quad K_2 \neq 0, \quad (3.15)$$

and the original functions

$$\psi_{K\ell m} = \frac{1}{(Kr)^{\frac{1}{2}}} J_{u_2}(Kr) Y_{\ell, m}(\theta, \varphi) \quad (3.16)$$

are properly normalized eigenfunctions of Eq. (3.5) in the sense of the integration of Eq. (3.16) which, together with Eq. (3.7), preserves the similarity with an integration over momenta as in the Thomas-Fermi basis.

4. The Quasi-Classical Density Matrix

The quasi-classical single-particle approximation to the density matrix now may be expressed in terms of the basis which has been described. Making use of Eqs. (2.12), (3.4), (3.8), (3.15) and (3.16), we obtain

$$\begin{aligned}
 [\rho_M(\vec{r}_1, \vec{r}_2)]_{f.c.} = & \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} \int_0^{\infty} dK K^2 \left\{ \theta\left(\lambda_M - \frac{\pi^2}{2m} K^2 - \frac{\pi^2}{8m} \frac{c}{r_1^2} - V(r_1)\right) \right. \\
 & \times \frac{J_{u_\ell}(Kr_1)}{(Kr_1)^{\frac{1}{2}}} Y_{\ell m}(\theta_1, \varphi_1) \Big\} \\
 & \times \left\{ \theta\left(\lambda_M - \frac{\pi^2}{2m} K^2 - \frac{\pi^2}{8m} \frac{c}{r_2^2} - V(r_2)\right) \frac{J_{u_\ell}(Kr_2)}{(Kr_2)^{\frac{1}{2}}} Y_{\ell m}(\theta_2, \varphi_2) \right\} \Bigg].
 \end{aligned} \quad (4.1)$$

With the aid of the Addition Theorem of Legendre polynomials, ⁽²²⁾ the previous equation reduces to

$$[\rho_M(\vec{r}_1, \vec{r}_2)]_{f.c.} = \frac{1}{4\pi} \sum_{\ell=0}^{\infty} (2\ell + 1) P_\ell\left(\frac{\vec{r}_1 \cdot \vec{r}_2}{r_1 r_2}\right) \int_0^{\infty} dK K \theta_1 \theta_2 \frac{J_{u_\ell}(Kr_1) J_{u_\ell}(Kr_2)}{(r_1 r_2)^{\frac{1}{2}}}, \quad (4.2)$$

where

$$\theta_j = \theta\left(\lambda_M - \frac{\pi^2 K^2}{2m} - \frac{\pi^2}{8m} \frac{c}{r_j^2} - V(r_j)\right).$$

The expression in Eq. (4.2) can be reduced further. However, we shall only do so for the diagonal elements of the density matrix, which correspond to the density. Making use of the properties of the θ -function, we obtain

$$\rho_M = \frac{1}{4\pi r} \sum_{\ell=0}^{\infty} (2\ell + 1) \int_0^{K_M} dK K J_{u_\ell}^2(Kr) \quad (4.3)$$

where

$$K_M^2 = \frac{2m}{\pi^2} (\lambda_M - V(r)) - c/4r^2 \geq 0. \quad (4.4)$$

Apart from the difference in the orders of the Bessel functions, the presence of the inverse square potential and a neglect of spin multiplicity, Eq. (4.3) is formally identical with the corresponding expression obtained in a

free-particle basis.⁽²³⁾ Thus the terms corresponding to each value of l represent the contribution to the density made by particles having a square angular momentum of $l(l+1)\hbar^2$.

The approximation to the density given in Eq. (4.3) can be seen to exhibit no divergence at the origin providing $V(r)$ exhibits no divergence as strong as $1/r^2$. In cases of atomic systems, in which Coulomb forces apply, this condition is satisfied. Because K_M vanishes then at some non-zero value of r the density vanishes for all smaller values. While this behavior is clearly unsatisfactory, it possibly represents a better approximation to a finite density than is afforded by the original Thomas-Fermi theory. For atomic systems we may anticipate that λ_M is negative. Thereupon, an asymptotic vanishing of $V(r)$ as $r \rightarrow \infty$ indicates the existence of a finite distance beyond which the density vanishes identically. Thus, the density given by Eq. (4.3) may be anticipated to have non-zero values in a limited region of space.⁽²⁴⁾

5. The Energy

The approximation which has been given for the quasi-classical density matrix enables one to obtain an approximation for the energy. The simplest way to do so is with the aid of the relation⁽²⁵⁾

$$\langle E \rangle_M = \int_0^M \lambda_M dM = \lambda_M M - \int_{\lambda_0}^{\lambda_M} M d\lambda, \quad (5.1)$$

where $\langle E \rangle_M$ is the sum of the first M eigenvalues. Since the quasi-classical approximation, Eq. (4.2), involves λ_M through the spectral operator, we need to examine

$$\int_{\lambda_0}^{\lambda_M} \theta \left(\lambda - \frac{\hbar^2 k^2}{2m} - \frac{\hbar^2}{8m} \frac{c}{r^2} - V(r) \right) d\lambda = \left\{ \lambda_M - \frac{\hbar^2 k^2}{2m} - \frac{\hbar^2}{8m} \frac{c}{r^2} - V(r) \right\} \\ \times \theta \left(\lambda_M - \frac{\hbar^2 k^2}{2m} - \frac{\hbar^2}{8m} \frac{c}{r^2} - V(r) \right).$$

When combined with Eqs. (4.2), (4.3) and (5.1), there results the approximation

$$\langle E \rangle_M = \int_0^\infty dr r \left[\sum_{\ell=0}^\infty (2\ell+1) \left\{ \frac{\hbar^2}{8m} \frac{c}{r^2} + V(r) \right\} \int_0^{K_M} dK K J_{u_\ell}^2(Kr) \right] \\ + \int_0^\infty dr r \left[\sum_{\ell=0}^\infty (2\ell+1) \frac{\hbar^2}{2m} \int_0^{K_M} dK K^3 J_{u_\ell}^2(Kr) \right]. \quad (5.2)$$

The last integral is the approximate expectation value of H_0 given by Eq. (3.4) for the first M eigenvalues of H . Both integrals can be reduced considerably with the aid of the Schafheitlin reduction formula for Bessel functions. (26)

The present approximation gives no indication of the reliability of the energy values, there being no extremum principle available to serve in this connection. However, since the eigenvalues satisfy the relation

$$\langle E^2 \rangle_1 = \langle E \rangle_1^2 \\ \langle E^2 \rangle_2 = \langle E \rangle_1^2 + \{ \langle E \rangle_2 - \langle E \rangle_1 \}^2,$$

some estimate of the precision of the approximate energy values can be made. For this purpose, we must have an expression for the sum of the squares of the first M eigenvalues. We have

$$\langle E^2 \rangle_M = \int_0^M \lambda_M^2 dM = \lambda_M^2 M - 2 \int_{\lambda_0}^{\lambda_M} \lambda_M d\lambda \quad (5.3)$$

Making use of Eq. (4.2) and carrying out the requisite operations, we obtain

$$\begin{aligned}
 \langle E^2 \rangle_M = & \int_0^\infty dr \, r \left[\sum_{\ell=0}^\infty (2\ell+1) \left\{ \frac{\hbar^2}{8m} \frac{c}{r^2} + V(r) \right\}^2 \int_0^{K_M} dK \, K \, J_{u_\ell}^2(Kr) \right] \\
 & - \int_0^\infty dr \, r \left[\sum_{\ell=0}^\infty (2\ell+1) \frac{\hbar^2}{8m} \left\{ \frac{\hbar^2}{8m} \frac{c}{r^2} + V(r) \right\} \int_0^{K_M} dK \, K^3 \, J_{u_\ell}^2(Kr) \right] \\
 & + \int_0^\infty dr \, r \left[\sum_{\ell=0}^\infty (2\ell+1) \frac{\hbar^4}{4m^2} \int_0^{K_M} dK \, K^5 \, J_{u_\ell}^2(Kr) \right]. \quad (5.4)
 \end{aligned}$$

With Eqs. (5.1) and (5.4) we are able to estimate the precision of the approximate energy values. Thus, for the lowest eigenvalue, $M = 1$ and $\Delta = \langle E^2 \rangle_1 - \langle E \rangle_1^2$ is a non-negative quality which vanishes for an eigenvalue. In the present approximation, Δ is a function of c , the strength of the inverse square potential. Hence, we may anticipate that some value of c may possibly render Δ a minimum. For such a value, the most precise value of $\langle E \rangle_1$ is obtained within the framework of the present theory.

6. Computational Techniques

The integration indicated in (4.3) may be performed explicitly, giving

$$\rho_M(\vec{r}) = \frac{r^2}{8\pi^3} \sum_{\ell=0}^\infty (2\ell+1) \left[J_{u_\ell}^2(X) - J_{u_\ell+1}(X) J_{u_\ell-1}(X) \right], \quad (6.1)$$

$X = K_M r,$

where K_M is given by (4.4) and u_ℓ by (3.14). This expression is convenient for computational purposes. Obvious alterations give density expressions for the various energy terms considered in the last section.

The definition of density thus derived was applied to the problem of the hydrogen atom, for which $V(r) = -e^2/r$. An IBM 7090 Data Processing System was programmed to evaluate (6.1), together with (4.4) and (3.14), for various values of λ_M and c . In all cases, summations over l were extended until further contribution was less than 1 part in 10^8 ($l \sim 7$). From the densities thus tabulated, normalization and energy integrals were evaluated, using Simpson's Rule. In each case, the expression under consideration was tabulated at a sufficient number of points so that further increase caused no significant change in the result of the integration. A typical mesh density was 50 points/Bohr radius. The energy expressions were evaluated and integrated in such a way as to allow the use of some of them as expectation values of inverse powers of r .

Several sets of calculations were performed. The equation $M(\lambda_M, c) = M_0$ was solved for λ_M at various values of c for $M_0 = 1$ and $M_0 = 5$, corresponding respectively to a filled 1s level, and filled 1s, 2s, and 2p levels, and energies and expectation values of powers of r were found for these values of λ_M and c . In addition, points on the curve $M(\lambda_M, c)$ vs. λ_M for $c = 0.5$ were found, so as to allow evaluation of

$$\int_0^{M_0} \lambda_M dM = \lambda_M M_0 - \int_{-\infty}^{\lambda_0} M d\lambda \quad (6.2)$$

for $M_0 = 1$, where λ_0 is given by $M(\lambda_0, c) = 1$, the integration again being by Simpson's Rule. This constitutes a check on the numerical procedures, as the energy found in this way must agree with that found from the integration of the energy density at λ_0 and c . The number of points on this curve, 40, was not sufficient to give a result comparable in precision to the integration of the energy density, these calculations being used mainly to demonstrate the internal consistency of the results.

7. Numerical Results and Discussion

Table IA presents the results of calculations of the energy for $M = 1$. The most interesting feature of these results is the variation of $\langle E \rangle$ with c . As can be seen, $\langle E \rangle$ has a definite, though broad, maximum rather close to the quantum-mechanical value, suggesting the possible existence of a statistical analogue of the Rayleigh-Ritz variational principle. Similar behavior can be seen in Table IB, which gives results for $M = 5$. These quantities are shown graphically in Fig. 1a and 1b. Table IA also gives $\Delta = \langle E^2 \rangle_1 - \langle E \rangle_1^2$, while a plot of Δ vs. c is shown in Fig. 2. There exists a value for c for which Δ is a minimum, though this does not coincide with the "best" value.⁽²⁷⁾ Also noteworthy is the great improvement in the energy over the value from the Fermi-Amaldi theory, with which these results are directly comparable, since the theories become identical for $c = 0$. Table IB also presents expectation values of various powers of r together with the Quantum Mechanical results and Fermi-Amaldi results. At $c = 0.6638$, corresponding to the "best" value of $\langle E \rangle$ there is a distinct improvement in all quantities. At $c = 0.95$, approximately the value corresponding to minimum Δ , the expectation values of the positive powers of r are estimated much less well. Presumably this is caused by the somewhat earlier cutoff in density attendant upon the use of $c > 0$. The removal of the infinity in the density at $r = 0$ is undoubtedly responsible for the improvement in the other quantities.

8. Future Applications

So far, we have constructed a generalized Thomas-Fermi single particle theory. Application to multiparticle systems may be achieved by invoking the idea of the self-consistent field. We may thus anticipate that the improvement in energies found here would be carried over into a

self-consistent field calculation. Such a series of calculations is in progress. Also, a molecular system of high symmetry is suitable for treatment in this way. This work is also in progress. Both will be reported on in due course.

Acknowledgements

We wish to acknowledge the following grants in support of this research: SG--Nonr - 1677-01, and GSH--AT(30-1) - 2968. We are also grateful for a grant of access to the 7090 Data Processing equipment at the M.I.T. Computation Center.

Table IA

Summary of Results for $M = 1$ (in Atomic Units)

c	λ	$\langle E^2 \rangle_1$	$\langle E^2 \rangle_1$	Δ	$\langle 1/r \rangle_1$	$\langle 1/r^2 \rangle_1$	$\langle r \rangle_1$	$\langle r^2 \rangle_1$
0.00	-.24038	-0.7211	∞	-	1.442	∞	1.541	3.205
0.50	-.24760	-0.5333	0.4458	.1614	1.067	2.414	1.543	3.117
0.60	-.25065	-0.5289	0.4170	.1373	1.058	2.239	1.528	3.048
0.6638	-.25310	-0.5280	0.4046	.1258	1.056	2.162	1.515	2.992
0.70	-.25470	-0.5283	0.3993	.1202	1.057	2.128	1.506	2.956
0.80	-.26040	-0.5325	0.3918	.1082	1.065	2.074	1.474	2.830
0.90	-.26940	-0.5455	0.3981	.1005	1.091	2.095	1.424	2.642
0.95	-.27705	-0.5596	0.4123	.0991	1.119	2.164	1.382	2.494
0.99	-.28920	-0.5857	0.4448	.1018	1.172	2.330	1.319	2.281
1.00	-.30105	-0.6139	0.4845	.1076	1.228	2.537	1.262	2.096
Quantum		-0.5000	0.2500	0	1.000	2.000	1.500	3.000

$\langle E \rangle_1$ at $c = 0.50$ from integration of (6.2) = -0.537.

Table IB


Summary of Results for $M = 5$ (in Atomic Units)


α	λ	$\langle E \rangle_5$	$\langle E^2 \rangle_5$
0.0	-0.08221	-1.2331	∞
0.3	-0.08248	-1.0711	0.6349
0.4	-0.08260	-1.0606	0.5647
0.5	-0.08273	-1.0559	0.5242
0.6	-0.08287	-1.0541	0.4933
0.7	-0.08303	-1.0578	0.4770
Quantum		-1.0000	0.3125


References

- (1) L. H. Thomas, Proc. Cambridge Phil. Soc. 23, 542 (1927).
- (2) E. Fermi, Z. Physik 48, 73 (1928).
- (3) See, for example, P. Gombas, Die Statistische Theorie des Atoms und Ihre Anwendungen (Springer-Verlag, Vienna, 1949).
- (4) C. F. von Weizsacker, Z. Physik 96, 431 (1935).
- (5) See, for example, R. A. Berg and L. Willets, Proc. Phys. Soc. (London) A68, 229 (1955).
- (6) See, for example, S. Golden, Phys. Rev. 105, 604 (1957); see, also, G. A. Baraff and S. Borowitz, Phys. Rev. 121, 1704 (1961).
- (7) An early attempt to deal with this problem is that of W. J. Swiatecki, Proc. Phys. Soc. A68, 285 (1955).
- (8) See, for example, J. M. C. Scott, Phil. Mag. 43, 859 (1952), and R. A. Ballinger and N. H. March, Phil. Mag. 46, 246 (1955).
- (9) See, for example, S. Golden in Ref. (6).
- (10) S. Golden, Phys. Rev. 107, 1283 (1957), referred hereinafter as (I).
- (11) This tacitly assumes that the energy spectrum is non-degenerate.
- (12) One should use the two-sided Laplace transform, but with the tacit assumption that the Hamiltonian is measured from the lowest eigenvalue, the one-sided Laplace transform may be employed in the formalism.
- (13) This tacitly assumes equal numbers of electrons in each spin state.
- (14) We are restricting the following considerations to a single-particle approximation. Thus, $\vec{r}$ refers to the position vector of the particle.
- (15) See, for example, L. D. Landau and E. M. Lifschitz, Quantum Mechanics (Addison-Wesley Publishing Co., Reading, 1958), pp. 118-121.
- (16) This basis may be termed the "barely-free-particle" basis.
- (17) See, for example, Ref. (15), page 87.
- (18) See, for example, G. N. Watson, A Treatise on the Theory of Bessel Functions (Cambridge University Press, Cambridge, 1948), p. 95.
- (19) When $c > 1$ the order of the Bessel function for $\ell = 0$ is imaginary and diverges at the origin.
- (20) Reference (18), p. 395.

- (21) Reference (18), p. 203.
- (22) Reference (15), p. 87.
- (23) See, for example, S. Golden, Phys. Rev. 110, 1349 (1958).
- (24) The same sort of behavior is imposed in an ad hoc manner in Ref. (6).
- (25) Indicated in (I) and in S. Golden, Rev. Mod. Phys. 32, 322 (1960).
- (26) Reference (18), p. 137.
- (27) One must be hesitant about use of such a term as "best," since in the absences of both prior information as to the correct answer and a variational principle, no particular confidence can be placed in this value. The quantity, defined above, is a measure of significance.







TECHNICAL REPORT DISTRIBUTION LIST

CONTRACTOR BRANDEIS UNIVERSITY

NR NO. 051-362

CONTRACT NUMBER Nonr 1677(01)

<u>NO. COPIES</u>	<u>NO. COPIES</u>
Commanding Officer Office of Naval Research Branch Office The John Crerar Library Building 86 East Randolph Street Chicago 1, Illinois (1)	Air Force Office of Scientific Research (SRC-E) Washington 25, D.C. (1)
Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, New York (1)	Commanding Officer Diamond Ordnance Fuze Laboratories Washington 25, D.C. Attn: Technical Information Office Branch 012 (1)
Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena 1, California (1)	Office, Chief of Research and Development Department of the Army Washington 25, D.C. Attn: Physical Sciences Division(1)
Commanding Officer Office of Naval Research Branch Office Box 39 Navy No. 100 Fleet Post Office New York, New York (7)	Chief, Bureau of Ships Department of the Navy Washington 25, D.C. Attn: Code 342C (2)
Director, Naval Research Laboratory Washington 25, D.C. Attn: Technical Information Officer(6) Chemistry Division (2)	Chief, Bureau of Naval Weapons Department of the Navy Washington 25, D.C. Attn: Technical Library (4)
Chief of Naval Research Department of the Navy Washington 25, D.C. Attn: Code 425 (2)	ASTIA Document Service Center Arlington Hall Station Arlington 12, Virginia (10)
DDR and E Technical Library Room 3C-128, The Pentagon Washington 25, D.C. (1)	Director of Research U.S. Army Signal Research and Development Laboratory Fort Monmouth, New Jersey (1)
Technical Director Research and Engineering Division Office of the Quartermaster General Department of the Army Washington 25, D.C. (1)	Naval Radiological Defense Laboratory San Francisco 24, California Attn: Technical Library (1)
Research Director Clothing and Organic Materials Division Quartermaster Research and Engineering Command U.S. Army Natick, Massachusetts (1)	Naval Ordnance Test Station China Lake, California Attn: Head, Chemistry Division (1)

REVISED 1 FEB 1962

TECHNICAL REPORT DISTRIBUTION LIST

CONTRACTOR BRANDEIS UNIVERSITY

Page 2

NR NO 051-362

CONTRACT NUMBER Monr 1677(01)

	<u>NO. COPIES</u>		<u>NO. COPIES</u>
Commanding Officer Army Research Office Box CM, Duke Station Durham, North Carolina Attn: Scientific Synthesis Office	(1)	Dr. J. T. Vanderslice Institute of Molecular Physics University of Maryland College Park, Maryland	(1)
Brookhaven National Laboratory Chemistry Department Upton, New York	(1)	Dr. Harrison Shull Department of Chemistry Indiana University Bloomington, Indiana	(1)
Atomic Energy Commission Division of Research Chemistry Programs Washington 25, D.C.	(1)	Dr. G. B. Kistiakowsky Department of Chemistry Harvard University Cambridge 38, Massachusetts	(1)
Atomic Energy Commission Division of Technical Information Extension Post Office Box 62 Oak Ridge, Tennessee	(1)	Dr. D. F. Hornig Department of Chemistry Princeton University Princeton, New Jersey	(1)
U.S. Army Chemical Research and Development Laboratories Technical Library Army Chemical Center, Maryland	(1)	Dr. H. P. Broida Heat and Power Division National Bureau of Standards Washington 25, D.C.	(1)
Office of Technical Services Department of Commerce Washington 25, D.C.	(1)	Dr. S. H. Bauer Department of Chemistry Cornell University Ithaca, New York	(1)
Commanding Officer ONR Branch Office 495 Summer Street Boston 10, Massachusetts	(2)	Dr. G. Barth-Wehrenalp, Director Inorganic Research Department Pennsalt Chemicals Corporation Post Office Box 4388 Philadelphia 18, Pennsylvania	(2)
Dr. Virginia Griffing Department of Chemistry Catholic University of America Washington, D.C.	(1)	Dr. E. B. Wilson, Jr. Department of Chemistry Harvard University Cambridge 38, Massachusetts	(1)
Dr. J. O. Hirschfelder Naval Research Laboratory University of Wisconsin Madison 6, Wisconsin	(1)		

TECHNICAL REPORT DISTRIBUTION LIST

CONTRACTOR BRANDEIS UNIVERSITY

Page 3

CONTRACT NUMBER Nonr 1677(01)

NR NO 051-362

<u>NO. COPIES</u>	<u>NO. COPIES</u>
Dr. Henry Eyring Graduate School University of Utah Salt Lake City, Utah (1)	Commanding Officer ONR Branch Office Box 39 Navy No.100 Fleet Post Office New York, New York (7)
Dr. B. L. Crawford, Jr. Department of Chemistry University of Minnesota Minneapolis, Minnesota (1)	FOR TRANSMITTAL TO: Dr. N.H. March Department of Physics Sheffield University Sheffield, England
Dr. J. H. Goldstein Department of Chemistry Emory University Emory, Georgia (1)	Dr. H.C. Longuet-Higgins Department of Theoretical Chemistry Cambridge University Cambridge, England
Dr. J. C. Slater Department of Physics Massachusetts Institute of Technology Cambridge 39, Massachusetts (1)	Dr. D. ter Haar Clarendon Laboratory Oxford University Oxford, England
Dr. R. S. Mulliken Department of Physics University of Chicago Chicago 37, Illinois (1)	Dr. C. A. Coulson Mathematical Institute Oxford University Oxford, England
Dr. Martin L. Sage Department of Chemistry University of Oregon Eugene, Oregon (1)	Dr. D. R. Bates Department of Applied Mathematics The Queen's University of Belfast Belfast, N. Ireland
Dr. I. Amdur Department of Chemistry Massachusetts Institute of Technology Cambridge 39, Massachusetts (1)	Dr. W. A. Bingel Max-Planck-Institute for Physics and Astrophysics Aumsterstrasse 6, Munich, Germany
Dr. D. A. Brown Department of Chemistry University College Dublin, Ireland (1)	Madame R.Herman Observatoire de Paris Section d'Astrophysique Meudon (S.-et-O), France
Dr. B. G. Anex Department of Chemistry Yale University New Haven, Connecticut (1)	