

**UNCLASSIFIED**

**AD** **414835**

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

FTD-TT- 62-978

CATALOGED BY DDC

414835

AS AD 111

414835

# TRANSLATION

PROBLEMS OF HEAT TRANSFER IN RAREFIED GASES

By


L. Ye. Kalikhman

FOREIGN TECHNOLOGY  
DIVISION

AIR FORCE SYSTEMS COMMAND

WRIGHT-PATTERSON AIR FORCE BASE

OHIO



## UNEDITED ROUGH DRAFT TRANSLATION

PROBLEMS OF HEAT TRANSFER IN RAREFIED GASES

BY: L. Ye. Kalikhman

English Pages: 3

SOURCE: Russian Periodical, Soveshchaniye po  
Teplo i Massoobmenu, Minsk, 1961,  
Tezisy Dokladov i Soobshcheniy, pp. 31-  
32.

T-110  
S/594-61-0-0-3-11

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:  
TRANSLATION DIVISION  
FOREIGN TECHNOLOGY DIVISION  
WP-AFB, OHIO.

## PROBLEMS OF HEAT TRANSFER IN RAREFIED GASES

L. Ye. Kalikhman

The problems of convective heat transfer in rarefied gases are the same as in ordinary gas-dynamic conditions complicated by additional effects.

A decrease in the gas pressure leads first of all to a change in the surface conditions; a temperature jump and slippage on the wall occurs.

The processes of interaction of the flow with the surface depend essentially on the accommodation coefficient  $\alpha$  and the coefficient of diffusion reflection  $\sigma$ .

The fundamental problem of convective heat transfer consists in determining the heat-transfer coefficients (the Stanton number  $St$  or the Nusselt number  $Nu$ ) as functions of the parameters  $M$ ,  $Re$ ,  $Pr$ ,  $\bar{T}_w$ ,  $\alpha$ ,  $\sigma$  and the Knudsen number  $Kn = M/\sqrt{Re}$ .

Taking into account the effect of a temperature jump in Lees' solution for planar and axisymmetrical gas flows allows us to determine the effect of the Knudsen number on the Nusselt number at the critical point, assuming thermodynamic equilibrium. With an increase in  $Kn$  the heat-transfer intensity decreases sharply.

In this case an increase in the enthalpy factor  $\bar{I}_w$  causes a re-  
duction, while a decrease in  $\bar{I}_w$  causes a growth, in the heat flows.  
A decrease in the accommodation coefficient from  $\alpha = 0.8$  to  $\alpha = 0.4$   
leads to an extremely sharp decrease in the heat-transfer intensity.  
These conclusions agree with Giedt's experiments carried out when  
 $\bar{I}_w = 0.3$  and  $0.7$ , when  $M_\infty = 1.32-5.7$ , and  $Re_H = 30-80$ .

In Giedt's experiments the Mach number has practically no effect;  
however, further increase of this number under low-pressure conditions  
should lead to freezing of the boundary layer. Consequently, the  
problem of heat-transfer due to recombination of atoms on the wall will  
be a critical one. This problem, which was sufficiently developed for  
a continuous medium in the works of Gulard and Rosner, has not yet  
been solved for rarefied gases, in view of the vagueness of the  
boundary conditions for an inhomogeneous gas.

We solved the problem of heat transfer of a longitudinally stream-  
lined flat plate by two methods: the method of linearization of  
equations and the method of series expansion in a small parameter  
inversely proportional to the free path. In the region of large  
 $Kn$  both solutions coincide; in the region of small  $Kn$ , as was to be  
expected, the second solution is unsuitable. The solution obtained  
is a generalization of Schaaf's solution, indicating that under  
equilibrium conditions  $StM$  depends not only on  $Kn$ , but on the density  
ratio  $\rho_s/\rho_{\omega_1}$ . Calculations showed that, when  $Kn$  is constant,  $StM$   
depends essentially on the Mach number and to a lesser degree on  
the enthalpy factor  $\bar{I}_w$ . Large temperature jumps on the surface  
correspond to high Mach numbers and  $\bar{I}_w$  values essentially different  
from unity.

... the effect of the slip velocity into account in the bound-  
ary condition for enthalpy stagnation and taking strict account of

the effect of the work of the friction force in the expression for the heat flow pose certain difficulties. Calculations taking the above factors into account allow us to conclude that at large Kn and Mach numbers these effects are extremely significant.

In this case the heat flow is not proportional to the enthalpy drop  $i_e - i_w$ . Calculations showed that the heat-transfer intensity for a longitudinally streamlined plate is sharply reduced with a decrease in  $\alpha$  and  $\sigma$ .

The extension of the results relating to planar flows of rarefied gas to the case of conical flows by means of a Stepanov transformation is not feasible, due to the changes in the boundary conditions. Direct calculations of the heat flows for this case yield results coinciding with the experiments of Drake and Maslach.

(Moscow)

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE	Nr. Copies	MAJOR AIR COMMANDS	Nr. Copies
		AFSC	
		SCFDD	1
		DDC	25
		TDBTL	5
HEADQUARTERS USAF		TDBDP	3
AFCIN-3D2	1	AEDC (ARY)	1
ARL (ARB)	1	BSD (BSF)	1
		AFFTC (FTY)	1
		ASD (ASYIM)	2
OTHER AGENCIES			
CIA	1		
NSA	6		
DIA	9		
AID	2		
OTS	2		
AEC	2		
PWS	1		
NASA	1		
ARMY (FSTC)	3		
NAVY	3		
NAFEC	1		
PGE	12		
RAND	1		

FTD-TT- 62-978/1+4