

UNCLASSIFIED

AD 4 3 8 7 4 5

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

157-61-58
64-13

QUARTERLY REPORT ON

(15) → CONTRACT NO DA 92-557-FEC-35580

INCLUSIVE DATES 1 October 1961 TO 31 December 1961

Quarterly rpt. 1 Oct-31 Dec 61,

SUBJECT OF INVESTIGATION

(6) →

BACTERIOLOGICAL, IMMUNOLOGICAL
AND
VIRAL STUDIES ON RECTAL MUCUS
IN
ENTERIC INFECTIONS

(SHIGELLOSIS, SALMONELLOSIS,
PATHOGENIC COLI INFECTIONS
AND VIRAL ENTERIC INFECTIONS),

RESPONSIBLE INVESTIGATOR

(10) →

by . Yasuzo Nakamizo ,

Chief Doctor, Medical Section

(5) 876350 →

Tokyo Metropolitan Ebura Infectious
Disease Hospital (JAPAN).
805, Yurigaya, Ota-ku, Tokyo, Japan

MAY 12 1964

U.S. Army Research & Development Group (9852) (Far East)

Office of the Chief of Research and Development
United States Army
APO 343

9

1 1 11.11

AR NO. 1
DDC FILE COPY

438745

438745

BACTERIOLOGICAL, IMMUNOLOGICAL AND
VIRAL STUDIES ON RECTAL MUCUS IN
ENTERIC INFECTIONS
(SHIGELLOSIS, SALMONELLOSIS, PATHO-
GENIC COLI INFECTIONS AND VIRAL
ENTERIC INFECTIONS)

The purpose of investigation

under Contact No DA-92-557-FEC-35580

1. Compare feces with mucus as the isolation source of the pathogenic microbe in enteric infections.
2. Estimate the serological potentialities of mucus in contrast with serum, bile and feces.
3. Measure antibacterial or antiviral resistance of mucus and reseaches for the agents which intensify antibacterial or antiviral resistance of mucus, such as antibiotic, gamma-globulin and vaccin, etc.,.
4. Trace source of immunological potentialities in mucus.

Serological findings on serum and rectal mucus in bacillary dysentery patients obtained during the period of 1 Oct. - 31 Dec. 1961 are as follows.

1. Agglutinin titers against live *Shigella* antigen

a. Agglutinin titers of serum and rectal mucus aspirated from rectal cavity using a romanoscope in bacillary dysentery patients reached peak mostly in the second or third week of illness, but in a few cases within the first week of illness.

b. It was observed that the agglutinin titers in *Shigella sonnei* infected cases were generally lower than those of other shigella infected cases. (Figure 1)

c. Remarkable differences in agglutinin titers were not recognized between the standard strains and the freshly isolated *Shigella* strains from bacillary dysentery patients.

d. Comparative studies of agglutinin titers in serum and mucus materials taken at the same time demonstrated that the serum agglutinin titer is higher than the mucus agglutinin titer in the first to third week of illness but the mucus titer is higher than the serum titer in the second to fourth week of illness.

(Table 1)

e. Attempts were made to see if the agglutinin activity of serum and mucus is affected when heated at 56°C 30 minutes or at 60°C for 3 minutes. The agglutinin titers remained intact even after heating. (Table 2)

2. Agglutin titers of serum and mucus against heated antigen

a. When *Shigella* antigen was heated at 100°C for 30 minutes, 60 minutes or 120 minutes, the agglutinin titer did not vary. (Table 2)

b. It was clarified that the agglutinin titers of serum and mucus in shigella flexneri 2a and 2b and *Shigella sonnei* infected cases are lower than those of *Shigella flexneri* 3a and v-x infected cases.

c. Throughout the process of Shigellosis, distinct differences in agglutinin titers were not demonstrated between serum and mucus.

3. Agglutinin titers of serum and mucus against *Escherichia coli* antigen (live or heated) isolated from *Shigella* positive dysentery cases were usually lower than those against *Shigella* antigen.

Figure 1. Serum agglutinin titer
against live *Shigella* antigen




Table 1

Comparison of agglutinin titer between serum and mucus against live <i>Shigella</i> antigen	Week of illness				
	1 st	2 nd	3 rd	4 th	5 th
serum = mucus	3	3	1	1	1
serum > mucus	5	7	11	1	1
serum < mucus	0	6	6	7	2

Table 2

Material <i>Shigella</i> <i>flexneri</i> 3 a infected case	Inactivation	<i>Shigella</i> antigen			
		live antigen	heated antigen		
			100° C 30 M.	100° C 60 M.	100° C 120 M.
serum	Not inactivated	320	80	80	80
	56° C 30 M.	640	160	160	320
	60° C 3 M.	640	320	320	640
mucus	Not inactivated	1280	640	640	640
	56° C 30 M.	1280	640	640	640
	60° C 3 M.	1280	640	640	640