

UNCLASSIFIED

AD 4 4 1 2 7 3

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

DISCLAIMER NOTICE

**THIS DOCUMENT IS BEST QUALITY
PRACTICABLE. THE COPY FURNISHED
TO DTIC CONTAINED A SIGNIFICANT
NUMBER OF PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

Stauffer

4 4 1 2 7 3



STAUFFER CHEMICAL COMPANY



HIGH ENERGY OXIDIZERS

CONTRACT Nonr-4019(00)

Project NR 093-035

**Richmond Research Center
Richmond, California**

STAUFFER CHEMICAL COMPANY
Richmond Research Center
Richmond, California

"HIGH ENERGY OXIDIZERS"

CONTRACT Nonr-4019(00)

Project NR 093-035

ARPA No. 399-62

OFFICE OF NAVAL RESEARCH

WASHINGTON, D. C.

Quarterly Technical Summary Report
for the Period February 1, 1964 to May 1, 1964

Investigators



Dr. K. O. Christie



Dr. A. E. Pavlath

Department Supervisor



Dr. E. G. Wallace

Richmond Research Center
Richmond, California

TABLE OF CONTENTS

	<u>Page No.</u>
Technical Report Distribution List	1
Summary	1
Abstract	2
Introduction	3
Discussion	3
Infrared spectra	3
Reaction of ClF_3 with AgNO_3	5
Reaction between ClF_3 and KClO_4	5
Reaction between ClF_3 and N_2O	5
Experimental	6
Vacuum line	6
Infrared Spectra	6
Reaction between ClF_3 and AgNO_3	6
Reaction between ClF_3 and KClO_4	6
Reaction between ClF_3 and N_2O	6

TECHNICAL REPORT DISTRIBUTION LIST

STAUFFER CHEMICAL COMPANY

Contract Nonr 4019(00)

NR 093-035

<u>Addressee</u>	<u>No. Copies</u>
Advanced Research Projects Agency Propellant Chemistry The Pentagon, Room 3D165 Washington 25, D. C.	6
Chemical Propulsion Information Agency Applied Physics Laboratory Johns Hopkins University Silver Spring, Maryland	3
Office of Naval Research Branch Office 1000 Geary Street San Francisco 1, California Attn: Dr. P. A. Miller	1
Office of Naval Research U. S. Navy Department Power Branch, Code 429 Washington 25, D. C.	2
American Oil Company Attn.: Dr. T. D. Nevitt Whiting Laboratories Whiting, Indiana	1
Astropower, Inc. Attn.: Dr. W. D. English 2069 Randolph Avenue Costa Mesa, California	1
Boston College Attn.: Dr. Robert F. O'Malley Chestnut Hill 67, Massachusetts	1
Georgetown University Attn.: Dr. William Clinton Department of Chemistry Washington 7, D. C.	1
Harshaw Chemical Company Attn.: Dr. D. R. Martin 1945 East 9th Street Cleveland 6, Ohio	1
Hercules Powder Company Attn.: Dr. R. S. Voris Wilmington, Delaware	1

Stauffer Chemical Company
Contract Nonr 4019(00)
NR 093-035

Page 11

TECHNICAL REPORT DISTRIBUTION LIST

<u>Addressee</u>	<u>No. Copies</u>
IIT Research Institute Attn.: Dr. Irvine J. Solomon 10 W. 35th Street Chicago 16, Illinois	1
King's College Attn.: Dr. Grihapati Mitra Wilkes-Barre, Pennsylvania	1
Midwest Research Institute Attn.: Dr. Florence Metz 425 Volker Boulevard Kansas City 10, Missouri	1
Monsanto Research Corporation Attn.: Dr. James W. Dale Boston Laboratories Everett 49, Massachusetts	1
National Bureau of Standards Attn.: Dr. David E. Mann Washington 25, D. C.	1
Ozark-Mahoning Company Attn.: Dr. Albert W. Jache 310 West Sixth Street Tulsa 19, Oklahoma	1
Royal College of Science & Technology Attn.: Dr. D. W. A. Sharp Chemistry Department Glasgow, C.1, Scotland	1
Shell Development Company Attn.: Dr. T. W. Evans Emeryville, California	1
Stanford Research Institute Attn.: Dr. Edwin S. Gould Menlo Park, California	1
Thiokol Chemical Corporation Attn.: Dr. A. R. Young Reaction Motors Division Denville, New Jersey	1

Stauffer Chemical Company
Contract Nonr 4019(00)
NR 093-035

Page 111

TECHNICAL REPORT DISTRIBUTION LIST

<u>Addressee</u>	<u>No. Copies</u>
Tracor, Inc. Attn.: Dr. Ray M. Hurd 1701 Guadalupe Street Austin 1, Texas	1
Union Carbide Corporation Attn.: Dr. R. G. Breckenridge P. O. Box 6116 Cleveland 1, Ohio	1
University of Washington Department of Chemistry Seattle 5, Washington Attn: Dr. G. H. Cady	1
Stanford Research Institute Propulsion Sciences Division Menlo Park, California	1
Dr. H. H. Selig Building 200 Argonne National Laboratory 9700 S. Cass Avenue Argonne, Illinois	1
University of Idaho Moscow, Idaho Attn: Dr. Jean'ne M. Shreeve	1

This report is distributed to the Chemical Propulsion Mailing List
of December 1963.

Summary

Using infrared spectroscopy the ionic structure of the chlorine-trifluoride complexes was proved in the solid form also. Conclusion was made concerning the structure of the ClF_2^+ cation. Investigations were started for the preparation of other complexes with anions of higher energy content.

Abstract

This is the first Quarterly Technical Summary Report of the second year on the investigation of chlorinetrifluoride complexes.

The IR spectra of gaseous and solid chlorinetrifluoride and arsenicpentafluoride were taken and compared with the spectrum of $\text{ClF}_2^+\text{AsF}_6^-$ and KAsF_6 . A comparison between the spectra indicated the presence of AsF_6^- in the complex. Analogous results were obtained for the $\text{ClF}_2^+\text{BF}_4^-$ complex. From the absorption of the ClF_2^+ cation it was concluded that it exists in an angular form and is not linear.

The reaction of AgNO_3 , KClO_4 and N_2O with chlorinetrifluoride was studied in an attempt to prepare ClF_2^+X^- where X^- was expected to be NO_3^- , ClO_4^- or N_2OF^- .

An all metal vacuum line and a Teflon IR cell were designed for future work.

A paper was presented on the results of the first year's study at the ARPA Conference, Chicago.

Introduction

It was established earlier that the chlorinetrifluoride complexes are dissociated in solvents such as halogenfluorides. Low temperature infrared study was needed to determine the structure in solid form.

Discussion

Infrared spectra

TABLE I

$\text{ClF}_2^+ \text{AsF}_6^-$	KAsF_6	$\text{ClF}_3(\text{g})$	$\text{ClF}_3(\text{s})$	$\text{AsF}_5(\text{g})$	$\text{AsF}_5(\text{s})$
405 s	413 ms		499 m 508 m	409 s 488 w	404 m
521 m		518 s 535 s			
558 m			604 vs 630 vs		
688 vs	685 vs	694 vs			
703 vs	705 vs	703 vs 713 vs			
729 vs	725 vs	741 s 761 s	765 m 770 w		725 vs 738 vs
				786 vs 811 vs 818 m 858 w 866 w	814 vs
					891 w
		957 m 1022 m		977 w 1023 w	
		1223 s 1273 w			1101 vw 1127 w 1201 w
1284 vw 1298 w		1451 s 1466 s 1488 s 1505 s			

The absorptions obtained for $\text{ClF}_2^+ \text{AsF}_6^-$, KAsF_6 , gaseous and solid ClF_3 and AsF_5 are tabulated in Table I.

It can be seen that the absorptions of the $\text{ClF}_2^+\text{AsF}_6^-$ cannot be due to those of ClF_3 or AsF_5 in either gaseous or solid form. The comparison of the spectra of $\text{ClF}_2^+\text{AsF}_6^-$ and KAsF_6 clearly indicates the presence of the AsF_6^- anion in the complex. Similar observations can be made in the case of the $\text{ClF}_2^+\text{BF}_4^-$ complex establishing the presence of the BF_4^- anion as it was already shown earlier by Selig and Shamir⁽¹⁾. Knowing that these complexes are formed from their components in a ratio of one:one, the presence of the AsF_6^- and BF_4^- anion leaves no doubt about the form of their chlorinetrifluoride part: it has to exist as a ClF_2^+ cation in solid phase also. In Table II the absorption of both complexes are given together with their assignment.


TABLE II

	$[\text{ClF}_2]^+[\text{AsF}_6]^-$	$[\text{ClF}_2]^+[\text{BF}_4]^-$
	729 cm^{-1}	1036 cm^{-1}
✓ anion	703	1005
	688	
✓ as ClF_2^+	557	536
✓ sym ClF_2^+	519	518

It can be seen that in each case there are two absorptions which do not belong to the anions; therefore, we must assign them to the ClF_2^+ cation. Since the highest frequency reported for a deformation vibration of the Cl-F bond is 434 cm^{-1} (2) it can be assumed that both observed absorptions are stretching modes.

Theoretically the ClF_2^+ cation could have the symmetry C_{2v} (bent form) or $\text{D}_{\infty h}$ (linear). The possible fundamentals for stretching vibrations are shown in Figure I.


FIGURE I



Since the (s) fundamental of symmetry $D_{\infty h}$ is IR inactive, a linear XY_2 molecule would show only one stretching vibration. Therefore, symmetry C_{2v} has to be favored, where both vibrations are IR active. Based on these results, it can be concluded that the ClF_2^+ cation has an angular form.

Reaction of ClF_3 with $AgNO_3$

This reaction was carried out in an attempt to prepare $ClF_2^+NO_3^-$ according to the following equation:




The reaction, however, did not result in $ClF_2^+NO_3^-$, or if it did form during the reaction it decomposed. Silverdifluoride and NO_2F were identified as the reaction products. Based on this observation, the following equation is suggested to describe this reaction:



Reaction between ClF_3 and $KClO_4$


The reaction which was expected to proceed in the following way:



did not result in any change of the $KClO_4$. X-ray analysis of the residue showed only unreacted $KClO_4$ and no KF.

Reaction between ClF_3 and N_2O

This reaction was described in a U. S. Patent as a way to prepare NF_3 (3). The reaction conditions, however, are too drastic ($300^\circ C$. under pressure). It was hoped that under milder conditions the following reaction could take place:



The N_2OF^- anion would be isoelectronic with N_2F_2 and therefore its existence might be possible. The reaction was investigated at different temperatures between -196° and $0^\circ C$., but the IR spectrum showed in each case only unreacted starting materials.

Experimental

Vacuum line

All reported reactions were carried out in a glass vacuum line.

Infrared Spectra

A Beckman IR-9 all grating instrument was used for this investigation covering a range from 400 to 4000 cm^{-1} . In addition to earlier investigations, the spectra of the complexes were taken using AgCl windows. The cell was made of glass, with an internal cold AgCl window.

Reaction between ClF_3 and AgNO_3

Chlorine trifluoride was condensed on top of powdered AgNO_3 at -196°C . The mixture was warmed up slowly to room temperature. The gases were analyzed by IR and the presence of NO_2F was established besides unreacted chlorine trifluoride. The dark solid residue was analyzed: Ag = 72.5%, F = 24.9%. This represents an atomic ratio of 1:1.95 between silver and fluorine. The X-ray powder diagram of the compound was also taken.

Reaction between ClF_3 and KClO_4

The reaction was carried in a similar way as with AgNO_3 . The vapor phase did not contain anything outside of ClF_3 according to the IR. The white solid residue was identified by X-ray analysis as unreacted starting material.

Reaction between ClF_3 and N_2O

The two components were condensed together on the cold AgCl window of the low temperature IR cell. The IR spectrum, at temperatures between -196° and 0°C ., showed only the absorptions of the starting materials. In another attempt the two components were condensed together in the vacuum line and were allowed to warm up to 0°C . No visible sign of any reaction was observed. When the liquid was warmed up to room temperature everything evaporated without any residue and the vapor phase again showed only ClF_3 and N_2O .

ls
May 21, 1964

- (1) Selig H., Shamir J., Inorg. Chem. 3, 294 (1964)
- (2) Classen, H. H., Weinstock, B., Malm, J. G., J. Chem. Phys. 28, 285 (1958)
- (3) Marsh F., U. S. Patent 3,032,400