

UNCLASSIFIED

AD 444518

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

444518

MEMORANDUM
RM-4104-PR
AUGUST 1964

ADP BY DDC
AD No.

OSCILLATORY AND UNIMODAL PROPERTIES OF SOLUTIONS OF SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS

Richard Bellman

PREPARED FOR:
UNITED STATES AIR FORCE PROJECT RAND

DDC
RECEIVED
AUG 21 1964
MILITARY DISTRICT OF COLUMBIA

The **RAND** Corporation
SANTA MONICA • CALIFORNIA

MEMORANDUM

RM-4104-PR

AUGUST 1964

OSCILLATORY AND UNIMODAL PROPERTIES
OF SOLUTIONS OF SECOND ORDER
LINEAR DIFFERENTIAL EQUATIONS

Richard Bellman

This research is sponsored by the United States Air Force under Project RAND--Contract No. AF 49(638)-700 monitored by the Directorate of Development Plans, Deputy Chief of Staff, Research and Development, Hq USAF. Views or conclusions contained in this Memorandum should not be interpreted as representing the official opinion or policy of the United States Air Force.

DDC AVAILABILITY NOTICE

Qualified requesters may obtain copies of this report from the Defense Documentation Center (DDC).

The RAND Corporation

1700 MAIN ST • SANTA MONICA • CALIFORNIA • 90406

PREFACE

Part of the Project RAND research program consists of basic supporting studies in mathematics. The mathematical research presented here concerns a technique for studying certain properties of solutions to various types of differential equations.

SUMMARY

In previous papers, we have shown how to derive fundamental properties of the solutions of

$$u'' + p(x)u = 0$$

directly from the fact that it is the Euler equation associated with the functional

$$J(u) = \int_0^a [u'^2 - p(x)u^2] dx.$$

Here we wish to demonstrate a Sturm oscillation theorem and the unimodal property of the solution in analogous fashion, under the assumption that $p(x) > 0$.

CONTENTS

PREFACE.....	iii
SUMMARY.....	v
Section	
1. INTRODUCTION.....	1
2. THE VARIATIONAL FORMULATION.....	1
3. THE STURMIAN PROPERTY.....	2
4. A UNIMODAL PROPERTY.....	4
5. DISCUSSION.....	7
REFERENCES.....	9

OSCILLATORY AND UNIMODAL PROPERTIES OF SOLUTIONS OF SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS

1. INTRODUCTION

In two previous papers, we have shown how to derive fundamental properties of the solutions of

$$(1.1) \quad u'' + p(x)u = 0$$

directly from the fact that it is the Euler equation associated with the functional

$$(1.2) \quad J(u) = \int_0^a [u'^2 - p(x)u^2] dx,$$

[1], [2]. Here we wish to demonstrate a Sturm oscillation theorem and the unimodal property of the solution in analogous fashion, under the assumption that $p(x) > 0$.

2. THE VARIATIONAL FORMULATION

Consider the problem of minimizing

$$(2.1) \quad K(u) = \frac{\int_0^a u'^2 dx}{\int_0^a p(x)u^2 dx}$$

over all functions $u(x)$ satisfying the conditions $u(0) = u(a) = 0$, and such that the numerator and denominator exist and are nonzero. This leads in the usual way to the equation

$$(2.2) \quad u'' + \lambda p(x)u = 0, \quad u(0) = u(a) = 0,$$

with the minimum equal to the smallest characteristic value. If the equation

$$(2.3) \quad u'' + p(x)u = 0, \quad u(0) = u(a) = 0,$$

with $p(x) > 0$, has a solution which is positive for $0 < x < a$, we know that $\lambda = 1$ is the smallest characteristic value and that $u(x)$ is, up to a normalization factor, the corresponding characteristic function. This result can be established without the use of Sturmian oscillation theorems, if we employ the Perron-Frobenius-Jentszch techniques of positive operators.

3. THE STURMIAN PROPERTY

Suppose that we are given the two equations

$$(3.1) \quad u'' + p(x)u = 0, \quad u(0) = 0,$$

$$v'' + q(x)v = 0, \quad v(0) = 0,$$

with $q(x) \geq p(x) > 0$. We wish to demonstrate that the first zero of $v(x)$ cannot occur after the first zero of u .

We assume that the situations were the reverse, as indicated below (see Fig. 1). Then, from what has preceded,




Fig. 1

$$(3.2) \quad 1 = \min_v \frac{\int_0^b v'^2 dx}{\int_0^b q(x)v^2 dx} < \frac{\int_0^b w'^2 dx}{\int_0^b q(x)w^2 dx},$$

where $w(x)$ is another function defined over $[0, b]$.
Choose $w(x)$ in the following way:

$$(3.3) \quad \begin{aligned} w(x) &= u(x), & 0 \leq x \leq a, \\ &= 0, & a \leq x \leq b. \end{aligned}$$

Then (3.2) yields

$$(3.4) \quad 1 < \frac{\int_0^a u'^2 dx}{\int_0^a q(x)u^2 dx} \leq \frac{\int_0^a u'^2 dx}{\int_0^a p(x)u^2 dx} = 1,$$

a contradiction.

4. A UNIMODAL PROPERTY

Let us now demonstrate a familiar property of the first characteristic function $u(x)$, its unimodal character. Suppose that $u(x)$, the solution of

$$(4.1) \quad u'' + p(x)u = 0, \quad u(0) = u(a) = 0,$$

with the property that $u(x) > 0$, $0 < x < a$, had the form in Fig. 2. Let us obtain from this function a function which yields a smaller value of $K(u)$, as defined in (2.1), in the fashion of Fig. 3. In the interval $[a_2, a_1]$, $v(x) = u(a_1)$, otherwise $v(x) = u(x)$.




Fig.2




Fig.3

We see that

$$(4.2) \quad \int_0^a v'^2 dx < \int_0^a u'^2 dx,$$
$$\int_0^a p(x)v^2 dx > \int_0^a p(x)u^2 dx.$$

Hence,

$$(4.3) \quad K(v) < K(u),$$

a contradiction to the minimizing property of u .

5. DISCUSSION

One value of the foregoing technique is that it can be applied to study nonlinear ordinary and partial differential equations as well. Furthermore, we can study the properties of the higher characteristic functions by using relative minima.

REFERENCES

1. Bellman, R., "On the Nonnegativity of Solutions of the Heat Equation," Boll. d'Unione Matematico, Vol. 12, 1957, pp. 520-523.
2. ———, "On Variation-diminishing Properties of Green's Functions," Boll. d'Unione Matematico, Vol. 16, 1961, pp. 164-166.

UNCLASSIFIED

UNCLASSIFIED