

UNCLASSIFIED

AD NUMBER
AD459384
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; FEB 1965. Other requests shall be referred to Army Biological Labs., Fort Detrick, MD.
AUTHORITY
BDRL D/A ltr, 27 Sep 1971

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD 4 5 9 3 8 4

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CREATED BY: DDC

AS AD NO. 459384

TECHNICAL MANUSCRIPT 200

A DEVICE
FOR CREATING DRY AEROSOLS
FOR EXPERIMENTAL INFECTIVITY STUDIES

4 5 9 3 8 4

FEBRUARY 1965

UNITED STATES ARMY
BIOLOGICAL LABORATORIES
FORT DETRICK

FOR ERRATA

AD 459384

THE FOLLOWING PAGES ARE CHANGES

TO BASIC DOCUMENT

AD 459384

459384

ERRATA SHEET

SUBJECT REPORT: Technical Manuscript 200

TITLE: A device for creating dry aerosols for
experimental infectivity studies

DATE ISSUED: February 1965

CORRECTION: Dr. Beebe's name should be corrected to read
James M. Beebe

U.S. ARMY BIOLOGICAL LABORATORIES
Fort Detrick, Frederick, Maryland

TECHNICAL MANUSCRIPT 200

A DEVICE FOR CREATING DRY AEROSOLS FOR
EXPERIMENTAL INFECTIVITY STUDIES

William S. Woodrow

James M. Bebee

Gordon L. Jessup, Jr.

William G. Roessler

Medical Bacteriology and Aerobiology Divisions
DIRECTORATE OF BIOLOGICAL RESEARCH
and
Biomathematics Division
DIRECTORATE OF TECHNICAL SERVICES

Project 1C522301A059

February 1965

This publication or any portion thereof may not be reproduced without specific authorization from the Commanding Officer, U. S. Army Biological Laboratories, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland. 21701. However, DDC is authorized to reproduce the publication for U. S. Government purposes.

The information in this publication has not been cleared for release to the public.

DDC AVAILABILITY NOTICE

Qualified requestors may obtain copies of this publication directly from DDC.

Foreign announcement and dissemination of this publication by DDC is limited.

ABSTRACT

Equipment is needed that will permit laboratory studies with experimental animals using aerosols of dusts, spores, microorganisms, or toxins. This report describes a machine capable of producing dynamic clouds and suitable for attachment to animal exposure boxes.

A dry preparation of Serratia marcescens with 68.0% of the particles less than 5 μ in diameter was used for calibration. Test material compacted in a small cartridge was forced by plunger action onto the serrated edge of a metering wheel, which then presented the powder to a dry air jet, creating a cloud. Samples were collected by impinging the particles in a liquid. Viability was determined by spreading dilutions of that liquid on agar plates. The distribution of particle diameters was determined by measuring at least 500 particles collected on microscope slides.

Plunger speed was found to have a highly significant effect on particle concentration in the aerosol at the slow speed of the metering wheel. An approximate maximum fivefold difference in cloud concentration (from 3.9×10^6 to 21.8×10^6 viable S. marcescens cells) was found by testing the various plunger and metering wheel agar combinations. Approximately 97% of the particles generated in the experimental aerosols were less than 5 μ in diameter.

A DEVICE FOR CREATING DRY AEROSOLS FOR EXPERIMENTAL INFECTIVITY STUDIES

Investigations of the toxicity of soluble substances and the infectivity and virulence of microorganisms by exposure of laboratory animals to aerosols of such materials have largely been confined to "clouds" generated from liquid suspensions. However, many diseases and pathological conditions result from the inhalation of dry microorganisms, dusts, and toxic materials. It would be highly desirable to have equipment that would permit laboratory studies with experimental animals using dusts, spores, microorganisms, or toxins prepared in an aerosol of finely divided dry powder. The lack of a suitable device for continuously generating an aerosol from dried materials has hindered such investigations. This report discusses a study of a small machine that is capable of producing dynamic clouds of dry materials; and this device can be attached to animal exposure boxes. Because the concentration of the cloud could be varied by means of gears, the experimental conditions, number of particles counted, and determination of viability of bacterial cells were replicated so that the results could be analyzed statistically.

The device disseminates the aerosol from a column of packed powder that is forced by a plunger onto the serrated edge of a metering wheel. The compacted powder is removed in small increments by the metering wheel and is presented to a high-velocity air jet that generates the aerosol at a rate of 8 liters per minute. The rate of plunger travel and speed of the metering wheel are controlled by separate sets of gears. In order to observe the behavior of clouds thus produced, the disseminating device was attached to a small animal holding box, one of several different kinds used with a Henderson apparatus in our laboratory for respiratory infection studies. Figure 1 is a photograph of the disseminator attached to the animal holding box with a Shipe aerosol sampler in position. The Shipe sampler collects the disseminated powder by impinging the particles in a liquid. Figure 2 is a close-up view of the gear trains that control the rate of plunger travel and speed of the metering wheel; the aerosol exit port and the connection for compressed air are shown in Figure 3.

A dry preparation of Serratia marcescens with 68.0% of the particles less than 5 μ in diameter was used to calibrate the disseminator. This powder was packed in a small (0.746cc) cartridge-type magazine by the use of weights. The magazine was held in alignment with a piston in a simple device that permitted dry powders to be compacted to various degrees by



Figure 1. Dry Aerosol Disseminator Attached to Animal Exposure Box.





Figure 2. Close-Up View of Dry Aerosol Disseminator Gear Trains.



METERING WHEEL HOUSING COMPRESSED AIR LINE AEROSOL PORT PLUNGER

Figure 3. Dry Aerosol Disseminator Showing Connection for Compressed Gas, Plunger Mechanism, and Aerosol Exit Port.

placing weights on top of the piston, as shown in Figure 4. Weight in excess of 700 grams packed the test material too tightly, but 200- to 700-gram weights provided a uniform cohesive column. A quantity of the powdered material sufficient for the entire investigation was kept under refrigeration, and an amount sufficient for one day's testing was removed on each of nine working days. Decimal dilutions of 0.1 gram of test material in tryptose saline diluent were spread on replicate tryptose agar plates at the beginning and end of each working day for viability determinations.

The cloud was initiated by manual movement of the plunger until the cloud became visible in a Tyndall beam. The machine was then operated mechanically for a one-minute equilibration period before a one-minute cloud sample was collected. After the impingers were changed, the machine was restarted and a second one-minute equilibration period was allowed before a second one-minute sample was collected. A third sample was collected in like manner. After 30 such runs, the data showed that counts of the first impinger sample in each run were somewhat higher than those of the second and third samples. Therefore, a two-minute equilibration period was employed for runs 31 through 54. Cloud samples were collected in Shipe impingers containing 25 ml of tryptose saline solution at a flow rate of 12.5 liters per minute. Impinging fluids were diluted and spread on tryptose agar plates to determine viable cell numbers.

Between the first and second impinger samples, a cloud sample was taken to determine particle size distribution. This was accomplished by placing a clean glass slide, bearing no adhesive, on the bottom inside a one-gallon paint can, passing the aerosol through the can for 3 minutes, and then allowing the aerosol to settle 48 hours. At least 500 particles per slide, which comprised 20 to 30 fields, were measured under high dry magnification by comparison with a series of calibrated circles on a Howard disc incorporated into the eyepiece of a microscope. Alterations were made in plunger and metering wheel speeds by changing gear settings prior to each run, in order to assess a range of cloud concentrations. Between runs the machine was partly disassembled and parts in contact with test material (and the animal holding box) were cleaned in 95% ethyl alcohol. After reassembly, an operating period of one minute without aerosol generation assured the complete evaporation of residual alcohol. Air was by-passed through the animal holding box at 20 liters per minute. Immediately before each run, a one-minute sample of that air was collected to make certain that no significant contamination from the previous run was present. This contamination, or carry-over from one run to the next, when present, had a concentration approximately four to six logarithms less than the experimental clouds and was ignored in the statistical analysis.



Figure 4. Device for Compacting Dry Test Materials.

One complete replication of this experiment required 18 runs; that is, 3 metering wheel speeds by 3 plunger speeds by either 200 or 700 grams used for compaction of test material. Only six runs per day could be accomplished efficiently. A split plot design was utilized, wherein wheel speed was considered as the whole plot, and the subplot was composed of the factorial arrangement of plunger speed and compaction weight. In this particular design, wheel speed is completely confounded, with day effects, which were assumed to be negligible. Wheel speeds were randomized within each replication, and combinations of plunger and compaction treatment were randomized within each day. Table 1 shows the order in which the runs were conducted.

Plunger speed was found to have a highly significant effect on cloud concentration at the slow metering wheel speed, but effects due to wheel speed and compaction weights were not detected. The interaction of plunger and wheel speeds indicated that the plunger effect was not the same with all wheel speeds. This is illustrated in Table 2, which shows that the plunger has its greatest effect at slow wheel speed, and that at the fast wheel speed the plunger had no effect. Plunger effect on cloud concentration at the intermediate wheel speed gave values between those found as the fast and slow wheel speeds. An approximate maximum fivefold difference in cloud concentration (from 3.9×10^6 to 21.8×10^6 viable Serratia marcescens cells) was found by testing these various combinations of plunger and metering wheel gear.

Plunger speed had a statistically significant linear effect on mean particle size and wheel speed had a quadratic effect. Figure 5 shows the mean relative frequency as per cent of each size classification by plunger speed.

The smallest comparison circle on the Howard ocular disc measured 1.4μ , which prevented recording particles smaller than this figure. The detection of very large particles would have required examination of many more than 20 to 30 fields for full representation.

This machine has a potential application in many areas of investigation involving aerosols of small dry particulates. Respiratory infectivity or toxicity could be studied for many biological materials, such as bacterial or fungal spores and various toxic materials. Inorganic chemical fumes, smokes, or dusts could be assayed in the dry state, provided that they were not too corrosive or abrasive for the machine. Patent rights are pending on the dry aerosol disseminator described.

TABLE 1. SPLIT PLOT DESIGN AND ORDER IN WHICH TRIALS WERE CONDUCTED

	Run	Pa/ Cb/	Run	P	C	Run	P	C
	Day 1 Wheel 7 ^c /		Day 2 Wheel 1			Day 3 Wheel 4		
Rep 1	1	1	2	7	1	2	13	3
	2	3	2	8	1	7	14	1
	3	1	7	9	3	7	15	3
	4	6	2	10	6	2	16	6
	5	3	7	11	6	7	17	6
	6	6	7	12	3	2	18	1
	Day 4 Wheel 1		Day 5 Wheel 4			Day 6 Wheel 7		
Rep 2	19	6	7	25	3	2	31	1
	20	3	7	26	1	2	32	6
	21	1	7	27	6	2	33	6
	22	1	2	28	1	7	34	3
	23	6	2	29	6	7	35	3
	24	3	2	30	3	7	36	1
	Day 7 Wheel 4		Day 8 Wheel 7			Day 9 Wheel 1		
Rep 3	37	1	2	43	3	7	49	6
	38	6	7	44	1	2	50	1
	39	3	2	45	6	2	51	3
	40	3	7	46	3	2	52	1
	41	6	2	47	6	7	53	6
	42	1	7	48	1	7	54	3

a. P = Plunger Gear no.

b. C = Compaction Weight, Hundred grams.

c. Wheel = Metering Wheel Gear no.

TABLE 2. AEROSOL CONCENTRATIONS ATTAINED WITH DISSEMINATOR USING
 DRY, SIZED SERRATIA MARCESCENS

Metering Wheel	Viable cells per liter of cloud, 10 ⁸		
	No. 1 Plunger Gear, Fast (1.54 mm/minute)	No. 3 Plunger Gear, Intermediate (0.64 mm/minute)	No. 6 Plunger Gear, Slow (0.19 mm/minute)
Fast (6.0 rpm)	9.6	5.7	7.3
Intermediate (1.8 rpm)	16.2	6.6	5.3
Slow (0.5 rpm)	21.8	5.0	3.9




Figure 5. Mean Relative Frequency (per cent) of each Size Classification by Plunger Speed.

DISTRIBUTION LIST

<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>
Assistant Scientific Director Building 812	1
Directorate of Biological Research Building 560	1
Directorate of Development Building 824	1
Directorate of Industrial Health & Safety Building 550	1
Chief, Program Coordination Office Building 825	1
Chief, Aerobiology Division Building 459	1
Chief, Medical Bacteriology Division Building 560	1
Chief, Medical Investigation Division Building 604	1
Chief, Physical Sciences Division Building 568	1
Chief, Biomathematics Division Building 1422	1
Chief, Process Development Division Building 469	1
Chief, Technical Evaluation Division Building 568	1
Chief, Virus & Rickettsia Division Building 539	10
Documents, Technical Library Building 426	2
Test Chamber Branch Technical Evaluation Division Building 1412	1

<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>
Editorial Branch Technical Information Division Building 816	1
Technical Releases Branch Technical Information Division Building 426	10
U.S. Army Medical Unit Ft. Detrick, Maryland Building 120	1
Liaison Representative Animal Disease Investigation Building 1301	5
Public Health Service Liaison Office Building 1301	9
Commanding Officer Naval Unit Building 125	3
Commanding General U.S. Army Edgewood Arsenal ATTN: SMUEA-CS Edgewood Arsenal, Maryland, 21010	1
Commanding Officer U.S. Army Chemical Research & Development Laboratories ATTN: Librarian Edgewood Arsenal, Maryland, 21010	2
Commanding General U.S. Army Munitions Command ATTN: AMSMU-CS Dover, New Jersey, 07801	1
Commanding General U.S. Army Munitions Command ATTN: AMSMU-RE-RR, Mr. G. Chesnov Dover, New Jersey, 07801	1
Commandant U.S. Army CBR Weapons Orientation Course Dugway Proving Ground Dugway, Utah, 84022	1

<u>ADDRESSEE</u>	<u>NUMBER OF COPIES</u>
Commanding General Deseret Test Center ATTN: Technical Library Fort Douglas, Utah, 84113	2
Commanding General U.S. Army Materiel Command Research Division, AMCRD-RC R&D Directorate Washington, D.C., 20315	1
Department of the Army Asst Chief of Staff/Force Operations ATTN: Technical Coordinator (B) CBR & N Directorate The Pentagon, Room 3A-480 Washington 25, D.C.	1
Defense Documentation Center Cameron Station Alexandria, Virginia, 22314	20
Detachment 4, RTD (ATCB) Eglin Air Force Base, Florida, 32542	1
APGC (PGBAP-1) Eglin Air Force Base, Florida, 32542	1
Dr. S. H. Madin Scientific Director Naval Biological Laboratory Naval Supply Center Oakland, 14, California, 94614	1
Commander (Code 4036) U.S. Naval Ordnance Test Station China Lake, California, 93557	1
Commanding Officer and Director U.S. Naval Applied Science Laboratory Naval Base, Code 9440 Brooklyn, New York, 11251	1
U.S. Army Medical R&D Command Office of the Surgeon General ATTN: MEDDH-C Main Navy Building, Room 2526 Washington, D.C., 20315	1

ADDRESSEENUMBER OF COPIES

Commandant
 USA CmlCen & Sch, ATTN: Bio Branch
 Ft. McClellan, Alabama, 36205

U.S. Army Standardization Group-Canada
 Office, Senior Standardization Rep.
 c/o Director of Equipment Policy
 Canadian Army Headquarters
 Ottawa 4, Canada

Munitions/TW
 Defence Research Staff
 British Embassy
 3100 Massachusetts Avenue, N.W.
 Washington 8, D.C.

Canadian Liaison Office (CER)
 Building 5101
 Edgewood Arsenal, Maryland, 21010

Australian Embassy
 ATTN: Lt. Col. P. D. Yonge
 Australian Army Staff (W)
 2001 Connecticut Avenue, N. W.
 Washington 7, D.C.

1

1

3

3

2