

UNCLASSIFIED

AD NUMBER
AD464264
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; FEB 1965. Other requests shall be referred to Office of Naval Research, Arlington, VA 22203.
AUTHORITY
ONR ltr, 4 May 1977

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD. 4 6 4 2 6 4

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

CATALOGED BY...

AS AD...

464264

4 6 4 2 6 4

DDC  
JUN 7 1965  
JISA B

AVAILABLE COPY WILL NOT PERMIT  
FULLY REPRODUCIBLE COPY.  
REPRODUCTION PERMITTED IF  
BY USER OF DDC.

**HYDRONAUTICS, incorporated**  
research in hydrodynamics

HYDRONAUTICS, Incorporated

TECHNICAL REPORT 231-5

ON INTERNAL GRAVITY WAVES  
GENERATED BY LOCAL DISTURBANCES

By

K. K. Wong

February 1965

Prepared for  
Office of Naval Research  
Department of the Navy  
Contract No. Nonr 3688(00)  
NR 220-016

TABLE OF CONTENTS

	Page
INTRODUCTION.....	1
THE GOVERNING EQUATIONS.....	1
ELEMENTARY WAVES.....	6
INTERNAL WAVES GENERATED BY AN OSCILLATING SINGULARITY.....	9
INTERNAL WAVES GENERATED BY AN INSTANTANEOUS SOURCE.....	17
REFERENCES.....	22

LIST OF FIGURES

- Figure 1 - Allowable Wave Numbers for Different Frequencies
- Figure 2 - Dependence of the Phase Velocity on the Direction of the Wave Number
- Figure 3 - Dependence of the Magnitude of the Group Velocity on the Direction of the Wave Number
- Figure 4 - Internal Motion Due to an Oscillating Source Whose Frequency is Less than the Vaisala Frequency
- Figure 5 - The Wave Front of an Instantaneous Disturbance in an Exponentially Stratified Fluid of Infinite Extent

## NOTATION

A	Source amplitude
b	Exponent of density stratification
c	Phase velocity
g	Acceleration due to gravity
G	Magnitude of group velocity
G <sub>x</sub> , G <sub>y</sub>	Components of group velocity
H <sub>v</sub> <sup>(1)</sup>	Hankel function of the first kind of order v
k	Magnitude of wave number
K <sub>v</sub>	Modified Bessel function of the second kind of order v
M	$M^2 = \frac{N^2}{\omega^2} - 1$
N	Vaisala frequency
P	$p^2 = - M^2$
p, p', $\bar{p}$	Total, perturbation, and mean pressure, respectively
q	Source distribution
t	Time
u', v'	Components of perturbation velocity
u, v	Components of total velocity

$V$	$V = \sqrt{\bar{\rho}} v'$
$x, y$	Cartesian coordinates
$\omega$	Frequency
$\alpha, \beta$	Components of wave number
$\rho, \rho', \bar{\rho}$	Total, perturbation, and mean density, respectively
$\theta$	Direction of wave number
$\Phi$	Amplitude of $V$
$\phi$	Green's function
$\delta(x)$	Dirac delta function

## INTRODUCTION

Love (see Lamb 1945), was the first to derive the linearized equations governing small oscillations about the state of hydrostatic equilibrium in a continuously stratified fluid. The propagation of free elementary waves in continuously stratified fluids were studied by Yih (1960), Eckart (1960), Yanowitch (1962) and Tolstoy (1963). Some of these authors also considered the effects of compressibility and rotation.

In this report the internal waves generated by a time dependent local disturbance in an incompressible continuously stratified fluid are examined. Following the Love theory of wave motion in heterogeneous liquids, the motion is regarded as a small perturbation about the state of hydrostatic equilibrium. For simplicity, the equilibrium density profile is taken to be exponential whereby the coefficients of the governing equations become constants. The complications of multiple reflections from physical boundaries are avoided by considering a fluid of infinite extent. The disturbance is taken to be two-dimensional and time dependent but non-moving.

## THE GOVERNING EQUATIONS

Consider the equations governing the two-dimensional motion of an incompressible, continuously stratified fluid under gravity. Let  $(x,y)$  be a Cartesian coordinate system oriented with gravity pointing in the negative  $y$  direction, and  $(u,v)$  be the velocity components in the  $x$  and  $y$  directions respectively.

Since the fluid is incompressible, then

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} = 2\pi q(x, y, t) \quad [1]$$

where  $2\pi q$  denotes the volume rate of outflow from a source distribution in the fluid. Conservation of mass demands that

$$\frac{\partial \rho}{\partial t} + \rho \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + u \frac{\partial \rho}{\partial x} + v \frac{\partial \rho}{\partial y} = 2\pi q(x, y, t) \rho_*(x, y, t) \quad [2]$$

where  $\rho$  is the density field and  $\rho_*$  is the density of the fluid being given out (in this case,  $\rho_*$  is left open) or taken in (in this case,  $\rho_*$  must be identically equal to the ambient density) by the source distribution. Subtracting [1] from [2] yields the following 'equation of state':

$$\frac{\partial \rho}{\partial t} + u \frac{\partial \rho}{\partial x} + v \frac{\partial \rho}{\partial y} = 2\pi q(\rho_* - \rho)H(q) \quad [3]$$

where  $H$  denotes the Heaviside function. If  $\rho_*$  is controlled such that it is always equal to the ambient density, then Equation [3] simplifies to

$$\frac{\partial \rho}{\partial t} + u \frac{\partial \rho}{\partial x} + v \frac{\partial \rho}{\partial y} = 0 \quad [4]$$

In what follows, this is assumed to be always the case.

Equations [1] and [4], together with the Euler equations

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} = - \frac{1}{\rho} \frac{\partial p}{\partial x} \quad [5]$$

$$\frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} = - \frac{1}{\rho} \frac{\partial p}{\partial y} - g \quad [6]$$

where  $p$  denotes the pressure and  $g$ , the acceleration due to gravity, form a system of four equations for the four unknowns  $(u, v, p, \rho)$ .

If the motion is weak, it may be regarded as a small perturbation about the state of hydrostatic equilibrium and, as a first approximation, second order effects may be neglected. Denoting the perturbations by  $(u', v', p', \rho')$ , the hydrostatic pressure by  $\bar{p}$  and the equilibrium density by  $\bar{\rho}$ , Equations [1], [4], [5] and [6], on neglecting second order terms, become

$$\frac{\partial u'}{\partial x} + \frac{\partial v'}{\partial y} = 2\pi q \quad [7]$$

$$\frac{\partial \rho'}{\partial t} + v \frac{d\bar{\rho}}{dy} = 0$$

$$\bar{\rho} \frac{\partial u'}{\partial t} = - \frac{\partial p'}{\partial x} \quad [9]$$

$$\bar{\rho} \frac{\partial v'}{\partial t} = - \frac{\partial p'}{\partial y} - g\rho' \quad [10]$$

The coefficients are seen to be functions of  $y$  only and the resulting system is linear. From this system of equations, the following equation for  $v'$  may be uncoupled:

$$\frac{\partial^2}{\partial t^2} \left\{ \frac{\partial^2}{\partial y^2} (\bar{\rho}v') + \frac{\partial^2}{\partial x^2} (\bar{\rho}v') + \frac{N^2}{g} \frac{\partial}{\partial y} (\bar{\rho}v') + \frac{2N}{g} \frac{dN}{dy} \bar{\rho}v' \right\} + N^2 \frac{\partial^2}{\partial x^2} (\bar{\rho}v') = 2\pi \frac{\partial}{\partial y} \left( \bar{\rho} \frac{\partial^2 q}{\partial t^2} \right) \quad [11]$$

where  $N$  is the Vaisala frequency defined by

$$N = \sqrt{- \frac{g}{\bar{\rho}} \frac{d\bar{\rho}}{dy}} \quad [12]$$

This important parameter serves to characterize the capability of the medium as an internal wave carrier.

In what follows, attention is focused entirely on the vertical velocity component only. It is convenient to transform it to a new variable  $V$  according to

$$V = \sqrt{\bar{\rho}} v' \quad [13]$$

The Equation for  $V$  is

$$\begin{aligned} \frac{\partial^2}{\partial t^2} \left\{ \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial x^2} + \left( \frac{N}{g} \frac{dN}{dy} - \frac{N^2}{4g^2} \right) V \right\} + N^2 \frac{\partial^2 V}{\partial x^2} \\ = \frac{2\pi}{\sqrt{\bar{\rho}}} \frac{\partial}{\partial y} \left( \bar{\rho} \frac{\partial^2 q}{\partial t^2} \right) \end{aligned} \quad [14]$$

For an exponential density profile, viz.

$$\bar{\rho} = \rho_0 e^{-2by} \quad [15]$$

where  $\rho_0$  and  $b$  are constants, Equation [14] simplifies to

$$\frac{\partial^2}{\partial t^2} \left\{ \frac{\partial^2 V}{\partial y^2} + \frac{\partial^2 V}{\partial x^2} - b^2 V \right\} + N^2 \frac{\partial^2 V}{\partial x^2}$$

$$= 2\pi \sqrt{\rho_0} e^{by} \frac{\partial}{\partial y} \left( e^{-2by} \frac{\partial^2 q}{\partial t^2} \right) \quad [16]$$

and the Vaisala frequency  $N = \sqrt{2gb}$  is constant throughout the fluid which means that the fluid is homogeneous in character as a wave carrier but anisotropic.

#### ELEMENTARY WAVES

In this section, certain results concerning the behaviour of elementary waves in a stratified fluid are discussed. For a more general treatment in this respect, the reader is referred to Eckart (1960).

The homogeneous character of the fluid suggests that plane waves of the form

$$V = A e^{i(\alpha x + \beta y - \omega t)}$$

where  $A$  is a complex constant,  $\alpha$ ,  $\beta$  and  $\omega$  are real constants, are possible homogeneous solutions of Equation [16]. Substitution into Equation [16] reveals that this is indeed so provided that the following characteristic equation is satisfied, viz

$$\omega^2(\alpha^2 + \beta^2 + b^2) - N^2\alpha^2 = 0 \quad [17]$$

In the first place, a necessary condition for Equation [17] to have real solutions is that  $\omega$  must be less than the Vaisala frequency, i.e.

$$\frac{N^2}{\omega^2} - 1 = M^2 > 0$$

Since the Vaisala frequency measures the stability or 'stiffness' of the fluid, the above condition implies that higher frequency waves require a stiffer medium; a result which is in accord with intuition. For each  $\omega$  that is less than the Vaisala frequency, Equation [17] is satisfied by all points lying on an hyperbola of two sheets in the wave number plane. In Figure 1 the hyperbolas in the first quadrant for various  $\omega$  are shown. The family of curves is symmetric about both the vertical and horizontal axes. It is seen that, for a given  $\omega/N$ , the magnitude,  $k$ , of the wave number depends on its inclination from the horizontal axis,  $\theta$ , which cannot exceed  $\tan^{-1} M$ .

The phase velocity or, in other words, the velocity of propagation of the individual crests and troughs has the same direction as the wave number and its magnitude,  $c$ , is given by

$$c = \frac{N}{b} \cdot \sqrt{\cos^2 \theta - \frac{\omega^2}{N^2}} \quad [18]$$

Figure 2 is a polar plot of  $c$  against  $\theta$  for various values of  $\omega$ . The fact that  $c$  is a function of  $\theta$  demonstrates the anisotropic character of the medium.

The group velocity, according to which energy is propagated, is given by

$$(G_x, G_y) = \left( \frac{\partial \omega}{\partial \alpha}, \frac{\partial \omega}{\partial \beta} \right) \quad [19]$$

The direction of the group velocity, then, is normal to that hyperbola in which the wave number is found at  $(\alpha, \beta)$  and is, in general, different from the direction of the phase velocity. This behaviour is characteristic of wave motions in anisotropic media (see Lighthill 1960). At  $\theta = 0$  the two velocities have the same direction, as  $\theta$  increases the difference in direction increases and tends to  $\pi/2$  as  $\theta$  tends to  $\tan^{-1} M$ . The magnitude of the group velocity,  $G$ , is found to be

$$G = \frac{c}{a \sin \theta} \frac{\omega^2}{N^2} \sqrt{M^2 + \tan^2 \theta} \quad [20]$$

Figure 3 is a polar plot of  $G$  against  $\theta$  for various values of  $\omega$ .

It is noted, that the amplitude of the actual velocity,  $v'$ , is not constant but varies with  $y$  as  $1/\sqrt{\rho}$ . It follows that the first order kinetic energy is conserved.

#### INTERNAL WAVES GENERATED BY AN OSCILLATING SINGULARITY

Consider the waves generated by a source of constant amplitude  $A$  and simple harmonic time dependency of frequency  $\omega$  located at the origin. After a sufficient time has elapsed for the transients to die away, the dependent variables would also acquire the same time dependency. Therefore,

$$V(x,y,t) = \Phi(x,y)e^{-i\omega t} \quad [21]$$

The equation governing the amplitude function  $\Phi(x,y)$  is

$$\frac{\partial^2 \Phi}{\partial y^2} - M^2 \frac{\partial^2 \Phi}{\partial x^2} - b^2 \Phi = 2\pi A \sqrt{\rho_0} \delta(x) [\dot{\delta}(y) - b\delta(y)] \quad [22]$$

where  $\delta$  denotes the Dirac delta function and a dot denotes a differentiation.

First, consider the Green's function,  $\phi$ , of Equation [22]. By definition,  $\phi$  satisfies the following equation:

$$\frac{\partial^2 \phi}{\partial y^2} - M^2 \frac{\partial^2 \phi}{\partial x^2} - b^2 \phi = \delta(x) \delta(y) \quad [23]$$

The two dimensional Fourier integral representation of  $\phi$  is easily found to be

$$\phi(x, y) = \frac{1}{(2\pi)^2} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{e^{i(\alpha x + \beta y)}}{M^2 \alpha^2 - (\beta^2 + b^2)} d\alpha d\beta \quad [24]$$

Consider first the case where  $\omega$  is larger than the Vaisala frequency, thus  $M^2$  is real and negative. Let  $M^2 = -P^2$  where  $P$  is real and positive. Rewrite Equation [24] as

$$\phi = - \frac{1}{(2\pi)^2} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{e^{i(\alpha x + \beta y)}}{\alpha^2 P^2 + (\beta^2 + b^2)} d\alpha d\beta \quad [25]$$

Integrate first with respect to  $\alpha$ . The integrand has simple poles at

$$\alpha_{\pm} = \pm \frac{i \sqrt{\beta^2 + b^2}}{P} \quad [26]$$

The contour is closed by a large semi-circle in the upper or lower half-plane depending on whether  $x$  is positive or negative and the integral evaluated by the method of residues; the contribution from the semi-circle vanishes as its radius tends to infinity. From here on, attention is focussed on  $x > 0$  only. For  $x > 0$

$$\begin{aligned}\phi &= -\frac{1}{4\pi P} \int_{-\infty}^{+\infty} \frac{e^{i\beta y - \sqrt{\beta^2 + b^2} \frac{x}{P}}}{\sqrt{\beta^2 + b^2}} d\beta \\ &= -\frac{1}{2\pi P} \int_0^{\infty} \frac{e^{-\frac{x}{P} \sqrt{\beta^2 + b^2}}}{\sqrt{\beta^2 + b^2}} \cos \beta y d\beta\end{aligned}$$

But

$$\int_0^{\infty} \frac{e^{-\frac{x}{P} \sqrt{\beta^2 + b^2}}}{\sqrt{\beta^2 + b^2}} \cos \beta y d\beta = K_0 \left[ b \left( \frac{x^2}{P^2} + y^2 \right) \right]$$

(see Watson 1958), where  $K_\nu$  denotes the modified Bessel function of the second kind of order  $\nu$ . Therefore,

$$\phi = - \frac{1}{2\pi P} K_0 \left( b \sqrt{\frac{x^2}{P^2} + y^2} \right) \quad [27]$$

Next, consider the case where  $\omega$  is less than the Vaisala frequency, thus  $M^2$  is real and positive. The poles now lie on the real  $\alpha$  axis and the integral is indeterminate. Under such circumstances it is usual to deform the path of integration slightly such that it goes around instead of through the poles. How the path should be deformed depends on additional physical information. In the present problem it is physically obvious that  $\phi$  must be symmetric in  $x$  and that all internal waves are outgoing waves from the source at the origin. With this considerations in mind it is found that the path of integration should be displaced above the negative pole and below the positive pole by small semi-circles. The integral is then evaluated as in the previous case to give

$$\begin{aligned} \phi &= \frac{i}{4\pi M} \int_{-\infty}^{+\infty} \frac{e^{i(\beta y + \sqrt{\beta^2 + b^2} \frac{x}{M})}}{\sqrt{\beta^2 + b^2}} d\beta \\ &= \frac{1}{2\pi M} \int_0^{\infty} \frac{e^{i \frac{x}{M} \sqrt{\beta^2 + b^2}}}{\sqrt{\beta^2 + b^2}} \cos \beta y d\beta \end{aligned}$$

But

$$\int_0^{\infty} \frac{e^{-\frac{x}{M}\sqrt{\beta^2+b^2}}}{\sqrt{\beta^2+b^2}} \cos \beta y \, d\beta = \frac{\pi 1}{2} H_0^{(1)} \left( b \sqrt{\frac{x^2}{M^2} - y^2} \right)$$

(see Watson 1958), where  $H_\nu^{(1)}$  denotes the Hankel function of the first kind of order  $\nu$ . Therefore,

$$\phi = -\frac{1}{4M} H_0^{(1)} \left( b \sqrt{\frac{x^2}{M^2} - y^2} \right) \quad [28]$$

The motion generated by the oscillating source can be expressed in terms of  $\phi$  as

$$v'(x,y,t) = e^{-i\omega t} 2\pi A e^{by} \left( \frac{\partial \phi}{\partial y} - b\phi \right) \quad [29]$$

First examine the case where  $\omega$  is greater than the Vaisala frequency. Substituting Equation [27] into [29] the following equation for  $v'$  is obtained

$$v' = e^{-i\omega t} \frac{Ab e^{by}}{P} \left[ \frac{y}{\sqrt{\frac{x^2}{P^2} + y^2}} K_1 \left( b \sqrt{\frac{x^2}{P^2} + y^2} \right) + K_0 \left( b \sqrt{\frac{x^2}{P^2} + y^2} \right) \right] \quad [30]$$

As the stratification tends to zero, i.e. as  $b \rightarrow 0$

$$v' \rightarrow e^{-i\omega t} A \frac{y}{x^2 + y^2} \quad [31]$$

This limit is the motion generated by an oscillating source in a homogeneous fluid. For finite  $b$ , the standing wave character of the motion is preserved because both  $K_0$  and  $K_1$  are real for real arguments. However, the purely antisymmetric character of the vertical velocity component with respect to  $y$  is destroyed. It is noted that no progressive internal wave is generated. This is in accord with the result of the last section which excludes the possibility of progressive waves whose frequency is greater than the Vaisala frequency.

Next consider the case where  $\omega$  is less than the Vaisala frequency. Substitution of Equation [28] into [29] results in the following equation for  $v'$ :

$$v' = -e^{-i\omega t} \frac{\pi A b e}{2M} \left[ \frac{y}{\sqrt{\frac{x^2}{M^2} - y^2}} H_1^{(1)} \left( b \sqrt{\frac{x^2}{M^2} - y^2} \right) - H_0^{(1)} \left( b \sqrt{\frac{x^2}{M^2} - y^2} \right) \right] \quad [32]$$

The solution behaves very differently depending on whether  $x^2/M^2 - y^2$  is positive or negative. Consider first the region which satisfies the inequality

$$\frac{x^2}{M^2} - y^2 < 0$$

Rewrite Equation [32] as

$$v' = -e^{-i\omega t} \frac{\pi A b e}{2M} \left[ \frac{y}{i \sqrt{y^2 - \frac{x^2}{M^2}}} H_1^{(1)} \left( i b \sqrt{y^2 - \frac{x^2}{M^2}} \right) - H_0^{(1)} \left( i b \sqrt{y^2 - \frac{x^2}{M^2}} \right) \right] \quad [33]$$

Using the following relation, (see Watson 1958)

$$K_\nu(\xi) = \frac{1}{2}\pi i e^{\frac{1}{2}\nu\pi i} H_\nu^{(1)}(i\xi)$$

where  $\xi$  is real and positive, Equation [33] becomes

$$v' = e^{-i(\omega t + \pi/2)} \frac{Ab e^{by}}{M} \left[ \frac{y}{\sqrt{y^2 - \frac{x^2}{M^2}}} K_1 \left( b \sqrt{y^2 - \frac{x^2}{M^2}} \right) + K_0 \left( b \sqrt{y^2 - \frac{x^2}{M^2}} \right) \right] \quad [34]$$

It is seen from the behaviour of  $K_0$  and  $K_1$  that the motion in this region is a standing wave also, but, here, the amplitude of the motion is singular on the lines  $y^2 = x^2/M^2$  instead of just at the origin where the source is located.

In the region

$$\frac{x^2}{M^2} - y^2 > 0$$

the arguments of the Hankel functions appearing in Equation [32] are real. From the known behaviours of the Hankel function for real arguments it is seen that internal gravity waves are present in this region.

This peculiar pattern of motion is related to the behaviour of elementary waves discussed in the previous section. The periodic source may be regarded as an integral of elementary waves of all wave numbers and of a given frequency  $\omega$ . It was seen that only those waves whose number lie on a particular hyperbola are progressive. The asymptotes of the hyperbola make an angle of  $\tan^{-1} M$  with the horizontal so that the direction of the wave number cannot exceed  $\tan^{-1} M$ . However, the group velocity or the velocity of propagation is normal to the hyperbola so that the region in which progressive waves are found are sectors bounded by the lines  $y^2 = x^2/M^2$  as shown in Figure 4. Except for the small wave number range the group velocities of the elementary waves are approximately in the direction of either one of these lines. Thus, there is a large concentration of energy in them.

#### INTERNAL WAVES GENERATED BY AN INSTANTANEOUS SOURCE

In this section, the transient problem of the motion due to a source at the origin with a time dependency of the form  $\delta(t)$  is considered. The appropriate forcing function for Equation [16] is

$$2\pi A \rho_0 \delta(x) \ddot{\delta}(t) [\dot{\delta}(y) - b\delta(y)]$$

Let  $\phi$  be the solution of the equation

$$\frac{\partial^2}{\partial t^2} \left\{ \frac{\partial^2 \phi}{\partial y^2} + \frac{\partial^2 \phi}{\partial x^2} - b^2 \phi \right\} + N^2 \frac{\partial^2 \phi}{\partial x^2} = \delta(x) \delta(y) \delta'(t) \quad [35]$$

therefore, the vertical velocity component  $v'$  is given by

$$v' = 2\pi A e^{by} \left( \frac{\partial \phi}{\partial y} - b\phi \right) \quad [36]$$

The Fourier integral representation of  $\phi$  is easily found to be

$$\phi(x, y, t) = \frac{1}{(2\pi)^3} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{e^{i(ax + \beta y + \omega t)}}{\left( \frac{N^2}{\omega^2} - 1 \right) \alpha^2 - (\beta^2 + b^2)} d\alpha d\beta d\omega \quad [37]$$

Integrate first with respect to  $\omega$ . The initial condition of the fluid is the state of hydrostatic equilibrium. With this consideration in mind, the path of integration is displaced slightly below the real axis. The contour is closed by a large semi-circle in the upper half plane and the integral evaluated by residues to give

$$\phi = \frac{\pi i}{(2\pi)^3} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{Na e^{i(\alpha x + \beta y)}}{(\alpha^2 + \beta^2 + b^2)^{3/2}} \left( e^{-i \frac{Nat}{\sqrt{\alpha^2 + \beta^2 + b^2}}} - e^{+i \frac{Nat}{\sqrt{\alpha^2 + \beta^2 + b^2}}} \right) d\alpha d\beta$$

$$= \frac{\pi i}{8\pi^2} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{Na e^{itf_1}}{(\alpha^2 + \beta^2 + b^2)^{3/2}} d\alpha d\beta -$$

$$\frac{\pi i}{8\pi^2} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \frac{Na e^{itf_2}}{(\alpha^2 + \beta^2 + b^2)^{3/2}} d\alpha d\beta = I_1 + I_2$$

where

$$f_1 = \alpha \frac{x}{t} + \beta \frac{y}{t} - \frac{Na}{\sqrt{\alpha^2 + \beta^2 + b^2}}$$

$$f_2 = \alpha \frac{x}{t} + \beta \frac{y}{t} + \frac{Na}{\sqrt{\alpha^2 + \beta^2 + b^2}}$$

The points of stationary phase of the integrals occur at

$$\frac{\partial f_1}{\partial \alpha} = 0, \quad \frac{\partial f_1}{\partial \beta} = 0 \quad [38]$$

As before we consider only  $x > 0$ . For  $x/t > 0$  only  $I_1$  can have points of stationary phase. Given a pair  $(x/t, y/t)$  the real solutions of the following simultaneous equations

$$\frac{x}{t} = \frac{N(\beta^2 + b^2)}{(\alpha^2 + \beta^2 + b^2)^{3/2}} \quad [39]$$

$$\frac{y}{t} = \frac{N\alpha\beta}{(\alpha^2 + \beta^2 + b^2)^{3/2}} \quad [40]$$

are points of stationary phase. These two equations may be combined to give

$$\left(\frac{y}{x}\right)^2 = \frac{\beta^2}{\beta^2 + b^2} \left\{ \left[ \frac{2gb}{\frac{x^2}{t^2} (\beta^2 + b^2)} \right]^{1/3} - 1 \right\} \quad [41]$$

For a given  $y/x$ , the above equation passes from a situation of no real solution to a situation of two real solutions as  $x/t$  tends to zero from a very large value. The transition where the equation has only one real solution, therefore, corresponds to the maximum distance from the origin at a given time where the disturbance is appreciable if we keep terms of  $O(1/t)$  only. Hence, the wave front is the locus of the transition points for all  $y/x$ . The equation of the wave front is obtained by eliminating  $\beta$  from Equation [41] and its derivative with respect to  $\beta$ . The resulting equation is

$$\left(\frac{x}{t} \sqrt{\frac{2g}{b}}\right)^{2/3} = \frac{B^4}{3\left(\frac{y}{x}\right)^2 (1 + B^2)^{1/3}} \quad [42]$$

where

$$B^2 = \frac{\sigma^2 + \sqrt{4\sigma^4 + 3\sigma^2}}{1 + \sigma^2}$$


$$\sigma = y/x$$

Since Equation [36] involves differentiation of  $\phi$  with respect to  $y$  only, the above asymptotic property of the wave front also applies to that of the internal waves generated by the instantaneous source.

Equation [42] is plotted in Figure 5. It is seen that the wave fronts for different times are geometrically similar.

REFERENCES

- Eckart, C., 1960, "Hydrodynamics of Oceans and Atmospheres," Pergamon Press.
- Lamb, H., 1945, "Hydrodynamics," Dover Publication.
- Lighthill, M. J., 1960, Trans. Royal Soc., 252A, 397-430.
- Tolstoy, I., 1963, Rev. of Mod. Phy. 35, 207-230.
- Watson, G. N., 1958, "Theory of Bessel Functions," Cambridge University Press.
- Yanowitch, M., 1962, Comm. of Pure and App. Math., XV, 45-61.
- Yih, C. S., 1960, J. Fluid Mech. 8, 481-508.



LEGEND

LOCUS OF THE ALLOWABLE WAVE NUMBERS FOR A GIVEN DIMENSIONLESS FREQUENCY  $\frac{\omega}{N}$

ASYMPTOTES OF THE HYPERBOLAS

INCLINATION OF THE WAVE NUMBER TO THE HORIZONTAL

$\frac{1}{b}$  x (MAGNITUDE OF THE WAVE NUMBER)


—————

- - - - -

POLAR ANGLE  $\theta$

RADIAL DISTANCE =

FIGURE 1 - ALLOWABLE WAVE NUMBERS FOR DIFFERENT FREQUENCIES



LEGEND


— THE MAGNITUDE OF THE PHASE VELOCITY FOR A GIVEN DIMENSIONLESS FREQUENCY  $\frac{\omega}{N}$

- - - MAXIMUM  $\theta$  FOR A GIVEN DIMENSIONLESS FREQUENCY

POLAR ANGLE  $\theta$  = INCLINATION OF THE WAVE NUMBER TO THE HORIZONTAL

RADIAL DISTANCE =  $\frac{b}{N} \times$  (MAGNITUDE OF THE PHASE VELOCITY)

FIGURE 2 - DEPENDENCE OF THE PHASE VELOCITY ON THE DIRECTION OF THE WAVE NUMBER



LEGEND

— THE MAGNITUDE OF THE GROUP VELOCITY FOR A GIVEN DIMENSIONLESS FREQUENCY,  $\frac{\omega}{N}$

- - - MAXIMUM  $\theta$  FOR A GIVEN DIMENSIONLESS FREQUENCY

POLAR ANGLE,  $\theta$  = INCLINATION OF THE WAVE NUMBER TO THE HORIZONTAL

$\frac{b}{N} \times$  (MAGNITUDE OR THE GROUP RADIAL DISTANCE = VELOCITY)

FIGURE 3 - DEPENDENCE OF THE MAGNITUDE OF THE GROUP VELOCITY ON DIRECTION OF THE WAVE NUMBER




FIGURE 4 - INTERNAL MOTION DUE TO AN OSCILLATING SOURCE WHOSE FREQUENCY IS LESS THAN THE VAISALA FREQUENCY




FIGURE 5—THE WAVE FRONT OF AN INSTANTANEOUS DISTURBANCE IN AN EXPONENTIALLY STRATIFIED FLUID OF INFINITE EXTENT

UNCLASSIFIED

## Security Classification

DOCUMENT CONTROL DATA - R&D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
HYDRONAUTICS, Incorporated, Pindell School Road, Howard County, Laurel, Maryland		UNCLASSIFIED
		2b. GROUP
3. REPORT TITLE		
ON INTERNAL GRAVITY WAVES GENERATED BY LOCAL DISTURBANCES		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
Technical Report		
5. AUTHOR(S) (Last name, first name, initial)		
Wong, K. K.		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
February 1965	27	7
8a. CONTRACT OR GRANT NO.	8a. ORIGINATOR'S REPORT NUMBER(S)	
Nonr 3688(00) NR 220-016	Technical Report 231-5	
b. PROJECT NO.	8b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
231		
c.		
d.		
10. AVAILABILITY/LIMITATION NOTICES		
Qualified requesters may obtain copies of this report from DDC		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
		Office of Naval Research Department of the Navy
13. ABSTRACT		
<p>The internal waves generated by a time dependent local disturbance in an incompressible continuously stratified fluid are examined. Following the Love theory of wave motion in heterogeneous liquids, the motion is regarded as a small perturbation about the state of hydrostatic equilibrium. For simplicity, the equilibrium density profile is taken to be exponential. It was found that, for a disturbance having a simple harmonic time dependency, the ratio of the disturbance frequency to the Vaisala frequency determines whether progressive internal waves are present or not and in which regions. The propagation of the wave front of the internal waves generated by an instantaneous source was also analyzed.</p>		

DD FORM 1473  
1 JAN 64UNCLASSIFIED  
Security Classification

## Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
1. Internal wave						
2. Stratified fluids						
3. Water waves						

## INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

HYDRONAUTICS, Incorporated

DISTRIBUTION LIST  
(Contract Nonr-3688(00))

Chief of Naval Research		Commanding Officer	
Department of the Navy		Office of Naval Research	
Washington, D. C. 20360		Branch Office	
Attn: Codes 438	2	1000 Geary Street	
416	1	San Francisco 9, California	1
461	1		
104	1	Commanding Officer and Director	
406T	1	David Taylor Model Basin	
427	1	Washington 7, D. C.	1
402	1		
446	1	Director	
466	1	U. S. Naval Research Laboratory	
		Washington, D. C. 20360	
Commanding Officer		Attn: Codes 2000	1
Office of Naval Research		5500	1
Branch Office		6100	1
495 Summer Street		2020	1
Boston 10, Massachusetts	1	2027	1
		5300	1
Commanding Officer		7300	1
Office of Naval Research		4000	1
Branch Office		5000	1
219 S. Dearborn Street		7350	1
Chicago, Ill. 60604	1		
		Chief, Bureau of Ships	
Commanding Officer		Department of the Navy	
Office of Naval Research		Washington, D. C. 20360	
Branch Office		Attn: Codes 305	1
207 West 24th Street		335	1
New York 11, New York	1	370	1
		374	1
Commanding Officer		421	1
Office of Naval Research			
Branch Office		Chief, Bureau of Naval Weapons	
1030 East Green Street		Department of the Navy	
Pasadena 1, California	1	Washington, D. C. 20360	
		Attn: Codes RU-2	1
		RUDC-43	1
		RRRE-4	1

HYDRONAUTICS, Incorporated

-2-

Chief of Naval Operations Department of the Navy Washington, D. C. 20360		Institute of Mathematical Sciences New York University 25 Waverly Place New York 3, New York	
Attn: Codes OP-71B	1	Attn: Prof. J. J. Stoker	1
OP-07C	1		
OP-07T	1		
OP-717	1	New York University Dept. of Oceanography University Heights New York 53, New York	
Commanding Officer U. S. Naval Ordnance Laboratory Corona, California	1	Attn: Prof. J.W. Pierson, Jr	1
Commander U. S. Naval Ordnance Test Station Pasadena Annex 3202 E. Foothill Boulevard Pasadena 8, California	1	The Johns Hopkins University Baltimore 18, Maryland Attn: Prof. O.M. Phillips	1
Superintendent U. S. Naval Postgraduate School Monterrey, California Attn: Library	1	California Institute of Tech. Pasadena 4, California Prof. T. Y. Wu	1
Commanding Officer and Director U. S. Naval Electronic Laboratory San Diego 52, California	1	Director Scripps Institution of Oceanography Applied Oceanography Group University of California La Jolla, California	1
Defense Documentation Center Cameron Station, 5010 Duke St. Alexandria, Virginia	20	Iowa Institute of Hydraulic Research State University of Iowa Iowa City, Iowa Attn: Prof. H. Rouse	1
Fluid Mechanics Section National Bureau of Standards Washington, D. C. 20360 Attn: Dr. O.B. Schubauer	1	Harvard University Cambridge 38, Massachusetts Attn: Prof. G.F. Carrier	1
National Research Council Montreal Road Ottawa 2, Canada Attn: Mr. E. S. Turner	1	Massachusetts Institute of Technology Cambridge 39, Massachusetts Attn: Prof. J.F. Kennedy	1

HYDRONAUTICS, Incorporated

-3-

Stevens Institute of Technology		Dr. A V. Hershey	
Davidson Laboratory		Computation and Exterior	
Hoboken, New Jersey		Ballistics Laboratory	
Attn: Mr. D. Savitsky	1	U. S. Naval Weapons Laboratory	
Dr. S. J. Lukasik	1	Dahlgren, Virginia	1
Director		Commander	
Woods Hole Oceanographic		U. S. Naval Ordnance Laboratory	
Institute		White Oak, Maryland	
Woods Hole, Massachusetts	1	Attn: Dr. S. J. Raff	
		(DU 5-224)	1
Dr. B. Zarwyn		Dr. E. Montrell	
Airborne Instruments Laboratory		Institute for Defense Analysis	
Division of Cutler-Hammer Inc.		1666 Connecticut Avenue, N.W.	
Walt Whitman Road, Melville,		Washington, D. C.	1
Long Island, New York	1		
Mr. Myron J. Block		Director, Special Projects Office	
Block Associates, Inc.		Department of the Navy	
385 Putnam Avenue		Washington, D. C. 20360	
Cambridge 39, Massachusetts	1	Attn: Code SP-001	1
Director		Dr. I. Katz	
Hudson Laboratories		Applied Physics Laboratory	
Dobbs Ferry, New York	1	Johns Hopkins University	
		Baltimore 18, Maryland	1
Dr. Savet		Dr. Harvey Brooks	
American Bosch Arma Corporation		School of Applied Science	
Arma Division		Harvard University	
Garden City, New York	1	Cambridge, Massachusetts	1
HYDRONAUTICS, Incorporated		Arthur D. Little, Inc.	
Pindell School Road		35 Acorn Park	
Howard County		Cambridge 40, Massachusetts	
Laurel, Maryland		Attn: Trident Library	1
Attn: Mr. P. Eiserberg, Pres.	1		
Mr. M.P. Tulin, V. Pres.	1		
Dr. L. L. Higgins			
Space Technology Lab. Inc.			
Canoga Park Facility			
8433 Fallbrook Avenue			
Canoga Park, California	1		

HYDRONAUTICS, Incorporated

-4-

Dr. R. W. Stewart c/o Pacific Naval Laboratory Esquimalt, Vancouver, British Columbia Canada	1
Professor M. J. Lighthill Imperial College London, England	1
Commanding Officer U. S. Naval Air Development Center Johnsville, Pennsylvania	1
Mr. Richard Burakat ITEK Corporation 10 Maguire Road Lexington 73, Massachusetts	1
Dr. J. Kotik Technical Research Group, Inc. Route 110 Melville, New York	1