

UNCLASSIFIED

AD NUMBER
AD465240
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; Apr 1965. Other requests shall be referred to U.S. Army Ballistic Research Laboratory, Aberdeen Proving Ground, MD.
AUTHORITY
USABRL ltr, 8 May 1967

THIS PAGE IS UNCLASSIFIED

CATALOGED BY: DDC

AS AD 117

465240 BRL R 1278

BRL

REPORT
NO. 1278

AD

BRL R 1278

ASYMPTOTIC SOLUTIONS FOR THE EXPANSION OF GAS INTO VACUUM

By E. H. Wedemeyer

APRIL 1965

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

DDC
RECEIVED
JUL 1 1965
DDC-IRA E

U. S. ARMY MATERIEL COMMAND
BALLISTIC RESEARCH LABORATORIES
ABERDEEN PROVING GROUND, MARYLAND

Destroy this report when it is no longer needed.
Do not return it to the originator.

DDC AVAILABILITY NOTICE

Qualified requesters may obtain copies of this report from DDC.

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

BALLISTIC RESEARCH LABORATORIES

REPORT NO. 1278

APRIL 1965

ASYMPTOTIC SOLUTIONS FOR THE EXPANSION OF GAS INTO VACUUM

E. H. Wedemeyer

Exterior Ballistics Laboratory

RDT & E Project No. 1A222901A201

ABERDEEN PROVING GROUND, MARYLAND

Previous page was blank, therefore not filmed.

BALLISTIC RESEARCH LABORATORIES

REPORT NO. 1278

EHWedemeyer/kel
Aberdeen Proving Ground, Md.
April 1965

ASYMPTOTIC SOLUTIONS FOR THE EXPANSION OF GAS INTO VACUUM

ABSTRACT

When a spherical or cylindrical gas cloud expands into vacuum, the resultant flow becomes asymptotically self-similar and independent of the original cloud diameter. An analytic solution for the asymptotic density profiles, given by Mirels and Mullen [4] is re-examined and a new solution proposed, which gives better agreement with numerical computations.

Some applications of the asymptotic solutions are discussed.

Previous page was blank, therefore not filmed.

TABLE OF CONTENTS

	PAGE
ABSTRACT	3
1. INTRODUCTION	7
2. ASYMPTOTIC NATURE OF THE FLOW	8
3. ANALYTICAL SOLUTION	10
4. LATERAL EXPANSION OF A HYPERSONIC JET BOUNDED BY VACUUM	19
5. MEASUREMENT OF PLASMA TEMPERATURE	21
APPENDIX A	23
REFERENCES	25

Previous page was blank, therefore not filmed.

1. INTRODUCTION

The investigations reported here were initiated by a paper "Translational Equilibration of Wires Exploded in Vacua," by O. H. Zinke et al [1]*. The authors of this paper made use of an unsteady spherical expansion of ionized gas in order to infer the temperature of the gas from the characteristics of the expansion. The method described by Zinke can be a powerful tool, provided that the unsteady flow resulting from an expanding gas cloud is understood theoretically. It is believed that the theoretical solution presented here straightens out some discrepancies observed by Zinke. Further remarks on this are given in section 5. The solution for the unsteady cylindrical expansion can also be applied to a related problem, the expansion of a hypersonic jet into vacuum.

In the above mentioned and other cases, the main interest is in the asymptotic flow field after long times. For large times the density distribution of the expanding gas cloud approaches an asymptotic similarity form, which is independent of the original diameter of the cloud.

It can be shown, however, that the shape of the asymptotic density profile depends on the initial density distribution of the gas before expansion. In the following an attempt is made to derive approximate solutions for the asymptotic density profiles for the case that the initial density distribution is uniform. Expansions with plane, cylindrical and spherical symmetry will be considered.

*Numbers in brackets denote references listed on page 25.

2. ASYMPTOTIC NATURE OF THE FLOW

In the following it is assumed that the flow is isentropic and that the gas can be treated as a continuum; i.e., the mean free path is very small compared with the initial diameter of the gas cloud. Then the equations of motion for flows of cylindrical ($\sigma = 1$) or spherical ($\sigma = 2$) symmetry are respectively:

$$\frac{\partial \bar{p}}{\partial \bar{t}} + \frac{\partial \bar{p}}{\partial \bar{r}} \bar{u} + \sigma \frac{\bar{p} \bar{u}}{\bar{r}} = 0 \quad (1a)$$

$$\frac{\partial \bar{u}}{\partial \bar{t}} + \bar{u} \frac{\partial \bar{u}}{\partial \bar{r}} + \frac{1}{\bar{p}} \frac{\partial \bar{p}}{\partial \bar{r}} = 0 \quad (1b)$$

$$(\bar{p}/p_0) = (\bar{\rho}/\rho_0)^\gamma \quad (1c)$$

ρ_0 and p_0 are taken as initial density and pressure if the density is initially uniform, or as the initial values at $r = 0$ if the density is not uniform. It is assumed that the gas is at rest initially and inclosed within a cylinder or sphere of radius R_0 .

For the following we introduce nondimensional quantities:

$$\rho = \frac{\bar{\rho}}{\rho_0} \quad u = \frac{\bar{u}}{a_0} \quad r = \frac{\bar{r}}{R_0} \quad t = \frac{\bar{t} a_0}{R_0} \quad (2)$$

where a_0 is the sound speed of the quiescent gas and found by

$$a_0^2 = \gamma \frac{p_0}{\rho_0} \quad (3)$$

With these nondimensional quantities the equations of motion are:

$$\frac{\partial \rho}{\partial t} + \frac{\partial \rho u}{\partial r} + \sigma \frac{\rho u}{r} = 0 \quad (4a)$$

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial r} + \rho^{\gamma-2} \frac{\partial \rho}{\partial r} = 0 \quad (4b)$$

For the case that the density is initially uniform, the boundary conditions are:

$$\text{at } t = 0; 0 < r < 1: \quad \rho = 1, u = 0$$

$$1 < r: \quad \rho = 0 \quad (5)$$

$$\text{at } t \geq 0, r = 0: \quad u = 0$$

For the case that $\sigma = 0$, the Equations (4) together with boundary conditions (5) describe the one-dimensional (plane) expansion of a slab of gas extending either from $r = -1$ to $r = +1$ with $r = 0$ a plane of symmetry or from $r = 0$ to $r = 1$ with a fixed wall at $r = 0$. In a few cases, exact solutions of Equations (4a, 4b) can be obtained. These solutions which we shall discuss in the next paragraph, will serve as a guide to construct approximate asymptotic solutions for the more general case.

Here we are concerned with the asymptotic nature of the flow after long times; i.e., for times large compared to R_0/a_0 . For times which are large in this sense, the original gas cloud has expanded many times and the pressure term $\rho^{\gamma-2} \frac{\partial \rho}{\partial r}$ in Equation (4b) becomes relatively small. After neglecting the pressure term Equation (4b) is solved by

$$u = r/t \quad (6)$$

With the solution (6) for u , Equation (4a) has the solution:

$$\rho t^{\sigma+1} = \text{const } f(r/t) \quad (7)$$

where $f(r/t)$ is an arbitrary function of the one variable r/t . It can be seen from Equation (6) and (7) that the flow becomes asymptotically self-similar; i.e., the density and velocity distributions at different times are similar and differ only by scaling factors. While this similarity form of Equations (6) and (7) means a certain simplification of the mathematical treatment, the problem of finding the particular solution for $f(r/t)$ is still

very difficult for the following reason. Since any solution of the form (6) and (7) solves the equations of motion asymptotically, any of these solutions represents a possible asymptotic flow, each one belonging to different initial conditions. Thus, one has to relate a given initial density distribution to a corresponding $f(r/t)$ and it seems that this can be done exactly only by computing the whole flow history from the beginning of the expansion up to large times where the flow becomes self-similar. A computation of this sort can be performed numerically, using the finite step method of characteristics. It must be considered, however, that any finite step computation may become increasingly inaccurate when carried out for too many steps, due to the accumulation of errors. The number of steps necessary in order to reach the asymptotic similarity-solution must be enormously large. In spite of this, numerical computations for uniform initial densities and various values of γ and σ have been carried out by several authors [2,3] and it appears that the results of these computations are surprisingly accurate.

3. ANALYTICAL SOLUTION

A first attempt toward an analytical solution has been made by Mirels and Mullen [4], however the solution does not show satisfactory agreement with numerical results [2,3].

The solution proposed here is obtained by only slight modification of the view of Mirels and Mullen [4] but it gives, nevertheless, far better agreement with numerical results. The procedure used to derive an analytical expression for the asymptotic flow consists in finding a valid generalization

of certain exact solutions. As has been mentioned before, in a few cases the Equations (4a), (4b) can be solved exactly. These cases are:

1. The one-dimensional ($\sigma = 0$) expansion of an initially uniform slab of gas.
2. The similarity flows for $\sigma = 0, 1, 2$. These flows are strictly self-similar at all times, but the initial density-distribution is not uniform.

The characteristics solution for one-dimensional expansion from uniform initial conditions (case 1) is given in [5] (Courant and Friedrichs) and is discussed in greater detail in [6,7]. In the asymptotic limit, this solution simplifies to:

$$\rho t = \frac{\gamma-1}{\sqrt{\pi}} \frac{\Gamma(N + \frac{1}{2})}{\Gamma(N)} \left[1 - \left(\frac{\gamma-1}{2} \frac{r}{t} \right)^2 \right]^{\frac{3-\gamma}{2(\gamma-1)}} \quad (8a)$$

$$u = r/t \quad (8b)$$

where $N = \frac{\gamma+1}{2(\gamma-1)}$ and Γ denotes the gamma-function.

The similarity flows have been investigated in References [8] and [9]. If $R(t)$ is the position of the leading edge of the expansion, one can define a similarity variable

$$y = r/R(t) \quad (9)$$

so that the gas is confined in $0 \leq y \leq 1$.

Assuming for the self-similar flow a form $\rho = R^{-(\sigma+1)} f(y)$; $v = \frac{dR}{dt} \cdot g(y)$ and substituting into Equations (1a, b) one obtains

$$\rho = R^{-(\sigma+1)} (1 - y^2)^{\frac{1}{\gamma-1}} \quad (10)$$

and $u = \frac{dR}{dt} \cdot y \quad (11)$

where R satisfies the differential equation

$$\frac{dR}{dt} = \frac{1}{\sqrt{\sigma+1}} \cdot \frac{2}{\gamma-1} [1 - R^{-(\sigma+1)(\gamma-1)}]^{\frac{1}{2}} \quad (12)$$

The constants of integration are so determined, that for

$$t = 0: R = 1; \frac{dR}{dt} = 0.$$

In the asymptotic limit one finds

$$R = \frac{1}{\sqrt{\sigma+1}} \cdot \frac{2}{\gamma-1} \cdot t \quad (13)$$

and Equations (10) and (11) become asymptotically:

$$\rho t^{\sigma+1} = \left(\sqrt{\sigma+1} \frac{(\gamma-1)}{2} \right)^{\sigma+1} \left[1 - \left(\sqrt{\sigma+1} \cdot \frac{\gamma-1}{2} \frac{r}{t} \right)^2 \right]^{\frac{1}{\gamma-1}} \quad (14a)$$

$$u = r/t \quad (14b)$$

A comparison of Equation (8) and Equation (14) shows that both formulas are quite similar and of the form:

$$\rho t^{\sigma+1} = D \left[1 - \left(\frac{r}{Vt} \right)^2 \right]^B \quad (15a)$$

$$u = r/t \quad (15b)$$

where D , V and B are constants. Thus, we assume, in accordance with Mirels and Mullen [4], that the asymptotic solution for uniform initial density and $\sigma \neq 0$ is also of the form (15); so that it remains only to determine the constants D , V and B . If one of the three constants is given, say V , then the other two can be determined from the conditions, that mass and total energy are conserved. In our nondimensional variables these two conditions can be expressed as:

$$\int_0^{Vt} \rho r^\sigma dr = \frac{1}{\sigma+1} ; \quad (16)$$

$$\int_0^{Vt} \frac{\rho}{2} u^2 r^\sigma dr = \frac{1}{\sigma+1} \cdot \frac{1}{\gamma(\gamma-1)} . \quad (17)$$

Equation (16) equates the mass of the expanded gas to the mass at $t = 0$ and Equation (17) does the same for the energy. Note, that for $t \rightarrow \infty$ the thermal energy vanishes so that only the integral over the kinetic energy density $\rho/2 u^2$ is taken at the left side of Equation (17), while for $t = 0$ the kinetic energy is zero and the thermal energy per unit mass is:

$$C_v T_0 = \frac{a_0^2}{\gamma(\gamma-1)} .$$

Introducing the new variable $y = \frac{r}{Vt}$ and inserting (15a, b) into (16) and (17) gives:

$$DV^{\sigma+1} \int_0^1 (1 - y^2)^B y^\sigma dy = \frac{1}{\sigma+1} ; \quad (18)$$

$$DV^{\sigma+3} \int_0^1 (1 - y^2)^B y^{\sigma+2} dy = \frac{2}{\sigma+1} \cdot \frac{1}{\gamma(\gamma-1)} . \quad (19)$$

The integrals in Equations (18) and (19) can be transformed into Eulerian integrals of the first kind and expressed in terms of gamma functions. It is, e.g.,

$$\int_0^1 (1 - y^2)^B y^\sigma dy = \frac{1}{2} \frac{\Gamma(B+1) \Gamma(\frac{\sigma+1}{2})}{\Gamma(B+1 + \frac{\sigma+1}{2})} . \quad (20)$$

Solving Equation (18) and (19) for B and D one obtains:

$$B = \left[\frac{\gamma(\gamma-1)}{2} V^2 - 1 \right] \left(\frac{\sigma+1}{2} \right) - 1 ; \quad (21)$$

$$D = \frac{1}{V^{\sigma+1}} \cdot \frac{2}{\sigma+1} \cdot \frac{\Gamma(B+1 + \frac{\sigma+1}{2})}{\Gamma(B+1) \Gamma(\frac{\sigma+1}{2})} \quad (22)$$

Now only one constant, V , remains to be determined and at this point the present analysis differs from Mirels and Mullen [4]. From Equation (15a) it can be seen, that V is the velocity of the leading edge of the expansion (in units of a_0).

Now Mirels and Mullen maintain that for uniform initial density the velocity of the leading edge should be the same as for one-dimensional flow, and therefore $V = \frac{2}{\gamma-1}$, regardless of the value of σ . In other words, the geometry, whether spherical or cylindrical, does not influence the leading edge velocity. The arguments given in favor of this are difficult and not fully convincing, but even if they were true they would not demand for V a value of $\frac{2}{\gamma-1}$. It is understood that Equation (15a) can be only an approximate formula which must not reflect every detail of an exact solution. It could be, that a minute fraction of the total mass "escapes" with the leading edge velocity $\frac{2}{\gamma-1}$ and that the bulk of the remaining mass expands with an "effective" leading edge velocity less than that. In order to have the best approximation, valid for the bulk of the mass, it would then be advisable to pick a smaller value for V . This is also suggested by Equation (14a) for the similarity flow.

Here one observes that the geometry does have an influence on the leading edge velocity expressed by the geometrical factor $\frac{1}{\sqrt{\sigma+1}}$. If the geometrical factor $\frac{1}{\sqrt{\sigma+1}}$ has an influence in the case of similarity flow, it is reasonable to assume that it has an analogous influence in the case of uniform initial density. This would suggest:

$$V = \frac{1}{\sqrt{\sigma+1}} \cdot \frac{2}{\gamma-1} , \quad (23)$$

for both similarity and uniform initial density flow.

A more rigorous derivation of (23) is given in Appendix A.

With (23) inserted into (21) and (22) the constants B and D are:

$$B = \left(\frac{1}{\gamma-1}\right) - \left(\frac{\sigma+1}{2}\right) , \quad (24)$$

$$D = \left(\frac{2}{\sigma+1}\right) \left(\sqrt{\sigma+1} \frac{(\gamma-1)}{2}\right)^{\sigma+1} \frac{\Gamma(B+1 + \frac{\sigma+1}{2})}{\Gamma(B+1) \Gamma(\frac{\sigma+1}{2})} , \quad (25)$$

and from Equation (15a),

$$\rho t^{\sigma+1} = D \left[1 - \left(\frac{\sqrt{\sigma+1}}{2} (\gamma-1) \frac{r}{t} \right)^2 \right]^B . \quad (26)$$

For $\sigma = 0$ Equation (8a) is recovered from Equation (26); i.e., Equation (26) is exact for $\sigma = 0$. For $\sigma \neq 0$ the formula (26) is different from Mirels and Mullen's, who obtain larger values for the exponent B and the factor D as a result of the larger V values they assumed. In fact, for $\sigma \geq 1$ the exponents B obtained by Mirels and Mullen are even larger than the exponent for similarity flow, which is $\frac{1}{\gamma-1}$ (Equation 14a). A consequence of this is that the resultant density distribution is less full than the corresponding distribution for similarity flow. This is an odd result, considering that the uniform density profile is fuller initially and one ought to expect that it also remains fuller than the similarity profile after expansion. The exponents obtained by Mirels and Mullen are so large (e.g. for $\sigma = 2$, $\gamma = 7/5$; $B = 8$ instead $B = 1$ following from Equation 24)

that they essentially cut off the density profile at a distance inside of the leading edge which is tantamount to our supposition that only a minute fraction of the mass actually achieves a velocity close to that of the leading edge. In other words, the large values of B tend to partly compensate for the prohibitively large leading edge velocity assumed by Mirels and Mullen.

The values here observed for B (Equation 24) are always smaller and therefore the density profiles fuller, than for the corresponding similarity flow.

Apparently the formula (26) for the density profile becomes nonsensical for $B < 0$, or

$$\frac{1}{\gamma-1} < \frac{\sigma+1}{2} \quad (27)$$

Now, it is interesting to note that inequality (27) describes just those cases which are physically meaningless. To see this, we recall that $\gamma = \frac{N+2}{N}$ where N is the number of internal degrees of freedom of the gas. Therefore, $\frac{1}{\gamma-1} = \frac{N}{2}$. On the other hand $(\sigma+1)$ is the number of dimensions of the space into which the gas expands. If this is $(\sigma+1) = n$, then inequality (27) states: $N < n$, which obviously cannot happen, since even a hypothetical gas must have at least as many internal as external degrees of freedom.

The density factor D of Equation (25) has a simple physical meaning: it gives, apart from the time factor $t^{\sigma+1}$, the density at the origin $r = 0$. The factor D is plotted in Figure 1 and Figure 2 for $\sigma = 1, 2$ versus γ . For comparison, the corresponding curves obtained by Mirels and Mullen [4]




FIGURE 1

Density factor for $\sigma=1$.




FIGURE 2

Density factor for $\sigma=2$.

and the results of numerical computations by Greifinger [2] are plotted in the same diagram. Also the curve for $\sigma = 0$ is shown in Figure 1. For this case the results of Mirels and Mullen [4], Greifinger [2] and the present analysis agree with each other and the exact solution. For $\sigma = 1, 2$ the results of the present analysis are in very good agreement with the numerical data [2].

Another check against numerical results is given in Figure 3, which shows the density profile for $\sigma = 1$, $\gamma = 5/3$ in comparison with the results of Mirels and Mullen [4] and numerical data computed at NRL [3]. All profiles are normalized by dividing through the density at $r = 0$. Again, the agreement between the present analysis and the numerical computation is very good. In Figure 2 the values for D according to Equation (24) are plotted only for $\gamma \leq 5/3$. For larger γ -values the exponent B (Equation 24) becomes negative and the formula (26) is no longer applicable. One might suspect that the approximate solution (Equation 26) becomes less accurate on approaching $\gamma = 5/3$.

4. LATERAL EXPANSION OF A HYPERSONIC JET BOUNDED BY VACUUM

If the lateral flow velocity u of the expanding jet is small compared with its axial velocity V_0 , then the equations of motion for steady flow reduce to Equations (1) with $\bar{t}$ replaced by $\frac{\bar{x}}{V_0}$, where $\bar{x}$ is the axial distance from the nozzle. Using the nondimensional quantities defined in Equation (2) and defining a nondimensional axial distance $x = \frac{\bar{x}}{R_0}$, one




FIGURE 3

Asymptotic density profile for $\sigma = 1$, $\gamma = \frac{5}{3}$

has to replace t in Equations (4) by $\frac{x a_0}{V_0} = \frac{x}{M_0}$. Thus, the solution for the density distribution of the jet is obtained from Equation (26) for $\sigma = 1$ and substituting $\frac{x}{M_0}$ for t . This solution is valid for $M_0 \gg 1$.

5. MEASUREMENT OF PLASMA TEMPERATURE

In their paper, "Translational Equilibration of Wires Exploded in Vacua," [1] the authors describe a method for measuring the temperature of ionized gas with the aid of a drift tube. The method, which is described in detail in [1], is essentially as follows: The plasma source, in this case an exploding wire, is located at one end of a long evacuated drift tube, while an electrode is placed at its opposite end. It is assumed that the wire is heated uniformly by a short current pulse. After the short heating period the gaseous, ionized wire material starts to expand without being heated subsequently. The ion flux, after reaching the electrode, gives rise to a time dependent current in the electrode circuit which can be observed. From the observed current distribution the temperature of the original plasma cloud can be inferred.

In evaluating the experimental data Zinke et al. [1] assumed, the plasma would expand like a collision-free gas; i.e., according to free molecular flow. On the basis of this assumption, it was impossible to fit the theoretical prediction to the experimental data unless a hypothetical "containment-time" was introduced. In other words, the shape of the observed current-time distribution was quite different from the shape of any of the possible theoretical distributions. The theoretical current distributions for free molecular flow are given by

$$j(\bar{t}) = \frac{K}{\bar{t}^4} \exp\left(-\frac{m}{2kT} \frac{\bar{r}^2}{\bar{t}^2}\right) \quad (28)$$

where $\bar{t}$ is the time elapsed from the explosion of the wire, $\bar{r}$ the distance between the explosion and the electrode, T the temperature of the original plasma cloud and K a constant related to the total number of ions.

The distributions Equation (28) form a two-parameter class of curves with parameters K and T . None of these curves fit the experimental data. A satisfactory fit could be achieved only after introducing a third parameter, H , the "containment-time." Assuming the plasma did not start to expand at time $t = 0$, i.e., immediately after heating, but at a later time, $t = H$, the distribution Equation (28) would be altered to:

$$j(\bar{t}) = \frac{K}{(\bar{t}-H)^4} \exp\left[-\frac{m}{2kT} \frac{\bar{r}^2}{(\bar{t}-H)^2}\right] \quad (29)$$

With properly chosen K , T , H , Zinke could obtain a reasonable fit to all experimental data. However, the physical meaning of the containment-time, H , remained completely unclear. Obviously, the assumption of free molecular flow is not correct for the high initial densities encountered in the wire explosion. In continuum expansion the energy of the gas molecules is redistributed as a result of collisions. This may significantly change the flow and the current distribution observed at the electrode of the drift tube. The theoretical current distribution for continuum flow is proportional to $\rho u = \rho \cdot \frac{\bar{r}}{\bar{t}}$ with ρ given by Equation (26) and $\sigma = 2$. Thus:

$$j = \frac{K \cdot \bar{r}}{\bar{t}^4} \left[1 - \left(\frac{\sqrt{3}(\gamma-1)}{2} \frac{\bar{r}}{a_n \bar{t}} \right)^2 \right]^B \quad (30)$$

Here $\bar{r}$ is the distance between the electrode and the plasma source (exploding wire) and $\bar{t}$ the time elapsed from the explosion.

When K and a_0 were chosen properly, a good fit could be obtained to the experimental data of Zinke without the assumption of a containment-time.

From a_0 the temperature can be determined by: $a_0^2 = \gamma \frac{kT}{m}$.

The values for the temperature obtained from this fit were about 10^5 °K. These values are more likely correct than those obtained by Zinke of about 10^6 °K.

APPENDIX A

Introducing a new dependent variable ρ^* by

$$\rho = (\rho^*)^{\sigma+1} \quad (31)$$

and substituting (31) into Equations (4a, b) gives the following set of equations for ρ^* :

$$\frac{\partial \rho^*}{\partial t} + \frac{\partial \rho^* u}{\partial r} + \frac{\sigma}{\sigma+1} \rho^* \left[\frac{u}{r} - \frac{\partial u}{\partial r} \right] = 0 \quad (32a)$$

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial r} + (\sigma+1) \rho^* (\gamma-2)(\sigma+1) \frac{\partial \rho^*}{\partial r} = 0 \quad (32b)$$

Let us now consider the factor $\left(\frac{u}{r} - \frac{\partial u}{\partial r} \right)$ in the last term of Equation (32a). This factor is exactly zero for the similarity solution, as can be seen from Equation (11).

For the case of uniform initial density this factor becomes asymptotically zero, since $u \sim \frac{r}{t}$. It is also zero near the center $r = 0$, which follows from the boundary conditions (5) according to which $u = c_1 r + c_2 r^2 + \dots$. We therefore assume, that the average effect of the last term of Equation (32a) is small and to a certain approximation it can be neglected altogether.

With the further transformations:

$$u^* = \frac{u}{\sqrt{\sigma+1}} ; \quad t^* = t\sqrt{\sigma+1} \quad (33)$$

$$(\gamma^* - 2) = (\gamma-2)(\sigma+1) + \sigma$$

we obtain by substituting into (32a,b):

$$\frac{\partial \rho^*}{\partial t^*} + \frac{\partial \rho^* u^*}{\partial r} = 0 \quad (34a)$$

$$\frac{\partial u^*}{\partial t^*} + u^* \frac{\partial u^*}{\partial r} + (\rho^*)^{(\gamma^*-2)} \frac{\partial \rho^*}{\partial r} = 0 \quad (34b)$$

From (34a,b) we observe, that ρ^* , u^* satisfy the equations of motion (4a,b) for $\sigma = 0$ with γ^* replacing γ . ρ^* , u^* also satisfy the same boundary conditions as ρ , u for the case of uniform initial density. The asymptotic solution, therefore, can be obtained from (8a,b) with ρ , u , t , γ replaced by ρ^* , u^* , t^* , γ^* ; i.e.,

$$\rho^* t^* \sim \left[1 - \left(\frac{\gamma^*-1}{2} \frac{r}{t^*} \right)^2 \right]^{\frac{3-\gamma^*}{2(\gamma^*-1)}}, \quad (35a)$$

$$u^* = r/t^* . \quad (35b)$$

Taking the $(\sigma+1)$ power of Equation (35a) and substitution of (31) and (33) into (35a,b) gives:

$$\rho t^{\sigma+1} = \text{const} \left[1 - \left(\sqrt{\sigma+1} \frac{(\gamma-1)}{2} \frac{r}{t} \right)^2 \right]^{\left(\frac{1}{\gamma-1} - \frac{\sigma+1}{2} \right)}, \quad (36a)$$

$$u = r/t . \quad (36b)$$

The constant in Equation (36a) can now be re-adjusted so as to satisfy the mass conservation. The solution (36a) then is identical with Equation (26).

REFERENCES

1. O. H. Zinke, R. B. Owen and C. K. Manka, "Translational Equilibration of Wires Exploded in Vacua." Exploding Wires 3, edited by W. G. Chace and H. K. Moore, Plenum Press, New York (1964).
2. C. Greifinger, Rand Corporation (private communication, 1965).
3. O. B. Langworthy, et al, "Electrically Exploded Wires, Experiments and Theory. Progress Report for March 1, 1958 to June 30, 1960," NRL Report 5489 (May 1961).
4. H. Mirels and J. F. Mullen, "Expansion of Gas Clouds and Hypersonic Jets Bounded by a Vacuum," AIAA Journal 1, 596 (1963).
5. R. Courant and K. O. Friedrichs, Supersonic Flow and Shock Waves, p. 191, Interscience Publishers, New York (1948).
6. C. Greifinger and J. Cole, "One-Dimensional Expansion of a Finite Mass of Gas into Vacuum." Rand Report P2008 (June 1960).
7. K. P. Stanyukovich, Unsteady Motion of Continuous Media, p. 147, 498-506, Pergamon Press, New York (1960).
8. Sedov, L. I., Similarity and Dimensional Methods in Mechanics, p. 271-281, Academic Press, New York (1959).
9. Keller, J. B., "Spherical, Cylindrical, and One-Dimensional Gas Flows," Quart. Appl. Math. 14, 171-184 (1956).

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) U. S. Army Ballistic Research Laboratories Aberdeen Proving Ground, Maryland		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE ASYMPTOTIC SOLUTIONS FOR THE EXPANSION OF GAS INTO VACUUM		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial) Wedemeyer, Erich H.		
6. REPORT DATE April 1965	7a. TOTAL NO. OF PAGES 31	7b. NO. OF REFS 9
8a. CONTRACT OR GRANT NO. b. PROJECT NO. 1A222901A201 c. d.		9a. ORIGINATOR'S REPORT NUMBER(S) Report No. 1278
		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
10. AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain copies of this report from DDC.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY U. S. Army Materiel Command Washington, D. C.
13. ABSTRACT When a spherical or cylindrical gas cloud expands into vacuum, the resultant flow becomes asymptotically self-similar and independent of the original cloud diameter. An analytic solution for the asymptotic density profiles, given by Mirels and Mullen ⁴ is re-examined and a new solution proposed, which gives better agreement with numerical computations. Some applications of the asymptotic solutions are discussed.		

DD FORM 1473
1 JAN 64

Unclassified
Security Classification

Unclassified
Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Unsteady expansion into vacuum Asymptotic gas flows Hypersonic jets Drift tube						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

Unclassified
Security Classification