

604154



ANALYTICAL APPROXIMATIONS

Volume 3

Cecil Hastings, Jr.

P-340

✓ gwl

30 October 1952

Approved for OTS release

6 p

COPY	7	OF	7
HARD COPY	\$. 1.00		
MICROFICHE	\$. 0.50		

6 p

DDC
RECEIVED
AUG 19 1964
DDC-IRA C

The RAND Corporation

• SANTA MONICA • CALIFORNIA

**Best
Available
Copy**

10-14-52

Analytical Approximation

Offset Circle Probability Function: We consider the function

$$q(R, x) = \int_R^{\infty} e^{-\frac{1}{2}(\rho^2 + x^2)} I_0(\rho x) \rho d\rho$$

in which $I_0(z)$ is the usual Bessel function. To better than .00014 over $(-\infty, \infty)$,

$$q(1, x) \doteq 1 - \frac{.3935}{[1 + .5968x^2 + .0047x^4 + .00028x^6]^4}$$

The parametric form used is convenient for approximating fixed-R cross-sections of the $q(R, x)$ surface for small values of R.

Cecil Hastings, Jr.
RAND Corporation
Copyright 1952

10-15-52

Analytical Approximation

Offset Circle Probability Function: We consider the function

$$q(R,x) = \int_R^\infty e^{-\frac{1}{2}(\rho^2+x^2)} I_0(\rho x) \rho d\rho$$

in which $I_0(z)$ is the usual Bessel function. To better than .0035 over $(-\infty, \infty)$,

$$q(2,x) \approx 1 - \frac{.865}{[1+.038x^2+.004x^4]^4}.$$

The parametric form used is convenient for approximating fixed-R cross-sections of the $q(R,x)$ surface for small values of R .

Cecil Hastings, Jr.
RAND Corporation
Copyright 1952

Analytical Approximation

Offset Circle Probability Function: We consider the function

$$q(R,x) = \int_R^\infty e^{-\frac{1}{2}(p^2+x^2)} I_0(px) p dp$$

in which $I_0(z)$ is the usual Bessel function. To better than .001 over $(-\infty, \infty)$,

$$q(2,x) \doteq 1 - \frac{.865}{[1+.0401x^2+.00309x^4+.000075x^6]^4}.$$

The parametric form used is convenient for approximating fixed- R cross-sections of the $q(R,x)$ surface for small values of R .

Cecil Hastings, Jr.
RAND Corporation
Copyright 1952

Analytical Approximation

Offset Circle Probability Function: We consider the function

$$q(R,x) = \int_R^{\infty} e^{-\frac{1}{2}(\rho^2+x^2)} I_0(\rho x) \rho d\rho$$

in which $I_0(z)$ is the usual Bessel function. To better than .0014 over $(-\infty, \infty)$,

$$q(1,x) \doteq 1 - \frac{.393}{[1+.093x^2 + .007x^4]^4}$$

The parametric form used is convenient for approximating fixed-R cross-sections of the $q(R,x)$ surface for small values of R.

Cecil Hastings, Jr.

RAND Corporation

Copyright 1952

10-29-52

Analytical Approximation

Offset Circle Probability Function: We consider the function

$$q(R,x) = \int_R^\infty e^{-\frac{1}{2}(\rho^2+x^2)} I_0(\rho x) \rho d\rho$$

in which $I_0(z)$ is the usual Bessel function.

To better than .0009 over $(0,\infty)$,

$$q(2,2+y) \doteq 1 - \frac{.397}{[1 + .236y + .066y^2 + .056y^3]^4}$$

The parametric form used is convenient for approximating fixed- R semi-cross-sections of the $q(R,R+y)$ surface for any $R \gg 0$ and for y ranging over $(0,\infty)$.

Cecil Hastings, Jr.
RAND Corporation
Copyright 1952