

THE COAGULATION OF CHARGED CLOUD-DROPLETS

by

L.M. Levin

Dok. Akad. Nauk SSSR, 94 (1954), 3, 467-470

and

SIZE DISTRIBUTION FUNCTION FOR CLOUD-DROPLETS AND RAINDROPS

by

L.M. Levin

Dok. Akad. Nauk 94 (1954), 6, 1045-1048

Translated by

E. R. Hope

Directorate of Scientific Information Service

DFB Canada

March, 1958

T 263 R

AD 610 961

Translated from Dok. Akad. Nauk SSSR, 94 (1954), 3, 467-470.

THE COAGULATION OF CHARGED CLOUD-DROPLETS

L. M. Levin

(Presented by O. Yu. Shmidt,
Member of the Academy, November 26, 1956.)

According to present theory of the formation of precipitations [1, 2], the growth of cloud-droplets in the range of diameters $d = 4 \sim 30 \mu$ takes place only by way of condensation of water vapor. The basis of this conclusion is that since there exists a critical value $k_{cr} = 1.214$ [2, 3] for the parameter of gravitational coagulation, then coagulation of $\sqrt{\text{two}}$ neutral droplets is possible only when the diameter of the larger drop exceeds 30μ .* But it seems to us that growth of cloud-droplets in the range $d = 4 \sim 30 \mu$ may take place through coagulation of electrically charged droplets. To confirm this hypothesis, let us calculate E , the coefficient of capture [3], for electrically charged droplets.

Langmuir [2] and Shishkin [1] unjustifiably applied the theory of deposition of an aerosol on obstacles to the theory of cloud-droplet growth. The coefficient of capture, in the deposition of an aerosol on a sphere, was calculated [2] under the assumption that the sphere is considerably larger than the aerosol particle. Consequently these calculations may be applied to the descent of droplets of raindrop size in a cloud, but not to the case of two droplets of comparable size. Langmuir, in calculating the growth of cloud-droplets, really considers all the cloud-droplets to be fixed in space, except one droplet (sometimes of the same dimensions as the others) which moves under the influence of gravitation and of rising air currents [2]. Shishkin, however, takes the coagulation of droplets as depending only on their relative velocity [1], although it is easy to see that the air velocity fields around the droplets, which determine the droplet interaction, depend on the velocities of the droplets relative to the air. Since it is not at present possible to account for the interaction of the aerodynamic fields of two droplets, it seems reasonable to us to perform the calculations for cases when one of the drops is considerably smaller than the other, and in doing so to take the aerodynamic field of the small droplet as not affecting the field of the large droplet. Then in a coordinate system tied to the large droplet, the small droplet will be acted upon by the force of gravity, by the electric forces, and by the aerodynamic Stokes force.

* For instance, with a rising air velocity of 10 cm/sec, the droplet diameter grows by condensation from 11 to 30μ in 7.5 hours, and then grows by gravitational coagulation to 1300μ in 15-20 minutes [1].

1) In the case of both droplets' bearing electric charges of different signs, the equation of motion of the small droplet will take the form

$$\frac{\pi d^3 \rho_k}{6} \frac{d v_1(r_1)}{dt} = -3\pi \mu d [u_1(r_1) - v_1(r_1)] + \frac{e_1 e_2}{r_1^2} r_1 + \frac{\pi d^3}{6} \rho_k g, \quad (1)$$

where r_1 is the radius vector to the center of the small droplet, $v_1 = dr_1/dt$ is its velocity vector, d is the droplet diameter, ρ_k its density, e_1, e_2 the charges on the droplets; $u_1(r_1)$ is the field of velocities of the air stream around the large droplet, μ is the viscosity coefficient of the air, and g is the vector acceleration of gravity.

Converting equation (1) to the dimensionless form [3] (as characteristic velocity we take the constant descent velocity of the large droplet, $u_\infty = \rho_k g D^2 / 18 \mu$, and as characteristic size $D/2$), we find *

$$k \frac{dv}{dt} + v = u + k \sigma_1 \frac{r}{r^2} + g_1, \quad (2)$$

where

$$k = \frac{d^3 \rho_k u_\infty}{9 \mu D}; \quad \sigma_1 = \frac{12 e_1 e_2}{\pi d^3 D u_\infty^2 \rho_k} < 0; \quad g_1 = \frac{g D}{2 u_\infty^2} k = \frac{d^2}{D^2}$$

are dimensionless scaling criteria for the phenomena.

It is obvious that for $|\sigma_1| \gg 1$ we may neglect the inertia term in equation (2), whereupon this equation takes the form (for $\alpha = -k\sigma_1 > 0$):

$$v = u - \alpha \frac{r}{r^2} + g_1. \quad (3)$$

Examination of equation (3) shows that for small values of α it has seven singular points (four saddles on the ρ axis, two centers and a dipole at the coordinate origin), and for large values of α , three singular points (two saddles on the ρ axis and a dipole). Figure 1 illustrates the topological configurations of the trajectories of equation (3) for different values of α . This figure shows that the separatrices A passing through a saddle ($\rho = 0, x > 1$) represent limiting trajectories determining the capture coefficient E. This means we have $E = \rho_0^2$ (ρ_0 being the distance of the said separatrix from the ρ axis at $x \rightarrow \infty$), since, as we shall show infra, these separatrices do not intersect the surface of the large droplet.

On the other hand, we can derive expressions for the trajectories described by equation (3) in finite form, if we note that the field corresponds to a Stokes stream having, in dimensionless (spherical) coordinates, the stream function $\psi(r, \theta) = \frac{\sin^2 \theta}{2} \left(r^2 - \frac{1}{2} r + \frac{1}{2r} \right)$. Then for the trajectories we get

$$\frac{dr}{r d\theta} = \frac{v_r}{v_\theta} = \frac{u_r - \alpha/r^2 + g_{1,r}}{u_\theta + g_{1,\theta}} = \frac{r^2 \sin \theta \frac{\partial \psi}{\partial \theta} - \frac{\alpha}{r^2} - g_1 \cos \theta}{-r \sin \theta \frac{\partial \psi}{\partial r} + g_1 \sin \theta}, \quad (4)$$

* In equation (2) we have the dimensionless quantities $v = v_1 / u_\infty$, $u = u_1 / u_\infty$, $r = 2r_1 / D$, $\alpha = 2u_\infty / D$ where D is the diameter of the large droplet.




Fig. 1

and consequently,

$$d\psi_1 = d\left(\psi - g_1 \frac{r^2 \sin^2 \theta}{2}\right) = \alpha \sin \theta d\theta. \quad (5)$$

From this it is easy to find the equation of the trajectories:

$$\psi_1 = \psi_{1,\pi} - \alpha(1 + \cos \theta), \quad (6)$$

where $\psi_{1,\pi}$ is the value of ψ_1 for the trajectory at $\theta = \pi$.

Noting that the separatrix we are seeking has, at $\theta = 0$, a finite value of r , we find for it the value of $\psi_{1,\pi}$ equal to $\psi_{1,\pi} = \frac{\rho_2^2(1-g_1)}{2} = 2\alpha$. Since according to equation (5) we have $d\psi_1/d\theta > 0$ along the trajectory, then the separatrix in question, having $\psi_1 = 0$ at $\theta = 0$, nowhere intersects the surface of the large droplet, for which $\psi_1 < 0$. Consequently,

$$E = \rho_2^2 = \frac{4\alpha}{1-g_1}. \quad (7)$$

- oOo -

2) Let us now consider that case when the large droplet is charged and the small droplets neutral. Then in the first approximation the charge of the large droplet will induce dipoles in the small droplets and attract them. The equation of motion of the small droplets, in the same coordinate system, takes the form (ref. [4]):

$$\frac{d\pi^2}{6} \rho_k \frac{dv_1(r_1)}{dt} = -3\pi_1 d |u_1(r_1) - v_1(r_1)| - \frac{1}{4} \frac{\epsilon-1}{\epsilon+2} d^3 \frac{\epsilon_1}{r_1^2} r_1 + \frac{\pi d^3}{6} \rho_k g, \quad (8)$$

where ϵ is the small droplet dielectric constant.

In dimensionless form equation (8) becomes:

$$k \frac{dv}{dt} + v = u - k\sigma_2 \frac{r}{r^2} + g_1, \quad (9)$$

where $\sigma_2 = \frac{24}{\pi} \frac{\epsilon-1}{\epsilon+2} \frac{r_1^3}{\rho_k D^2 u_\infty^2} > 0$.

If $\sigma_2 \gg 1$, then equation (9) is simplified:

$$v = u - k\sigma_2 \frac{r}{r^2} + g_1, \quad (10)$$

and the trajectories of motion may be determined from an equation similar to (5):

$$d\psi_1 = \frac{\alpha \sigma_2 \sin \theta d\theta}{r^2}. \quad (11)$$

Equation (10) has eight singular points. Their positions and the topological configurations of the trajectories are similar to the cases illustrated in Figure 1.*

- 000 -

3) Tables 1 and 2 present the results of our calculations of E ; likewise values of σ_1 and σ_2 , enabling one to estimate the applicability of the computation methods as developed (the calculations were carried out for $\sigma_1, \sigma_2 > 5$ and $g_1 \leq 0.25$). The order of magnitude of the charge was calculated according to Ya. I. Frenkel' [5], and for the absolute magnitude of the charge in the CGSE system we accepted the expression

$$e = 0.5 \cdot 10^{-7} d [\mu], \quad (12)$$

corresponding to a value of 0.3 V for the electrokinetic potential. It seems to us that the magnitude of the charge accepted by us according to expression (12) has been selected reasonably as regards order of magnitude, and does not contradict the experimental findings of Gunn [6], who measured the gross charge of cloud-droplets of diameter $d > 10 \mu$.

TABLE 1

$D [\mu]$	$d [\mu]$	σ_1	E	$D [\mu]$	$d [\mu]$	σ_1	E
5	1	15000	65,2	15	6	5,4	2,8
5	2	3900	74,7	20	1	61	0,98
7,5	1	3080	18,8	20	3	6,8	1,00
7,5	4	192	25,9	25	1	25	0,50
10	1	975	7,9	25	2	6,2	0,50
10	5	39	10,4	30	1	12	0,29
15	1	192	2,3	30	1,5	5,4	0,29

TABLE 2

$D [\mu]$	σ_1	E		
		$\epsilon_1 = 0,25$	$\epsilon_1 = 0,1$	$\epsilon_1 = 0,04$
5	1250	2,32	1,48	0,89
7,5	110	1,06	0,69	0,40
10	20	0,57	0,38	0,22
12,5	5,1	0,34	0,23	0,13

* The difference amounts to the fact that the dipole reduces to two nodes, a stable node, at the coordinate origin, and an unstable node ($\rho = 0, 0 < x < 1$), while the centers are converted into unstable foci or unstable nodes. As previously, the separatrix A passing through a saddle ($\rho = 0, x > 1$) does not intersect the surface of the large droplet (equation (11)) and constitutes a limit trajectory. Numerical calculations have shown that equation (11) is convenient for constructing this separatrix (calculation of E).

Tables 1 and 2 show that under these assumptions the coefficient of capture for cloud-droplets, in the range with which we are dealing, is high enough to guarantee a considerable coagulation of droplets, while for neutral droplets of the same sizes there is no coagulation. The amount of coagulation will be rather great even if the actual size of the charges is an order of magnitude lower than that which we have taken.

- oOo -

4) From the expressions for σ_1 and σ_2 it follows that at large values of D (when u_∞ also is large) the quantities $\sigma_1, \sigma_2 \ll 1$ and, consequently, the effect of the electric charges on coagulation may be neglected. This accounts for the experimental results of Gunn and Hitschfeld [7], in which $D = 3.2$ mm, $d > 5 \mu$ and $e_1 = 0.2$ CGSE. Thus in these experiments $u_\infty = 825$ cm/sec and $\sigma_1 = 4 \cdot 10^{-5}$, $\sigma_2 < 7 \cdot 10^{-3}$. Consequently these authors did not find any electric charge effect on the coagulation of a large droplet with small droplets.

In the same way, by calculating the quantities σ_1 and σ_2 it may be shown that in cloud-droplet catchers the precipitation of droplets onto the receiver is in practice independent of the electric charges of the droplets and of the droplet receiver.

Similar results are indeed found for charged droplets.

In conclusion I express my cordial thanks to V. M. Bovsheverov for fruitful discussion of the results of this work.

Geophysical Institute,
Academy of Sciences of the USSR.

Received October 21, 1953.

REFERENCES

- 1) N.S. SHISHKIN, *Usp. Fiz. Nauk* [Progress of the Physical Sciences], 45, 3, 313 (1951).
- 2) I. Langmuir, *J. Meteor.*, 5, 175 (1948).
- 3) L.M. LEVIN, *Dok. Akad. Nauk*, 91, No. 6 (1952).
- 4) R. Cochet, *Ann. de Géophys.*, 8, 33 (1952).
- 5) Ya. I. FRENKEL', *Teoriya yavlenii atmosfernogo elektrichestva* [Theory of atmospheric electric phenomena], 1949.
- 6) R. Gunn, *J. Meteor.*, 9, 397 (1952).
- 7) K. Gunn, W. Hitschfeld, *ibid.*, 8, 7 (1951).

SIZE DISTRIBUTION FUNCTION FOR CLOUD-DROPLETS AND RAINDROPS

L. M. Levin

(Presented by G. A. Hambrug'ev,
Member of the Academy, December 4, 1953.)

Modern theories of the formation and development of clouds and forms of precipitation are essentially concerned with empirical size distribution functions for cloud-droplets and raindrops. In chronological order, four distribution functions have been used in these theories: the Smolukhovski-Schumann theory [1-3], the Marshall-Palmer theory [4], the Best theory [5, 6] and the Kargian-Mazin theory [7]. The first of these, a theoretical formula, was derived by Smolukhovski for colloids, and the same scheme for deriving the distribution function was, without any sufficient justification, carried over by Schumann to the cloud-droplet case. The other distribution functions are empirical.

In 1941, A.N. Kolmogorov proved [8] that if we take a rather general scheme of random break-up of particles, then in the limit we obtain a logarithmically normal size-distribution of particles. It is natural to suppose that a logarithmically normal distribution might likewise be found for the random coagulation of cloud-size droplets, in the limiting case of a great number of particle-merger events.

In 1952-1953, a large amount of statistical data was collected by the Elbruz expedition of the Geophysical Institute, Academy of Sciences of the USSR, data which made it possible to examine the question of cloud-droplet distribution function and to state the premises for the raindrop distribution function.*

Figure 1a shows "rectified diagrams" [10] for the size-distribution of cloud-droplets collected from individual clouds. Plotted as abscissae are the logarithms of droplet diameter d in microns; as ordinates, values of $\Psi^{-1} [F(d)]$, where $F(d)$ is the selection distribution function, and

$$\Psi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt$$
 (here Ψ^{-1} is the inverse function of Ψ). According to

the logarithmically normal distribution for which the density of distribution will be given by the formula:

$$n(x) = \frac{1}{\sigma\sqrt{2\pi}x} e^{-\left[\ln \frac{x}{x_0}\right]^2 / 2\sigma^2}, \quad (1)$$

* The experimental procedure followed by the expedition in its research on the microstructure of clouds is described in reference [9]. Rain spectra were studied by a dyed filter paper method.

the experimental points must, in these coordinates, lie on a straight line. The point of intersection of this straight line and the axis of the abscissae is $\ln x_0$, and the tangent of the straight line's angle of slope is $1/\sigma$ [10].

Figure 1e shows that the experimental findings for clouds are satisfactorily described by a logarithmically normal distribution function. At the present time, the expedition has accumulated over thirty samplings of cloud-droplets, with the numbers of droplets in the sample ranging from 2000 to 20,000. They all support the above conclusion.

As for the raindrop size distribution, we are not at the present time in possession of any similar material collected from individual rainfalls. From the material for 1952 it was possible to determine only the average spectra for a number of rainfalls of one particular intensity. These spectra are shown, in the same coordinates, in Figure 2. The graphs of Figure 2 indicate that rain-droplets likewise have a logarithmically normal distribution law. Since this last conclusion is drawn only for the average spectra, it is of great importance to collect statistical material on the spectra of individual rainfalls.

Moreover, to answer precisely the question of the distribution function it is essential to expand the range of captured droplets, both for clouds and for rain. In the case of clouds, it would appear essential to establish a procedure for capturing fine droplets ($d < 4 \mu$) and a procedure for collecting a greater amount of the large droplets ($d > 50-60 \mu$). In the case of rain, it would be desirable to effect the capture of drops of diameter $d > 400 \mu$.

It seems to us that the logarithmically normal law of size-distribution of cloud-droplets in the diameter-range $d = 4 \sim 50 \mu$ proves that coagulation of cloud-droplets occurs over the whole of this range. This fact indirectly confirms our working hypothesis of coagulation of electrically charged cloud-droplets in the diameter-range $d = 4 \sim 30 \mu$, as set forth in reference [11].

In reference [7] it was shown that the size-distribution of cloud-droplets is well approximated by the formula

$$n(x) = ax^2 e^{-bx}. \quad (2)$$

This formula was derived from the fact that over a large range of diameters the empirical quantity $\lg \frac{n(d)}{d^2}$ was found to be linearly related to d . But this relationship holds with fair accuracy for a great range of x , in the case of a logarithmically normal distribution.

In Figure 3a we show graphs of $\lg \frac{n(x)}{x^2}$, where $n(x)$ is given by formula (1).^{*} These graphs show that for the range $\sigma = 0.25 \sim 0.5$ (that is, a range practically covering all the values of σ encountered in the clouds investigated), relationship (2) is fairly well satisfied for $x > x_1 = 0.5 \sim 1.0 x_0$. In the region $x < x_1$ there is a characteristic bend in the curve of

^{*} The quantity x_0 was taken for the scale-unit on the x-axis of Figure 3. We note that x_0 is the median of the distribution.




Fig. 1.

- 1 - Sept. 3, 1953, at 18:00 hrs:
 $N = 27,800$, $\sigma = 0.35$, $d_0 = 8.6 \mu$.
- 2 - Sept. 3, 1953, at 18:30 hrs:
 $N = 16,000$, $\sigma = 0.35$, $d_0 = 9.3 \mu$.
- 3 - Sept. 3, 1953, at 19:00 hrs:
 $N = 7,500$, $\sigma = 0.31$, $d_0 = 10.7 \mu$.
- 4 - Sept. 19, 1953, at 18:45 hrs:
 $N = 4,352$, $\sigma = 0.30$, $d_0 = 8.8 \mu$.
- 5 - Sept. 19, 1953, at 19:00 hrs:
 $N = 3,900$, $\sigma = 0.27$, $d_0 = 8.8 \mu$.




Fig. 2.

- 1 - $I = 5.6-7.5 \text{ mm/hr}$, $N = 28,700$,
 $\sigma = 0.52$, $d_0 = 0.57 \text{ mm}$.
- 2 - $I = 3.6-5.5 \text{ mm/hr}$, $N = 37,300$,
 $\sigma = 0.46$, $d_0 = 0.64 \text{ mm}$.
- 3 - $I = 2.6-3.5 \text{ mm/hr}$, $N = 27,700$,
 $\sigma = 0.48$, $d_0 = 0.56 \text{ mm}$.
- 4 - $I = 1.6-2.5 \text{ mm/hr}$, $N = 24,000$,
 $\sigma = 0.46$, $d_0 = 0.56 \text{ mm}$.
- 5 - $I = 0.5-1.5 \text{ mm/hr}$, $N = 22,300$,
 $\sigma = 0.41$, $d_0 = 0.59 \text{ mm}$.
- 6 - $I = 0.5-7.5 \text{ mm/hr}$, $N = 140,600$,
 $\sigma = 0.48$, $d_0 = 0.58 \text{ mm}$.




Fig. 3.

$\lg \frac{n(x)}{x^2}$. an inflexion which is seen in nearly all the experimental curves of $\lg \frac{n(d)}{d^2}$ (see Figure 1b) corresponding to the distributions shown in Figure 1a.

Similarly it may be shown that the size-distribution of raindrops as found empirically by Marshall and Palmer [4]:

$$n(x) = ae^{-bx} \quad (3)$$

is, over a large part of the curve, an approximation to the logarithmically normal distribution. This fact is illustrated in Figure 3b, which is a graph of $\lg n(x)$ for $n(x)$ as given by formula (1).

Here, incidentally, we must note that the experimental data obtained by us on cloud-droplets do not satisfy the Smolukhovski-Schumann distribution, for which

$$n(x) = ax^2 e^{-bx^3} \quad (4)$$

This follows from the fact that in the coordinates of Figure 1b this distribution (4) should appear as a cubic parabola. As may be seen from Figure 1b this is not the case.

In conclusion we express our cordial thanks to A.M. Obukhov, Member-Correspondent of the Academy of Sciences of the USSR, for his valuable advice.

Geophysical Institute,
Academy of Sciences of the USSR.

Received December 1, 1953.

REFERENCES

- 1) M. SMOLUKHOVSKI, Sborn. Koaguliaciya kolloidov [Collection of papers: Coagulation of Colloids], 1936.
- 2) T.Q. Schumann, Quart. J. Roy. Met. Soc., 66, 195 (1940).
- 3) N.S. SHISHKIN, Usp. Fiz. Nayk [Progress of the Physical Sciences] 45, 3, 313 (1951).
- 4) J.S. Marshall, W. McK. Palmer, J. Meteor., 3, 165 (1948).
- 5) A.C. Best, Quart. J. Roy. Met. Soc., 76, 16 (1950).
- 6) A.C. Best, *ibid.*, 77, 418 (1951).
- 7) A. Kh. KHRGIAN, I.I. MAZIN, Tr. CAO [Tr. Central Aerological Observatory], No. 7, 56 (1952).
- 8) A.N. KOLMOGOROV, Dok. Akad. Nauk, 31, No. 2 (1941).

- 9) L.M. LEVIN, R.F. STAROSTINA, Dok. Akad. Nauk, 92, No. 1 (1953).
- 10) N. Arley, K.R. Buch, Introduction to the Theory of Probability and Mathematical Statistics [Russian translation], 1951, pp. 140-142.
- 11) L.M. LEVIN, Dok. Akad. Nauk, 94, No. 3 (1954).