

AD 624 626


MAXIMUM VOLUNTARY VENTILATION
AFTER $+G_x$ IMPACT IN HUMANS

Peter G. Hanson, Captain, USAF, BSC

CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION		
Hardcopy	Microfiche	
1.00	0.50	15 pp. 15
ARCHIVE COPY		

Code 1

November 1965



6571st Aeromedical Research Laboratory
Aerospace Medical Division
Air Force Systems Command
Holloman Air Force Base, New Mexico

This report is made available for study with the understanding that proprietary interests in and relating thereto will not be impaired. In case of apparent conflict or any other questions between the Government's rights and those of others, notify the Judge Advocate, Air Force Systems Command, Andrews Air Force Base, Washington, D.C. 20331.

This report may be reproduced to satisfy need of U.S. Government agencies. No other reproduction is authorized except with permission of the 6571st Aeromedical Research Laboratory, Holloman AFB, NMex.

Do not return this copy. Retain or destroy.

MAXIMUM VOLUNTARY VENTILATION
AFTER $+G_x$ IMPACT IN HUMANS

Peter G. Hanson, Captain, USAF, BSC

FOREWORD

This study was conducted during February 1965 by the Biodynamics Branch of the 6571st Aeromedical Research Laboratory.

The author wishes to acknowledge the cooperation and assistance of all members of Dynalectron Corporation, Land-Air Division Aeromed Sub-Group under the supervision of Mr. Mac Connell.

* * *

This technical report has been reviewed and is approved.


CLYDE H. KRATOCHVIL
Lt Colonel, USAF, MC
Commander

ABSTRACT

Eighteen volunteer male subjects were exposed to 20 +G_x impact on the Daisy Decelerator. Measurements of maximum voluntary ventilation (MVV) obtained 10 minutes prior to, immediately after and 20 minutes after impact were compared with previously determined baseline MVV values. The results indicate that MVV performance is elevated immediately after impact. It is suggested that this response is related to subject anxiety with accompanying sympathicotonia.

TABLE OF CONTENTS

	Page
I. INTRODUCTION	1
II. METHODS.	1
A. Experimental Design.	1
B. Subject Protocol	1
C. Determination of Maximum Voluntary Ventilation.	2
D. Treatment of Data.	4
III. RESULTS.	4
IV. DISCUSSION	6
REFERENCES	11

LIST ILLUSTRATIONS

Figure

1. Subject and Equipment During MVV Determination	3
2. Combined Group Data for MVV Measurements Prior to and After Impact.	7

Table

I. Baseline and Predicted MVV Values.	5
II. Statistical Summary of Experimental Results.	8

I

INTRODUCTION

Exposure to sustained acceleration in the $+G_x$ axis results in pulmonary dysfunction in human and laboratory animal subjects. Significant diminutions in pulmonary function as measured by tidal volume, vital capacity, maximum voluntary ventilation (MVV) and pulmonary diffusion have been demonstrated in humans (Ref. 1). Roentgenographic studies in humans and post-acceleration autopsy findings in dogs also suggest that the anterior-posterior displacement of the pulmonary structures within the thoracic cavity results in the engorgement of pulmonary vessels and atelectasis (Ref. 2, 3). These effects develop after 3 to 10 minutes of acceleration at 6 to 12 $+G_x$ (Ref. 4).

Impact acceleration in the $+G_x$ axis produces a marked anterior-posterior compression of the thoracic cavity and inertial displacement of the cardio-pulmonary organs (Ref. 5) which could result in serious lung damage at high impact levels. Accumulated experience with human test subjects on the Daisy Decelerator (Ref. 6) indicate that no clinically perceptible impairment of pulmonary function occurs after $+G_x$ impact at levels up to 25 G; however, these observations have not been verified with quantitative measurements. Therefore, it was of interest to measure MVV before and after $+G_x$ impact to more clearly evaluate the physiological effects of impact on pulmonary function in humans.

II

METHODS

A. Experimental Design

Experiments were conducted during the months of January and February, 1965. A total of 18 volunteer subjects were exposed to impact on the Daisy Decelerator in the $+G_x$ vector (0-5-180).¹ The conditions of deceleration were: brake entrance velocity 30 fps, onset of deceleration 1000 G/sec. with peak of 20 ± 7 G. Each subject was exposed once to the impact conditions.

B. Subject Protocol

A baseline MVV was determined on each subject prior to the actual impact experiments. The measurement technique will be discussed below.

¹ 0-5-180 refers to the orientation of the subject in degrees of roll, pitch and yaw on a polar coordinate system. In the neutral position of 0-0-0 the subject is seated upright facing forward toward the point of impact. In the 0-5-180 orientation the subject is seated upright, 5 degrees back from perpendicular with the back facing the point of impact. Inertial forces in this orientation are directed from anterior to posterior.

On the day of an experiment, the subject reported to the Daisy Decelerator facility approximately one hour prior to the scheduled impact. After a medical examination and placement of vectorcardiogram leads, a pre-impact MVV was determined. The subject was then taken to the Daisy sled for the impact run. A second pre-impact MVV was determined while the subject sat in the Daisy sled prior to attachment of the restraint harness. Sled firing and impact followed within 10 minutes of this pre-impact MVV.

Immediately after impact, the restraint harness was released and the subject was examined briefly to ascertain the state of consciousness and whether any obvious physical injury had occurred. The immediate post-impact MVV was then measured. An average time of 40 seconds elapsed from impact to the beginning of the MVV determination. Following a post-run physical examination, a final MVV measurement was made. This measurement was approximately 20 minutes post-impact.

C. Determination of Maximum Voluntary Ventilation

Maximum voluntary ventilation is defined as the maximum volume of air which can be moved in and out of the lungs per unit time (Ref. 7). It is usually accomplished by having the subject breathe into a spirometer through a low-resistance respiratory valve. The subject is urged to breathe rapidly and yet deeply to produce the maximum possible ventilation volume. Several trial sessions are required to attain consistent results. Since the over-breathing process produces respiratory alkalosis, the collection period is limited to 10 or 15 seconds.

The conditions of this study required a portable gas collection system so that MVV could be determined on the Daisy Decelerator before and after impact. A 100-liter capacity Douglas bag (Thomas) was fitted with a plastic two-way respiratory valve (Collins No. P-304). This system was supported by a metal frame so that the bag would hang freely and the valve remain upright for presentation to the subject (Fig. 1). Prior to a MVV determination, the bag was evacuated with a Collins motor blower (No. P-533) and the bag neck clamped below the point of respiratory valve insertion with a quick release hose clamp. The valve was then inserted and secured with a hose clamp.

A MVV determination was carried out with the subject seated, nose clamp in place, and the respiratory valve mouthpiece held firmly to prevent expulsion. At the end of a normal expiration, the subject was instructed to begin maximum effort breathing which continued over a timed 15-second period. At the end of 15 seconds the mouthpiece was withdrawn from the subjects' mouth and the bag neck quickly clamped below the valve.

The volume of expired air collected in the Douglas bag was then measured by connecting the bag to the intake end of a Collins motor blower which was in turn connected to a dry gas meter (American Meter Co.) so



Figure 1. Subject and Equipment During MVV Determination.

that the contents of the bag were expelled through the gas meter. Gas temperature readings were taken when approximately half of the collected volume had passed through the meter. Barometric pressure was obtained from a mercury manometer situated in the room where the gas volume was measured.

D. Treatment of Data

MVV is usually expressed in liters per minute BTPS. Because of the 4,100 foot altitude of Holloman Air Force Base, volumes measured (ATPS) were relatively large compared to clinical values established near sea level (Ref. 8). The data in this paper, therefore, are expressed in liters per minute (BTPS) corrected to 760 mm Hg. Calculation of MVV was accomplished by the following method:

$$MVV = \dot{V}_E \times 4 \times K \times \frac{(P_B - P_{H_2O})}{760 \text{ mm Hg} - 47 \text{ mm Hg.}} \times \frac{273^\circ + T_B}{273^\circ + T}$$

Where:

MVV = maximum voluntary ventilation (liters/minute BTPS)

$\dot{V}_E$ = expired volume for 15 seconds (liters ATPS)

K = dry gas meter correction factor

P_B = ambient barometric pressure (mm Hg.)

T = gas temperature ($^\circ\text{C}$) in gasometer

P_{H_2O} = vapour tension of water at temperature of gasometer.

T_B = body temperature (37°)

III

RESULTS

The baseline and predicted MVV values along with the physical characteristics of each subject are presented in Table I. With the exception of several severe discrepancies, the measured baseline values are within 15 percent of the predicted values. The comparatively lower volumes obtained in the baseline determinations may be partially attributed to the sitting position of the subjects in this experiment. Predicted values were determined in the standing position (Ref. 9),

Table I. BASELINE AND PREDICTED MVV VALUES

						MVV (L/MIN BTFS)	
	Subject	Age	Ht. (cm)	Wt. (kg)	S.A. (M ²)	Baseline + S $\bar{x}$	Predicted*
1.	J.L.A.	30	170	71	1.81	161 + 1.1	148
2.	C.V.A.	27	170	59	1.68	129 + 5.7	141
3.	J.S.A.	23	183	88	2.10	145 + 1.1	181
4.	W.H.A.	35	183	89	2.14	133 + 8.1	171
5.	J.R.B.	24	178	59	1.74	153 + 5.3	148
6.	C.T.B.	28	190	84	2.20	162 + 3.5	183
7.	W.C.C.	25	170	87	1.99	164 + 1.9	168
8.	W.E.	26	183	83	2.10	136 + 3.7	176
9.	J.O.E.	23	181	75	1.94	126 + 4.3	165
10.	J.F.H.	30	178	79	1.98	133 + 7.8	162
11.	K.D.H.	33	185	96	1.98	133 + 8.2	158
12.	C.W.H.	24	183	74	1.99	174 + 6.0	168
13.	A.A.P.	31	183	87	2.15	174 + 6.7	173
14.	J.L.P.	22	178	79	1.99	148 + 3.2	170
15.	J.W.P.	27	178	87	2.15	178 + 4.3	178
16.	J.W.R.	23	181	75	1.98	118 + 3.7	170
17.	L.M.S.	24	178	71	1.88	162 + 0.1	160
18.	J.P.W.	33	183	79	2.20	153 + 2.4	178

* Motley, H.L.: J. Chest. Dis: 24, 378, 1953.

$$MVV = (97 - 0.50 \text{ Age (yrs)}) \times S.A.(M^2) \text{ L/MIN (BTFS)}.$$

which is known to yield greater MVV volumes than the recumbent or sitting position. Reproducibility of the MVV measurement normally increases with repeated trials (Ref. 10). In our determinations an average of five trials were necessary to establish an acceptable baseline which was set at $|\bar{S}\bar{x}| \leq 10$ L/min. This figure agrees with obtained values in multiple determinations as discussed by Bartels et al. (Ref. 10).

The results of the experimental conditions are summarized in Figure 2. These data represent three groups of MVV measurements obtained; one preceding and two following impact, each plotted as a function of the baseline MVV. Comparison of the groups is made to illustrate the major trends observed in these experiments. In general, there is an increased MVV 10 minutes prior to impact compared to baseline measurements (B_2 vs B_1). Immediately after impact the elevation in MVV is more pronounced (P_1 vs B_1). Twenty minutes after impact there is a diminished MVV in seven subjects, no change in five and continued elevation in the remaining six.

The changes observed for individual subjects are tabulated in Table II along with a statistical summary of the group data. Interindividual variability clearly accounts for the rather small alteration in MVV when expressed as group averages. A statistical analysis based on Wilcoxon's matched-pairs signed-ranks test (Ref. 11) has been included to evaluate the magnitude and direction of the observed differences. The increased MVV immediately after impact (P_1) is significantly ($P < .05$) greater than the baseline (B_1). Changes observed in the other comparisons are not significant.

IV

DISCUSSION

Several factors contribute to subject performance during MVV measurement: the muscular force available, the compliance of the lungs and thoracic cage, and resistance of the airway and pulmonary thoracic tissues (Ref. 8). Therefore, the MVV measurement does provide a useful overall evaluation of breathing mechanics. However, changes in MVV also may reflect a combination of these factors which precludes a definitive diagnosis without ancillary studies of pulmonary function.

The results of this study indicate that $+G_x$ impact ($20 + 1 G_x$) produces no pulmonary dysfunction as measured by MVV. The data actually suggest that prior to and immediately after the impact exposure subject MVV is elevated above baseline values. This phenomenon may be caused by progressive anxiety which is clearly observable in most human subjects prior to impact testing. After the impact episode, subjects tend to relax and anxiety subsides. Increased secretion of epinephrine, which accompanies anxiety, could produce bronchodilation and increased voluntary




Figure 2. Combined Group Data for MVV Measurements Prior to and After Impact

Table II. STATISTICAL SUMMARY OF EXPERIMENTAL RESULTS

Changes in MVV L/Min BTFS

Subject	$(B_2 - B_1)$		$(P_1 - B_1)$		$(P_2 - B_1)$	
	d	rank	d	rank	d	rank
1. J.L.A.	1	1.5	0		-26	-14.0
2. C.V.A.	13	13.5	12	10.0	20	11.0
3. J.S.A.	5	4.0	-2		-6	-4.0
4. W.N.A.	6	5.0	4	4.5	8	6.0
5. J.R.B.	12	12.0	5	6.5	11	9.0
6. G.T.B.	16	16.0	20	14.5	7	5.0
7. W.C.C.	8	7.5	3	3.0	2	1.5
8. W.E.	10	-10.0	-12	-10.0	-10	-7.5
9. J.O.E.	-3	-3.0	19	13.0	0	
10. J.F.H.	7	6.0	20	14.5	22	12.5
11. K.D.H.	13	13.5	12	10.0	4	3.0
12. C.W.H.	1	1.5	5	6.5	0	
13. A.A.P.	10	10.0	-4	-4.5	-2	-1.5
14. J.L.P.	10	10.0	16	12.0	-10	-7.5
15. J.W.P.	-20	-16.0	-28	-17.0	-28	-15.0
16. J.W.R.	15	15.0	25	16.0	22	12.5
17. L.M.S.	-8	-7.5	-2	-1.5	-19	-11.0
18. J.P.W.	-25	-17.0	7	8.0	-13	-10.0

$\bar{X} = 2.8$ $T = -53.5$ $\bar{X} = 6.1$ $T = -34.5$ $\bar{X} = -1.0$ $T = -70.5$

$S\bar{X}_D = \pm 11.5$ $P = N.S.$ $S\bar{X}_D = \pm 12.1$ $P \leq 0.05$ $S\bar{X}_D = \pm 15.5$ $P = N.S.$

muscle activity and thus improve the MVV performance (Ref. 10). The greatest increase in MVV occurs immediately after impact exposure when the subject has sustained the maximum anxiety. Urinary excretion of catecholamines, as measured by vanilmandelic acid (VMA), has been studied under similar impact conditions and is known to increase in urine specimens collected 10 minutes prior to impact (Ref. 12). The combination of these observations suggest that increased adrenergic activity in the subjects prior to impact exposure may account for the elevated MVV. Decreased MVV which occurs 20 minutes after impact is not significant and probably reflects the more relaxed attitude of the subjects.

Transient displacement of the pulmonary tree which occurs during $+G_x$ impact does not produce a measureable change in MVV in contrast to a significant decrease in pulmonary function observed with short-term $+G_x$ centrifugation. The latter changes are thought to be associated with radical alterations in pulmonary blood flow and alveolar ventilation produced by prolonged hydrostatic pressure imbalance within the lungs and thorax (Ref. 2). The average duration of impact G in this experiment was 80 to 100 msec. Hydrostatic pressure changes which occur within this time interval probably do not alter pulmonary circulation sufficiently to produce areas of uneven perfusion.

It is concluded that $+G_x$ impact at the magnitude imposed in these experiments does not produce significant diminution in maximum voluntary ventilation performance. Changes which do occur may tentatively be explained by subject anxiety attendant to impact exposure.

REFERENCES

1. Hyde, A. S. Physiological Effects of Acceleration on Respiration and Protective Measures. In: Bio-Assay Techniques for Human Centrifuges and Physiological Effects of Acceleration. New York: Pergamon Press, Chapter IX_B. p. 101 (1961).
2. Banchem, N., L. Cronin, A. C. Nolan and E. H. Wood. Blood Oxygen Changes Induced by Forward (+G_x) Acceleration. AMRL-TR-64-132, 1964.
3. Hershgold, E. J. "Roentgenographic Study of Human Subjects During Transverse Accelerations." Aerospace Med. 31 No. 3, 213, 1960.
4. Mueller, G. C. E. Cardiovascular Effects of Forward Acceleration. In: Bio-Assay Techniques for Human Centrifuges and Physiological Effects of Acceleration. New York: Pergamon Press, Chapter XI. p. 119 (1961).
5. McDonald, R. K., V. C. Kelly and R. Kaye. "Etiology of Pulmonary Hemorrhage in Cats Exposed to Abrupt Deceleration." Aerospace Med. 19, No. 1, 138, 1948.
6. Chandler, R. F. The Daisy Decelerator. 6571st Aeromedical Research Laboratory Technical Report (in press).
7. Consolazio, C. F., R. E. Johnson and L. J. Pecora. Physiological Measurements of Metabolic Functions in Man. New York: McGraw-Hill, Inc. p. 225 (1963).
8. Comroe, J. H. (Jr.), R. E. Forster II, A. B. DuBois, W. A. Briscoe and E. Carlsen. The Lung - Clinical Physiology and Pulmonary Function Tests. Chicago: Year Book Med. Pub. p. 202 (1962).
9. Motley, H. L. "Pulmonary Function Measurements." Am. J. Surg. 88, 103, 1954.
10. Bartels, H., E. Bucherl, C. W. Hertz, G. Rodewald and M. Schwab. Methods in Pulmonary Physiology. New York: Hafner Co. pp. 69-78 (1963).
11. Siegel, S. Nonparametric Statistics, New York: McGraw-Hill, Inc. p. 75 (1956).
12. Hanson, P. G. Unpublished Observations, 1965.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author) 6571st Aeromedical Research Laboratory		2a REPORT SECURITY CLASSIFICATION Unclassified
		2b GROUP
3 REPORT TITLE MAXIMUM VOLUNTARY VENTILATION AFTER +G _x IMPACT IN HUMANS		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates) Interim February 1965		
5 AUTHOR(S) (Last name, first name, initial) Hanson, Peter G., Captain, USAF, BSC		
6 REPORT DATE November 1965	7a TOTAL NO OF PAGES 10	7b NO OF REFS 12
8a CONTRACT OR GRANT NO	8a ORIGINATOR'S REPORT NUMBER(S) ARL-TR-65-22	
b PROJECT NO		
c	8b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d		
10 AVAILABILITY/LIMITATION NOTICES		
11 SUPPLEMENTARY NOTES		12 SPONSORING MILITARY ACTIVITY 6571st Aeromedical Research Laboratory Holloman AFB, New Mexico
13 ABSTRACT Eighteen volunteer male subjects were exposed to 20 +G _x impact on the Daisy Decelerator. Measurements of maximum voluntary ventilation (MVV) obtained 10 minutes prior to, immediately after and 20 minutes after impact were compared with previously determined baseline MVV values. The results indicate that MVV performance is elevated immediately after impact. It is suggested that this response is related to subject anxiety with accompanying sympathicotonia.		

DD FORM 1 JAN 64 1473

Unclassified
Security Classification

KEY WORDS

Humans
Impact
Maximum voluntary ventilation
Anxiety

LINK A

ROLE

WT

LINK B

ROLE

WT

LINK C

ROLE

WT

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS) (S) (C) or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.