AFCRL-66-153

AD 634 070

NUMERICAL INVESTIGATION OF ELECTROMAGNETIC SCATTERING AND DIFFRACTION BY CONVEX OBJECTS

Nelson A. Logan

Lockheed Missiles & Space Company
A Group Division of Lockheed Aircraft Corporation
Sunnyvale, California

Contract No. AF19(628)-4393

Project No. 4600

Task No. 460004

FINAL REPORT

Period Covered: 16 Nov 1964 through 15 Nov 1965

Date of Report

December, 1965

Prepared
for

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES
OFFICE OF AEROSPACE RESEARCH
UNITED STATED AIR FORCE
BEDFORD, MASSACHUSETTS

Distribution of this document is unlimited.

UNITED STATES AIR FORCE
AFCL Research Library

AFCRL-66-153
AD 634 070

NUMERICAL INVESTIGATION OF ELECTROMAGNETIC SCATTERING AND DIFFRACTION BY CONVEX OBJECTS

Nelson A. Logan

Lockheed Missiles & Space Company
A Group Division of Lockheed Aircraft Corporation
Sunnyvale, California

Contract No. AF19(628)-4393

Project No. 4600

Task No. 460004

FINAL REPORT

Period Covered: 16 Nov 1964 through 15 Nov 1965

Date of Report

December, 1965

Prepared
for

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES
OFFICE OF AEROSPACE RESEARCH
UNITED STATES AIR FORCE
BEDFORD, MASSACHUSETTS

Distribution of this document is unlimited.

LOCKHEED MISSILES & SPACE COMPANY

ABSTRACT

The solutions $D(x,y,y_o,q)$ of the partial differential equation $D_{yy} + iD_x + yD = 0$ are employed as a mathematical model for the fields reflected from and diffracted around an opaque convex surface. This solution involves the Airy function $w_1(t)$ which is a solution of $w_1''(t) - tw_1(t) = 0$. FORTRAN computer programs and tables are given for the appropriate Airy functions. A considerable portion of the report is devoted to discussions of the representations for the roots of the equation $w_1'(t_s) - qw_1(t_s) = 0$ which play a fundamental role in the theory. As a function of q, these roots satisfy the Riccati equation $(t - q^2)(dt/dq) = 1$. The Riccati equation is used to develop several representations for these roots. FORTRAN programs are presented for the evaluation of these constants. Tables of the roots are given for certain cases which have been shown to be of practical significance. Asymptotic expansions are developed for the height gain function $F(y)$ which is a solution of $F''(y) + (k^2f(y) - \lambda_s)F(y) = 0$. The theory of representing functions by a series of Chebyshev polynomials is reviewed and a discussion is given of the significance of these expansions for future work in diffraction theory.

ACKNOWLEDGMENTS

Many of the results contained in this report had their beginnings in the period of several years which preceded the one-year period of Contract AF 19(628)-4393. The author wishes to express his gratitude to the three department managers under whom he served during these years when the research was supported by the Lockheed Independent Research Program. Without the encouragement and understanding of Messrs. E. A. Blasi, A. S. Dunbar and N. J. Gamara this work would never have been carried out to the degree of completeness at which is now stands. During the past year (while the work has been supported by the contract with the Air Force Cambridge Research Laboratories), the author has had the privilege of working under the management of Mr. A. S. Dunbar.

Although the work of the last year has been carried out solely by the author, many of the results reported herein are the fruits of the labors of Mr. R. S. Mason and Dr. K. S. Yee.

Some of the equations in this report were typed by Miss Ann Youngs and the author gratefully acknowledges his endebtedness to her for her diligence in transcribing handwritten equations into typescript. The author is also endebted to Mrs. Bette DeGand for the typing of several of the tables and to Mrs. Marion Hegeman for assistance in administrative details.

RELATED CONTRACTS AND PUBLICATIONS

1. Logan, N.A., "Survey of Some Early Studies of the Scattering of Plane Waves by a Sphere," Proc. of IEEE, Vol. 3, Aug 65, pp. 773-785. Contract No. AF 19(628)-4393

PREFACE

Although this report is officially entitled a "final" report upon Contract AF 19(628)-4393, the report is in fact an "interim" report upon a subject which has attracted the author's attention for over ten years. Prior to the one-year period during which the research has been carried out under the present contract, the author undertook work along these lines as part of the Independent Research Program of the Lockheed Missiles and Space Company. The attainments made to date fall far short of the author's goals. However, the author hopes to continue to strive towards filling some of the gaps that exist between the results in both the asymptotic theory of diffraction by convex surfaces and the numerical analysis which is required in order to translate the esoteric mathematical results into curves and tables that will meet the needs of the practical engineer. The author readily admits that the opaque convex surface is an over-idealized model for any except the simplest of propagation and diffraction problems. However, the actual problems that confront most engineers in this field are so complex that there is a great demand for information concerning the phenomena associated with a smooth convex surface. The present research efforts have been directed towards extending the algorithms and the computer programs that will permit us to **obtain numerical results when the scatterer is an opaque surface which is characterized by an impedance boundary condition.**

The problem of the diffraction of electromagnetic waves by convex surfaces has been the subject of hundreds of papers since the appearance over 100 years ago of a classic paper by A. Clebsch which can be considered to be the "first" paper upon this subject. The

author recently had the honor of having a survey of some of the early history of this problem to appear as the lead article in a "Special Issue on Radar Reflectivity" which was published as the August, 1965, issue of the Proceedings of the Institute of Electrical and Electronics Engineers (Proc. IEEE, Vol. 53, No. 8, Aug., 1965). The author would like to encourage his readers to read this historical survey. If the reader is not already familiar with this special issue, the author is certain that the very fact that the reader has been interested enough in this report to have read this far is a very sure indication that the "Special Issue on Radar Reflectivity" will provide the reader with a "gold mine" of information.

The theory of diffraction by convex surfaces cannot be readily separated from a large body of work in applied mathematics. Therefore, the engineer who finds himself facing a practical problem involving diffraction by a convex surface finds that he is confronted with an almost chaotic mass of literature in which there are many gaps in the practical information to say nothing of the fact that there is a lack of a coherent and effective general theory. It is with regrets that the present author finds himself in the position of adding to the confusion and the frustration of the "newcomers" by releasing the present material in a form such that a considerable background is required in order to pick out results which may be helpful in a particular practical problem.

The author would like to illustrate his sympathy for the reader who is a "newcomer" to the diffraction field by relating an experience which occurred during the course of this contract. The author is frequently asked to recommend books, reports, and papers which can provide an engineer with a means of learning what results are currently available for him to use in connection with problems associated with diffraction by a convex surface.

For several years, the author has been telling his colleagues that the book that they were seeking was about to be published as Vol. 1, Diffraction by Convex Surfaces in the International Series of Monographs on Electromagnetic Waves which was begun several years ago by Pergamon Press. The author of this book was to have been Academician V. A. Fock of Leningrad University. The present author* had been considerably influenced by a series of papers by Fock which he had discovered in the mid-1950's. With the cooperation of P. Blacksmith, M. D. Friedman, and others, the author had made available to a limited circle of readers a collection** of 13 papers by Fock, together with a summary of Fock's contributions by Academician V. I. Smirnov and an appendix which contained a pertinent paper by Academician M. A. Leontovich. It was the understanding of the present author that when Fock visited the United States in the late 1950's that he had refused permission for the collection of papers to be issued in book-form on the grounds that the papers were incoherent and that he intended to prepare a monograph in which a unified presentation of his research would be presented. However, even behind the "Iron Curtain" the scientists apparently find the days and nights passing

---

*Although the present author has since become sufficiently acquainted with the literature to realize that much that he learned from Fock's papers could have been learned from papers by White, Pekeris, Pryce, Furry, and others, the author recognizes the fact that the clarity of presentation and the thoroughness of the investigations which were found in Fock's papers were such that he can best describe Fock's influence upon his own work by asserting that Fock was his "teacher."

**Air Force Cambridge Research Center, Diffraction, Refraction, and Reflection of Radio Waves: Thirteen Papers by V. A. Fock, N.A. Logan and P. Blacksmith, eds., Report AFCRC-TN-57-102, ASTIA Document No. AD117276, Bedford, Mass., June, 1957

in such swift processions that one has to sometimes face reality and admit that the goal which one has set for oneself is too ambitious. Anyway, it was a great disappointment to the present author when in mid-1965 the long-promised Pergamon Press volume appeared in print.* This monograph is nothing more than 17 papers by Fock (with several having been written with co-authors) which (except for minor corrections) appear in their original form. Except for the inclusion of papers published since 1957, the book which is now available is essentially the same collection of "raw" reference material which the present author and his associates had sought to provide to scientists in the United States when the 391-page paperbound AFCRC report was distributed in 1957. The author strongly urges the serious student of diffraction theory to acquaint himself with these papers by Fock. However, since these are research papers and the material has not been put together in "textbook style," the reader may find Fock's book a difficult source from which to acquire a basic knowledge and background upon which to build the ability to read the hundreds of papers which are available in this subject area. Furthermore, since no effort is made to relate the material in Fock's book to the more recent publications of other authors, the reader will have to "forage" for himself to learn what other results are available to him. Making a literature survey requires considerable "detective work" since a number of the modern authors have worked independently of one another and cross-references to each other's works are often conspicuous by their absence.

There are sufficient gaps in our present state of knowledge in

---

*V. A. Fock, Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965

this subject area that any book or monograph published today would be virtually out-of-date by the time it appeared in print. This is particularly true at the present time when the availability of the electronic computer is making it possible for us to attempt to obtain numerical results which were "impossible" during World War II when the widespread use of high frequency radio waves created a need for results from diffraction theory for those cases in which classical geometrical optics failed to provide an explanation of the observed phenomena. As an example of the progress which has been made in the computer field, it is interesting to cite the fact that Aiken* made the observation in 1945 that "...if it had been possible to run the computation without interruption forty-five days would have been sfficient time to complete the tables..." of Airy functions which Furry and other scientists needed in their studies of wave propagation. One of the FORTRAN IV programs which is given in this report could be used on the UNIVAC 1107 to produce these identical tables in approximately thirty minutes. The author can also cite a more recent example based upon his own experience. Upon some output sheets from the IBM-650 which were used to prepare the curves for the scattering properties of perfectly conducting spheres which appear in the appendix of the monograph of King and Wu** there appear some

---

*The Staff of the Computation Laboratory, Tables of the Modified Hankel Functions of Order One-Third and of Their Derivatives, Cambridge, Mass., Harvard University Press, 1945

**R. W. P. King and T. T. Wu, The Scattering and Diffraction of Waves, Cambridge, Mass., Harvard University Press, 1959. (Graphs in appendix illustrate current distributions and bistatic cross sections for perfectly conducting spheres for ka = 1.1(0.6)9.5, 10, and 20.)

penciled notes to indicate that these computations required forty-five minutes for a set of E-plane and H-plane curves for a single value of the parameter ka. It is now possible to generate this same data on a computer such as the UNIVAC 1107 in less than forty-five seconds.

However, as the present author looks back at his computing experience, his respect for the amazing speed of the modern electronic computer does not leave him standing in awe as much as does the realization that programming languages such as FORTRAN IV offer the modern scientist a powerful tool which was virtually beyond his wildest dream less than 20 years ago. One has to have grown-up with the "computer age" to really appreciate what a change has been made possible by the development of mathematically-oriented programming languages which enenable a scientist who is not a "computer specialist" to perform millions of mathematical manipulations at the rate of thousands per second.

The author's first major computation job involved some diffraction patterns for a circular cylinder and the work was carried out in the late 1940's by a group of ladies who worked for months at Friden and Marchant desk machines. Even when two ladies would work in tandem on worksheets which were laid out in identical manners, the detection of errors was a very frustrating task. With human hands and minds at work, it was virtually impossible to eliminate blunders caused by an error in lookup in a table, in transfering a number into the keyboard, copying the number from the register onto the work sheets, an error in placing a decimal point, and, ABOVE ALL, one had to contend with frequent replacement of the computer personnel so that with even a moderate size group there was usually always a "new girl." Anyway, much of this was drastically changed a few years later when the author

set outto compute the scattering diagrams which are contained in the appendix of the monograph by King and Wu. The author had started some of these computations on desk machines of the Friden and Marchant variety, but he was bogged down in trying to detect the errors so as to be able to construct the scattering patterns. At this point, the services of G. E. Reynolds and M. R. Hoes were enlisted and they came up with a set of instructions which made it possible to obtain the desired computations-free of human error- on the IBM-650 computer. Although the manner in which these programmers managed to get the computer to perform was, to the author, a sort of "black magic," the author does recall the excessive handling of thousands of IBM cards as different portions of the computation were handled by the limited "memory" of the computer and the final results obtained by having hundreds of cards read back into the machine and the final answers assembled. This was a far cry from the tedious and frustrating manner of trying to obtain the results by hand, but the specialists trained in programming the computer were an essential part of the team.

The development of the FORTRAN, ALGOL, MAD, and similar "languages" has made it possible for the scientist, if he so desires (and budget conditions can sometimes make desire into a necessity), to dispense with the programmer for many of the day-to-day type of calculations. The introduction of these languages have had an important effect in that they are generally languages which are independent of a particular computer and hence a scientist who has developed a program for his computer (which might be a UNIVAC 1107, for example) may be able to share the program with another scientist who has acess to a totally different computer (such as an IBM 7094). This is a remarkable new aspect of cooperation between scientists and makes it possible for one to build upon the experience of other workers. In the days of hand computation

many of the present concepts of programming were in use. A worksheet properly laid out with all the instructions explicitly displayed at the headings of the proper columns constituted an algorithm (a set of explicit rules for carrying out a computation) for the numerical problem which was under consideration. However, except for the layout for certain problems such as an harmonic analysis of a set of data, it was very uncommon for one scientist to pass on to another his "set-up" for the arriving at the numbers which he presented in his tables and graphs.

The situation has now markedly changed with the development of the new programming languages. The Communications of the Association for Computing Machinery have been supplying a medium for the publication of algorithms written in the ALGOL language for the past several years. However, the largest cooperative effort in making programs available for use by persons or groups other than the initiators is the IBM SHARE system. The rapid advances in computers in the past several years has made it almost mandatory that a program which is to be effectively shared with another user be written in a standard language. For example, the use of a machine-oriented language should be avoided whenever possible. The present author knows of one extremely large program which was coded in the FORTRAN ASSEMBLY PROGRAM (FAP) language and the need has now arisen to convert this program to be used on a computer which uses a different assembly language. The net result is that a team of programmers are engaged in translating a FAP-coded program into FORTRAN IV so as to be able to have the program compile upon other computers than the IBM-7090-7094 for which it was originally prepared. Because compilers can be expected to change relatively rapidly over the next few years, the use of a standard language such as FORTRAN IV is highly recommended. Although the different compilers may accept only certain "dialects" of FORTRAN IV, the changes which must be made in going from one compiler to another are rather minor if one restricts oneself to

the use of features which are available on the widest class of compilers. A useful reference for FORTRAN IV is the report* of the American Standards Association (ASA) FORTRAN Working Group. Their report lists side-by-side a set of standards for FORTRAN (which is FORTRAN IV) and BASIC FORTRAN (which is FORTRAN II). McCracken** has attempted to collect in tabular form some of the characterisitics of over two dozen programming languages which are being referred to as FORTRAN IV. Unfortunately, in many compilers FORTRAN II is not solely a subset of FORTRAN IV and hence not all FORTRAN II programs can be compiled on software which compiles FORTRAN IV programs. The language situation promises to be a bit more confused in mid-1966 when the manuals and the compilers will be available for IBM Operating System/360. Although this new language has been called FORTRAN VI, NPL (New Programming Language), and MPPL (Multi-Purpose Programming Language), the most recent announcements apparently refer to it as PL/I (Programming Language/One). However, some of the early indications are that FORTRAN IV as it is now defined will most likely be a subset of the more versatile IBM programming language.

The carrying out of certain portions of the work of the present contract has been hampered by the fact that much of the work prior to this contract had involved the use of a set of FAP-coded SHARE routines known as NPREC. The conventional FORTRAN II compilers did not provide means for the input and output of double precision numbers, and the floating point numbers were required to lie within the range 10^{-38} and 10^{38}. We had not only

---

*W. P. Heising, "History and Summary of FORTRAN Standardization Development for the ASA," Comm. Assoc. Comput. Mach., Vol. 7 Oct., 1964, pp. 590-625

**D. D. McCracken, "How to Tell If It's FORTRAN IV," Datamation, Vol. 11, Oct., 1965, pp. 38-41

the need, in certain algorithms, to greatly exceed the range of the numbers permitted in FORTRAN, but we also wanted to have the ability to input and output more than the 8-digits that were maximum number of digits which the FORTRAN compiler would handle. The author was confronted with several problems where the cancellation of numbers of comparable size was so complete that answers to even several significant figures could not be obtained with the relatively simple algorithm unless many significant figures could be manipulated. Nearly a decade ago the author showed that the use of high precision could be used to sum certain divergent series in diffraction theory and this led to the development at the Air Force Cambridge Research Center of the capability of working with quadruple precision numbers which consisted of floating point numbers of 37 significant figures.* However, the capability developed at AFCRC was limited to the computer available to them because this was a machine-language coded program. It was not until 1961 that the author learned of the existence of a FAP-coded SHARE program which would enable a FORTRAN programmer to compute in double precision floating point arithmetic. This very ingenious program had been prepared by R. I. Berggren, J. C. Cysbers, R. Hafner, and L. Sonneville of the Rocketdyne Division of North American Aviation, Inc. The program used three consecutive 36-bit words to store a single number, with one word being used for the exponent and two words for the fraction. This arrangement made it possible to work with floating point numbers which consisted of more than 21 decimal digits, and with exponents whose absolute value could not exceed $(2^{35}-1) \approx 3.4 \times 10^{10}$. Although this set of subroutines could be employed in a FORTRAN II program, its use was rather clumsy. When it was announced in the Fall of 1964 that the UNIVAC 1107 which was to be installed at Lockheed in the Spring of 1965 was to have a double-precision capability of

*Air Force Cambridge Research Center, Cambridge Computer Interpretive Routine for Quadruple Precision Numbers, Report TN-59-155, 1959

21-significant decimal digits, the author resolved to attempt to convert to FORTRAN IV for the UNIVAC 1107 and carry out much of the previous work which had involved the use of the NPREC subroutines by employing the relatively simple double-precision of the new computer. However, upon delivery it was found that the UNIVAC 1107 performed double-precision by software operations which limited the precision of the numbers to 17 decimal digits. Nevertheless, the greater simplicity of the use of FORTRAN IV as compared with the use of FORTRAN II supplemented by the NPREC subroutines (together with the constant anxiety to the effect that a systems change on the IBM 7094 might result in the NPREC subroutine, coded in FAP, no longer being compatible with the system tapes being installed in the more advanced system monitors being made available by IBM) led the author to seek to continue to strive towards the use of the FORTRAN IV compiler for the UNIVAC 1107. Although the loss of 4 to 5 significant figures when the FORTRAN IV program was coded to replace the FORTRAN II-NPREC program proved to limit a few of the problems that could be handled, two far more important limitations appeared. Since the manipulations had been almost solely performed internally in the computer and the magnitude of the numbers involved in many of the algorithms had not been known to the author, it came as quite a shock to realize that many of the algorithms which had worked successfully when using the NPREC subroutines could not be used at all on the UNIVAC 1107 (or, on the FORTRAN IV compiler for the IBM 7094) because the exponents exceeded the magnitude of 10^{38} which is an inherent property of both the IBM and the UNIVAC compilers. Most compiler designers have apparently decided that the need for the manipulation of such large numbers is restricted to such special applications that only the compilers for the Control Data Corporation (CDC 3000 and 6000 series) and Philco (2000 series) computers take into consideration these needs, and even here the restrictions are to 10^{308} and 10^{616}, respectively.

A second surprise that came with the attempts to use the more sophisticated UNIVAC 1107 computer was the receipt from the compiler of messages reading "...UNRESOLVABLE AMBIGUITY CAUSED BY SOURCE LANGUAGE ERROR..." when attempts were made to run certain FORTRAN II-NPREC programs on the UNIVAC 1107 after the conversion process (consisting mainly of changing read and write statements and the making of allowances for some minor changes such as the replacement of COSF(X) by COS(X) for single-precision cosines and DPCOS(X) by DCOS(X) for double-precision cosines) had been carried out. The difficulty can be traced to several new sophistications of the newer software and hardware of the new generation of electronic computer of which the UNIVAC 1107 is a typical example. With the newer compilers, one is dealing with both new hardware and new software. One of the hardware difficulties apparently arises from the fact that greatly increased operation speed is achieved by using a set of film registers to store the subscripts which are used during the execution of a DO-loop. This means that for most programs a very significant saving in computer time is saved since the subscripts are not computed, stored in memory, and then retrieved as was the case in the earlier hardware. However, if one has made excessive use of subscripts in a DO-loop, the system apparently becomes saturated and informs the programmer that an unresolvable ambiguity has been encountered. Therefore, in such cases the original algorithm has to be recoded and broken down into smaller units. A not-too-dissimilar situation can occur with rather long arithmetic expressions that involve a great deal of multiplying, dividing, raising to a power, etc., and the user has been lax in the use of parentheses. The compilers have apparently been designed to try to "look ahead" and optimize an expression and it is possible that a lengthy expression which was compiled by "brute force" on an earlier FORTRAN II compiler may result in a message regarding an unresolvability in the source language

when submitted to a more sophisticated compiler. This latter type of difficulty is readily avoided by using parenthesis wherever they help the "human reader" to understand what is required and the result will generally be that the "inhuman compiler" will also encounter less difficulty in understanding what it is that the programmer wishes to have the computer do with his lengthy set of instructions. Although it is discouraging to find that a program that already works on a less sophisticated compiler and computer may need to be drastically modified to be acceptable upon the newer equipment, the greater flexibility of the FORTRAN IV language and the greater speed of compilation and of execution of the generation of computers more than offsets the inconveniences.

However, the difficulties associated with the fact that the number of decimal digits in a double-precision number is 16 for the IBM 7094 and 17 for the UNIVAC 1107, while the magnitude of the numbers may not exceed 10^{38}, remains a serious limitation for a number of computing algorithms. In spite of the fact that its use is somewhat clumsy, the author would like to see the old NPREC subroutines revised so as to be compatible with FORTRAN IV and be acceptable to the "systems tape" on which is "located" the FORTRAN IV compiler. Although the author knows of no efforts which will lead to this capability for the UNIVAC 1107 or 1108, the author knows that a group at the Ames Research Center of the National Aeronautics and Space Administration are in the process of preparing a FORTRAN IV version of the NPREC subroutines which will go several steps beyond the original NPREC subroutines and which may permit the use of triple- and quadruple-precision. When this set of subroutines has been debugged and submitted to the IBM SHARE library, the author looks foward to employing it for many problems which have been currently stymied because the algorithm known to the author would require drastic revision before the quantity can be computed with the normal range of precision and magnitude that the FORTRAN compilers have been designed to

cope with in their everyday use. That this class of problems is not purely academic and confined to only a few problems such as those encountered by the present author can be seen by observing that there have recently appeared several papers that deal with "arbitrary precision" floating-point computations and the associated input-output conversion techniques.*

Further evidence of the fact that this type of increased facility for the electronic computer is more widespread than one might imagine at a first consideration of the problem can be seen from the fact that R. K. Holmberg of the National Bureau of Standards, Boulder, Colorado, has prepared a special set of IBM 7090 subroutines which are known as the EPREC or "Extended Precision Arithmetic Package." Whereas the NPREC package from North American Aviation used two words for the fraction and one word for the exponent, the EPREC package requires just one word for the fraction and one word for the exponent. The result is that the precision of the number is essentially the same as in conventional FORTRAN, but overflow and underflow problems are postponed until much larger numbers are encountered since the routine will handle numbers as large as 10^{4920}.

The drawback of the use of software such as the NPREC and EPREC subroutines is that a considerable amount of the core storage (it takes nearly 4000 "words" to store the NPREC subroutines) is consumed in making the subroutines available. The author hopes that in the near future that the computer manufacturers

---

*A. H. Stroud and D. Secrest, "A Multiple-Precision Floating-Point Interpretive Program for the Control Data 1604," Computer J., Vol. 7, Jan., 1964, pp. 62-66

Y. Ikebe, "Note on Triple-Precision Floating Point Arithmetic with 132-Bit Numbers," Comm. Assoc. Comput. Mach., Vol. 8, March, 1965, pp. 175-177

will become more aware of the special needs of scientific computation and build in the ability to handle numbers of greater precision and larger range in the hardware itself. At least one manufacturer seems to have made a good start in this direction, namely the Honeywell Series 200. The programmer can specify by a parameter card at the beginning of the source program an integer N which will determine the number of "characters" to be employed for each quantity which is the appear in the program that follows. Each "character" consists of 6 binary bits and hence the fact that any number of "characters" between 3 to 12 can be specified means that the programmer can specify by means of a control card whether the program is to use floating point numbers that range from an accuracy of 5 digits to 20 digits. The range of normalized floating point values is given as 10^{-616} to 10^{+616} in a recent brochure on the Honeywell Model 4200. It is interesting to observe that because of the variable word length made available to the programmer, that the "dialect" of FORTRAN IV which is used on this Honeywell computer does not include the type declarations COMPLEX and DOUBLE PRECISION. It is obvious that the need for double precision has been eliminated because of the choice of word length that is made available to the programmer, but the denial of the complex arithmetic ability found in other FORTRAN compilers means that the programmer must work out the real and imaginary parts of all quantities. However, the author can forsee the day when a later generation of this compiler will have the complex arithmetic built into the compiler. The author hopes that other compiler designers will see the advantages to the scientific computer of having the ability to control the precision by means of a "control card" and that the need for clumsy subroutines (written in assembly languages such as FAP for the IBM-7094 and in SLEUTH for the UNIVAC 1107) to perform calculations of high precision with "large" numbers will be eliminated in a future generation of FORTRAN IV compilers.

The time "ran out" on the present contract before the author had the opportunity to employ the Stromberg-Carlson 4020 film plotter which is available as part of the computer facilities at LMSC. A subroutine has been prepared by M. H. Seigel* for a customer on another contract who wished to make contour plots (similar to the familiar geological contour maps) of the radar reflectivity of certain targets as a function of the polar angles θ and the azimutal angles φ, when the angles are treated in the contour plot as a number pair (θ,φ) in a complex plane. It is hoped that in future work that this mode of presentation will be frequently employed in the author's efforts to depict to his readers the behavior of diffraction phenomena as a function of the complex-valued impedance parameter. This new plotter opens up a broad new field of applications for the visual presentation of data. The reader is urged to consult Fig. 1 of a recent paper by Chezem** where it is shown that the "beautiful" three-dimensional presentations of the behavior of functions which have long intrigued physicists who thumb through the pages of Jahnke and Emde have now been brought within the reach of any programmer who has access to a versatile electronic plotter such as the SC 4020.

The author hopes to continue the work which is reported upon in this document. He looks foward to the day when sufficient knowledge of the means by which the required functions can be evaluated will have been acquired so that the user need only follow a "cook-book" type of set of instructions to obtain numerical results for a practical problem. Just as the housewife seldom

---

*Lockheed Missiles and Space Company, Digital Contour Plotting Techniques, by M. H. Seigel, Report ISSA-507, Sunnyvale, Calif., June, 1965

**C. G. Chezem, "Note on "Three-Dimensional" Plotting As a Technique for Finding the Zeros of Functions in the Complex Plane," Computer J., Vol. 8, Oct., 1965, p. 288

has any knowledge about the complicated chemistry taking place in her cooking utensils, so also the electronics engineer seldom has a need to actually know the fine details of asymptotic expansions, Airy functions, etc., in order to arrive at the guidelines which he requires from theory provided he can have before him the proper "cook-book" type of instructions as to how to obtain the numerical information which will describe his physical phenomena. There has already been an enormous amount of work reported upon by investigators who have found diffraction theory a fertile area from which to pluck a subject to develop into a paper for publication. However, in spite of the large amount of tables and graphs which are available, the investigator who seeks to obtain some results "...for an engineering study which must be completed by a week from Friday..." will usually find his situation quite hopeless. When he goes to the library to do his literature survey the chances are quite good that he will find scores of papers which are quite confusing because although the authors are treating essentially the same problems by essentially the same methods, they persist in using their own personal preferences for the notation for the various functions involved and hence a comparison of the results of the papers will be quite difficult. Then, after he has mastered the papers (long after the "due date" for the engineering study) he will almost certainly find that certain difficult aspects of the theory have been glossed over and that the filling in of these gaps requires considerable experience, mathematical ingenuity, and competence in numerical analysis and computer programming.

The author has worked at this subject area for long enough, and he is aware of his own personal limitations, and therefore his dreams of providing a "relatively simple" set of rules and programs for the practical engineer have become less ambitious with the passing of several years. However, the dream persists and the author hopes that his next reports will come closer to helping his engineering colleagues than does this present report.

CONTENTS

Section		Page
	ABSTRACT	iii
	ACKNOWLEDGMENTS	v
	RELATED CONTRACTS AND PUBLICATIONS	vi
	PREFACE	vii
	LIST OF ILLUSTRATIONS	xxix
	LIST OF TABLES	xxxi
	LIST OF FORTRAN PROGRAMS	xxxvii
	LIST OF FORTRAN BLOCKS*	xxxix
1	THE DIFFRACTION FUNCTIONS	1-1
	References for Section 1	1-34
2	THE ROOTS t_s OF $w_1'(t_s) - q\, w_1(t_s) = 0$	
	2.1 Notation for the Root Defining Equation	2-1
	2.2 The Riccati Equation	2-7
	2.3 The Root Locus for the Ground Wave Problem	2-36
	2.4 Numerical Values for the Classical Expansions	2-56
	2.5 Programs for Step-by-Step Integration	2-94
	References for Section 2	2-101
3	EXPANSIONS IN THE LIGHTED REGION OF A CIRCULAR CYLINDER	3-1
	References for Section 3	3-35

*BLOCKS are FORTRAN subprograms which are used to enter data into labeled COMMON. They contain no executable statements and are essentially just tables of data which are read into the computer storage at the time of making the compilation.

Section		Page
4	ASYMPTOTIC EXPANSIONS FOR THE HEIGHT GAIN FUNCTION FOR THE CASE OF NORMAL REFRACTION	
	4.1 Introduction	4-1
	4.2 Problems Leading to the Height Gain Function	4-1
	4.3 Survey of Some Previous Work on the Height Gain Function	4-6
	4.4 The Research Reported by Fock	4-12
	4.5 The Contributions of Friedlander	4-15
	4.6 Determination of the $H_n(u,\tau)$	4-19
	4.7 Determination of $\eta(u,t)$	4-25
	4.8 Application to Bessel Functions	4-41
	4.9 Numerical Aspects of Employing These Expansions	4-47
	References for Section 4	4-48
5	ASYMPTOTIC EXPANSIONS FOR THE PROPAGATION CONSTANTS AND THE NORMALIZATION INTEGRAL	
	5.1 The Eigenvalue Problem	5-1
	5.2 Another Expansion for λ_s	5-6
	5.3 The Turning Points	5-11
	5.4 The Propagation Constants	5-13
	5.5 The Characteristic Wave Numbers	5-14
	5.6 The Normalization Integral	5-19
	5.7 Direct Integration of N_s	5-31
	5.8 Eigenvalues Derived from the Langer Approximation	5-34
	References for Section 5	5-38
6	HEIGHT GAIN FUNCTIONS FOR LARGE HEIGHTS	
	6.1 The Langer Asymptotic Estimate	6-1
	6.2 The W.K.B. Approximation	6-4
	6.3 Some Further Forms for $F_{1,2}(y,\lambda)$	6-7
	References for Section 6	6-15

Section		Page
7	ASYMPTOTIC ESTIMATES FOR $U(x,y;x_o,y_o)$	
	7.1 The Diffraction Functions	7-1
	7.2 The Reflection Formula	7-3
	7.3 The Diffraction Formula	7-15
	7.4 A Digression on Hankel Functions	7-23
	7.5 The Extended Height Gain Approximation	7-30
	7.6 Extension of Bremmer's Results	7-32
	References for Section 7	7-39
8	CHEBYSHEV POLYNOMIAL REPRESENTATIONS	
	8.1 Introduction	8-1
	8.2 Polynomial Expansions	8-4
	8.3 Properties of Chebyshev Polynomials	8-9
	8.4 Summing Series of Chebyshev Polynomials	8-11
	8.5 Further Remarks on the Clenshaw Algorithm	8-17
	8.6 Determination of Coefficients	8-22
	8.7 The Future for Diffraction Theory	8-30
	References for Section 8	8-41
Appendix		
A	FORTRAN AND NUMERICAL ANALYSIS	A-1
B	TABLES OF THE AIRY FUNCTION ON THE REAL AXIS	B-1
C	TABLES TO FACILITATE HAND COMPUTATION OF THE ROOT t_1 FOR THE CASE OF A REAL EARTH	C-1
D	TABLES OF VALUES OF ROOTS OF $w_1'(t_s) - qw_1(t_s)$	D-1
E	THE HEIGHT GAIN FUNCTIONS FOR $q = 0$ AND $q = \infty$	E-1
	References for Appendix E	E-50

LIST OF ILLUSTRATIONS

Figure		Page
1-1	Variation of the Field Strength Near The Horizon	1-5
1-2	Source on the Surface of a Spherical Interface	1-6
1-3	Plane Wave Incident Upon a Convex Surface	1-7
1-4	Behavior of $V_1(0,q)$	1-9
1-5	Reflection from a Convex Surface	1-11
1-6	Behavior of $V_2(0,q) = \lim_{\xi \to C} \left\{ \frac{\exp(i\pi/4)}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t) \frac{v'(t) - q\,v(t)}{w_1'(t) - q\,w_1(t)}\, dt - \frac{\exp(i\pi/4)}{2\sqrt{\pi}\,\xi} \right\}$	1-19
1-7	Attenuation Function $u(\xi)$ (Real and Imaginary Parts)	1-25
1-8	Attenuation Function $u(\xi)$ (Modulus)	1-26
1-9	Attenuation Function $v(\xi)$ (Real and Imaginary Parts)	1-27
1-10	Attenuation Function $v(\xi)$ (Modulus)	1-28
2-1	The Logarithmic Derivative $q(\tau) = w_1'(\tau)/w_1(\tau)$	2-30
2-2	The Logarithmic Derivative $q(\tau) = w_1'(\tau)/w_1(\tau)$	2-31
2-3	Boundaries for Regions of Applicability of Eqs. (2-21), (2-22) and (2-26)	2-35
2-4	The Logarithmic Derivative $L(t) = Ai'(t)/Ai(t)$	2-37
2-5	The Root Locus of $\tau_1[Q\exp(i45^\circ)]$ for $0 < Q < 0.4$	2-46
2-6	Loci of the Roots in Vicinity of $\tau_1^\infty = 1.169 + i2.025$ for Propagation Around the Earth's Surface	2-48
3-1	Geometry of the Reflection Problem	3-1
3-2	The Geometry as Defined by Keller, Lewis and Seckler (Ref. 8)	3-17

Figure		Page
7-1	Ray Paths in Inhomogeneous Medium	7-21
8-1	Behavior of the Airy Functions	8-33
B-1	The Auxiliary Functions $\chi(x)$ and $\psi(x)$	B-76
B-2	The Behavior of $\chi'(t)$ and $\psi'(t)$	B-77

LIST OF TABLES

Table		Page
1-1	Values of A_n and B_n Occurring in Asymptotic Expansion of the Logarithmic Derivatives of Ai(α) and Ai′(α)	1-30
1-2	The Attenuation Function $u(\xi)$	1-31
1-3	The Attenuation Function $v(\xi)$	1-32
2-1	Roots and Turning Values for Ai(-α)	2-9
2-2	Roots and Turning Values for Ai′(-β)	2-10
2-3	Coefficients for Polynomial Representation for $A_n(\tau)$	2-12
2-4	Coefficients for Polynomial Representation for $B_n(\tau)$	2-19
2-5	The Critical Points τ_s^c and Corresponding Values of q_c	2-32
2-6	The Root τ_1 As Calculated by Admiralty Computing Service	2-44
2-7	Table of Roots $\tau_1(q)$ Obtained by Johler, Walters and Lilley	2-45
2-8a	Table of Values of the Parameter log n Showing Dependence upon Physical Nature of the Soil and on the Wavelength λ	2-51
2-8b	The Function $q(\alpha,n) = in^{5/6}/\sqrt{i + \alpha n}$	2-52
2-9	Coefficients in the Expansion of $\tau_s(q)$ in Series of Powers of q	2-58
2-10	Coefficients in the Expansion of $\tau_s(q)$ in Series of Powers of $\frac{1}{q}$	2-61
2-11	The Coefficients $C_n(\tau_1^o)$ As Computed in Single Precision	2-66

Table		Page
2-12	The Coefficients $D_n(\alpha_1)$ and $F_n(\alpha_1)$	2-72
2-13	The Coefficients $C_n(\beta_1)$ and $E_n(\beta_1)$	2-78
2-14	Coefficients for Polynomial Representation for $F_n(\tau)$	2-79
3-1	Leading Terms in the Asymptotic Expansion of $A_n(p,0)$	3-6
3-2	Leading Terms in the Asymptotic Expansion of $A_n(p,\infty)$	3-7
3-3	Coefficients B_{jt} and C_{jt} for the Dirichlet Problem	3-19
3-4	Coefficients D_{jt} for the Dirichlet Problem	3-20
3-5	Coefficients E_{jt} for the Dirichlet Problem	3-20
3-6	Coefficients F_{jt} for the Dirichlet Problem	3-22
3-7	Coefficients G_{jt} for the Dirichlet Problem	3-23
3-8	Coefficients B_{jt} and C_{jt} for the Neumann Problem	3-26
3-9	Coefficients D_{jt} for the Neumann Problem	3-27
3-10	Coefficients E_{jt} for the Neumann Problem	3-27
3-11	Coefficients F_{jt} for the Neumann Problem	3-28
3-12	Coefficients G_{jt} for the Neumann Problem	3-30
4-1	Particular Solutions of $\frac{d^2y}{dx^2} - xy = f(x)$	4-21
8-1	Chebyshev Coefficients for exp(x)	8-30
B-1	The Airy Functions ($10 < x < 25$)	B-29
B-2	Derivative of the Airy Function ($10 < x < 25$)	B-33
B-3	The Airy Functions ($0.001 < \frac{1}{x} < 0.100$)	B-37
B-4	Derivative of the Airy Functions ($0.001 < \frac{1}{x} < 0.100$)	B-40
B-5	The Auxiliary Functions Q(x) and M(x)	B-43
B-6	The Auxiliary Functions P(x) and A(x)	B-46

Table		Page
B-7	The Auxiliary Functions R(z) and S(z) for $0.001 < \frac{1}{x} < 0.100$	B-49
B-8	The Auxiliary Functions W(z) and X(z) for $0.001 < \frac{1}{x} < 0.100$	B-50
B-9	The Airy Functions Ai(-x) and Bi(-x) for $0 < x < 10$	B-52
B-10	Airy Function Derivatives Ai'(-x) and Bi'(-x) for $0 < x < 10$	B-58
B-11	The Auxiliary Functions F(-x) and $\chi(-x)$ for $0 < x < 10$	B-51
B-12	The Auxiliary Functions G(-x) and $\psi(-x)$ for $0 < x < 10$	B-64
B-13	The Airy Functions Ai(-x) and Bi(-x) for $x = 10.0(0.1)25.0$	B-67
B-14	The Airy Derivatives Ai'(-x) and Bi'(-x) for $x = 10.0(0.1)25.0$	B-71
B-15	The Auxiliary Functions F(-x) and $\chi(-x)$ for $x = 10.0(0.1)25.0$	B-75
B-16	The Auxiliary Functions G(-x) and $\psi(-x)$ for $x = 10.0(0.1)25.0$	B-79
B-17	The Functions $\exp(\pm\xi)$ with $\xi = \frac{2}{3} x^{3/2}$	B-83
B-18	The Functions $\cos(\xi + \pi/4)$ and $\sin(\xi + \pi/4)$ with $\xi = \frac{2}{3} x^{3/2}$	B-84
B-19	The Coefficients $A_n(x)$ in the Classical Asymptotic Expansions	B-85
B-20	The Coefficients $A_n(x)$ in the Classical Asymptotic Expansions	B-87
C-1	Values of q_o, t_o and A_n	C-4
C-2	Values of q_o, t_o and A_n	C-31
D-1	The Roots $t_1[Q\exp(i145^\circ)]$	D-5

Table		Page
D-2	The Roots $t_2[Q\exp(i145^\circ)]$	D-8
D-3	The Roots $t_3[Q\exp(i145^\circ)]$	D-11
D-4	The Roots $t_4[Q\exp(i145^\circ)]$	D-14
D-5	The Roots $t_5[Q\exp(i145^\circ)]$	D-17
D-6	The Roots $t_1[Q\exp(i10^\circ)]$	D-20
D-7	The Roots $t_2[Q\exp(i10^\circ)]$	D-23
D-8	The Roots $t_3[Q\exp(i10^\circ)]$	D-26
D-9	The Roots $t_4[Q\exp(i10^\circ)]$	D-29
D-10	The Roots $t_5[Q\exp(i10^\circ)]$	D-32
D-11	The Roots $t_1[Q\exp(i190^\circ)]$	D-35
D-12	The Roots $t_2[Q\exp(i190^\circ)]$	D-38
D-13	The Roots $t_3[Q\exp(i190^\circ)]$	D-41
D-14	The Roots $t_4[Q\exp(i190^\circ)]$	D-44
D-15	The Roots $t_5[Q\exp(i190^\circ)]$	D-47
D-16	The Roots $t_1[Q\exp(i30^\circ)]$	D-50
D-17	The Roots $t_2[Q\exp(i30^\circ)]$	D-53
D-18	The Roots $t_3[Q\exp(i30^\circ)]$	D-56
D-19	The Roots $t_4[Q\exp(i30^\circ)]$	D-59
D-20	The Roots $t_5[Q\exp(i30^\circ)]$	D-62
D-21	The Roots $t_1[Q\exp(i300^\circ)]$	D-65
D-22	The Roots $t_1[Q\exp(i305^\circ)]$	D-66
D-23	The Roots $t_1[Q\exp(i310^\circ)]$	D-67
D-24	The Roots $t_1[Q\exp(i315^\circ)]$	D-68
D-25	The Roots $t_1[Q\exp(i330^\circ)]$	D-69

Table		Page
D-26	The Roots $t_1[Q\exp(i345^\circ)]$	D-70
D-27	The Roots $t_1[Q\exp(i350^\circ)]$	D-71
D-28	The Roots $t_1[Q\exp(i355^\circ)]$	D-72
D-29	The Roots $t_1[Q\ \exp(i1^\circ)]$	D-75
D-30	The Roots $t_1[Q\ \exp(i5^\circ)]$	D-78
D-31	The Roots $t_1[Q\exp(i10^\circ)]$	D-81
D-32	The Roots $t_1[Q\exp(i15^\circ)]$	D-84
D-33	The Roots $t_1[Q\exp(i20^\circ)]$	D-87
D-34	The Roots $t_1[Q\exp(i44^\circ)]$	D-90
D-35	The Roots $t_1[Q\exp(i46^\circ)]$	D-92
D-36	The Roots $t_1[Q\exp(i60^\circ)]$	D-94
D-37	The Roots $t_1[Q\exp(i85^\circ)]$	D-96
D-38	The Roots $t_1[Q\exp(i120^\circ)]$	D-99
E-1	Table of Coefficients $a_n(0)$ and $b_n(0)$ for $u(x,0)$ and $v(x,0)$	E-14
E-2	Samples of Tables of $f_s(y)$ and $f_s'(y)$ Published by Soviet Authors	E-20
E-3	Samples of Tables of $g_s(y)$ and $g_s'(y)$ Published by Soviet Authors	E-21
E-4	The Height Gain Function $f_s(y)$	E-31
E-5	The Height Gain Function $f_s'(y)$	E-35
E-6	The Height Gain Function $g_s(y)$	E-39
E-7	The Height Gain Function $g_s'(y)$	E-43

LIST OF FORTRAN PROGRAMS

Program		Page
2-1	SUBRØUTINE CØEFSQ(BET,N,C)	2-64
2-2	SUBRØUTINE LARGEQ(ALP,N,D,F)	2-70
2-3	SUBRØUTINE LARGEQ(ALP,D,F)	2-73
2-4	SUBRØUTINE CMRTSM(QO,TO,Q,T,EPSI,CØNVRG)	2-90
2-5	SUBROUTINE CMRTLG(XO,TO,X,T,EPSI,CØNVRG)	2-94
2-6	SUBRØUTINE RØØTSM(QOR,QOI,TOR,TOI,QR,QI, TR,TI,EPSI,CØNVRG)	2-95
2-7	SUBRØUTINE RØØTLG(XOR,XOI,TOR,TOI,XR,XI, TR,TI,EPSI,CØNVRG)	2-97
8-1	SUBRØUTINE CHEBYS(X,SUM,AO,A,N)	8-12
8-2	SUBRØUTINE CLENSW(X,SUM,AO,A,N)	8-13
8-3	SUBRØUTINE TNSTAR(X,SUM,CO,C,N)	8-14
8-4	SUBRØUTINE TNEVEN(X,SUM,EO,E,N)	8-15
8-5	SUBRØUTINE TNØDDS(X,SUM1,SUM2,FO,F,N)	8-16
8-6	SUBRØUTINE CBS2PS(N,A,B)	8-23
8-7	SUBROUTINE PS2CBS(N,A,B)	8-25
8-8	SUBRØUTINE CLNSHW(FUNCTN,A,B,N,CØEF)	8-28
8-9	Main Program to Test CLNSHW	8-29
B-1	SUBRØUTINE AIBIZ(Z,AI,BI,AIP,BIP)	B-23
B-2	SUBRØUTINE ASYPOS(X,AL,AM,AI,AIP,BL,BM, BI,BIP,ERRØR,FLØVER)	B-26

Program		Page
E-1	SUBRØUTINE STØKES(N,A,B,C,D,U,V)	E-12
E-2	SUBRØUTINE TAYSUM(T,X,WT,WPT,WTPX,N)	E-13
E-3	SUBRØUTINE TERMS(N,AR,AI,BR,BI,XR,XI,TR,TI)	E-16
E-4	Main Program to Test TERMS	E-17
E-5	SUBRØUTINE HGFY(N,Y,FYR,FYI)	E-23
E-6	SUBRØUTINE HGFPY(N,Y,FPYR,FPYI)	E-25
E-7	SUBRØUTINE HGGY(N,Y,GYR,GYI)	E-27
E-8	SUBRØUTINE HGGPY(N,Y,GPYR,GPYI)	E-29
E-9	SUBRØUTINE VTWTZ(Z,VT,WT,VTP,WTP)	E-47

LIST OF FORTRAN BLOCKS

Block		Page
2-1	CØMMØN/BREMM/BREM(18,9),BR(20,3)	2-74
B-1	The Coefficients for Asymptotic Expansion of $\theta(x)$ and $\varphi(x)$	B-14
B-2	CØMMØN/EQUADP/EAIRYF(100),EBIRYF(100),EAIRYD(100), EBIRYD(100)	B-17

Section 1
THE DIFFRACTION FUNCTIONS

This research program is concerned with the theory and application of the Green's function $D(x,y,y_o,q)$ defined by

$$\text{Differential Equation} \quad \frac{\partial^2 D}{\partial y^2} + i\frac{\partial D}{\partial x} + yD = -\delta(x)\,\delta(y - y_o) \qquad \begin{matrix} 0 \le y < \infty \\ 0 < x < \infty \end{matrix} \tag{1.1}$$

$$\text{Boundary Condition} \quad \left[\frac{\partial D}{\partial y} + qD\right]_{y=o} = 0 \tag{1.2}$$

$$\text{Radiation Condition} \quad \lim_{y \to \infty} \left[\frac{1}{\sqrt{y}}\left(\frac{\partial D}{\partial y} - i\sqrt{y}\,D\right)\right] = 0 \tag{1.3}$$

The solution of this problem involves the Airy functions

$$\left(\frac{d^2}{dt^2} - t\right) w_{1,2}(t) = 0 \tag{1.4}$$

$$w_{1,2}(t) = \sqrt{\pi}\left[Bi(t) \pm i\,Ai(t)\right] = \frac{1}{\sqrt{\pi}}\int_o^\infty \exp\left(-\frac{1}{3}x^3 + xt\right)dx \pm \frac{i}{\sqrt{\pi}}\int_o^\infty \exp\left[-i\left(\frac{1}{3}x^3 + xt\right)\right]dx \tag{1.5}$$

$$v(t) = \frac{1}{2i}\left[w_1(t) - w_2(t)\right] = \sqrt{\pi}\,Ai(t) = \frac{1}{\sqrt{\pi}}\int_o^\infty \cos\left(\frac{1}{3}x^3 + xt\right)dx \tag{1.6}$$

It is possible to express this Green's function in the form of a Fourier integral or in the form of a "normal mode" expansion which involves roots t_s which are solutions of (1.8).

$$D(x, y, y_o, q) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp(ixt) \left[w_1(t - y_>)\, v(t - y_<) - \frac{v'(t) - q v(t)}{w'_1(t) - q w_1(t)}\, w_1(t - y_o)\, w_1(t - y) \right] dt$$

$$= -i \sum_{s=1}^{\infty} \frac{\exp(ixt_s)\, w_1(t_s - y_o)\, w_1(t_s - y)}{\left[w'_1(t_s)\right]^2 - t_s \left[w_1(t_s)\right]^2} \tag{1.7}$$

where

$$w'_1(t_s) - q\, w_1(t_s) = 0 \tag{1.8}$$

In some applications, it is convenient to renormalize $D(x,y,y_o,q)$ in order to employ a function $V(x,y,y_o,q)$ which has the property $V(0,0,0,0) = 2$.

$$V(x, y, y_o, q) = 2\sqrt{-i\pi x}\; D(x, y, y_o, q) \tag{1.9}$$

$$\frac{\partial^2 V}{\partial y^2} + i \frac{\partial V}{\partial x} + \left(y - \frac{i}{2x}\right) V = 0 \tag{1.10}$$

$$\left(\frac{\partial V}{\partial y} + qV\right)_{y=0} = 0 \tag{1.11}$$

When $y_o \to \infty$, it is convenient to express $D(x,y,y_o,q)$ in the form

$$D(x, y, y_o, q) \xrightarrow[y_o \to \infty]{} \frac{\exp\left(i \frac{2}{3} y_o^{3/2} + i \frac{\pi}{4}\right)}{2\sqrt{\pi}\, \sqrt[4]{y_o}}\, E(x - \sqrt{y_o}, y, q) \tag{1.12}$$

$$E(\xi, y, q) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t) \left[v(t - y) - \frac{v'(t) - q v(t)}{w'_1(t) - q w_1(t)}\, w_1(t - y) \right] dt \tag{1.13}$$

$$\frac{\partial^2 E}{\partial y^2} + i \frac{\partial E}{\partial \xi} + yE = 0 \quad , \quad \left(\frac{\partial E}{\partial y} + qE\right)_{y=0} = 0 \tag{1.14}$$

It is often convenient to decompose $E(\xi,y,q)$ into a "plane wave" field $E_o(\xi,y)$ and a reflected field $P(\xi,y,q)$.

$$E_o(\xi, y) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t)\, v(t - y)\, dt$$

$$= \exp\left(i\xi y - i\frac{\xi^3}{3}\right) \qquad (1.15)$$

$$P(\xi, y, q) = -\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t)\, \frac{v'(t) - q\,v(t)}{w_1'(t) - q\,w_1(t)}\, w_1(t - y)\, dt \qquad (1.16)$$

1.1 Applications

The function $V(x,y,y_o,q)$ is well known in the theory of the propagation of the ground wave around the earth's surface under the conditions of "normal refraction" (i.e., in the absence of ducts and atmospheric inhomogeneities). As an example, we cite the problem of finding the Hertz potential for a vertically-directed electric dipole located at $z = r_o$ on the polar axis in the region exterior to a sphere of radius a. Let the Hertz potential in free space be described by

$$U_o = \exp(ikR)/(4\pi R) \quad , \qquad r = \sqrt{\rho^2 + (z - r_o)^2} \qquad (1.17)$$

The field in the presence of the sphere must satisfy the equations

$$\left(\nabla^2 + k^2\right)U = -\frac{\delta(\Theta)\,\delta(r - r_o)}{2\pi r_o^2 \sin\Theta} \qquad (1.18)$$

$$\frac{\partial U}{\partial r} + k\left[\left(\frac{2}{ka}\right)^{1/3} q\right]U = 0 \quad , \quad r = a \qquad (1.19)$$

The solution is generally given in the form

$$U \sim \frac{e^{iks}}{4\pi s}\sqrt{\frac{\theta}{\sin\theta}}\; V(x, y, y_o, q) \tag{1.20}$$

$$x = \left(\frac{k}{2a^2}\right)^{1/3} s, \quad y = \left(\frac{2k^2}{a}\right)^{1/3} h, \quad y_o = \left(\frac{2k^2}{a}\right)^{1/3} h_o \tag{1.21}$$

$$s = a\theta \quad , \qquad h = r - a \quad , \quad h_o = r_o - a \tag{1.22}$$

The attenuation function $V(x,y,y_o,q)$ is often written in a form which employs a so-called "height-gain" function $f_s(y)$

$$V(x, y, y_o, q) = 2\sqrt{i\pi x}\sum_{s=1}^{\infty} \frac{\exp(ixt_s)}{t_s - q^2}\, f_s(y)\; f_s(y_o) \tag{1.23}$$

$$f_s(y) = \frac{w_1(t_s - y)}{w_1(t_s)} = 1 - qy + \frac{1}{2}t_s y^2 + \cdots \tag{1.24}$$

$$f_s(y) \xrightarrow[y \gg |t_s|]{} \frac{\exp\left[i\left(\frac{2}{3}y^{3/2} - \sqrt{y}\, t_s - \frac{\pi}{4}\right)\right]}{\sqrt[4]{y}\; w_1(t_s)}\left\{1 + i\frac{t^2}{4\sqrt{y}} + \cdots\right\} \tag{1.25}$$

In the pioneering work of van der Pol and Bremmer (Ref. 1) in the late 1930's, the attenuation function $V(x,y,y_o,q)$ was quite adequate because of the low antenna heights and the relatively long radio wave lengths which were of interest . However, after the development of radar in the early 1940's it became very important to treat the case of very short wavelengths and very great heights. The calculations which were made by van der Pol and Bremmer from regions deep in the shadow to just within the lighted region had been made on the basis of (1.7). Fig. 1-1 is taken from one of the classic papers by these authors and shows how by using an increasing number of terms that the variation of the field can be followed from its region of deep shadow below the horizon to points within the lighted region where the dashed lines in Fig. 1-1 show the envelop as predicted by optical




Fig. 1-1 Variation of Field Strength Near the Horizon

theory. When the asymptotic form for the height-gain factor given by (1.25) was introduced into (1.23), the residue series, or "normal mode" expansion, failed to converge except for points below the optical horizon. Since this situation corresponded to the very practical problem of predicting the high altitude coverage of ground-based radars, the problem was very vigorously attacked by scientists in the United States, England, and in Russia. In 1941, Burrows and Gray (Ref. 2) (in a notation which is quite different from that which we are using) expressed the results of the van der Pol-Bremmer theory for this case in the form

$$U \sim \frac{\exp(i\,ka\,\theta)}{4\pi a\Theta}\sqrt{\frac{\Theta}{\sin\Theta}}\,\exp\left(i\,\frac{2}{3}y^{3/2}\right)V_1(x-\sqrt{y}\,,\,q) \tag{1.26}$$

$$x = \left(\frac{ka}{2}\right)^{1/3}\theta\,,\quad y = \left(\frac{2}{ka}\right)^{1/3}k(r-a)\,,\quad q = i\left(\frac{ka}{2}\right)^{1/3}\frac{k}{k_2}\sqrt{1-\frac{k^2}{k_2^2}} \tag{1.27}$$





Fig. 1-2 Source on the Surface of a Spherical Interface

In 1945, Fock (Ref. 3) published a similar result (in the same notation we have used) and gave a table of $V_1(\xi,0)$ for $\xi = 4.5(0.1)4.5$. It was long overlooked that **Burrows** and Gray (Ref. 2) gave a curve for $V_1(\xi,0)$ and $V_1(\xi,\infty)$ for $\xi>0$ in their original paper. The quantity plotted in their graph was defined by

$$F_L(q) = \left| \frac{\left(t_1 - q^2\right) w_1(t_1)}{\sqrt[3]{4}\; w'_1\left(t_1^{\infty}\right)} V_1\left(\frac{\left|t_1^{\infty}\right|}{|t_1|} \frac{L}{\sqrt[3]{2}}, q \right) \right| \qquad (1.28)$$

The curves given are for $q = 0$ and $q = \infty$, with $L \geq 0$.

These results of Burrows-Gray and Fock are only valid for natural units of height y of the order of two to five. For larger heights, it is desirable to use the reciprocity principle and use the fields on the surface which are induced by a plane wave. This problem was studied by Fock (Ref. 3). Fock described his results as "The Principle of the Local Field in the Penumbra Region." According to this principle: "The transition from light to shadow on the surface of the body takes place in a narrow strip along the boundary of the geometrical shadow. The width of this strip is of the order of $(\lambda a^2/\pi)^{\frac{1}{3}}$, where $\underline{a}$ is the radius of curvature of the normal section of the body in the plane of incidence. If the geometry is as illustrated in Fig. 1-3, then the principle



$$\xi = \left(\frac{k}{2a^2}\right)^{1/3} x$$

Fig. 1-3 Plane Wave Incident Upon a Convex Surface

is expressed in the following formulae.

$$H_y = H_y^o \left[\exp\left(ikx + \frac{\xi^3}{3}\right)\right] V_1(\xi, q) \tag{1.29}$$

$$H_x = i(ka/2)^{-\frac{1}{3}} H_z^o \left[\exp\left(ikx + \frac{\xi^3}{3}\right)\right] f(\xi) \tag{1.30}$$

The current distribution functions $V_1(\xi,q)$ and $f(\xi)$ are defined as limiting forms of Eq. (1-23) which results from a use of Eq. (1-25)

$$V_1(\xi, q) = \lim_{y_o \to \infty} \left\{ \frac{\sqrt[4]{y_o}}{\sqrt{\xi + \sqrt{y_o}}} \exp\left(-i\frac{2}{3}y_o^{3/2}\right) V(\xi + \sqrt{y_o}, 0, y_o, q) \right\}$$

$$= i2\sqrt{\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{(t_s - q^2) w_1(t_s)}$$

$$= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t) \frac{1}{w_1'(t) - q\, w_1(t)}\, dt \tag{1.31}$$

and

$$f(\xi) = \left\{ \lim_{q \to \infty} - q\, V_1(\xi, q) \right\} = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \frac{\exp(i\xi t)}{w_1(t)}\, dt \tag{1.32}$$

For $x < 0$ these formulae transform into the equations of geometric optics. This property can be readily verified by noting that for ξ tending to large negative values that the special functions have the behavior

$$V_1(\xi, q) \to \frac{2 \exp(-i\xi^3/3)}{1 + i(q/\xi)}$$

$$f(\xi) \to i2\xi \exp(-i\xi^3/3)$$

For the special case $\xi = 0$ (i.e., the horizon) we know that $V_1(0,0) = 1.399$ and $f(0) = 0.766 \exp(-i\pi/3)$. The behavior for the for other values of q is depicted in Fig. 1-4.




Fig. 1-4 Behavior of $V_1(0,q)$

The principle which was enunciated by Fock in 1945, was extended to the case when both the source and the receiver were above the surface by the papers of Pekeris (Ref. 4) in 1947 and Fock (1951). However, let us first review the manner in which the elevated source and receiver were handled in the work of van der Pol and Bremmer (Ref. 1).

The starting point for the discussion will be the exact solution for the Hertz potential for a vertical electric dipole. The problem can be discribed as follows:

Differential Equation

$$\nabla^2 U + k^2 U = -\frac{\delta (r - r_o)\, \delta (\theta)}{2\pi r^2 \sin \theta} \tag{1.33}$$

$$k^2 = \omega^2 \epsilon_o \mu_o \text{ for } r>a \; ; \quad k_1^2 = \omega^2 \epsilon_1 \mu_o + i\omega \mu_o \sigma_1 \text{ for } r<a \tag{1.34}$$

Boundary Condition

$$\frac{\partial}{\partial r} (r\,U) \text{ and } k^2 U \text{ continuous across } r = a \tag{1.35}$$

Periodic Condition

$$U (\theta \pm 2m\pi) = U (\theta) \tag{1.36}$$

Radiation Condition

$$\lim_{r \to \infty} \left\{ r \left[\frac{\partial U}{\partial r} - i k U \right] \right\} = 0 \tag{1.37}$$

The classical methods for the solution of a problem of this type involves the expansion of the field in a series of spherical harmonics. The resulting representation for $U(\theta, r, r_o, \Gamma)$ involves a complex combination of spherical Hankel functions. If we let $r_>$ and $r_<$ represent, respectively, the larger and the smaller of the two radial distances r and r_o, then the exact solutions take the form of the following series:

$$U(\Theta, r, r_o, \Gamma) = \frac{i\,k}{8\pi} \frac{1}{(kr)(kr_o)} \sum_{n=0}^{\infty} (2n+1) \left| \zeta_n^{(1)}(kr_>)\, \zeta_n^{(2)}(kr_<) - \frac{\zeta_n^{(2)'}(ka) + \Gamma \zeta_n^{(2)}(ka)}{\zeta_n^{(1)'}(ka) + \Gamma \zeta_n^{(1)}(ka)} \zeta_n^{(1)}(kr)\, \zeta_n^{(1)}(kr_o) \right| P_n(\cos\Theta) \qquad (1.38)$$

$$\zeta_n^{(1,2)}(x) = \sqrt{\frac{\pi x}{2}}\, H_{n+1/2}^{(1,2)}(x) \qquad \psi_n(x) = \sqrt{\frac{\pi x}{2}}\, J_{n+1/2}(x) \qquad (1.39)$$

$$\Gamma = -\frac{k}{k_1} \frac{\psi_n'(k_1 a)}{\psi_n(k_1 a)} \sim i \frac{k}{k_1} \sqrt{1 - \left(\frac{k}{k_1}\right)^2} \qquad (1.40)$$

The classical procedure involves the transformation of the sum into an integral. From the integral, one extracts an integral which can be evaluated by the method of stationary phase to yield the reflected wave. Before we consider the form taken by the result, we must make some geometrical definitions by introducing Fig. 1-5.




Fig. 1-5 Reflection from a Convex Surface

$$\mathcal{U}(\Theta, r, r_o, \Gamma) \sim \frac{\exp(ikR)}{4\pi R} - \frac{ik}{4\pi} \frac{1}{(kr)(kr_o)} \int_{-\infty+io}^{\infty+io} (n + 1/2) \frac{\zeta_n^{(2)'}(ka) + \Gamma \zeta_n^{(2)}(ka)}{\zeta_n^{(1)'}(ka) + \Gamma \zeta_n^{(1)}(ka)}$$

$$\zeta_n^{(1)}(kr)\, \zeta_n^{(1)}(kr_o) \left[P_n(\cos\Theta) - i\frac{2}{\pi} Q_n(\cos\Theta) \right] dn$$

$$\sim \frac{\exp(ikR)}{4\pi R} + \frac{1}{4\pi} \left\{ \sqrt{\frac{a \sin\alpha}{r r_o \sin\Theta}} \frac{\cos\alpha + i\Gamma}{\cos\alpha - i\Gamma} \right.$$

$$\left. \sqrt{\frac{a \cos\alpha}{2R_o R_1 + a(R_o + R_1)\cos\alpha}} \right\} \exp\left[ik(R_o + R_1) \right] \qquad (1\text{-}41)$$

Van der Pol and Bremmer showed that this formula agrees with the approximation based upon the attenuation function Eq. (1-20) provided the comparison is made in the vicinity of the first and second lobe just above the horizon. However, for points well above the horizon, the attentuation function fails to agree with the stationary phase (geometrical optics) result.

It would be interesting to seek a formula which agrees with the classic result involving $V(x,y,y_o,q)$ for points near the horizon, but which is in better agreement with the stationary phase result for points well above the horizon. Such a formula has been found in the course of the studies made by the present author. It involves the reflected field $R(x,y,y_o,q)$ which can be defined by separating the Green's function $D(x,y,y_o,q)$ into a free space field $D_o(x,y,y_o)$ and a secondary field which for points above the horizon can be interpreted as a reflected field. The decomposition is made as follows:

$$D(x,y,y_o,q) = D_o(x,y,y_o) + R(x,y,y_o,q) \qquad (1\text{-}41)$$

$$D_o(x,y,y_o) = \frac{1}{2\pi}\int_{-\infty}^{\infty} \exp(ixt)\, w_1(t-y)\, v(t-y_o)\, dt$$

$$= \frac{\exp\left(i\frac{\pi}{4}\right)}{2\sqrt{\pi x}} \exp\left[-i\frac{x^3}{12} + i\frac{x}{2}(y+y_o) + i\frac{(y-y_o)^2}{4x}\right] \qquad (1\text{-}42)$$

$$R(x,y,y_o,q) = -\frac{1}{2\pi}\int_{-\infty}^{\infty} \exp(ixt)\,\frac{v'(t) - q\,v(t)}{w_1'(t) - q w_1(t)}\; w_1(t-y)\, w_1(t-y_o)\, dt \qquad (1\text{-}43)$$

The method of stationary phase can be used to show that

$$R(\zeta+\zeta_o,\ \zeta^2+2\mu\zeta,\ \zeta_o^2+2\mu\zeta_o,\ q) \xrightarrow[\mu>>1]{}$$

$$-\frac{\exp(i\pi/4)}{2\sqrt{\pi}}\,\frac{q-i\mu}{q+i\mu}\sqrt{\frac{\mu}{2\zeta\zeta_o+\mu(\zeta+\zeta_o)}}\,\exp(i\omega^*) \qquad (1\text{-}44)$$

where

$$\omega^* = \frac{2}{3}(\zeta^3+\zeta_o^3) + 2\mu(\zeta^2+\zeta_o^2) + \mu^2(\zeta+\zeta_o) \qquad (1\text{-}45)$$

It will often be more convenient to absorb the phase factor in Eq. (1-45) into the definition of the reflection function and express the results in terms of the modified reflection function $S(\mu,\zeta,\zeta_o,q)$

$$S(\mu,\zeta,\zeta_o,q) = \exp(-i\omega^*)\; R(\zeta+\zeta_o,\ \zeta^2+2\mu\zeta,\ \zeta_o^2+2\mu\zeta_o,\ q)$$

$$\xrightarrow[\mu\to\infty]{} -\frac{1}{2}\sqrt{\frac{i}{\pi}}\,\frac{q-i\mu}{q+i\mu}\sqrt{\frac{\mu}{2\zeta\zeta_o+\mu(\zeta+\zeta_o)}} \qquad (1\text{-}46)$$

The new formula for the region of direct visibility then takes the form

$$U(\Theta,r,r_o,\Gamma) = \frac{\exp(ikR)}{4\pi R}$$

$$+\sqrt{\frac{-ik}{8\pi}}\left(\frac{2}{ka}\right)^{1/3}\sqrt{\frac{a\sin\alpha}{rr_o\sin\Theta}}\,\exp\left[ik(D_o+D)\right]\,S(\mu,\zeta,\zeta_o,q) \qquad (1\text{-}47)$$

where

$$\zeta = \left(\frac{ka}{2}\right)^{1/3} \frac{D}{a} \qquad \mu = \left(\frac{ka}{2}\right)^{1/3} \cos\alpha$$

$$\zeta_o = \left(\frac{ka}{2}\right)^{1/3} \frac{D_o}{a} \qquad q = \left(\frac{ka}{2}\right)^{1/3} \Gamma \tag{1-48}$$

For points well above the horizon, $\mu \to \infty$ as $ka \to \infty$ since $\cos\alpha > 0$. Therefore, we observe that with the above definitions of ζ_o, ζ, and μ our new formulation agress with the classical results obtained from a stationary phase evaluation of the exact solution. The new formulation has the advantage, however, of leading to results which are useful up to the horizon. In fact, it can even be used in the region just beyond the horizon.

The asymptotic forms that appear in Eq. (1-44) and Eq. (1-46) are only valid provided the parameter μ is large. From Eq. (1-48) we see that since α tends to 90° as one approaches the horizon that the value of μ may become small (and even vanish) in this region regardless of how large the radius of curvature may be in comparison with the wavelength (i.e., no matter how large $ka = 2\pi a/\lambda$ may be). Therefore, if we continue to require that both ζ and ζ_o be very large, but leave μ unrestricted, we will find that the asymptotic properties of the field will have to be described in terms of a function $V_{11}(\xi,q)$ which we will refer to as the reflection coefficient function. If we use the leading term in the asymptotic expansion described by Eq. (1-25) we find that

$$V_{11}(\xi,q) = \lim_{\substack{y_o \to \infty \\ y \to \infty}} \left\{ \frac{\sqrt[4]{y_o y}}{\sqrt{\xi + \sqrt{y} + \sqrt{y_o}}} \exp\left[-i\frac{2}{3}\left(y^{3/2} + y_o^{3/2}\right)\right] \right.$$

$$\left. V\left(\xi + \sqrt{y} + \sqrt{y_o},\, y,\, y_o,\, q\right) \right\} \tag{1-49}$$

We can represent $V_{11}(\xi,q)$ by means of a residue series (involving the roots t_s defined by Eq. (1-8)) or by means of a Fourier integral.

$$V_{11}(\xi,q) = \exp\left(i\frac{3\pi}{4}\right) 2\sqrt{\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{\left(t_s - q^2\right)\left[w_1(t_s)\right]^2}$$

$$= -\frac{\exp(i\pi/4)}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t)\, \frac{v'(t) - q\, v(t)}{w_1'(t) - q\, w_1(t)}\, dt \qquad (1\text{-}50)$$

However, $V_{11}(\xi,q)$ is singular when $\xi \to 0$ and hence it is convenient to define a closely related funotion $V_2(\xi,q)$ which is an entire function of ξ.

$$V_2(\xi,q) = V_{11}(\xi,q) - (2\sqrt{\pi}\,\xi)^{-1} \exp(i\pi/4) \qquad (1\text{-}51)$$

If we use the properties which we have just cited, we can show that the function $S(\mu,\zeta,\zeta_o,q)$ defined by Eq. (1-46) has the property

$$S(\mu,\zeta,\zeta_o,q) \xrightarrow[\substack{\zeta_o > 1 \\ \mu > 0}]{\zeta > 1} \sqrt{\frac{1}{2\pi}}\, \frac{\exp\left(-i\frac{2}{3}\mu^3\right)}{\sqrt{2\zeta\zeta_o + \mu(\zeta+\zeta_o)}}\, V_{11}(-2\mu, q) \qquad (1\text{-}52)$$

If we reconsider the function $R(x,y,y_o,q)$, we see that these properties lead to

$$R\left(\zeta+\zeta_o,\ \zeta^2 + 2\mu\zeta,\ \zeta_o^2 + 2\mu\zeta_o,\ q\right) \xrightarrow[\zeta,\ \zeta_o \gg 1]{}$$

$$\frac{\exp\left(i\frac{\pi}{4}\right)}{\sqrt{2\pi}\sqrt{2\zeta\zeta_o + \mu(\zeta+\zeta_o)}}\, V_{11}(-2\mu, q) \exp\left(i\omega^* - i\frac{2}{3}\mu^3\right) \qquad (1\text{-}53)$$

For points well above the horizon (i.e., for large positive values of μ, the reflection coefficient function $V_{11}(-2\mu,q)$ has the asymptotic behavior

$$V_{11}(-2\mu, q) \xrightarrow[\mu \gg 1]{} -\frac{q - i\mu}{q + i\mu}\sqrt{\frac{\mu}{2}}\exp\left(i\frac{2}{3}\mu^3\right) \tag{1-54}$$

$$V_{11}(-2\mu, q) \xrightarrow[q \to \infty]{\mu \gg 1} -\sqrt{\frac{\mu}{2}}\exp\left[i\left(\frac{2}{3}\mu^3 - \frac{2\mu}{q}\right)\right] \tag{1-55}$$

When the asymptotic estimates for $V_{11}(-2\mu, q)$ are used in Eq. (1-52) and Eq. (1-47), we are led back to the result which was given in Eq. (1-41). However, Eq. (1-41) was valid only for points well above the horizon and we have now laid the groundwork for an asymptotic estimate which agrees with the predictions of Eq. (1-41) when the parameter μ is very large and positive but which has the advantage of holding for points in the vicinity of the horizon. Before we proceed to discuss the behavior on the horizon itself, let us first obtain a result which is valid in the vicinity of the horizon and which will contain the familiar Fresnel integral which we should expect to encounter in this region on the basis of the considerations of physical optics. For our purposes it is convenient to introduce the modified Fresnel integral $K(\tau)$ which was employed by Fock (Ref. 5).

$$K(\tau) = \exp\left(-i\tau^2 - i\frac{\pi}{4}\right)\frac{1}{\sqrt{\pi}}\int_{\tau}^{\infty}\exp(is^2)\,ds$$

$$\xrightarrow[\tau \to \infty]{} \frac{\exp\left(i\frac{\pi}{4}\right)}{2\sqrt{\pi}\,\tau} \tag{1-56}$$

From the property described by Eq. (1-51), we observe that

$$\mu K(\mu\xi) + V_2(\xi, q) \xrightarrow[\xi > 0]{\mu \to \infty} V_{11}(\xi, q) \tag{1-57}$$

We will define the "horizon" to be the locus of points for which

$x = (\sqrt{y} + \sqrt{y_o})$ and we will introduce the parameter

$$\xi = x - \sqrt{y} - \sqrt{y_o} \tag{1-58}$$

which will be positive below the horizon, zero on the horizon, and negative above the horizon. We also define the parameters μ and τ

$$\mu = \sqrt{\frac{\sqrt{y}\sqrt{y_o}}{\sqrt{y} + \sqrt{y_o}}} \quad , \quad \tau = \mu\xi \tag{1-59}$$

Let $F(x,y,y_o)$ be the Fresnel diffraction field

$$F(x,y,y_o) = \sqrt[4]{\frac{x^2}{y y_o}} \exp\left[i \frac{2}{3}\left(y^{3/2} + y_o^{3/2}\right)\right] \mu K(\tau), \quad \xi > 0 \tag{1-60a}$$

$$= \sqrt[4]{\frac{x^2}{y y_o}} \exp\left[i \frac{2}{3}\left(y^{3/2} + y_o^{3/2}\right)\right]\left[-\mu K(-\tau)\right]$$

$$+ \exp\left[i\left(-\frac{x^3}{12} + \frac{x}{2}(y + y_o) + \frac{(y - y_o)^2}{4x}\right)\right] \quad \xi < 0 \tag{1-60b}$$

We can then show that

$$V(x,y_o,y,q) \xrightarrow[\substack{y \to \infty \\ y_o \to \infty}]{} F(x,y,y_o) + \sqrt[4]{\frac{x^2}{y y_o}} \exp\left[i \frac{2}{3}\left(y^{3/2} + y_o^{3/2}\right)\right] V_2(\xi,q) \tag{1-61}$$

If we let μ tend to infinity we obtain from (1-61) the result

$$V(x,y,y_o,q) - \frac{\sqrt{x}}{\sqrt{\sqrt{y}\sqrt{y_o}}} V_{11}(\xi,q) \xrightarrow[\substack{y_o \to \infty \\ y \to \infty \\ \mu \to \infty}]{} G(x,y,y_o) \tag{1-61}$$

where $G(x,y,y_o) = 0$ when $\xi > 0$ and

$$G(x,y,y_o) = \exp[i(-\frac{x^3}{12} + \frac{x}{2}(y + y_o) + \frac{(y - y_o)^2}{4x})]$$

when $\xi < 0$. The result in Eq. (1-61) reveals that the function $V_{11}(\xi,q)$ describes the reflected field in the lighted region (i.e., $\xi < 0$) and the same function describes the diffracted field in the shadow region (i.e., $\xi > 0$). In the vicinity of the horizon we need to employ the function $V_2(\xi,q)$. On the horizon (i.e., $\xi = 0$) we find that


$$V(\sqrt{y} + \sqrt{y_o}, y, y_o, q) \xrightarrow[\substack{y \to \infty \\ y_o \to \infty}]{} [\frac{1}{2} + \frac{1}{\mu} V_2(0,q)]\exp(i\Lambda) \qquad (1\text{-}62)$$

where

$$\Lambda = \frac{2}{3}(y^{3/2} + y_o^{3/2})$$

In Fig. 1-6 we present an illustration which shows the behavior*

*Fig. 1-6 was previously presented by the author in Ref.6. Data has been generated and a much more detailed plot has been begun. However, the author became "side-tracked" on this task when it was found that as $q \to Q$, where Q is real and positive, that the data revealed an irregular behavior. The author became intrigued with this phenomena and investigated the behavior of the analogue of $V_2(0,q)$ in the problem of diffraction by a circular cylinder. Computations made with the exact solution also showed similar results to those found with the data for $V_2(0,q)$. The author strongly suspects that the erratic behavior is caused by errors introduced into the numerical work because of the fact that the expression $w_1'(t) - qw_1(t)$ which appears in the denominator of the integral defining $V_2(\xi,q)$ vanishes for a point close to the contour of integration. The author hopes to be able to devote time in the immediate future to the task of completing this very detailed contour plot of $V_2(0,q)$.


A=0.2
A=0.1
A=0.5
A=1.5
A=0.2
φ=0
φ=π
φ=0
φ=0
φ=0

of $V_2(0,q)$ as a function of q. The locus for which $\varphi = \arg(q) = 45^\circ$ was previously presented in a paper by Wait and Conda (Ref. 7). The reader will observe that the form taken by q in Fig. 1-6 has been taken to be

$$q = 2^{\frac{1}{3}} A \exp(i\varphi) \tag{1-63}$$

This form was chosen in order that the curves might supplement the curve presented by Wait and Conda who also used A in the manner displayed in Eq. (1-63).

If we apply the result in Eq. (1.61) to the function $S(\mu,\zeta,\zeta_o,q)$ defined in Eq. (1-46) we find that

$$S(\mu,\zeta,\zeta_o,q) \xrightarrow[\substack{\zeta_o \to \infty \\ \mu \to 0}]{\zeta \to \infty} \frac{1}{2}\sqrt{\frac{i}{\pi}}\ \frac{1}{\sqrt{\zeta\zeta_o}} \exp\left(-i\,\frac{2}{3}\,\mu^3\right)\left[-\eta\, K(-\tau) + V_2(-2\mu, q)\right] \tag{1-64}$$

where

$$\tau = -2\mu\eta \qquad , \qquad \eta = \sqrt{\frac{\zeta\zeta_o}{\zeta+\zeta_o}} \tag{1-65}$$

If Eq. (1-64) is used in Eq. (1-47) we obtain a useful extension of the classical reflection formulae. On the horizon, $\mu = 0$, and the theory reveals that we have the approximation

$$U(\theta,r,r_o,\Gamma) \approx \frac{\exp(ikR)}{4\pi R} + M(\theta,r,r_o)\ V_2(0,q) \tag{1-66a}$$

where

$$M(\theta,r,r_o) = \frac{1}{2\pi}\left(\frac{ka}{2}\right)^{1/3} \frac{1}{\sqrt{2kDD_o}} \sqrt{\frac{a}{rr_o \sin\theta}}\ \exp\left[ik(D+D_o)\right] \tag{1-66b}$$

If we let r and r_o tend to infinity, we obtain the result

$$U(\theta,r,r_o,\Gamma) \xrightarrow[\substack{r_o \to \infty \\ r \to \infty}]{} \left\{\frac{1}{2} + 2\left(\frac{ka}{2}\right)^{1/3} \sqrt{\frac{D+D_o}{2kDD_o}}\ V_2(0, q)\right\} \frac{\exp(ikR)}{4\pi R} \tag{1-67}$$

The factor "1/2" which appears as the first term inside the braces of Eq. (1-67) is the familiar Fresnel field factor for the field on the shadow boundary. The effects of the curvature of the surface which is tangent to the line joining the source and the receiver is contained in the term which contains the function $V_2(0,q)$.

The function $U(\theta,r,r_o,\Gamma)$ arose in the problem of an axially symmetric source above a spherical surface. We can show that very similar results can be obtained when the diffracting surface is a circular cylinder. For this case, we can again use Fiq. 1-5 and define μ, ζ,ζ_o, and q as in the case of the sphere. Let us denote the Green's function for the cylinder problem by $C(\theta,r,r_o,\Gamma)$. The optical results for this function are of the form

$$C(\theta, r, r_o, \Gamma) \xrightarrow[\mu\to\infty]{} \frac{\exp[i(kR+\pi/4)]}{\sqrt{8\pi kR}} + \left\{\sqrt{\frac{1}{8\pi k}}\ \frac{\cos\alpha + i\Gamma}{\cos\alpha - i\Gamma}\right.$$
$$\left.\sqrt{\frac{a\cos\alpha}{2DD_o + a(D+D_o)\cos\alpha}}\ \exp\left[ik(D+D_o)\right]\right\} \quad (1\text{-}68)$$

For points near the horizon, we can use use the same arguments which led to Eq. (1-67) to show that

$$C(\theta, r, r_o, \Gamma) \xrightarrow[\substack{\mu\to 0\\ r\to\infty\\ r_o\to\infty}]{} \left\{\frac{1}{2} + 2\left(\frac{ka}{2}\right)^{1/3}\sqrt{\frac{D+D_o}{2kDD_o}}\ V_2(0,q)\right\}\frac{\exp[i(kR+\pi/4)]}{\sqrt{8\pi kR}} \quad (1\text{-}69)$$

The case $r_o \to \infty$ corresponds to the problem of reflection of a plane wave from a circular cylinder. This problem follows from the results given in Eqs. (1-68) and (1-69) is we use the approximation

$$\frac{\exp(ikR)}{\sqrt{8\pi kR}} \sim \frac{\exp(ikr_o)}{\sqrt{8\pi kr_o}}\exp(-ikr\cos\theta) \quad (1\text{-}70)$$

and factor out the term $\exp(ikr_o)/(\sqrt{8\pi kr_o})$. We discuss the case of a cylinder illuminated by a plane wave in more detail in

Section 3.

Let us turn now to consider the two parameter functions which are the limiting forms of $V(x,y,y_o,q)$. These have been discussed in considerable detail by Logan (Ref. 8).

When $y = y_o = 0$ we obtain the function $V_o(\xi,q)$ which possesses the representations

$$V_o(\xi, q) = \frac{1}{2} V(\xi, 0, 0, q)$$

$$= \sqrt{\pi\xi}\ \exp(i\pi/4) \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{t_s - q^2}$$

$$= \frac{\exp(-i\pi/4)}{2}\sqrt{\frac{\xi}{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t)\ \frac{w_1(t)}{w_1'(t) - q w_1(t)}\ dt \qquad (1\text{-}71)$$

For $\xi > 0$ the residue series in Eq. (1-71) converges fast enough to be a practical means of computing $V_o(\xi,q)$. For $\xi \to 0$ it becomes completely impractical to sum the residue series and it becomes more attractive to follow Pekeris (Ref. 4) and express the integral in Eq. (1-71) in the form

$$V_o(\xi,q) = \frac{1}{2}\sqrt{\frac{-i\xi}{\pi}} \int_{\infty \exp(i\xi\pi)}^{\infty} \exp(i\xi t)\ \frac{w_1(t)}{w_1'(t) - q w_1(t)}\ dt \qquad (1\text{-}72)$$

Pekeris then used an asymptotic representation for the integrand which, in our notation, is of the form

$$\frac{w_1(t)}{w_1'(t) - q w_1(t)} \approx \frac{1}{\sqrt{t} - q} \qquad (1\text{-}73)$$

to suggest that the integrand should be broken into two components as in the following equation:

$$\frac{w_1(t)}{w_1'(t) - qw_1(t)} = \frac{1}{\sqrt{t} - q} + \frac{\sqrt{t}\, w_1(t) - w_1'(t)}{(\sqrt{t} - q)[w_1'(t) - qw_1(t)]} \tag{1-74}$$

We can then use the property of the Airy function

$$w_1[t\exp(i\tfrac{2}{3}\pi)] = \exp(i\tfrac{1}{3}\pi)w_2(t) \tag{1-75}$$

to express $V_0(\xi,q)$ in the form

$$\begin{aligned} V_0(\xi,q) = {} & \frac{1}{2}\sqrt{\frac{-i\xi}{\pi}} \int_{\infty\exp(i\frac{2}{3}\pi)}^{\infty} \frac{\exp(i\xi t)}{\sqrt{t} - q}\, dt \\ & + \frac{1}{2}\sqrt{\frac{-i\xi}{\pi}} \int_0^{\infty} \exp(i\xi t)\, F_1(t,q)dt \\ & + \frac{1}{2}\sqrt{\frac{-i\xi}{\pi}} \int_0^{\infty} \exp\left(-\frac{\sqrt{3} + i}{2}\,\xi t\right) F_2(t,q^*)dt \end{aligned} \tag{1-76}$$

where

$$F_1(t,q) = \frac{\sqrt{t}\, w_1(t) - w_1'(t)}{(\sqrt{t} - q)[w_1'(t) - qw_1(t)]} \tag{1-77a}$$

$$F_2(t,q) = \exp(i\tfrac{1}{3}\pi)\frac{\sqrt{t}\, w_2(t) + w_2'(t)}{(\sqrt{t} + q^*)[w_2'(t) - q^* w_2(t)} \tag{1-77b}$$

Pekeris observed that the first integral in Eq. (1-76) leads to the Weyl-van der Pol formula, i.e.,

$$\frac{1}{2}\sqrt{\frac{-i\xi}{\pi}}\int_{\infty\exp(i\frac{\pi}{2})}^{\infty}\frac{\exp(i\xi t)}{\sqrt{t}-q}\,dt = 1 - 2\sigma\exp(-\sigma^2)\int_{i\infty}^{\sigma}\exp(\alpha^2)\,d\alpha \qquad (1\text{-}78)$$

where σ is the Sommerfeld "numerical distance"

$$\sigma = q\sqrt{-i\xi} \qquad (1\text{-}79)$$

Pekeris evaluated the integrals which contain the factors $F_1(t,q)$ and $F_2(t,q)$ by means of numerical integration. The author knows of no later writers who have employed these techniques to evaluate Eq. (1-76). The technique of numerical integration is well known for the case of the functions $V_1(\xi,q)$ and $V_2(\xi,q)$, but the decomposition discovered by Pekeris has escaped the attention of those writers whose papers the author has consulted. The integral in Eq. (1-78) leads to the error function of complex argument.

The function $V_o(\xi,q)$ can be shown to be a solution of the integral equation

$$V_o(\xi,q) = \exp\left(-i\frac{1}{12}\xi^3\right) - \frac{\exp\left(-i\frac{\pi}{4}\right)}{2}\sqrt{\frac{\xi}{\pi}}\int_0^{\xi} V_o(x,q)\exp\left[-i\frac{1}{12}(\xi-x)^3\right]$$

$$\left|(\xi-x)-2iq\right|\frac{dx}{\sqrt{x(\xi-x)}} \qquad (1\text{-}80)$$

In Ref. 8 the author has discussed representations for the two limiting cases, namely

$$v(\xi) = V_o(\xi,0) = \frac{\exp(-i\,\pi/4)}{2}\sqrt{\frac{\xi}{\pi}}\int_{-\infty}^{\infty}\exp(i\xi t)\,\frac{w_1(t)}{w_1'(t)}\,dt \qquad (1\text{-}81)$$

$$u(x) = \lim_{q\to\infty}\left|2i\xi q^2 V_o(\xi,q)\right| = \frac{\exp(-i\,3\pi/4)}{\sqrt{\pi}}\,\xi^{3/2}\int_{-\infty}^{\infty}\exp(i\xi t)\,\frac{w_1'(t)}{w_1(t)}\,dt \qquad (1\text{-}82)$$

In Fig. 1.7 we illustrate the behavior of the real and the




Fig. 1.7 Attenuation Function $u(\xi)$ (Real and Imaginary Parts)

imaginary parts of $u(\xi)$. In Fig. 1.8 we illustrate the behavior of the modulus of $u(\xi)$.

In Fig. 1.9 we show the behavior of the real and imaginary parts of $v(\xi)$. In Fig. 1.10 we illustrate the behavior of the modulus of $v(\xi)$.

For small values of ξ we can compute $u(\xi)$ and $v(\xi)$ from series which progress in successive powers of $\xi^{3/2}$. The theory behind these representations has been developed in Ref. 3. The series are of the form

(Text continues on p. 1-28)




Fig. 1.8 Attenuation Function $u(\xi)$ (Modulus)




Fig. 1-9 Attenuation Function v(ξ) (Real and Imaginary Parts)

LOCKHEED MISSILES & SPACE COMPANY

$$u(\xi) = \left\{ 1 + \frac{1}{4} \frac{\Gamma\left(-\frac{1}{2}\right)}{\Gamma(2)} \exp\left(-i\frac{\pi}{4}\right)\xi^{3/2} - \frac{5}{32} \frac{\Gamma\left(-\frac{1}{2}\right)}{\Gamma\left(\frac{5}{2}\right)} \exp\left(-i\frac{\pi}{2}\right)\xi^{3} + \cdots\cdots \right\}$$

$$= -2\sqrt{-i\pi\xi^3} \sum_{n=0}^{\infty} A_n \frac{\left[e^{-i\pi/6}\xi\right]^{\frac{3n-3}{2}}}{\Gamma\left(\frac{3n-1}{2}\right)} \qquad (1\text{-}83)$$

$$v(\xi) = \left\{ 1 - \frac{1}{4} \frac{\Gamma\left(\frac{1}{2}\right)}{\Gamma(2)} \exp\left(-i\pi/4\right)\xi^{3/2} + \frac{7}{32} \frac{\Gamma\left(\frac{1}{2}\right)}{\Gamma\left(\frac{7}{2}\right)} \exp\left(-i\frac{\pi}{2}\right)\xi^{3} + \cdots \right\}$$

$$= -\exp\left(-i\frac{\pi}{3}\right) \sum_{n=0}^{\infty} B_n \frac{\left[e^{-i\pi/6}\xi\right]^{\frac{3n-1}{2}}}{\Gamma\left(\frac{3n+1}{2}\right)} \qquad (1\text{-}84)$$




Fig. 1.10 Attenuation Function v(ξ) (Modulus)

In Eq. (1-83) the coefficients A_n are the coefficients of $\alpha^{(-3n+\frac{1}{2})}$ in the asymptotic expansion of the logarithmic derivative of the Airy function

$$\frac{Ai'(\alpha)}{Ai(\alpha)} = \sum_{n=0}^{\infty} A_n \alpha^{(3n-1/2)} = -\sqrt{\alpha} - \frac{1}{4}\frac{1}{\alpha} + \frac{5}{32}\frac{1}{\alpha^{5/2}} - \frac{15}{64}\frac{1}{\alpha^4} + \frac{1105}{2048}\frac{1}{\alpha^{11/2}}$$
$$- \frac{1695}{1024}\frac{1}{\alpha^7} + \frac{414125}{65536}\frac{1}{\alpha^{17/2}} - \frac{59025}{2048}\frac{1}{\alpha^{10}}$$
$$+ \frac{12820\,31525}{8388608}\frac{1}{\alpha^{21/2}} - \frac{2421\,83775}{262144}\frac{1}{\alpha^{13}} + \cdots \quad (1\text{-}85)$$

In Eq. (1-84) the coefficients B_n are the coefficients of $\alpha^{(-3n+\frac{1}{2})}$ in the asymptotic expansion of the logarithmic derivative of the derivative of the Airy function

$$\frac{d}{d\alpha}\log Ai'(\alpha) = \frac{Ai''(\alpha)}{Ai'(\alpha)} = \alpha\frac{Ai(\alpha)}{Ai'(\alpha)} = \sum_{n=0}^{\infty} B_n \frac{1}{\alpha^{\frac{3n-1}{2}}}$$

$$= -\sqrt{\alpha} + \frac{1}{4}\frac{1}{\alpha} - \frac{7}{32}\frac{1}{\alpha^{5/2}} + \frac{21}{64}\frac{1}{\alpha^4} - \frac{1463}{2048}\frac{1}{\alpha^{11/2}}$$
$$+ \frac{2121}{1024}\frac{1}{\alpha^7} - \frac{495271}{65536}\frac{1}{\alpha^{17/2}} + \frac{136479}{4096}\frac{1}{\alpha^{10}}$$
$$- \frac{14457\,13003}{8388608}\frac{1}{\alpha^{23/2}} + \frac{268122561}{262144}\frac{1}{\alpha^{13}} - \cdots \quad (1\text{-}86)$$

In Table 1-1 we present numerical values for the coefficients A_n and B_n.

In Tables 1-2 and 1-3 we present numerical values for the functions $u(\xi)$ and $v(\xi)$.

The author is preparing a revision of Ref. 8 in which computer programs for these functions, as well as for $V_1(\xi,q)$ and $V_2(\xi,q)$, will be given.

Table 1-1

VALUE OF A_n AND B_n OCCURING IN ASYMPTOTIC EXPANSIONS OF THE LOGARITHMIC DERIVATIVES OF Ai(α) AND Ai'(α)

n	A_n	B_n
0	-0.10000000000000000000D 01,	-0.10000000000000000000D 01,
1	-0.25000000000000000000D+00,	+0.25000000000000000000D+00,
2	+0.15625000000000000000D+00,	-0.21875000000000000000D+00,
3	-0.23437500000000000000D+00,	+0.32812500000000000000D+00,
4	+0.53955078125000000000D+00,	-0.71435546875000000000D+00,
5	-0.16552734375000000000D 01,	+0.20712890625000000000D 01,
6	+0.63190460205078125000D 01,	-0.75572357177734375000D 01,
7	-0.28820800781250000000D 02,	+0.33320068359375000000D 02,
8	+0.15283006727695465087D 03,	-0.17234241998195648193D 03,
9	-0.92385778427124023437D 03,	+0.10228064002990722656D 04,
10	+0.62714540283195674419D 04,	-0.68477664783634245395D 04,
11	-0.47242087572813034057D 05,	+0.51038504238367080688D 05,
12	+0.39109379739530413644D 06,	-0.41901341755007504252D 06,
13	-0.35296291884854435920D 07,	+0.37563697358470559120D 07,
14	+0.34492359422529131052D 08,	-0.36507330917304253858D 08,
15	-0.36285905416732421144D 09,	+0.38230366458564009144D 09,
16	+0.40887481953623399013D 10,	-0.42911919679893578455D 10,
17	-0.49132931466161909011D 11,	+0.51394354289130102764D 11,
18	+0.62719845918866603263D 12,	-0.65417344539770204956D 12,
19	-0.84761139845881194130D 13,	+0.88182840831890162595D 13,
20	+0.12089738905565412566D 15,	-0.12549617754663210762D 15,
21	-0.18149699900371916926D 16,	+0.18802478828410565850D 16,
22	+0.28607122279051596739D 17,	-0.29582925127951160480D 17,
23	-0.47233560971778173036D 18,	+0.48765744586404374051D 18,
24	+0.81527705736479572024D 19,	-0.84048864895675398507D 19,
25	-0.14683164515496690264D 21,	+0.15116986400379079975D 21,
26	+0.27544975719777650174D 22,	-0.28324098625029985502D 22,
27	-0.53737875698420739846D 23,	+0.55195734973972829322D 23,
28	+0.10886622365991780594D 25,	-0.11170370989316488415D 25,
29	-0.22871046142438023036D 26,	+0.23444644959247598676D 26,
30	+0.49762975408878700819D 27,	-0.50965613963939228718D 27,
31	-0.11200533064150971566D 29,	+0.11461721236159604725D 29,
32	+0.26049620884159831713D 30,	-0.26636489757168919513D 30,
33	-0.62537903246900463830D 31,	+0.63900634447083650325D 31,
34	+0.15482495363475704390D 33,	-0.15809161838403698000D 33,
35	-0.39490815816136472322D 34,	+0.40298415692702386368D 34,
36	+0.10368921179090231495D 36,	-0.10574644297528990525D 36,
37	-0.28002659450985263641D 37,	+0.28542133273123163778D 37,
38	+0.77724602947290274652D 38,	-0.79179892468084064963D 38,

Table 1-2

THE ATTENUATION FUNCTION $u(\xi)$

ξ	Real	Imag.	Modulus	Argument
0	1.0000	0	1.0000	0
0.1	0.9802	0.0192	0.9804	1.12
0.2	0.9440	0.0528	0.9455	3.20
0.3	0.8974	0.0922	0.9021	5.86
0.4	0.8429	0.1334	0.8534	9.00
0.5	0 7822	0.1738	0.8013	12.51
0.6	0.7156	0.2123	0.7465	16.52
0.7	0.6487	0.2434	0.6929	20.57
0.8	0.5792	0.2700	0.6390	24.97
0.9	0.5090	0.2902	0.5859	29.68
1.0	0.4403	0.3035	0.5348	34.58
1.1	0.3740	0.3102	0.4859	39.67
1.2	0.3112	0.3105	0.4396	44.94
1.3	0.2526	0.3050	0.3960	50.36
1.4	0.1991	0.2944	0.3554	55.93
1.5	0.1510	0.2796	0.3178	61.63
1.6	0.1086	0.2615	0.2831	67.45
1.7	0.0719	0.2409	0.2514	73.38
1.8	0.0409	0.2187	0.2225	79.40
1.9	0.0154	0.1957	0.1963	85.50
2.0	-0.0051	0.1726	0.1727	91.68
2.1	-0.0209	0.1501	0.1515	97.93
2.2	-0.0326	0.1285	0.1326	104.24
2.3	-0.0407	0.1083	0.1157	110.61
2.4	-0.0458	0.0898	0.1008	117.02
2.5	-0.0483	0.0730	0.0875	123.48
2.6	-0.0487	0.0581	0.0759	129.98
2.7	-0.0476	0.0452	0.0656	136.50
2.8	-0.0453	0.0341	0.0567	143.06
2.9	-0.0421	0.0247	0.0488	149.63
3.0	-0.0384	0.0169	0.0420	156.23
3.1	-0.0345	0.0106	0.0361	162.85
3.2	-0.0304	0.0056	0.0309	169.49
3.3	-0.0264	0.0018	0.0265	176.13
3.4	-0.0226	-0.0011	0.0226	182.79
3.5	-0.0190	-0.0032	0.0193	189.46
3.6	-0.0158	-0.0046	0.0165	196.13
3.7	-0.0129	-0.0054	0.0140	202.81
3.8	-0.0104	-0.0059	0.0119	209.50
3.9	-0.0082	-0.0060	0.0101	216.19
4.0	-0.0063	-0.0059	0.0086	222.88

Table 1-3

THE ATTENUATION FUNCTION $\tau(\xi)$

ξ	Real	Imag.	Modulus	Argument
0.1	0.9901	0.0098	0.9901	0.57
0.2	0.9720	0.0271	0.9724	1.59
0.3	0.9486	0.0484	0.9498	2.92
0.4	0.9210	0.0721	0.9238	4.47
0.5	0.8899	0.0969	0.8952	6.22
0.6	0.8559	0.1221	0.8646	8.12
0.7	0.8195	0.1469	0.8325	10.16
0.8	0.7810	0.1707	0.7995	12.33
0.9	0.7410	0.1929	0.7657	14.59
1.0	0.6999	0.2134	0.7317	16.96
1.1	0.6581	0.2317	0.6977	19.40
1.2	0.6159	0.2477	0.6638	21.91
1.3	0.5737	0.2612	0.6304	24.48
1.4	0.5317	0.2723	0.5976	27.11
1.5	0.4908	0.2809	0.5655	29.79
1.6	0.4506	0.2870	0.5342	32.50
1.7	0.4115	0.2908	0.5039	35.25
1.8	0.3739	0.2924	0.4747	38.03
1.9	0.3379	0.2920	0.4466	40.83
2.0	0.3036	0.2897	0.4196	43.66
2.1	0.2710	0.2857	0.3938	46.51
2.2	0.2404	0.2801	0.3692	49.37
2.3	0.2117	0.2733	0.3457	52.24
2.4	0.1850	0.2653	0.3265	55.12
2.5	0.1602	0.2564	0.3024	58.01
2.6	0.1373	0.2468	0.2824	60.91
2.7	0.1163	0.2365	0.2636	63.81
2.8	0.0972	0.2258	0.2458	66.72
2.9	0.0747	0.2148	0.2291	69.63
3.0	0.0640	0.2035	0.2134	72.55
3.1	0.0499	0.1923	0.1986	75.46
3.2	0.0372	0.1810	0.1848	78.38
3.3	0.0260	0.1698	0.1718	81.30
3.4	0.0161	0.1589	0.1597	84.22
3.5	0.0074	0.1482	0.1484	87.14
3.6	-0.0002	0.1378	0.1378	90.06
3.7	-0.0067	0.1277	0.1279	92.98
3.8	-0.0122	0.1180	0.1187	95.90
3.9	-0.0169	0.1088	0.1101	98.82
4.0	-0.0208	0.0999	0.1021	101.74

Table 1-3 (Cont'd)

ξ	Real	Imag.	Modulus	Argument
4.1	-0.0240	0.0915	0.0946	104.66
4.2	-0.0265	0.0836	0.0877	107.58
4.3	-0.0285	0.0761	0.0812	110.50
4.4	-0.0299	0.0690	0.0752	113.42
4.5	-0.0309	0.0624	0.0697	116.34
4.6	-0.0315	0.0562	0.0645	119.25
4.7	-0.0318	0.0505	0.0597	122.17
4.8	-0.0318	0.0452	0.0552	125.09
4.9	-0.0315	0.0402	0.0511	128.01
5.0	-0.0310	0.0357	0.0472	130.93
5.1	-0.0303	0.0315	0.0437	133.85
5.2	-0.0294	0.0276	0.0404	136.77
5.3	-0.0285	0.0241	0.0373	139.69
5.4	-0.0274	0.0209	0.0345	142.61
5.5	-0.0263	0.0180	0.0319	145.52
5.6	-0.0251	0.0154	0.0295	148.44
5.7	-0.0239	0.0130	0.0272	151.36
5.8	-0.0226	0.0109	0.0251	154.28
5.9	-0.0214	0.0090	0.0232	157.20
6.0	-0.0202	0.0073	0.0214	160.12
6.1	-0.0189	0.0058	0.0198	163.04
6.2	-0.0177	0.0044	0.0183	165.96
6.3	-0.0165	0.0033	0.0169	168.87
6.4	-0.0154	0.0022	0.0156	171.79
6.5	-0.0143	0.0013	0.0144	174.71
6.6	-0.0132	0.0005	0.0132	177.63
6.7	-0.0122	-0.0001	0.0122	180.55
6.8	-0.0113	-0.0007	0.0113	183.47
6.9	-0.0103	-0.0012	0.0104	186.39
7.0	-0.0095	-0.0015	0.0096	189.30
7.1	-0.0086	-0.0019	0.0088	192.22
7.2	-0.0079	-0.0021	0.0082	195.14
7.3	-0.0072	-0.0023	0.0075	198.06
7.4	-0.0065	-0.0025	0.0069	200.98
7.5	-0.0058	-0.0026	0.0064	203.90
7.6	-0.0053	-0.0027	0.0059	206.82
7.7	-0.0047	-0.0027	0.0054	209.73
7.8	-0.0042	-0.0027	0.0050	212.65
7.9	-0.0038	-0.0027	0.0046	215.57
8.0	-0.0033	-0.0026	0.0043	218.49

REFERENCES FOR SECTION 1

1. B. van der Pol and H. Bremmer, "Diffraction of Electromagnetic Waves from an Electrical Point Source Round a Finitely Conducting Sphere with Applications of Radio Telegraphy and the Theory of the Rainbow," Part I: Phil. Mag., Vol. 24, 1937, pp. 141-176; Part II: Phil. Mag., Vol. 24, 1937, pp. 825-864; Part III, Phil. Mag., Vol. 25, 1938, pp. 817-837.

2. C. R. Burrows and M. C. Gray, "The Effect of the Earth's Curvature on Ground-Wave Propagation," Proc. IRE, Vol. 29, 1941, pp. 16-24

3. V. A. Fock, "The Distribution of Currents Induced by a Plane Wave on the Surface of a Conductor," J. Phys. USSR, Vol. 10, 1946, pp. 130-136. (V. A. Fock, Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965, pp. 10-22.)

4. C. L. Pekeris, "The Field of a Microwave Antenna in the Vicinity of the Horizon," J. Appl. Phys., Vol. 18, 1947, pp. 667-680, 1025-1027

5. V. A. Fock, "Fresnel Diffraction From Convex Bodies," Prog. Phys. Sci. (Uspekhi), Vol. 43, pp. 587-599, 1951. (V. A. Fock, Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965, pp. 134-146.)

6. N. A. Logan and K. S. Yee, "A Mathematical Model for Diffraction by Convex Surfaces," Electromagnetic Waves, R.E. Langer, ed., Madison, University of Wisconsin Press, 1962, pp. 139-180

7. J. R. Wait and A. M. Conda, "Diffraction of Electromagnetic Waves by Smooth Obstacles for Grazing Angles," J. Res. NBS, Vol. 63D, 1959, pp. 181-197

8. Lockheed Missiles and Space Co., General Research in Diffraction Theory, Vol. 1, by N. A. Logan, LMSC Report No. 288087, Sunnyvale, Calif., Dec., 1959

UNITED STATES AIR FORCE
AFGL Research Library

Section 2

THE ROOTS t_s OF $w_1'(t_s) - qw_1(t_s) = 0$

2.1 Notation for the Root Defining Equation

In electromagnetic diffraction problems which involve the propagation of waves around a convex surface, one often has to solve for the roots $\underline{t}$ of the equation

$$w_1'(t) - qw_1(t) = 0 \tag{2-1}$$

where $w_1'(t) = dw_1(t)/dt$ and q is a given "constant" which can generally be related to the physical quantities in such a manner that we can speak of q as being related to the "surface impedance." The function $w_1(t)$ is the Airy function introduced by Fock (Ref. 1) and used extensively by him, by Wait* (Ref. 2), by Logan (ref. 3), by Logan and Yee (Ref. 4), and by others. The function $w_1(t)$ satisfies the Airy differential equation

$$\frac{d^2y}{dt^2} - ty = 0 \tag{2-2}$$

and can be defined by the integral representation

$$w_1(t) = \frac{1}{\pi}\int_{-\infty}^{\infty e^{i\pi/3}} \exp\left(\frac{1}{3}z^3 - zt\right)dz \tag{2-3}$$

For real values of $\underline{t}$ it is convenient to resolve $w_1(t)$ into its real and imaginary parts by defining

$$w_1(t) = u(t) + iv(t) \tag{2-4}$$

where, for real values of t, both u(t) and v(t) are real.

---

*Wait denotes our $w_1(t)$ by $w_2(t)$, and our $w_2(t)$ by $w_1(t)$.

The integral definitions of u(t) and v(t) for real $\underline{t}$ can be taken to be

$$v(t) = \frac{1}{\sqrt{\pi}} \int_0^\infty \cos\left(\frac{x^3}{3} + tx\right) dx \qquad (2\text{-}4a)$$

$$u(t) = \frac{1}{\sqrt{\pi}} \int_0^\infty \left| \sin\left(\frac{x^3}{3} + tx\right) + \exp\left(-\frac{x^3}{3} + xt\right) \right| dx \qquad (2\text{-}4b)$$

The Airy functions are so fundamental in the mathematical discussion of a variety of physical phenomena that it is almost unexplainable that such an important function had not received a universally accepted notation prior to the beginning of World War II. The author has observed that this function is now finding its way into more and more textbooks, but the notation which seems to be "catching on" the most firmly is that which is due to Jeffreys and Miller (Ref. 5). The functions which were tabulated by Miller were denoted by Ai(x) and Bi(x) and defined by the integrals

$$Ai(x) = \frac{1}{\pi} \int_0^\infty \cos\left(\frac{t^3}{3} + xt\right) dt \qquad (2\text{-}5a)$$

$$Bi(x) = \frac{1}{\pi} \int_0^\infty \left[\sin\left(\frac{t^3}{3} + xt\right) + \exp\left(-\frac{t^3}{3} + xt\right) \right] dt \qquad (2\text{-}5b)$$

Since Eqs. (2-4) and (2-5) differ by only a factor of the form $\sqrt{\pi}$, it is very easy to employ both these notations since the conversion between them is

$$v(t) = \sqrt{\pi}\, Ai(t) \quad , \quad u(t) = \sqrt{\pi}\, Bi(t)$$

$$w_1(t) = \sqrt{\pi}\left[Bi(t) + i\, Ai(t)\right] \quad , \quad w_2(t) = \sqrt{\pi}\left[Bi(t) - i\, Ai(t)\right]$$

The need for the Airy functions in the theory of radiowave propagation during World War II was so great that each research group adopted its own notation. In the classic work published just before the start of the War, van der Pol and Bremmer (Ref. 6) had employed approximations for the Hankel functions $H_\nu^{(1)}(kr)$ which involved the Hankel functions

$$H_{\frac{1}{3}}^{(1)}(z) \quad , \quad H_{-\frac{1}{3}}^{(1)}(z) \quad , \quad H_{\frac{2}{3}}^{(1)}(z) \quad , \quad H_{-\frac{2}{3}}^{(1)}(z)$$

and much of the computational effort associated with their work revolved around finding the roots τ_s which were determined by the equation

$$\frac{\exp(-i\,\pi/3)\, H_{2/3}^{(1)}\left|\frac{1}{3}(-2\tau_s)^{3/2}\right|}{H_{1/3}^{(1)}\left|\frac{1}{3}(-2\tau_s)\right|^{3/2}} = -\frac{1}{\delta_e\sqrt{-2\tau_s}} \tag{2-6}$$

When Furry and his co-workers at the Radio Research Laboratory at Harvard University became interested in these problems they observed that by defining

$$\xi = \frac{2}{3}x^{3/2}$$

that they could simplify the theory by working with a set of modified Hankel functions of order one-third which were defined by means of the relations

$$\left.\begin{aligned} h_j(x) &= \left(\frac{2}{3}\right)^{1/3} x^{1/2} H_{1/3}^{(j)}\left(\frac{2}{3}x^{3/2}\right) = \xi^{1/3} H_{1/3}^{(j)}(\xi) \\ \frac{dh_j(x)}{dx} &= \left(\frac{2}{3}\right)^{1/3} x\, H_{-2/3}^{(j)}\left(\frac{2}{3}x^{3/2}\right) = \left(\frac{3}{2}\right)^{1/3} \xi^{2/3} H_{-2/3}^{(j)}(\xi) \end{aligned}\right\} j = 1,2 \tag{2-7}$$

Tables of these functions (Ref. 7) were computed on the famous Automatic Sequence Controlled Calculator (or "Mark I" as it came to be affectionately called by its users) which had been presented to Harvard University by the International Business Machines Corporation in 1944. In the introduction to the tables, Furry points out that the functions are solutions of

$$\frac{d^2 h_j}{dx^2} + x\, h_j = 0$$

and that they are related to the Airy functions being used by Miller and the English workers by means of the relations

$$h_1(x) = k\,[Ai(-x) - iBi(-x)]$$
$$h_2(x) = k^*[Ai(-x) + iBi(-x)]$$

where

$$k = (12)^{1/6}\exp(-i\pi/6) \quad , \quad k^* = (12)^{1/6}\exp(+i\pi/6)$$

The digression which we have made above to point out some of the differences in notation for the Airy functions is far from complete. For a more complete discussion, the reader is referred to a report by Logan (Ref. 3). However, in order to proceed with the study of the roots of $w_1'(t) - qw_1(t) = 0$, the author feels that the reader should have before him the following relations:

$$w_1(t) = -\sqrt{\pi}(12)^{-1/6}\exp(-i\tfrac{1}{6}\pi)h_1(-t) \qquad (2\text{-}8a)$$

$$= \sqrt{-\tfrac{1}{3}\pi t}\,\exp(i\tfrac{2}{3}\pi)H_{\frac{1}{3}}^{(1)}[\tfrac{2}{3}(-t)^{3/2}] \qquad (2\text{-}8b)$$

$$= u(t) + iv(t) \qquad (2\text{-}8c)$$

$$= \sqrt{\pi}\,[Bi(t) + iAi(t)] \qquad (2\text{-}8d)$$

$$= 2\sqrt{\pi}\,\exp(i\pi/6)Ai[t\exp(i\tfrac{2}{3}\pi)] \qquad (2\text{-}8e)$$

From Eq. (2-8e) we readily see that the solutions a_s of

$$\mathrm{Ai}'(-a_s) + q \exp(i\tfrac{1}{3}\pi)\, \mathrm{Ai}(-a_s) = 0 \qquad (2\text{-}9)$$

are the same as the solutions of Eq. (2-1), namely,

$$w_1'(t_s) - q w_1(t_s) = 0$$

provided we observe that

$$t_s = a_s \exp(i\tfrac{1}{3}\pi) \qquad (2\text{-}10)$$

If we use Eq. (2-7) and Eq. (2-8b), we find that

$$q = \frac{w_1'(t_s)}{w_1(t_s)} = -\frac{\xi_s^{\frac{1}{3}} H^{(1)}_{-\frac{2}{3}}(\xi_s)}{(\frac{2}{3})^{\frac{1}{3}} H^{(1)}_{\frac{1}{3}}(\xi_s)} = \frac{\sqrt{-t_s}\exp(-i\frac{1}{3}\pi) H^{(1)}_{\frac{2}{3}}(\xi_s)}{H^{(1)}_{\frac{1}{3}}(\xi_s)} \qquad (2\text{-}11)$$

where

$$\xi_s = \frac{2}{3}(-t_s)^{3/2} = \frac{1}{3}(-2\tau_{s-1})^{3/2} \qquad (2\text{-}11)$$

Therefore, we observe from a comparison of Eq. (2-6) and (2-11) that the roots t_s which were studied by Fock are essentially the same as the roots τ_{s-1} which were studied by van der Pol and Bremmer* and by Bremmer (Ref. 8).

Since van der Pol and Wyngaarden (Ref. 9) were influenced by the work and the tables of Furry (Ref. 7), their Eq. (4), namely

$$h(\mu_s) = b\, h'(\mu_s) \qquad (2\text{-}12a)$$

can be shown to be

$$h_2(\mu_s) = b\, h_2'(\mu_s)\,. \qquad (2\text{-}12b)$$

Since the modified Hankel functions possess the properties

---

*The relationship is $t_s = 2^{\frac{1}{3}}\tau_{s-1}$ since Fock lets $s = 1,2,3,\ldots$ whereas van der Pol and Bremmer let $s = 0,1,2,\ldots$.

$$h_1(z) = [h_2(z^*)]^* \quad , \quad h_1'(z) = [h_2'(z^*)]^*$$

(where the asterisks denote that the complex conjugate quantities are to be employed), we observe from the property

$$q = \frac{w_1'(t_s)}{w_1(t_s)} = - \frac{h_1'(-t_s)}{h_1(-t_s)} \tag{2-13}$$

that the roots t_s are related to the roots μ_s (of Ref. 9) by the simple property

$$t_s = \text{complex conjugate } (-\mu_s)$$

Further confusion has been fostered by the fact that a number of authors are employing an Airy function A(z) defined by

$$A(z) = \int_0^\infty \cos(\tau^3 - z\tau)d\tau \tag{2-14a}$$

which is related to the more conventional Airy function Ai(x) by means of the relations

$$Ai(x) = \frac{3^{\frac{1}{3}}}{\pi} A(-3^{\frac{1}{3}}x) \quad , \quad Ai'(x) = - \frac{3^{\frac{2}{3}}}{\pi} A'(-3^{\frac{1}{3}}x) \tag{2-14b}$$

From Eq. (2-9) we see that

$$q = \frac{w_1'(t_s)}{w_1(t_s)} = \frac{\exp(i\frac{2}{3}\pi)Ai'(-a_s)}{Ai(-a_s)} = \frac{3^{\frac{1}{3}}\exp(-i\frac{1}{3}\pi)A(3^{\frac{1}{3}}a_s)}{A(3^{\frac{1}{3}}a_s)} \tag{2-15}$$

where t_s and a_s are related as in Eq. (2-10).

The remarks which we have made above are sufficient to show the reader that he may expect to find the equivalent of Eq. (2-1) appearing in many disguises as he looks through the literature.

From the property

$$q(t) = \frac{d}{dt}\{\log[w_1(t)]\} = \frac{w_1'(t)}{w_1(t)} \tag{2-16}$$

we observe that the inverse problem to the finding of the roots of Eq. (2-1) can be described as "studying the behavior of the logarithmic derivative of the Airy function $w_1(t)$."

The occurrence of equations such as Eq. (2-1) is not limited to the radiowave propagation problems. For example, Jeffreys (Ref. 10) considered an elastic wave propagation problem which led him to study the problem of obtaining the roots ξ of the equation

$$\frac{Ai'(-\xi)}{Ai(-\xi)} = K$$

where k is a constant.

2.2 The Riccati Equation

The roots τ_s of Eq. (2-1), i.e., the roots of

$$w_1'(\tau_s) - q\, w_1(\tau_s) = 0 \qquad (2\text{-}17)$$

can often be evaluated by observing that the Airy equation

$$w_1''(\tau) = \tau\, w_1(\tau)$$

reveals that

$$\frac{dq(\tau)}{d\tau} = \frac{d}{d\tau}\,\frac{w_1'(\tau)}{w_1(\tau)} = \frac{w_1''(\tau)}{w_1(\tau)} - \left[\frac{w_1'(\tau)}{w_1(\tau)}\right]^2$$

or

$$\frac{dq}{dt} = t - q^2 \qquad \text{or} \qquad \frac{dt}{dq} = \frac{1}{t - q^2} \qquad (2\text{-}18)$$

Eq. (2-18) is a common type of non-linear first-order differential equation which is known in the literature as a Riccati equation. It is a simple matter to obtain the initial conditions under which Eq. (2-18) is to be integrated. Eq. (2-8e) reveals that $w_1(\tau)$ is merely a rotation of the function Ai(z). The roots of

the Airy function and its derivatives are well known. Interest in these roots goes back to as early as 1838 when Airy showed that the "rainbow integral"

$$W(m) = \int_0^\infty \cos\left[\frac{\pi}{2}(w^3 - mw)\right] dw$$

could be used to describe the maxima and minima of the system of spurious (or supernumerary) rainbows which are observed near the inner edge of the primary bow or the outer edge of the secondary bow. Numerical results for these quanties depended upon finding the roots of $W(m) = 0$ and $W'(m) = 0$. Airy's first numerical calculations led him to a determination of the first minimum and the first two maxima. By 1848 he had extended the results to find the location of the second zero. Then, in a classic paper which appeared in 1850, Stokes gave values (computed to 4 decimals) of the first 50 zeros and the first 10 maxima. During World War II, Miller (Ref. 5) obtained values (to 8 decimals) of the first 50 zeros of $Ai(x)$ and $Ai'(x)$. These results proved to be inadequate for some applications being made by the present author, and in 1958 he received 15-decimal values of these zeros from G. F. Miller and P. H. Haines at the National Physical Laboratory (Teddington, England). In the following year, Sherry (Ref. 11) published values for the first 50 zeros which are accurate to 25 decimals.

Let us now return to Eq. (2-17) and denote the roots for $q = 0$ by τ_s^0 and the root for $q \to \infty$ by τ_s^∞. Then we have

$$w_1(\tau_s^\infty) = 0 \quad , \quad w_1'(\tau_s^0) = 0 \qquad (2\text{-}19)$$

where

$$\tau_s^\infty = \alpha_s \exp(i\tfrac{1}{3}\pi) \quad , \quad \tau_s^0 = \beta_s \exp(i\tfrac{1}{3}\pi) \qquad (2\text{-}20)$$

where α_s and β_s are the roots of the Airy function and its derivative

$$Ai(-\alpha_s) = 0 \quad , \quad Ai'(-\beta_s) = 0$$

Values for these constants are given in Tables 2-1 and 2-2. Values are also given for the "turning values" of the functions.

Table 2-1

ROOTS AND TURNING VALUES OF Ai(-α)

s	α_s	Ai'(- α_s)
1	2. 33810 74104 59767	+0. 70121 08227 20691
2	4. 08794 94441 30971	-0. 80311 13696 54864
3	5. 52055 98280 95551	+0. 86520 40258 94152
4	6. 78670 80900 71759	-0. 91085 07370 49602
5	7. 94413 35871 20853	+0. 94733 57094 41568
6	9. 02265 08533 40980	-0. 97792 28085 69499
7	10. 04017 43415 58086	+0. 00437 01226 60312
8	11. 00852 43037 33263	-1. 02773 86888 20786
9	11. 93601 55632 36263	+1. 04872 06485 88189
10	12. 82877 67528 65757	-1. 06779 38591 57428
11	13. 69148 90352 10718	+1. 08530 28313 50700
12	14. 52782 99517 75335	-1. 10150 45702 77497
13	15. 34075 51359 77997	+1. 11659 61779 32656
14	16. 13268 51569 45771	-1. 13073 23104 93188
15	16. 90563 39974 29943	+1. 14403 66732 73553
16	17. 66130 01056 97057	-1. 15660 98491 16566
17	18. 40113 25992 07115	+1. 16853 47844 87525
18	19. 12638 04742 46952	-1. 17988 07298 70146
19	19. 83812 98917 21500	+1. 19070 61311 58776
20	20. 53733 29076 77567	-1. 20106 07915 19823
21	21. 22482 99436 42097	+1. 21098 75148 68287
22	21. 90136 75955 85131	-1. 22052 33738 97260
23	22. 56761 29174 96503	+1. 22970 07015 09681
24	23. 22416 50011 21681	-1. 23854 78753 29632
25	23. 87156 44555 35918	+1. 24708 99452 59407
26	24. 51030 12365 89678	-1. 25534 91404 75735
27	25. 14082 11661 48964	+1. 26334 52827 50799
28	25. 76353 14009 82756	-1. 27109 61262 18604
29	26. 37880 50521 37232	+1. 27861 76388 24258
30	26. 98698 51116 06368	-1. 28592 42371 22704
31	27. 58838 78098 82445	+1. 29302 89834 49956
32	28. 18330 55026 38645	-1. 29994 37525 11048
33	28. 77200 91652 37435	+1. 30667 93729 32094
34	29. 35475 05587 66288	-1. 31324 57481 80648
35	29. 93176 41190 86556	+1. 31965 19603 77514
36	30. 50326 86114 18505	-1. 32590 63598 38441
37	31. 06946 85851 83756	+1. 33201 66426 47702
38	31. 63055 56580 12659	-1. 33798 99181 42291
39	32. 18670 96529 52051	+1. 34383 27678 48983
40	32. 73809 96090 00269	-1. 34955 12971 47445
41	33. 28488 46819 01402	+1. 35515 11807 15907
42	33. 82721 49495 08652	-1. 36063 77026 40532
43	34. 36523 21338 63659	+1. 36601 57919 26784
44	34. 89907 02503 45312	-1. 37129 00540 34239
45	35. 42885 61927 47888	+1. 37646 47989 60084
46	35. 95471 02618 98629	-1. 38154 40663 17105
47	36. 47674 66443 74809	+1. 38653 16477 85955
48	36. 99507 38469 94501	-1. 39143 11072 66471
49	37. 50979 50920 05016	+1. 39624 57990 06725
50	38. 02100 86772 55254	-1. 40097 88839 49769
51	38. 52880 83050 94249	+1. 40563 33445 05322
52	39. 03328 33832 72514	-1. 41021 19979 25998
53	39. 53451 93007 23018	+1. 41471 75084 44110
54	40. 03259 76807 54176	-1. 41915 23983 05068
55	40. 52759 66138 89718	+1. 42351 90578 16189
56	41. 01959 08723 32490	-1. 42781 97545 15052

Table 2-2

ROOTS AND TURNING VALUES OF $A1'(-\beta)$

s	β_s	$A1(-\beta_s)$
1	1. 01879 29716 47471	+0. 53565 66560 15700
2	3. 24819 75821 79837	-0. 41901 54780 32564
3	4. 82009 92111 78736	+0. 38040 64686 28153
4	6. 16330 73556 39487	-0. 35790 79437 12292
5	7. 37217 72550 47770	+0. 34230 12444 11624
6	8. 48848 67340 19722	-0. 33047 62291 47967
7	9. 53544 90524 33547	+0. 32102 22881 94716
8	10. 52766 03969 57407	-0. 31318 53909 78682
9	11. 47505 66334 80245	+0. 30651 72938 82777
10	12. 38478 83718 45747	-0. 30073 08293 22645
11	13. 26221 89616 65210	+0. 29563 14810 01913
12	14. 11150 19704 62995	-0. 29108 16772 03539
13	14. 93593 71967 20517	+0. 28698 07069 99202
14	15. 73820 13736 92538	-0. 28325 27361 25021
15	16. 52050 38254 33794	+0. 27983 93053 60411
16	17. 28469 50502 16437	-0. 27669 44450 68930
17	18. 03234 46225 04393	+0. 27378 13856 46685
18	18. 76479 84376 65955	-0. 27107 02785 76971
19	19. 48322 16565 67231	+0. 26853 65782 82176
20	20. 18863 15094 63373	-0. 26615 98682 15709
21	20. 88192 27555 16738	+0. 26392 29929 60829
22	21. 56388 77231 98975	-0. 26181 14056 94794
23	22. 23523 22853 43913	+0. 25981 26701 51466
24	22. 89658 87388 74619	-0. 25791 60753 32572
25	23. 54852 62959 28802	+0. 25611 23337 79654
26	24. 19155 97095 26354	-0. 25439 33426 46825
27	24. 82615 64259 21155	+0. 25275 19925 76574
28	25. 45274 25617 77650	-0. 25118 20133 88409
29	26. 07170 79351 73912	+0. 24967 78484 21125
30	26. 68341 03283 22450	-0. 24823 45513 98365
31	27. 28817 91215 23985	+0. 24684 77011 60296
32	27. 88631 84087 68461	-0. 24551 33306 87119
33	28. 47810 96831 02278	+0. 24422 78676 45060
34	29. 06381 41626 38199	-0. 24298 80342 90143
35	29. 64367 48146 32016	+0. 24179 10550 23721
36	30. 21791 81244 68575	-0. 24063 41202 44844
37	30. 78675 56480 12503	+0. 23951 48554 15564
38	31. 35038 53790 83035	-0. 23843 10444 66267
39	31. 90899 29584 30463	+0. 23738 06563 33468
40	32. 46275 27462 38480	-0. 23636 18275 53143
41	33. 01182 87766 34287	+0. 23537 28399 36488
42	33. 55637 56097 89422	-0. 23441 21104 38024
43	34. 09653 90948 09138	+0. 23347 81753 92842
44	34. 63245 70546 35866	-0. 23256 96793 53833
45	35. 16425 99025 53408	+0. 23168 53648 03788
46	35. 69207 11985 10469	-0. 23082 40630 53231
47	36. 21600 81523 35199	+0. 22998 46861 64426
48	36. 73618 20799 46803	-0. 22916 62197 06428
49	37. 25269 88178 54148	+0. 22836 77166 46283
50	37. 76565 91005 38871	-0. 22758 82910 18357
51	38. 27515 89047 30879	+0. 22682 71133 87890
52	38. 78128 97640 80369	-0. 22608 34059 36628
53	39. 28413 90572 98596	+0. 22535 64383 68475
54	39. 78379 02724 68233	-0. 22464 55241 61432
55	40. 28032 32499 03719	+0. 22395 00171 79277
56	40. 77381 44056 64866	-0. 22326 93086 02552

There are two series which have played such a role in the calculating of the roots $\tau_s(q)$ of Eq. (2-17) that we shall refer to then as the classical expansions. For small values of q, we can express $\tau_s(q)$ in the form

$$\tau_s(q) = \tau_s^o + \sum_{n=1}^{\infty} A_n(\tau_s^o)\, q^n \tag{2-21a}$$

where

$$A_1(\tau) = \frac{1}{\tau} \tag{2-21b}$$

$$A_2(\tau) = -\frac{1}{2\tau^3} \tag{2-21c}$$

$$A_3(\tau) = \frac{1}{3\tau^2} + \frac{1}{2\tau^5} \tag{2-21d}$$

$$A_4(\tau) = -\frac{7}{12\tau^4} - \frac{5}{8\tau^7} \tag{2-21e}$$

$$A_5(\tau) = \frac{1}{5\tau^3} + \frac{21}{20\tau^6} + \frac{7}{8\tau^9} \tag{2-21f}$$

$$A_6(\tau) = -\frac{29}{45\tau^5} - \frac{77}{40\tau^8} - \frac{21}{16\tau^{11}} \tag{2-21g}$$

$$A_7(\tau) = \frac{1}{7\tau^4} + \frac{76}{45\tau^7} + \frac{143}{40\tau^{10}} + \frac{33}{16\tau^{13}} \tag{2-21h}$$

$$A_8(\tau) = -\frac{97}{140\tau^6} - \frac{163}{40\tau^9} - \frac{429}{64\tau^{12}} - \frac{429}{128\tau^{15}} \tag{2-21i}$$

$$A_9(\tau) = \frac{1}{9\tau^5} + \frac{13661}{5670\tau^8} + \frac{6769}{720\tau^{11}} + \frac{2431}{192\tau^{14}} + \frac{715}{128\tau^{17}} \tag{2-21j}$$

$$A_{10}(\tau) = -\frac{2309}{3150\tau^7} - \frac{820573}{113400\tau^{10}} - \frac{37961}{1800\tau^{13}} - \frac{46189}{1920\tau^{16}} - \frac{2431}{256\tau^{19}} \tag{2-21k}$$

$$A_{11}(\tau) = \frac{1}{11\tau^6} + \frac{1057}{330\tau^9} + \frac{263653}{13200\tau^{12}} + \frac{4641}{100\tau^{15}} + \frac{29393}{640\tau^{18}} + \frac{4199}{256\tau^{21}} \tag{2-21l}$$

$$A_{12}(\tau) = -\frac{31907}{41580\tau^8} - \frac{3089423}{267300\tau^{11}} - \frac{18592951}{356400\tau^{14}} - \frac{48347}{480\tau^{17}} - \frac{676039}{7680\tau^{20}} - \frac{29393}{1024\tau^{23}} \tag{2-21m}$$

(Text continues on p. 2-16)

Table 2-3

COEFFICIENTS FOR POLYNOMIAL REPRESENTATION FOR $A_n(\tau)$

m	A_n^m
	n = 1
1	.10000000000000000000E 1
	n = 2
1	-.50000000000000000000E +0
	n = 3
1	.50000000000000000000E +0
2	.33333333333333333333E +0
	n = 4
1	-.62500000000000000000E +0
2	-.58333333333333333333E +0
	n = 5
1	.87500000000000000000E +0
2	.10500000000000000000E 1
3	.20000000000000000000E +0
	n = 6
1	-.13125000000000000000E 1
2	-.19250000000000000000E 1
3	-.64444444444444444444E +0
	n = 7
1	.20625000000000000000E 1
2	.35750000000000000000E 1
3	.16888888888888888888E 1
4	.14285714285714285714E +0
	n = 8
1	-.33515625000000000000E 1
2	-.67031250000000000000E 1
3	-.40750000000000000000E 1
4	-.69285714285714285714E +0

m	A_n^m
	n = 9
1	.55859375000000000000E 1
2	.12661458333333333333E 2
3	.94013888888888888888E 1
4	.24093474426807760141 0E 1
5	.11111111111111111111E +0
	n = 10
1	-.94960937500000000000E 1
2	-.24056770833333333333E 2
3	-.21089444444444444444E 2
4	-.72360934744268077600 9E 1
5	-.73301587301587301587 1E +0
	n = 11
1	.16402343750000000000E 2
2	.45926562500000000000E 2
3	.46410000000000000000E 2
4	.19973712121212121212E 2
5	.32030303030303030303E 1
6	.90909090909090909090E -1
	n = 12
1	-.28704101562500000000E 2
2	-.88025911458333333333E 2
3	-.10072291666666666666E 3
4	-.52168773849607 82 98E 2
5	-.11557886270108492330 6E 2
6	-.76736411736411736411 7E +0
	n = 13
1	.50784179687500000000E 2
2	.16928059895833333333E 3
3	.21631614583333333333E 3
4	.13103160213243546576 6E 3
5	.37219566030677141787 8E 2
6	.40628634328634328634 3E 1
7	.76923076923076923076 9E -1

Table 2-3 (Cont'd)

m	B_n^m
	n = 14
1	-.906860351562500000000E 2
2	-.326469726562500000000E 3
3	-.460779687500000000000E 3
4	-.319649399350649350649E 3
5	-.111057272727272727272E 3
6	-.171803429903429903429E 2
7	-.797393083107368821651E +0
	n = 15
1	.144823496328125000000E 3
2	.[illegible]7430468750000000E 3
3	.[illegible]10111111111111111E 3
4	.762357128096440596418E 3
5	.313665989738656405314E 2
6	.498300897691373881846E 1
7	.498300897691373881846E 1
8	.666666666666666666666E -1
	n = 16
1	-.295863189697265625000E 3
2	-.122290118408203124996E 4
3	-.205247083550347222214E 4
4	-.178573479202365921108E 4
5	-.849846271053791887095E 3
6	-.212935579693625989917E 3
7	-.242352048901451282237E 2
8	-.824081474081474081474E +0
	n = 17
1	.539515228271484375000E 3
2	.237386700439453125000E 4
3	.430097192382812500000E 4
4	.412182204241071428552E 4
5	.222876115625000000000E 4
6	.668571732311805841197E 3
7	.100931804534401173054E 3
8	.595861628567510920447E 1
9	.588235294117647058823E -1

m	B_n^m
	n=18
1	-.989111251831054687500E 3
2	-.461585250854492187478E 4
3	-.897884212239583333333E 4
4	-.939836243851273148097E 4
5	-.569359025006430041122E 4
6	-.199300806213484631231E 4
7	-.377697704147928064645E 3
8	-.328471008737893269479E 2
9	-.848107085361987322768E +0
	n = 19
1	.182204704284667968750E 4
2	.898876541137695312500E 4
3	.186842506510416666666E 5
4	.212095927933304398133E 5
5	.142338772545653292171E 5
6	.570228663209453289764E 4
7	.130627541158318984506E 4
8	.152719292019624894291E 3
9	.698560301504426522849E 1
10	.526315789473684210526E -1
	n = 20
1	-.337078702926635742187E 4
2	-.175280925521850585926E 5
3	-.387724038574218750000E 5
4	-.474436188085937500000E 5
5	-.349459286718750000000E 5
6	-.157807166173975840326E 5
7	-.425166397738944062452E 4
8	-.630600872299870751863E 3
9	-.431344333467210955041E 2
10	-.869959153333766336057E +0

Table 2-3 (Cont'd)

m	B_n^m	
	n = 21	
1	.6260033054351806640 63E	4
5	.8448603061728395060 99E	5
9	.2218077197117825103 85E	3
2	.3422151403045654296 60E	5
6	.4248380433795307474 51E	5
10	.8060488264894588827 58E	1
3	.8026308104858398436 80E	5
7	.1318548625859510418 79E	5
11	.4761904761904761904 76E	-1
4	.1053188800577153604 40E	6
8	.2384208215879374681 08E	4
		+0.0
	n = 22	
1	-.1166425237655639648 4E	5
5	-.2015671744015802869 76E	6
9	-.1002914686651305546 74E	4
2	-.6688750469589233397 88E	5
6	-.1117446747904662580 42E	6
10	-.5521009986912292778 71E	2
3	-.1657990242614746093 59E	6
7	-.3931012466451850840 99E	5
11	-.8900026202024513310 69E	+0
4	-.2322402438420430119 43E	6
8	-.8418900546689349840 12E	4
	n = 23	
1	.2181114283561706542 97E	5
5	.4753906782589285713 78E	6
9	.4118863696318483013 27E	4
2	.1308668570137023925 66E	6
6	.2881429796860103131 86E	6
10	.3115461864535694739 20E	3
3	.3418431038818359374 62E	6
7	.1134101971405246395 97E	6
11	.9180268281714526948 53E	1
4	.5091160556489226105 31E	6
8	.2814566463616388245 24E	5
12	.4347826086956521739 13E	-1

m	B_n^m	
	n = 24	
1	-.4089589281678199766 07E	5
5	-.1109913976446132173 52E	7
9	-.1569320875227546051 38E	5
2	-.2562809283185005187 73E	6
6	-.7303590114526841810 48E	6
10	-.1532568416104050736 58E	
3	-.7036216669680277505 68E	6
7	-.3182245460278726068 52E	6
11	-.6918208724831558394 96E	2
4	-.1110277282165463490 07E	7
8	-.8996775458733614048 22E	5
12	-.9085171034625873769 71E	+0
	n = 25	
1	.7688427849555015563 97E	5
5	.2568277146142914331 75E	7
9	.5629170331357282904 07E	5
13	.4000000000000000000 00E	-1
2	.5023106195042610167 90E	6
6	.1823716386407737636 20E	7
10	.6799504680329346884 07E	4
3	.1446087111213658650 53E	7
7	.8719020517620402367 17E	6
11	.4255472816403201881 48E	3
4	.2410033228662574228 26E	7
8	.2769977767267925470 06E	6
12	.1034232341022393810 71E	2
	n = 26	
1	-.1448972940877676010 13E	6
5	-.5895651322685469192 98E	7
9	-.1921327900582214284 22E	6
13	-.9257216662211640311 95E	+0
2	-.9853015997968196867 73E	6
6	-.4494136755798536345 76E	7
10	-.2782214729234670985 08E	5
3	-.2967952939136810302 31E	7
7	-.2340057211579372922 19E	7
11	-.2264795558103938238 08E	4
4	-.5209506357072434858 04E	7
8	-.8261052446791436680 50E	6
12	-.8515396873185961325 86E	2

Table 2-3 (Cont'd)

m	B_n^m	m	B_n^m
	n = 27		n = 29
1	.2736948888324499130 25E 6	1	.9825376858948171138 76E 6
5	.1343745052897279238 37E 8	5	.6850943515271948822 06E 8
9	.6290086692278624900 990E 6	9	.6090023906904920405 79E 7
13	.1154434753487504671 16E 2	13	.7420691907149865681 01E 3
2	.1934110547749312718 46E 7	2	.7467286412800610064 36E 7
6	.1094580110439052486 59E 8	6	.6298963707272994268 84E 8
10	.1066401623393842007 20E 6	10	.1343772838613880646 18E 7
14	.3703703703703703703 70E -1	14	.1278429377751860167 28E 2
3	.6083904486954583061 09E 7	3	.2548051036623215674 84E 8
7	.6167710699078581623 33E 7	7	.4091205425833363573 59E 8
11	.1080344264887086929 99E 5	11	.1929412802928261681 73E 6
4	.1121831410446184143 15E 8	15	.3448275862068965517 24E -1
8	.2397150602466441785 31E 7	4	.5150479916213488259 83E 8
12	.5676810329416671074 11E 3	8	.1884612355300069038 40E 8
		12	.1661290901159638915 64E 5
	n = 28		n = 30
1	-.5180653252899944782 26E 6	1	-.1866821603200152516 36E 7
5	-.3043002454067744677 54E 8	5	-.1534222270891570181 62E 9
9	-.1987424104619707412 27E 7	9	-.1816951204459332905 57E 8
13	-.1032253244285711400 90E 3	13	-.4558268662250215244 36E 4
2	-.3799145718793292839 79E 7	2	-.1468566327850786645 97E 8
6	-.2638166404205204921 00E 8	6	-.1491217395152050564 88E 9
10	-.3872527378667851936 45E 6	10	-.4485299160414389257 83E 7
14	-.9417911856226953432 81E +0	14	-.1234921005203712849 47E 3
3	-.1245715149352592891 88E 8	3	-.5206998071263422964 85E 8
7	-.1599843273195582559 99E 8	7	-.1032974580453623131 38E 9
11	-.4727863922411735950 10E 5	11	-.7430681516960052013 08E 6
4	-.2407506577771342618 51E 8	15	-.9568675277247121355 95E +0
8	-.6792098046351854354 52E 7	4	-.1098710133994354361 76E 9
12	-.3252775048027061904 41E 4	8	-.5133231417191659008 25E 8
		12	-.7747383261241003798 36E 5

Numerical values for the coefficients which appear in Eqs. (2-21b) through (2-21m) are given in Table 2-3. We have expressed the explicit dependence of the $A_n(\tau)$ upon τ in the following manner:

$$A_n(\tau) = \frac{1}{\tau^{2n-1}} \sum_{m=1}^{M} A_n^m \, \tau^{m-1} \qquad (2\text{-}21n)$$

where M is the "integral part" of $[(n + 1)/2]$. The coefficients in Table 2-3 have been obtained by means of the NPREC subroutines which have been discussed in the Preface. The reader is advised to be very careful if he employs these tables for $n > 20$ since the format employed in the computer output has resulted in a table in which m successively takes on the values $1,5,9,\cdots$ in the order listed under the heading "m".

When the magnitude of τ is very large, the dominant terms in the expansion will be the terms

$$A_1(\tau) = \frac{1}{\tau}$$

$$A_3(\tau) = \frac{1}{3\tau^2} + \cdots$$

$$A_5(\tau) = \frac{1}{5\tau^3} + \cdots$$

We recognize these coefficients to be the coefficients in the well known expansion

$$\frac{\tanh^{-1}x}{x} = \frac{\coth^{-1}(1/x)}{x} = \frac{1}{2x}\left[\log(1 + x) - \log(1 - x)\right]$$

$$= 1 + \frac{1}{3}x^2 + \frac{1}{5}x^4 + \frac{1}{7}x^6 + \frac{1}{9}x^8 + \cdots, \quad x^2 < 1$$

Therefore, it appears that for very large values of τ_s^o that the series behaves like the series expansion of

$$t_s^o + (1/\sqrt{t_s^o})\tanh^{-1}(q/\sqrt{t_s^o})$$

Therefore, in order for the series in Eq. (2-21) to converge, we must require that the magnitude of $(q/\sqrt{\tau_s^o})$ be less than unity.* More precise definitions of the radius of convergence of the series in Eq. (2-21) will be possible after we discuss the roots t_s^c for which $\tau_s(q_c) = q_c^2$. We will postpone this discussion until after we have considered another classical expansion for $\tau_s(q)$

For large values of q, we can express $\tau_s(q)$ in the form

$$\tau_s(q) = \tau_s^\infty + \sum_{n=1}^{\infty} B_n(\tau_s^\infty)\, q^{-n} \tag{2-22a}$$

where

$$B_1(\tau) = 1 \tag{2-22b}$$

$$B_2(\tau) = 0 \tag{2-22c}$$

$$B_3(\tau) = \frac{\tau}{3} \tag{2-22d}$$

$$B_4(\tau) = \frac{1}{4} \tag{2-22e}$$

$$B_5(\tau) = \frac{\tau^2}{5} \tag{2-22f}$$

$$B_6(\tau) = \frac{7}{18}\tau \tag{2-22g}$$

$$B_7(\tau) = \frac{1}{7}\tau^3 + \frac{5}{28} \tag{2-22h}$$

$$B_8(\tau) = \frac{29}{60}\tau^2 \tag{2-22i}$$

$$B_9(\tau) = \frac{1}{9}\tau^4 + \frac{41}{81}\tau \tag{2-22j}$$

---

*The author has not found the time to investigate the possible advantages of rearranging the series in Eq. (2-21) so that it is of the form

$$\tau_s(q) = \tau_s^o + (1/\sqrt{\tau_s^o})\tanh^{-1} z + \sum_{n=2}^{\infty} C_n(\tau_s^o)\, z^n$$

where $z = q/\sqrt{\tau_s^o}$.

$$B_{10}(\tau) = \frac{97}{175}\tau^3 + \frac{423}{2520} \qquad (2\text{-}22k)$$

$$B_{11}(\tau) = \frac{1}{11}\tau^5 + \frac{947}{990}\tau^2 \qquad (2\text{-}22l)$$

$$B_{12}(\tau) = \frac{2309}{3780}\tau^4 + \frac{18842}{27216}\tau \qquad (2\text{-}22m)$$

$$B_{13}(\tau) = \frac{1}{13}\tau^6 + \frac{247464}{163800}\tau^3 + \frac{11781}{65520} \qquad (2\text{-}22n)$$

Numerical values for the coefficients which appear in Eqs. (2-22b) through (2-22n) are given in Table 2-4. We have expressed the explicit dependence of the $B_n(\tau)$ upon τ in the following manner:

$$B_n(\tau) = \sum_{m=1}^{M} B_n^m \tau^{m-1} \qquad (2\text{-}22o)$$

where M is the "integral part" of $[(n + 1)/2]$. This definition is quite "wasteful" since the series actually progress in successive power of τ^3. However, the "book-keeping" is somewhat easier if we use the form in Eq. (2-22o) so as to have a single form which is useful for all n instead of having one definition for $n = 1,4,7,10,\ldots$, another for $n = 2,5,8,11,\ldots$ and a third definition for $n = 3,6,9,12,\ldots$.

When the magnitude of τ is very large, the dominant terms in the expansion will be the terms

$$B_1(\tau) = \tau$$

$$B_3(\tau) = \frac{1}{3}\tau^3 + \cdots$$

$$B_5(\tau) = \frac{1}{5}\tau^5 + \cdots$$

$$B_7(\tau) = \frac{1}{7}\tau^7 + \cdots$$

(Text continues on p. 2-29)

Table 2-4

COEFFICIENTS FOR POLYNOMIAL REPRESENTATION FOR $B_n(\tau)$

m	B_n^m
	n = 1
1	.10000000000000000000E 1
	n = 2
1	+0.0
	n = 3
1	+0.0
2	.33333333333333333333E +0
	n = 4
1	.25000000000000000000E +0
2	+0.0
	n = 5
1	+0.0
2	+0.0
3	.19999999999999999999E +0
	n = 6
1	+0.0
2	.38888888888888888888E +0
3	+0.0
	n = 7
1	.17857142857142857142E +0
2	+0.0
3	+0.0
4	.14285714285714285714E +0
	n = 8
1	+0.0
2	+0.0
3	.48333333333333333333E +0
4	+0.0

m	B_n^m
	n = 9
1	+0.0
2	.50617283950617283950E +0
3	+0.0
4	+0.0
5	.11111111111111111111E +0
	n = 10
1	.16785,14285714285714E +0
2	+0.0
3	+0.0
4	.55428571428571428571E +0
5	+0.0
	n = 11
1	+0.0
2	+0.0
3	.95656565656565656565E +0
4	+0.0
5	+0.0
6	.90909090909090909090E -1
	n = 12
1	+0.0
2	.69231334509112286889E +0
3	+0.0
4	+0.0
5	.61084656084656084656E +0
6	+0.0
	n = 13
1	.17980769230769230769E +0
2	+0.0
3	+0.0
4	.15107692307692307692E 1
5	+0.0
6	+0.0
7	.76923076923076923076E -1

Table 2-4 (Cont'd)

m	B_n^m
	n = 14
1	+0.0
2	+0.0
3	.17569441581346343251OE 1
4	+0.0
5	+0.0
6	.65774067202638631209 9E +0
7	+0.0
	n = 15
1	+0.0
2	.97418626298255927885 3E +0
3	+0.0
4	+0.0
5	.21548606701940035273 3E 1
6	+0.0
7	+0.0
8	.66666666666666666666E -1
	n = 16
1	.20800137362637362637 3E +0
2	+0.0
3	+0.0
4	.35348667998667998667 9E 1
5	+0.0
6	+0.0
7	.69771894771894771894 6E +0
8	+0.0
	n = 17
1	+0.0
2	+0.0
3	.31039137315981386943 0E 1
4	+0.0
5	+0.0
6	.28782871532451364384 0E 1
7	+0.0
8	+0.0
9	.58823529411764705882 3E -1

m	B_n^m
	n = 18
1	+0.0
2	.13963268045778333844 1E 1
3	+0.0
4	+0.0
5	.61883153526857230560 8E 1
6	+0.0
7	+0.0
8	.73251686585019918353 0E +0
9	+0.0
	n = 19
1	.25314591423613980004 9E +0
2	+0.0
3	+0.0
4	.75875390524262704713 6E 1
5	+0.0
6	+0.0
7	.36727998818224382134 0E 1
8	+0.0
9	+0.0
10	.52631578947368421052 6E -1
	n = 20
1	+0.0
2	+0.0
3	.53545064003615201872 1E 1
4	+0.0
5	+0.0
6	.98694558376989749538 5E 1
7	+0.0
8	+0.0
9	.76329637682578859049 2E +0
10	+0.0

Table 2-4 (Cont'd)

m	B_n^m
	n = 21
1	+0.0
2	.20274541706387119449 4E 1
3	+0.0
4	+0.0
5	.15744038390442270512 7E 2
6	+0.0
7	+0.0
8	.45317800701020655669 0E 1
9	+0.0
10	+0.0
11	.47619047619047619047 6E -1
	n = 22
1	.31940253120939211164 7E +0
2	+0.0
3	+0.0
4	.15405278998205498949 2E 2
5	+0.0
6	+0.0
7	.14721449760752161138 1E 2
8	+0.0
9	+0.0
10	.79087195757615121532 7E +0
11	+0.0
	n = 23
1	+0.0
2	+0.0
3	.90897819138906711885 0E 1
4	+0.0
5	+0.0
6	.29190960007067913805 3E 2
7	+0.0
8	+0.0
9	.54497996829442451169 2E 1
10	+0.0
11	+0.0
12	.43478260869565217391 3E -1

m	B_n^m
	n = 24
1	+0.0
2	.29721955453724352443 2E 1
3	+0.0
4	+0.0
5	.36962435617442178415 9E 2
6	+0.0
7	+0.0
8	.20879477575203889121 2E 2
9	+0.0
10	+0.0
11	.81583573518558038681 7E +0
12	+0.0
	n = 25
1	.41401298828051647600 3E +0
2	+0.0
3	+0.0
4	.30083761057423716745 6E 2
5	+0.0
6	+0.0
7	.49840265636392173893 4E 2
8	+0.0
9	+0.0
10	.64223253038046170609 9E 1
11	+0.0
12	+0.0
13	.40000000000000000000 0E -1
	n = 26
1	+0.0
2	+0.0
3	.15253422079139393079 5E 2
4	+0.0
5	+0.0
6	.78116664039712413466 4E 2
7	+0.0
8	+0.0
9	.28471645596760314946 9E 2
10	+0.0
11	+0.0
12	.83863117242700373259 1E +0
13	+0.0

Table 2-4 (Cont'd)

m	B_n^m
	n = 27
1	+0.0
2	.43894641905880423827E 1
3	+0.0
4	+0.0
5	.81938991564840618568 2E
6	+0.0
7	+0.0
8	.79893157459162654695 4E 2
9	+0.0
10	+0.0
11	.74455120828592138079 6E 1
12	+0.0
13	+0.0
14	.37037037037037037037 0E -1
	n = 28
1	.54799472744605210655 3E +0
2	+0.0
3	+0.0
4	.57080994606336861204 8E 2
5	+0.0
6	+0.0
7	.15024012962178463023 3E 3
8	+0.0
9	+0.0
10	.37619767348190709130 5E 2
11	+0.0
12	+0.0
13	.85959869006250945754 1E +0
14	+0.0
	n = 29
1	+0.0
2	+0.0
3	.25374443779996556589 0E 2
4	+0.0
5	+0.0
6	.19442183362985648709 5E 3
7	+0.0
8	+0.0
9	.12183408358726292434 3E 3

m	B_n^m
	n = 29 (Cont'd)
10	+0.0
11	+0.0
12	.85160562692332108236 5E 1
13	+0.0
14	+0.0
15	.34482758620689655172 4E -1
	n = 30
1	+0.0
2	.65206931454947902457 7E 1
3	+0.0
4	+0.0
5	.17398639482411075419 4E 3
6	+0.0
7	+0.0
8	.26860342405694292144 9E 3
9	+0.0
10	+0.0
11	.48440030945067326399 0E 2
12	+0.0
13	+0.0
14	.87900510658118232039 9E +0
15	+0.0
	n = 31
1	.73756049654103755057 0E +0
2	+0.0
3	+0.0
4	.10593133715486414229 6E 3
5	+0.0
6	+0.0
7	.41591719006262661809 7E 3
8	+0.0
9	+0.0
10	.17842497041130901766 9E 3
11	+0.0
12	+0.0
13	.96310870261945895502 9E 1
14	+0.0
15	+0.0
16	.32258064516129032258 0E -1

Table 2-4 (Cont'd)

m	B_n^m n = 32
1	+0.0
2	+0.0
3	.41924823166894404386E 2
4	+0.0
5	+0.0
6	.45795864028248810588 0E 3
7	+0.0
8	+0.0
9	.45300040334300482172 5E 3
10	+0.0
11	+0.0
12	.61043568909858093206 0E 2
13	+0.0
14	+0.0
15	.89706330724191762712 4E +0
16	+0.0

n = 33

1	+0.0
2	.97330541062746599701 7E 1
3	+0.0
4	+0.0
5	.35724643093490827057 4E 3
6	+0.0
7	+0.0
8	.82051430433305492515 8E 3
9	+0.0
10	+0.0
11	.25269916625353609904 1E 3
12	+0.0
13	+0.0
14	.10788086532295894860 51E 2
15	+0.0
16	+0.0
17	.30303030303030303030 3E -1

m	B_n^m n = 34
1	.10063566436218606220 3E 1
2	+0.0
3	+0.0
4	.19315907481137165490 2E 3
5	+0.0
6	+0.0
7	.10786215300780753327 1E 4
8	+0.0
9	+0.0
10	.72837770410598695873 6E 3
11	+0.0
12	+0.0
13	.75536946274867368754 6E 2
14	+0.0
15	+0.0
16	.91394578031068786930 1E +0
17	+0.0

n = 35

1	+0.0
2	+0.0
3	.66893075841245259571 4E 2
4	+0.0
5	+0.0
6	.10330010103579762234 9E 4
7	+0.0
8	+0.0
9	.15164566894807756879 6E 4
10	+0.0
11	+0.0
12	.34795562755234278393 1E 3
13	+0.0
14	+0.0
15	.11984821953929999787 2E 2
16	+0.0
17	+0.0
18	.28571428571428571428 5E -1

Table 2-4 (Cont'd)

m	B_n^m
	n = 36
1	+0.0
2	.14585597218200322572 3E 2
3	+0.0
4	+0.0
5	.71407245709032556048 0E 3
6	+0.0
7	+0.0
8	.23274687595838646698 3E 4
9	+0.0
10	+0.0
11	.11254665999561945498 4E 4
12	+0.0
13	+0.0
14	.92022580754239325776 5E 2
15	+0.0
16	+0.0
17	.92979417388023448187 5E +0
18	+0.0
	n = 37
1	.13888224533024508197 8E 1
2	+0.0
3	+0.0
4	.34720860260156469737 3E 3
5	+0.0
6	+0.0
7	.26562110220082787902 0E 4
8	+0.0
9	+0.0
10	.26560060070938662611 1E 4
11	+0.0
12	+0.0
13	.46775326438395905726 5E 3
14	+0.0
15	+0.0
16	.13219309368743013203 3E 2
17	+0.0
18	+0.0
19	.27027027027027027027 0E -1

m	B_n^m
	n = 38
1	+0.0
2	+0.0
3	.11270330719017557894 6E 3
4	+0.0
5	+0.0
6	.22500381822571520775 0E 4
7	+0.0
8	+0.0
9	.46780990430299819193 6E 4
10	+0.0
11	+0.0
12	.16814154734981769673 0E 4
13	+0.0
14	+0.0
15	.11059910646436408781 4E 3
16	+0.0
17	+0.0
18	.94472619[illegible]7637E +0
19	+0.0
	n = 39
1	+0.0
2	.21930519062148916077 9E 2
3	+0.0
4	+0.0
5	.13961917816464701605 9E 4
6	+0.0
7	+0.0
8	.62232672750812864943 8E 4
9	+0.0
10	+0.0
11	.44467614148456389324 7E 4
12	+0.0
13	+0.0
14	.61590549495678032320 0E 3
15	+0.0
16	+0.0
17	.14489763670667863963 9E 2
18	+0.0
19	+0.0
20	.25641025641025641025 6E -1

Table 2-4 (Cont'd)

m	B_n^m
	n = 40
1	.19351727878111440988 8E 1
2	+0.0
3	+0.0
4	.61675311241431911550 3E 3
5	+0.0
6	+0.0
7	.62718863707938563340 4E 4
8	+0.0
9	+0.0
10	.88655875060148221130 1E 4
11	+0.0
12	+0.0
13	.24404216200858197366 2E 4
14	+0.0
15	+0.0
16	.13136169059234590780 4E 3
17	+0.0
18	+0.0
19	.95884069005271082749 8E +0
20	+0.0
	n = 41
1	+0.0
2	+0.0
3	.18368565429039014605 8E 3
4	+0.0
5	+0.0
6	.47616252996672314511 8E 4
7	+0.0
8	+0.0
9	.13515674102211869166 8E 5
10	+0.0
11	+0.0
12	.71644691399523686205 8E 4
13	+0.0
14	+0.0
15	.79647502137684760467 2E 3
16	+0.0
17	+0.0
18	.15794574233326871784 9E 2
19	+0.0
20	+0.0
21	.24390243902439024390 2E -1

m	B_n^m
	n = 42
1	+0.0
2	.33068322232248549881 7E 2
3	+0.0
4	+0.0
5	.26801453655965849944 1E 4
6	+0.0
7	+0.0
8	.15855640311338162472 8E 5
9	+0.0
10	+0.0
11	.15988303429049228207 0E 5
12	+0.0
13	+0.0
14	.34543614208246498394 4E 4
15	+0.0
16	+0.0
17	.15440231066429496538 5E 3
18	+0.0
19	+0.0
20	.97222144791602305073 0E +0
21	+0.0
	n = 43
1	.27187711430588036620 5E 1
2	+0.0
3	+0.0
4	.10846458541918446407 1E 4
5	+0.0
6	+0.0
7	.14301952636546437693 0E 5
8	+0.0
9	+0.0
10	.27556850668841872724 7E 5
11	+0.0
12	+0.0
13	.11167351476234589447 1E 5
14	+0.0
15	+0.0
16	.10137688287582756663 3E 4
17	+0.0
18	+0.0
19	.17132279115304294497 7E 2
20	+0.0

Table 2-4 (Cont'd)

m	B_n^m
	n = 43 (Cont'd)
21	+0.0
22	.23255813953488372093E -1
	n = 44
1	+0.0
2	+0.0
3	.29842768707165112129E 3
4	+0.0
5	+0.0
6	.98358447297616784590E 4
7	+0.0
8	+0.0
9	.37002283004106219886E 5
10	+0.0
11	+0.0
12	.27635640602956603457E 5
13	+0.0
14	+0.0
15	.47834181603441058576E 4
16	+0.0
17	+0.0
18	.17980999864115441754E 3
19	+0.0
20	+0.0
21	.98494009697136232240E +0
22	+0.0
	n = 45
1	+0.0
2	.49985713308389152933E 2
3	+0.0
4	+0.0
5	.50652496291481823612E 4
6	+0.0
7	+0.0
8	.38804379714157803155E 5
9	+0.0
10	+0.0
11	.53258787501649236658E 5
12	+0.0
13	+0.0

m	B_n^m
	n = 45 (Cont'd)
14	.16911979790884366973E 5
15	+0.0
16	+0.0
17	.12723334037134198246E 4
18	+0.0
19	+0.0
20	.18501544949988377649E 2
21	+0.0
22	+0.0
23	.22222222222222222222E -1
	n = 46
1	.38470442447016547165E 1
2	+0.0
3	+0.0
4	.16912685421920507407E 4
5	+0.0
6	+0.0
7	.31668806544128928040E 5
8	+0.0
9	+0.0
10	.80764903463533501820E 5
11	+0.0
12	+0.0
13	.46045267360116575535E 5
14	+0.0
15	+0.0
16	.64965706943407468289E 4
17	+0.0
18	+0.0
19	.20767105692490304263E 3
20	+0.0
21	+0.0
22	.99705833771653053506E +0
23	+0.0

Table 2-4 (Cont'd)

m	B_n^m
	n = 47
1	+0.0
2	+0.0
3	.48352986543615327878 6E 3
4	+0.0
5	+0.0
6	.19903165494288617313 3E 5
7	+0.0
8	+0.0
9	.96841883630022724794 8E 5
10	+0.0
11	+0.0
12	.98311304716378584343 4E 5
13	+0.0
14	+0.0
15	.24970712271482414369 4E 5
16	+0.0
17	+0.0
18	.15769501647889175136 3E 4
19	+0.0
20	+0.0
21	.19901150334180934699 0E 2
22	+0.0
23	+0.0
24	.21276595744680851063 8E -1
	n = 48
1	+0.0
2	.75720607391468194466 7E 2
3	+0.0
4	+0.0
5	.94456446054455646743 7E 4
6	+0.0
7	+0.0
8	.91789589786253221807 9E 5
9	+0.0
10	+0.0
11	.16654014710207427159 5E 6
12	+0.0
13	+0.0
14	.74293630691747083771 2E 5
15	+0.0
16	+0.0

m	B_n^m
	n = 48 (Cont'd)
17	.86728962998604754984 3E 4
18	+0.0
19	+0.0
20	.23806925048172202113 0E 3
21	+0.0
22	+0.0
23	.10086296880292310263 0E 1
24	+0.0
	n = 49
1	.54776591737586558132 4E 1
2	+0.0
3	+0.0
4	.32734891236421663967 4E 4
5	+0.0
6	+0.0
7	.68385242972101050863 6E 5
8	+0.0
9	+0.0
10	.22530951882085337895 5E 6
11	+0.0
12	+0.0
13	.17437510690100233934 8E 6
14	+0.0
15	+0.0
16	.36050713897125017746 8E 5
17	+0.0
18	+0.0
19	.19326310955378847705 6E 4
20	+0.0
21	+0.0
22	.21329971984050328919 2E 2
23	+0.0
24	+0.0
25	.20408163265306122448 9E -1

Table 2-4 (Cont'd)

m	B_n^m n = 50
1	
2	+0.0
3	+0.0
4	.78159920789608696296 4E 3
5	+0.0
6	+0.0
7	.39566156684222485711 3E 5
8	+0.0
9	+0.0
10	.24394206931435437812 3E 6
11	+0.0
12	+0.0
13	.32698848098248773876 8E 6
14	+0.0
15	+0.0
16	.11652356438185860918 9E 6
17	+0.0
18	+0.0
19	.11400826171147381531 4E 5
20	+0.0
21	+0.0
22	.27108597823534667023 6E 3
23	+0.0
24	+0.0
25	.10197008622318090952 6E 1
	+0.0

[Note added in proof:- The user who wishes to conserve storage in an electronic computer may find it convenient to replace Eq. (2-22c) by

$$B_n(\tau) = \tau^{\mu} \sum_{j=1}^{K} C_j^n \tau^{3(j-1)}$$

where, if we let [...] be interpreted as "the integral part of",

$$\mu = (2n + 1) - 3[(2n + 1)/3] \quad , \quad K = [(2n + 1)/3] = [n/2]$$

For example, if n = 10, we would have $\mu = 0$, K = 2; if n = 11, we would have $\mu = 2$, K = 2; and if n = 12, we would have $\mu = 1$, K = 2.]

As in the discussion following Eq. (2-21), we recognize the coefficients in the expansion of $x^{-1}\tanh^{-1}x$ and observe that the behavior of $\tau_s(q)$ for the cases in which the magnitude of τ_s^∞ is large is given by

$$\tau_s(q) \approx \tau_s^\infty + \frac{1}{\sqrt{\tau_s^\infty}} \tanh^{-1}(\sqrt{\tau_s^\infty}/q) \quad , \quad \left|\sqrt{\tau_s^\infty}/q\right| < 1$$

Since this result suggests that we must require that the magnitude of $\sqrt{\tau_s^\infty}/q$ be less than unity in order to employ the expansion in Eq. (2-22), we see that as the magnitude of τ_s^∞ increases, the region in the q-plane for which this expansion can be employed "shrinks" until it coalesces upon the "point at infinity." This is in sharp contrast with the expansion in Eq. (2-21) since it can be employed in ever enlarging regions in the q-plane as the magnitude of τ_s^o increases.

We observe that the roots for $q = 0$ and $q \to \infty$ lie along the line $\arg(q) = 60^o$, and lie in the first quadrant of the τ-plane. Much attention has been given in the literature to the study of vertically-polarized waves propagated over the convex interface between free space and a highly conducting homogeneous sphere. In this case, the impedance parameter has a phase of the order of 45^o. We will refer to these as the "Watson modes" since this was the type of problem in which Watson (and later van der Pol and Bremmer) was interested nearly fifty years ago. Since World War II there has developed an interest in the propagation of vertically-polarized electromagnetic waves over the convex surface of a perfect conductor which is either

a) slightly rough
b) covered with a thin dielectric layer, or
c) corrugated.

These modes were discussed in a paper by Elliott (Ref. 12) and we shall therefore refer to these as the "Elliot modes." The very important problem of absorbing layers upon a convex surface can be

often shown to correspond to the case in which the impedance parameter q has a phase of the order of 90°. Therefore, the applications which are being encountered are such that it becomes quite important to be able to find the roots $\tau_s(q)$ for an arbitrary value of q. Therefore, it has become important to study the behavior of the logarithmic derivative $q(\tau)$ in order to understand the mapping specified by the functions $\tau_s(q)$ and $q(\tau)$. The mapping is found to possess certain critical values $\tau_s^c(q)$ for which

$$\tau_s^c = q_c^2 = [w_1'(\tau_s^c)/w_1(\tau_s^c)]^2 \qquad (2\text{-}23)$$

We often refer to these points as the "saddle points." It will be observed that these points are singularities for the differential equation for dt/dq given in Eq. (2-18)




Fig. 2-1 The Logarithmic Derivative $q(\tau) = w_1'(\tau)/w_1(\tau)$
(——— phase , - - - - - modulus)

In Fig. 2-1, we have attempted to illustrate some of these properties of the logarithmic derivative function $q(\tau)$, with emphasis upon illustrating the saddle point, the Watson modes, and the Elliott modes. A more detailed illustration is given in Fig. 2-2




Fig. 2-2 The Logarithmic Derivative $q(\tau) = w_1'(\tau)/w_1(\tau)$
(——— phase , – – – – – modulus)

The critical points τ_s^c correspond to points at which two modes coalesce. They are very important in determining the radius of convergence (in the q-plane) of the series which we have given in Eqs. (2-21) and (2-22). The first two of these points have been located at

$$q_c \approx 1.731\exp(i19.29^\circ) \quad , \qquad q_c \approx 2.135\exp(i23.52^\circ)$$

The approximate location of the first 20 saddle points (or critical points) is given in Table 2-5. For $s > 3$, these critical points can be found from an asymptotic representation. From the definition given in Eq. (2-23), we see that the location of the critical points is equivalent to the problem of finding the roots of the function $f(\tau) = [w_1'(\tau)]^2 - \tau[w_1(\tau)]^2$. We can show that

Table 2-5

THE CRITICAL POINTS τ_s^c AND CORRESPONDING VALUES OF q_c

s	Real τ_s^c	Imag τ_s^c	Mod q_c	Arg q_c (deg.)
1	2.343	1.869	1.731	19.29
2	3.118	3.347	2.135	23.52
3	3.771	4.573	2.433	25.25
4	4.353	5.658	2.672	26.21
5	4.898	6.658	2.875	26.83
6	5.409	7.591	3.053	27.264
7	5.894	8.472	3.213	27.586
8	6.358	9.311	3.358	27.836
9	6.804	10.114	3.491	28.036
10	7.234	10.888	3.616	28.200
11	7.651	11.636	3.732	28.337
12	8.056	12.361	3.841	28.453
13	8.450	13.066	3.945	28.553
14	8.835	13.752	4.043	28.640
15	9.212	14.423	4.137	28.717
16	9.580	15.078	4.227	28.785
17	9.941	15.719	4.313	28.845
18	10.295	16.348	4.396	28.900
19	10.643	16.966	4.475	28.949
20	10.986	17.572	4.552	28.994

$f(\tau)$ is a solution of the differential equation

$$\frac{d^3f}{d\tau^3} - 4\tau \frac{df}{d\tau} + 2f = 0$$

This differential equation can be used to generate a Taylor series for $f(\tau)$. If we then use the fact that $f(\tau_s^c) = 0$, we see that the finding of the roots is reduced to the problem of inverting the series

$$f(\tau_s^c) = 0 = f(\tau) + f'(\tau)(\tau_s^c - \tau) + \sum_{n=2}^{\infty} \frac{f^{(n)}(\tau)}{n!}(\tau_s^c - \tau)^n$$

Since $f'(\tau) = -[w_1(\tau)]^2$ and $f''(\tau) = -2w_1(\tau)w_1'(\tau)$, it is a rather straight foward calculation to find a more accurate value of τ_s^c once we know $w_1(\tau)$ and $w_1'(\tau)$ at a neighboring point τ.

Our study of the asymptotic location of the τ_s^c for $s \to \infty$ has indicated that they can be represented in the form

$$\tau_s^c = S(s)\exp[i(\tfrac{1}{3}\pi - T(s)] \qquad (2\text{-}24a)$$

where

$$S(s) = (3/4)(X^2 + Y^2)^{\frac{1}{3}} \quad , \quad T(s) = \tfrac{2}{3}\tan^{-1}(Y/X)$$

where $X = X(s)$ and $Y = Y(s)$ possess the asymptotic behavior

$$X(s) \xrightarrow[s \to \infty]{} 2\pi s - \frac{\log(12\pi s)}{2\pi s} \qquad (2\text{-}24b)$$

$$Y(s) \xrightarrow[s \to \infty]{} \log[12\pi s - \frac{3}{\pi s}\log(12\pi s)] \qquad (2\text{-}24c)$$

If we study the behavior of the lines of constant phase in Fig. 2-2 we observe that when $30^\circ < \arg q < 210^\circ$ that all the root loci which start from τ_s^0 move along paths that end up at τ_s^∞. However, for $-150^\circ < \arg q < 30^\circ$ there is a type of root loci which starts from τ_s^0 but which "slips away" and runs off to infinity. We will designate this root as $\tau_0(q)$. From Fig. 1, we see that the root $\tau_0(q)$ corresponds to $\tau_1(q)$ for $-139.2^\circ < \arg q < 19.2^\circ$, to $\tau_2(q)$ for $-143.4^\circ < \arg q < -139.2^\circ$ and $19.2^\circ < \arg q < 23.4^\circ$, etc.

For the special case of the surface wave modes which we have designated as the "Elliott modes," the impedance parameter is characterized by arg $q = 0^\circ$. For this case we can compute $\tau_0(q)$ for $q >> 2$ by means of the relations

$$\text{Real } \tau_0(q) \approx q^2 + \frac{1}{2}\frac{1}{q} + \frac{1}{8}\frac{1}{q^4} + \frac{5}{32}\frac{1}{q^7} + \frac{11}{32}\frac{1}{q^{10}} + \frac{539}{512}\frac{1}{q^{13}} + \frac{8337}{2048}\frac{1}{q^{16}}$$

$$+ \frac{9659}{512}\frac{1}{q^{19}} + \frac{416349}{4096}\frac{1}{q^{22}} + O(q^{-25}) \qquad (2\text{-}25a)$$

$$\text{Imag } \tau_0(q) \approx 2q^2 \exp\left[-\frac{4}{3}q^3 - 1 - \frac{7}{12q^3} - \frac{31}{48q^6}\right.$$

$$\left. - \frac{397}{288q^9} - \frac{6427}{1536q^{12}} - \frac{248025}{15360q^{15}} - \cdots\right] \qquad (2\text{-}25b)$$

The asymptotic expansions which we have given in Eq. (2-25) for the case of q real (i.e., arg $q = 0^\circ$) can also be employed to obtain asymptotic series for $\tau_0(q)$ for $-150^\circ < \arg q < 30^\circ$ if we write

$$\tau_0(q) \approx \left[q^2 + \frac{1}{2}\frac{1}{q} + \cdots\right] + i2q^2 \exp\left[-\frac{4}{3}q^3 - 1 - \cdots\right] \qquad (2\text{-}26)$$

where further terms in the expansion can be obtained from Eqs. (2-25a) and (2-25b) where now the values of q are complex.

In Fig. 2-3 we have indicated the regions in which one can employ the representations for $\tau_s(q)$ which we have given in Eqs. (2-21), (2-22) and (2-26). The boundary for Eq. (2-26) has been drawn with dashed lines because this is an asymptotic representation and the expansion does not possess an exact boundary outside of which it can be employed. It is important to observe that there




Fig. 2-3 Boundaries for Regions of Applicability of Eqs. (2-21), (2-22) and (2-26)

are root loci (computed on the basis of Eq. (2-21)) which start from $q = 0$ at τ_2^0 and which cannot be "joined" to the root loci (computed on the basis of Eq. (2-22)) which start from $q = \infty$ at τ_2^∞. Since the expansions given by Eqs. (2-21) and (2-22) become useless (from a practical point of view) before the theoretical boundaries are reached, there are considerable "gaps" in the complex q-plane which cannot be reached by employing these classical expansions.

2.3 The Root Locus for the Ground Wave Problem

Before we continue with the discussion of methods for the numerical evaluation of the roots $\tau_s(q)$, let us consider in more detail the root locus which we have referred to as the "Watson modes." Up until about a decade ago the only interest in these roots was that which arose out of the problem of predicting the diffraction effects associated with the propagation of the ground wave around the earth's surface. The rapidly advancing technology associated with the development of missiles and satellites has opened up a broad new range of problems in which the microwave engineer requires the ability to predict the diffraction effects when radiowaves are radiated from (or received by) an antenna mounted upon a convex surface. Another broad new area of interest is that associated with the radar reflection characteristics of bodies whose shape includes include portions that come within the scope of the problem of reflection from and diffraction around convex surfaces. The development of the laser has provided physicists with a source of coherent electromagnetic waves and now (Ref. 20) there exists an interest in this diffraction theory even in the range of the optical wavelengths. The recent paper by Streifer and Kodis (Ref. 21) is probably only the precusor of the flood of papers which will appear within the next decade in which the logarithmic derivative of the Airy function will play a prominent role.

From the rotational property of the Airy functions, we know that

$$q(t) = \frac{w_1'(t)}{w_1(t)} = \exp(i\tfrac{2}{3}\pi)\ \frac{v'[t\exp(i\tfrac{2}{3}\pi)]}{v[t\exp(i\tfrac{2}{3}\pi)]}$$

$$= \exp(i\tfrac{3}{3}\pi)\ \frac{Ai'[t\exp(i\tfrac{2}{3}\pi)]}{Ai[t\exp(i\tfrac{2}{3}\pi)]} \qquad (2\text{-}27)$$

Therefore, by rotating the contour plot given in Fig. 2-2 and

by relabeling the lines of constant phase, we obtain the contour plot of the logarithmic derivative $L(t) = Ai'(t)/Ai(t)$ which is given in Fig. 2-4.

Let us consider briefly a problem in which both the logarithmic derivative $q(t) = w_1'(t)/w_1(t)$ and the logarithmic derivative $L(t) = v'(t)/v(t)$ occur. In the study of diffraction by circular cylinders we have to solve the two dimensional differential equation

$$\left[\frac{\partial^2}{\partial\rho^2} + \frac{1}{\rho}\frac{\partial}{\partial\rho} + \frac{1}{\rho^2}\frac{\partial^2}{\partial\varphi^2} + k^2\right] \Psi(\rho,\varphi) = 0 \qquad (2\text{-}28)$$

Let the electrical properties of the cylinder be characterized by k_1 while that of the outside is described by k. The cylinder has a radius a. We will use an $\exp(-i\omega t)$ time dependence. Since the wave must be outgoing at infinity and finite at $\rho = 0$, we choose




Fig. 2-4 The Logarithmic Derivative $L(t) = Ai'(t)/Ai(t)$
(———— phase , – – – – – modulus)

a solution proportional to

$$\exp(\pm i\upsilon\varphi)\ H_{\upsilon}^{(1)}(k\rho) \quad , \ \rho > a$$

$$\exp(\pm i\upsilon\varphi)\ J_{\upsilon}(k_1\rho) \quad , \ \rho < a$$

where

$$k^2 = \omega^2\varepsilon\mu \quad , \ \rho > a \tag{2-29a}$$

$$k_1^2 = \omega^2\varepsilon_1\mu_1 + i\omega\mu_1\sigma_1 \quad , \ \rho < a \tag{2-29b}$$

In electromagnetic diffraction problems which involve a circular cylinder, there are generally two cases of interest:

(a) $\Psi = E_z$, electric field parallel to the axis of the cylinder

(b) $\Psi = H_z$, magnetic field parallel to the axis of the cylinder

The continuity requirements on the electric and magnetic field at the interface $\rho = a$ lead to an eigenvalue problem for the determination of the roots υ_s defined by

$$\mu_1 k\, J_{\upsilon_s}(k_1 a) H_{\upsilon_s}^{(1)\prime}(ka) - \mu k_1 J_{\upsilon_s}'(k_1 a) H_{\upsilon_s}^{(1)}(ka) = 0 \ , \ \Psi = E_z \tag{2-30a}$$

$$\mu k_1\, J_{\upsilon_s}(k_1 a) H_{\upsilon_s}^{(1)\prime}(ka) - \mu_1 k J_{\upsilon_s}'(k_1 a) H_{\upsilon_s}^{(1)}(ka) = 0 \ , \ \Psi = H_z \tag{2-30b}$$

If the magnitude of k_1 is much greater than that of k, we can show that the magnitude of $k_1 a$ is much greater than the magnitude of υ_s (for small values of s if we order the υ_s and let υ_1 be the root with the smallest imaginary part, υ_2 be the root with the next smallest imaginary part, etc.) and then we can use the approximation

$$\frac{J_{\upsilon_s}'(k_1 a)}{J_{\upsilon_s}(k_1 a)} \approx -i\sqrt{(k_1 a)^2 - \upsilon_s^2}\ /(k_1 a) \tag{2-31}$$

Therefore, we find that we have to solve the problem defined by

$$\frac{H_{\upsilon_s}^{(1)\prime}(ka)}{H_{\upsilon_s}^{(1)}(ka)} \approx \begin{cases} -i(\mu k_1/\mu_{1k})\sqrt{(k_1 a)^2 - \upsilon_s^2}\,/(k_1 a)\,, & \Psi = E_z \quad (2\text{-}32a) \\ -i(\mu_1 k/\mu k_1)\sqrt{(k_1 a)^2 - \upsilon_s^2}\,/(k_1 a)\,, & \Psi = H_z \quad (2\text{-}32b) \end{cases}$$

When ka is very large, there are roots υ_s which are in the vicinity of $\upsilon_s = ka$. Therefore, we can use the Lorenz-Nicholson asymptotic approximation

$$H_{\upsilon}^{(1)}(ka) \approx -(i/\sqrt{\pi})(ka/2)^{-\frac{1}{3}} w_1(t) \qquad (2\text{-}33)$$

where $w_1(t)$ is the Airy function and t is defined by

$$\upsilon = ka + (ka/2)^{\frac{1}{3}} t \qquad (2\text{-}34)$$

Therefore, we can employ the approximation

$$\frac{H_{\upsilon_s}^{(1)\prime}(ka)}{H_{\upsilon_s}^{(1)}(ka)} \approx -(ka/2)^{-\frac{1}{3}} \frac{w_1'(t)}{w_1(t)} \qquad (2\text{-}35)$$

to obtain the following problem which is to be solved for the roots t_s

$$t_s = (ka/2)^{-\frac{1}{3}}(\upsilon_s - ka) \qquad (2\text{-}36)$$

defined by

$$\frac{w_1'(t_s)}{w_1(t_s)} \approx i(ka/2)^{\frac{1}{3}} \begin{cases} \dfrac{\mu k_1}{\mu_1 k} \dfrac{\sqrt{(k_1 a)^2 - \upsilon_s^2}}{(k_1 a)}\,, & \Psi = E_z \quad (2\text{-}37a) \\ \dfrac{\mu_1 k}{\mu k_1} \dfrac{\sqrt{(k_1 a)^2 - \upsilon_s^2}}{(k_1 a)}\,, & \Psi = H_z \quad (2\text{-}37b) \end{cases}$$

If we make the further approximation of replacing υ_s by (ka) in the radicals of Eq. (2-37) we can express the eigenvalue problem in the form

$$q(t_s) = \frac{w_1'(t_s)}{w_1(t_s)} = \begin{cases} q_v \ , & \Psi = E_z \quad (2\text{-}38a) \\ q_h \ , & \Psi = H_z \quad (2\text{-}38b) \end{cases}$$

where the impedance parameters are

$$q_v = i(ka/2)^{\frac{1}{3}}(k/k_1)\sqrt{1 - (k/k_1)^2} \quad (2\text{-}39a)$$

$$q_h = i(ka/2)^{\frac{1}{3}}(k_1/k)\sqrt{1 - (k/k_1)^2} = (k_1/k)^2 q_v \quad (2\text{-}39b)$$

Although we have derived Eq. (2-38) for the case of a circular cylinder, the analysis for the lossy dielectric sphere leads to an identical equation in the limit as $ka \to \infty$. If the diffracting obstacle is lossless, we observe that

$$\arg(q_v) = \arg(q_h) = 90^\circ$$

However, if $\sigma_1 >> \omega\varepsilon_1$, we find that

$$\arg(q_v) = 45^\circ$$

$$\arg(q_h) = 135^\circ$$

Therefore, we observe that

$$45^\circ < \arg(q_v) < 90^\circ$$

$$90^\circ < \arg(q_h) < 135^\circ$$

Most of the papers published to date have dealt with the eigenvalue problem defined by Eq. (2-38). This approximation is generally adequate for the problem of studying the propagation of the ground wave around the earth's surface. However, in problems which arise in the applications of microwaves, the condition that $k_1 a$ be much larger than ka is not satisfied in many of the situations for which data on the propagation phenomena is desired. Let us sketch a method which can be employed under much less stringent conditions than those which must be satisfied in order to have the problem reduce to Eq. (2-38).

We need an asymptotic approximation for the Bessel functions which is valid when both ka and $k_1 a$ are large, but the relation $|k_1 a| >> |ka|$ need not hold. A suitable asymptotic estimate is that which was employed in the mid-1930's by Langer and which has been employed in the more recent studies made by Olver (Ref. 22). In order to simplify the asymptotic formulae, let us make a few definitions. Let $\zeta = \zeta(z)$ be defined by

$$\frac{2}{3}\zeta^{3/2} = \log[(1 + \sqrt{1 - z^2})/z] - \sqrt{1 - z^2} \tag{2-40a}$$

and then define

$$\Phi(\zeta) = [4\zeta/(1 - z^2)]^{1/4} \tag{2-40b}$$

$$\Psi(\zeta) = 2/[z\Phi(\zeta)] \tag{2-40c}$$

The Langer-Olver asymptotic estimates are then given by

$$\sqrt{\pi}\, H_\upsilon^{(1)}(\upsilon z) \approx -i\Phi(\zeta)\upsilon^{-\frac{1}{3}} w_1(\upsilon^{\frac{2}{3}}\zeta) \tag{2-41}$$

$$\sqrt{\pi}\, H_\upsilon^{(1)\prime}(\upsilon z) \approx i\Psi(\zeta)\upsilon^{-\frac{2}{3}} w_1'(\upsilon^{\frac{2}{3}}\zeta) \tag{2-42}$$

$$\sqrt{\pi}\, J_\upsilon(\upsilon z) \approx \Phi(\zeta)\upsilon^{-\frac{1}{3}} v(\upsilon^{\frac{2}{3}}\zeta) \tag{2-43}$$

$$\sqrt{\pi}\, J_\upsilon'(\upsilon z) \approx -\Psi(\zeta)\upsilon^{-\frac{2}{3}} v'(\upsilon^{\frac{2}{3}}\zeta) \tag{2-44}$$

In order to use Eqs. (2-41) through (2-44) in Eq. (2-30), we define the quantities z and z_1 and ζ and ζ_1 by means of the relations

$$z = ka/\upsilon_s \quad , \quad z_1 = k_1 a/\upsilon_s$$
$$\zeta = \zeta(z) \quad , \quad \zeta_1 = \zeta(z_1)$$

and then observe that the logarithmic derivatives of the Bessel functions which occur in Eq. (2-30) can be expressed in the form

$$\frac{H_\upsilon^{(1)\prime}(ka)}{H_\upsilon^{(1)}(ka)} \approx -2\upsilon^{-\frac{1}{3}}(\upsilon/ka)\sqrt{[1-(ka/\upsilon)^2/(4\zeta)]}\ \frac{w_1'(\upsilon^{\frac{2}{3}}\zeta)}{w_1(\upsilon^{\frac{2}{3}}\zeta)} \qquad (2\text{-}45)$$

$$\frac{J_\upsilon'(k_1a)}{J_\upsilon(k_1a)} \approx -2\upsilon^{-\frac{1}{3}}(\upsilon/k_1a)\sqrt{[1-(k_1a/\upsilon)^2/(4\zeta_1)]}\ \frac{v'(\upsilon^{\frac{2}{3}}\zeta_1)}{v(\upsilon^{\frac{2}{3}}\zeta_1)} \qquad (2\text{-}46)$$

The subscript s on υ_s has been suppressed in writing Eqs. (2-45) and (2-46). We can then replace Eq. (2-30a) by the asymptotic approximation

$$\frac{w_1'(\upsilon_s^{\frac{2}{3}}\zeta)}{w_1(\upsilon^{\frac{2}{3}}\zeta)} = \Gamma\,\frac{v'(\upsilon_s^{\frac{2}{3}}\zeta_1)}{v(\upsilon_s^{\frac{2}{3}}\zeta_1)}\left[\frac{\zeta}{\zeta_1}\,\frac{\upsilon_s^2-(k_1a)^2}{\upsilon_s^2-(ka)^2}\right]^{\frac{1}{2}} \qquad (2\text{-}47)$$

where

$$\Gamma = \frac{\mu k_1}{\mu_1 k}$$

We can replace Eq. (2-30b) by a similar equation in which the only difference is that Γ is replaced by Γ^{-1}.

Eq. (2-47) provides an example of the direction in which future work will undoubtedly be directed. In Eq. (2-38) the right hand side is a constant, whereas in Eq. (2-47) we not only have a right hand side which is variable, but it has the interesting property of involving the logarithmic derivative of the Airy function v(t). In some situations we may expect to find roots for which $\upsilon_s \approx k_1a$ and k_1a and ka are widely enough separated that we can use an asymptotic estimate for the left hand side of Eq. (2-47) and then find that the problem to be solve is

$$\frac{v'(\upsilon_s^{\frac{2}{3}}\zeta_1)}{v(\upsilon^{\frac{2}{3}}\zeta_1)} = \frac{Ai'(\upsilon^{\frac{2}{3}}\zeta_1)}{Ai(\upsilon^{\frac{2}{3}}\zeta_1)} = F(\upsilon_s, k_1a, ka, \Gamma) \qquad (2\text{-}48)$$

where $F(v_s, k_1 a, ka, \Gamma)$ may in some problems be so slowly varying that we can consider it to be a constant and therefore find that the problem to be solved is

$$v'(t) - F\, v(t) = 0 \qquad (2\text{-}49a)$$

which is equivalent to

$$w_1'[t\exp(i\tfrac{4}{3}\pi)] - F\exp(-i\tfrac{4}{3}\pi)\; w_1[t\exp(i\tfrac{4}{3}\pi)] = 0 \qquad (2\text{-}49b)$$

We want now to turn away from this more general problem and look more closely at the classical ground wave propagation problem. However, we hope that the reader sees from the discourse above that there exists many more reasons to study the roots of

$$w_1'(\tau) - q\, w_1(\tau) = 0$$

than merely the fact that this equation plays a fundamental role in the problem of propagation of radio waves around the earth's surface.

The calculations made by van der Pol and Bremmer (Ref. 6), Norton (Ref. 13), Wyngaarden and van der Pol (Ref. 9), and Johler, et. al. (Ref. 15) have all been made for specific frequencies and electrical properties of the earth. Such calculations play an important role in specific engineering applications. However, from the point of view of making calculations which can be adapted by the potential user to a class of problems, there are advantages to be had by following the procedures which were employed by Domb* (Ref. 19) and Belkina (Ref. 17).

---

*Although the reference is to a paper by Domb, the author would like to remind the reader that this work was primarily under the technical direction of M. H. L. Pryce. The calculations were made in 1942-4 by the Admiralty Computing Service and the contributors included J. C. P. Miller, L. Fox, D. H. Sadler, R. H. Corkan, P. H. Haines, R. G. Taylor, and E. M. Wilson.

In the tables of Ref. 19, it is assumed that the impedance parameter q is of the form

$$q = Q\exp(i45^{\circ})$$

where Q is real and positive. Calculations were made for the first five roots for

$$Q = 0.0(0.1)1.0 \quad , \text{ and } \quad \frac{1}{Q} = 0.0(0.1)1.0$$

In Table 2-5 we list the values that were obtained for the first root $\tau_1(q)$ by Domb and his co-workers. The "tic" marks that appear on the locus for 45° (the "Watson modes") correspond to the entries in Table 2-6.

The present author feels that there is a definite advantage in the presenting of the results in terms of dimensionless parameters in the manner employed by Pryce and Domb (Ref. 25), Pryce (Ref. 24) and Domb (Ref. 19). In order to emphasize this point, we present in Table 2-7 the values for the root $\tau_1(q)$ which were obtained by Johler, Walters and Lilley for the case of propagation over typical land ($\varepsilon_1 = 15$, $\sigma_1 = 0.005$ mho/m) and over typical sea ($\varepsilon_1 = 80$, σ_1 5 mho/m) for frequencies between 0 and 2000 kc.

Table 2-6

THE ROOT τ_1 AS CALCULATED BY ADMIRALTY COMPUTING SERVICE

Q	Real τ_1	Imag τ_1	1/Q	Real τ_1	Imag τ_1
0.0	0.509	0.882	0.0	1.169	2.025
0.1	0.604	0.862	0.1	1.240	1.953
0.2	0.698	0.850	0.2	1.311	1.877
0.3	0.790	0.848	0.3	1.383	1.790
0.4	0.880	0.854	0.4	1.449	1.685
0.5	0.967	0.868	0.5	1.497	1.559
0.6	1.050	0.891	0.6	1.510	1.420
0.7	1.128	0.921	0.7	1.486	1.293
0.8	1.201	0.958	0.8	1.439	1.189
0.9	1.267	1.002	0.9	1.383	1.110
1.0	1.326	1.051	1.0	1.326	1.051

Table 2-7

TABLE OF ROOTS $\tau_1(q)$ OBTAINED BY JOHLER, WALTERS AND LILLEY

Frequency (kc)	$\varepsilon_1 = 15$, $\sigma = 0.005$ mho/m		$\varepsilon_1 = 80$, $\sigma = 5$ mho/m	
	Real τ_1	Imag τ_1	Real τ_1	Imag τ_1
0	0.5094	0.8823	0.5094	0.8823
0.1	0.5115	0.8811	0.5095	0.8823
0.2	0.5131	0.8813	0.5095	0.8823
0.5	0.5173	0.8802	0.5096	0.8822
1.0	0.5235	0.8786	0.5098	0.8822
2	0.5346	0.8759	0.5102	0.8821
3	0.5447	0.8735	0.5105	0.8820
4	0.5542	0.8714	0.5108	0.8819
5	0.5634	0.8694	0.5111	0.8818
6	0.5722	0.8676	0.5114	0.8818
7	0.5808	0.8659	0.5117	0.8817
8	0.5892	0.8643	0.5119	0.8816
9	0.5974	0.8628	0.5122	0.8816
10	0.6055	0.8614	0.5124	0.8815
20	0.6801	0.8519	0.5148	0.8809
30	0.7477	0.8483	0.5170	0.8803
50	0.8696	0.8542	0.5210	0.8793
60	0.9254	0.8625	0.5229	0.8788
70	0.9782	0.8739	0.5248	0.8783
80	1.0281	0.8879	0.5266	0.8778
90	1.0753	0.9045	0.5284	0.8774
100	1.1197	0.9233	0.5301	0.8770
200	1.4111	1.1944	0.5463	0.8731
300	1.4628	1.4552	0.5612	0.8698
400	1.4573	1.6096	0.5752	0.8669
500	1.4173	1.6992	0.5886	0.8644
600	1.3835	1.7538	0.6016	0.8620
700	1.3569	1.7906	0.6142	0.8600
800	1.3361	1.8171	0.6264	0.8581
900	1.3193	1.8373	0.6384	0.8564
1000	1.3055	1.8531	0.6502	0.8549
1100	1.2940	1.8660	0.6618	0.8536
1200	1.2843	1.8766	0.6732	0.8524
1300	1.2759	1.8856	0.6843	0.8513
1400	1.2687	1.8933	0.6954	0.8504
1500	1.2623	1.8999	0.7062	0.8497
1600	1.2566	1.9057	0.7170	0.8490
1700	1.2516	1.9108	0.7276	0.8485
1800	0.2471	1.9154	0.7380	0.8481
1900	1.2430	1.9195	0.7484	0.8479
2000	1.2393	1.9232	0.7586	0.8477




Fig. 2-5 The Root Locus of $\tau_1[Q\exp(i145^\circ)]$ for $0 < Q < 0.4$

In Fig. 2-5 we have presented (in strip form) a greatly magnified version of the root locus for arg(q) = 45° which has been illustrated in Figs. 2-1 and 2-2. Over this range it is not possible to distinguish between this locus and the locus of points which are obtained from the tables of Ref. 15* for frequencies up to

*Ref. 15 can be used to extend Table 2-6 to 10,000 kc.

50 kc over "typical land" and up to 3000 kc over "typical sea." The portion of the locus which is given in this figure is an interesting region because it contains the "minimum" in the imaginary part of $\tau_1[Q\exp(i145^\circ)]$ which occurs between $Q = 0.25$ and 0.30. This occurs for a frequency of the order of 30 kc over "typical land" and of the order of 2000 kc over "typical sea."

The data given in Ref. 15 reveals that when the frequency is 10,000 kc that the corresponding value of Q is of the order of unity in the case of "typical sea." If one plots the data of Table 2-6 for the case of "typical land" on a scale such as employed in Fig. 2-5, it will be observed that as the frequency continues to increase that the locus pulls slightly away from the locus for $\arg(q) = 45^\circ$. The reader who is interested in the relation of the impedance parameter to the frequency will find tables in Ref. 15 for quantities denoted by K_e and ψ_e. The relation to q is

$$q = \frac{1}{2^{\frac{1}{3}}K_e} \exp[i(\frac{\pi}{4} + \psi_e)]$$

In Fig. 2-6 we have plotted several sets of data which are related to the behavior of the root locus for high frequencies when $\tau_1(q)$ approaches $\tau_1^\infty = 1.169 + i2.025$ as the magnitude of q tends to infinity. The lowest set of points (labeled NBS DATA) is taken from Ref. 15 for the case of "typical land." The highest frequency considered was 10 Mc. The highest set of data in this figure is the root locus for $\arg(q) = 45^\circ$. Over the range covered by this illustration these two loci are virtually parallel. The reader will observe that the magnitude of q on the locus based on the data of Ref. 15 is of the order of 10 at a frequency of 3 Mc.

The C.C.I.R. atlas of ground wave curves which was prepared by Wyngaarden and van der Pol does not overlap with the data of Ref. 15 since the frequency range is taken to be that from 30 Mc to 300 Mc. Furthermore, the electrical parameters are not the




Fig. 2-6 Loci of Roots in Vicinity of $\tau_1^\infty = 1.169 + i2.025$ for Propagation Around the Earth's Surface

same in the two sets of calculations. The parameters employed were as indicated below.

Source	Johler, Walters and Lilley Ref. 15	Wyngaarden and van der Pol Ref. 9
"Typical Land"	$\varepsilon_1 = 15$ $\sigma_1 = 0.005$ mho/m	$\varepsilon_1 = 10$ $\sigma_1 = 0.001$ mho/m
"Typical Sea"	$\varepsilon_1 = 80$ $\sigma_1 = 5$ mho/m	$\varepsilon_1 = 80$ $\sigma_1 = 4$ mho/m

The curve for "typical sea" in Fig. 2-6 is virtually parallel over to the curve for arg(q) = 45°. The frequency has not been taken to be large enough for us to see how the locus of the roots for propagation over the "typical sea" approaches the limiting point at τ_1^∞. However, the curve for "typical land" in the C.C.I.R. atlas do show the manner in which that locus turns and heads into the limiting point along the locus for arg(q) = 90°. The atlas prepared by the Radio Research Laboratory (Ref. 23) covers the frequency range from 30 to 10,000 Mc , but the roots which were employed to make the calculations have not been listed in the report.

Except for the "tic" mark which is used to mark the location of the root for $q = 0.1\exp(i45^\circ)$, all the "tic" marks on the locus for arg(q) = 45° are taken from the data published by Belkina (ref. 17). The method employed by Belkina deserves further study because it is perhaps the most useful method of presenting data which has been developed to date. In Eq. (2-39) we have given expressions for the impedance parameters for vertical and horizontal polarization. They depend upon a, ω, ε_1, μ_1, and σ_1. In the ground wave problem the radius a is fixed (although it also

varies if we take into account the fact that the concept of an equivalent earth's radius is employed to take into account the rate of change of the index of refraction at the earth's surface.) but we still have the three variables ω, ε_1, and σ_1 even in those cases for which $\mu_1 = 1$. Following a suggestion make by Fock, Belkina used the approximation

$$\sqrt{1 - (k/k_1)^2} \approx \frac{1}{\sqrt{1 + (k/k_1)^2}}$$

which is valid when the magnitude of (k/k_1) is small. Belkina then observed the manner in which the frequency ω enters the expression for q_v of Eq. (2-39a) and assumed that q_v could be represented by an expression of the form

$$q = \frac{in^{5/6}}{\sqrt{1 + \alpha n}} \qquad (2\text{-}50)$$

This expression has the advantage of containing only two parameters and therefore it is possible to conceive of making computations for a range of values of α and n which will serve to cover a range of values of ω, ε_1, and σ_1. This is a very clever transformation since we have two adjustable parameters at our disposal in Eq. (2-50) and the physical problem consists of two basic variables, the wavelength and the physical nature of the earth's crust over which the wave is being propagated. The fact that the properties of the earth's crust requires two real parameters, ε_1 and σ_1, for its description forces us to consider the diffraction phenomena to depend upon at least three variables (three only if we assume μ_1 to be constant and equal to the value of μ in free space). The advantage of the form for q given in Eq. (2-50) is that we have only 2 parameters to vary and hence it becomes feasible to attempt to compute some universal data which can be used in a variety of physical problems. In Table 2-8a we have reproduced Belkina's table which shows the combination of

Table 2-8a

TABLE OF VALUES OF THE PARAMETER log n SHOWING DEPENDENCE UPON PHYSICAL NATURE OF THE SOIL OR SEA AND ON THE WAVELENGTH λ

Physical Nature of Soil	Very Salty Sea Water	Slightly Salty Sea Water	Slightly Salty Sea Water	Marsh	Damp Soil and Meadow Soil	Damp Soil and Meadow Soil
ε	80	80	80	15	15	15
$\frac{\sigma_0}{\sigma}$	2.10^5	$1,10^6$	2.10^6	1.10^7	1.10^8	2.10^8
α	0,010	0,018	0,024	0,010	0,023	0,031
λ (meters)	log n					
10	0,425	0,844	1,021	1,44	2,04	2,22
20	0,124	0,543	0,720	1,143	1,744	1,924
30	−0,052	0,367	0,544	0,967	1,568	1,748
40	−0,177	0,242	0,419	0,842	1,443	1,623
50	−0,274	0,145	0,322	0,745	1,346	1,526
60	−0,353	0,066	0,243	0,666	1,267	1,447
70	−0,420	−0,001	0,176	0,599	1,200	1,380
80	−0,478	−0,059	0,118	0,541	1,142	1,322
90	−0,520	−0,110	0,067	0,490	1,091	1,271
100	−0,575	−0,156	0,021	0,444	1,045	1,225
200	−0,876	−0,457	−0,280	0,143	0,744	0,924
300	−1,052	−0,633	−0,456	−0,033	0,568	0,748
400	−1,177	−0,758	−0,581	−0,158	0,443	0,623
500	−1,274	−0,855	−0,678	−0,255	0,346	0,526
600	−1,353	−0,934	−0,757	−0,334	0,267	0,447
700	−1,420	−1,001	−0,824	−0,401	0,200	0,380
800	−1,478	−1,059	−0,882	−0,459	0,142	0,322
900	−1,529	−1,110	−0,933	−0,510	0,091	0,271
1000	−1,575	−1,156	−0,979	−0,556	0,045	0,225
1100	−1,616	−1,198	−1,020	−0,597	0,004	0,184
1200	−1,654	−1,235	−1,058	−0,635	−0,034	0,146
1300	−1,689	−1,270	−1,093	−0,670	−0,069	0,111
1400	−1,721	−1,302	−1,125	−0,702	−0,101	0,079
1500	−1,751	−1,332	−1,155	−0,732	−0,131	0,049
1600	−1,779	−1,360	−1,183	−0,760	−0,159	0,021
1700	−1,806	−1,386	−1,209	−0,786	−0,185	−0,005
1800	−1,830	−1,411	−1,234	−0,811	−0,210	−0,030
1900	−1,854	−1,435	−1,258	−0,835	−0,234	−0,054
2000	−1,876	−1,457	−1,280	−0,857	−0,256	−0,076

Note: Belkina expressed the conductivity of the earth in the form σ_0/σ where σ is the conductivity of the earth and σ_0 is the conductivity of mercury which was taken to be 10440 $(\Omega^{\circ}cm)^{-1}$.

Table 2-8b

THE FUNCTION $q(\alpha,n) = \frac{ln^{\frac{5}{6}}}{\sqrt{1+\alpha n}}$

lg$n$	$\alpha = 0{,}00$	$\alpha = 0{,}01$	
	Req = Imq	Req	Imq
−1,0	0,1037891	0,1037371	0,1038409
−0,9	0,1257434	,1256642	0,1258225
−0,8	0,1523415	0,1522206	0,1524621
−0,7	0,1845660	0,1843816	0,1847498
−0,6	0,223607	0,223325	0,223887
−0,5	0.270906	0,270476	0,271333
−0,4	0,328210	0,327555	0,328861
−0,3	0,397635	0,396635	0,398628
−0,2	0,481746	0,480219	0,483259
−0,1	0,583649	0,581317	0,585953
0	0,707107	0,703545	0,710616
0,1	0,856679	0,851236	0,862020
0,2	1,037891	1,029569	1,046016
0,3	1,257434	1,244705	1,269788
0,4	1,523415	1,503929	1,542180
0,5	1,845660	1,815804	1,874132
0,6	2,23607	2,19027	2,27921
0,7	2,70906	2,63873	2,77429
0,8	3,28210	3,17393	3,38050
0,9	3,97635	3,80968	4,12430
1,0	4,81746	4,56015	5,03891
1,1	5,83649	5,43841	6,16599
1,2	7,07107	6,45394	7,55738
1,3	8,56679	7,60847	9,27653
1,4	10,37891	8,88952	11,39862
1,5	12,57434	10,26164	14,00751
1,6	15,23415	11,65625	17,18642
1,7	18,45660	12,9648	20,9998
1,8	22,3607	14,0434	25,4659
1,9	27,0906	14,7388	30,5303
2,0	32,8210	14,9365	36,0599
2,1	39,7635	14,6074	41,8749
2,2	48,1746	13,8223	47,8098
2,3	58,3649	12,7195	53,7664
2,4	70,7107	11,4522	59,7293
2,5	85,6679	10,1478	65,7465

Table 2-8b (Cont'd)

THE FUNCTION $q(\alpha,n) = \frac{in^{\frac{5}{6}}}{\sqrt{1+\alpha n}}$

lg n	α = 0.02		α = 0,03	
	Re q	Im q	Re q	Im q
-1,0	0,1036851	0,1038927	0,1036330	0,1039444
-0,9	0,1255848	0,1259014	0,1255018	0,1259836
-0,8	0,1520995	0,1525824	0,1519780	0,1527024
-0,7	0,1841967	0,1849331	0,1840111	0,1851159
-0,6	0,223043	0,224166	0,222760	0,224444
-0,5	0,270045	0,271758	0,269612	0,272182
-0,4	0,326896	0,329509	0,326232	0,330152
-0,3	0,395628	0,399613	0,394613	0,400591
-0,2	0,478678	0,484757	0,477123	0,486240
-0,1	0,578958	0,588229	0,576573	0,590476
0	0,699931	0,714070	0,696268	0,717469
0,1	0,845695	0,867256	0,840058	0,872384
0,2	1,021061	1,053939	1,012373	1,061652
0,3	1,231620	1,281748	1,218199	1,293298
0,4	1,483770	1,560182	1,462987	1,577381
0,5	1,784680	1,901119	1,752413	1,926529
0,6	2,14211	2,31944	2,09188	2,35658
0,7	2,56400	2,83385	2,48563	2,88730
0,8	3,05767	3,46777	2,93526	3,54298
0,9	3,62843	4,25036	3,43735	4,35271
1,0	4,27717	5,21731	3,98016	5,34946
1,1	4,99680	6,41087	4,53962	6,56711
1,2	5,76723	7,87812	5,07600	8,03403
1,3	6,54979	9,66574	5,53436	9,76282
1,4	7,28329	11,80968	5,85280	11,73901
1,5	7,88683	14,31989	5,98001	13,91601
1,6	8,27453	17,16533	5,89537	16,22415
1,7	8,3824	20,2709	5,61860	18,59365
1,8	8,1948	23,5356	5,2007	20,9779
1,9	7,7519	26,8668	4,7034	23,3633
2,0	7,1316	30,2098	4,1810	25,7645
2,1	6,4198	33,5565	3,6717	28,2121
2,2	5,6878	36,9341	3,1979	30,7423
2,3	4,9835	40,3885	2,7700	33,3909
2,4	4,3338	43,9710	2,3908	36,1908
2,5	3,7502	47,7310	2,0588	39,1718

α and n which are required to describe the variables ε_1, σ_1 and λ which enter into certain diffraction problems. In Table 2-8b we reproduce the table given by Belkina which presents the values of q which result from use of the combinations of α and n which are contained in the tables. The reader will observe that negative values of log n (i.e., $n < 1$) lead to combinations of the real and imaginary parts of q which are almost equal and therefore the argument of q is of the order of 45°. However, for large positive values of n (and $\alpha > 0$), we see that the imaginary part of q grows much faster than does the real part. Therefore, these data reveal that the family of root loci which are based upon Eq. (2-50) will parallel the locus $\arg(q) = 45^\circ$ for small values of n, but then will pull away from this locus and turn to approach τ_1^∞ along a line which is parallel to the imaginary axis.

Diffraction theory needs the ingenuity and inventiveness which leads to the recognition of the importance of models such as that contained in Eq. (2-50). Since the electrical properties to be assigned to an obstacle are often not known exactly, it would be a great advantage to have the calculations which are published have their basis in a relationship similar to that of Eq. (2-50) which will permit the construction of a family of <u>universal</u> curves which can be relatively easily adapted to a wide variety of approximate physical models.

The importance of seeking universal curves is clearly indicated by the C.C.I.R. atlas (Ref. 9). It is not possible to plot the roots for horizontal polarization on the graph given in Fig. 2-5 because they are all closely clustered around τ_1^∞. This suggests that the approximations suggested in Appendix E (see Eq. (E-7) and (E-8)) could be used to obtain the data for horizontal polarization from the universal functions computed for $q = \infty$. The importance of these concepts was apparently appreciated by Pryce and Domb (Ref. 25), but later writers have apparently not been

influenced by this relatively unknown work which was carried out in England during World War II.

The papers of Norton (Ref. 13) and Pryce and Domb (Ref. 25) are seldom referred to by recent writers. However, these papers are a "must" on the reading lists of those who undertake numerical work in diffraction theory. These men were not writing a paper just to add another paper to their list of publications; they were writing to try to guide engineers towards the obtaining of numerical results for the practical problems that they might encounter.

Our discussion of the root loci for the ground wave propagation problems have led us away from our development of numerical methods for solving for the roots and associated functions. However, we are entering an era in which we can expect an increase of interest in diffraction phenomena. Even if scores of atlases are prepared which will illustrate the diffraction phenomena associated with dielectric spheres and circular cylinders, the chances are quite large that the engineer who wants an answer will have difficulty in interpolating when the exisiting data have been obtained for a set of values of the electrical parameters, the radius a, and the wavelength. Even with the ability to employ the electronic computer to make calculations, it is still worth keeping in mind that if one can find a means of computing a "universal" function that one might not need to get back on the computer to make a completely new computation when one is faced with the problem of changing the parameters of the problem.

In Appendix C we have given some coefficients which will enable the user to calculate (on a desk calculator) the roots for arg(q) in the vicinity of 45° and for arg(q) in the vicinity of 0°. These two cases appear to be the ones in which there is the most interest at present.

2.4 Numerical Values for the Classical Expansions

The expansions which are designated as Eq. (2-21) and Eq. (2-22) have been employed in the papers of many writers since the first several terms were given in the classic papers of van der Pol and Bremmer (Ref. 6). Norton (Ref. 13) gave the first several terms with numerical values for the coefficients. Howe (Ref. 14) discussed the obtaining of numerical values of A_n and B_n on an electronic computer. Johler, Walters, and Lilley (Ref. 15) have given (in a slightly different notation) the explicit dependence of the coefficients upon τ for terms up to and including $A_{10}(\tau)$ and $B_{11}(\tau)$.

If one wants specific numerical values for the $A_n(\tau)$ and $B_n(\tau)$, the use of the explicit forms for these coefficients as a function of τ is to be discouraged since the coefficients (as has been shown by Howe(Ref. 14) and by a group at Leningrad University (Ref. 16)) can be readily obtained by means of recursion formulae. For $n > 2$, the recursion formula for the A_n is

$$nA_oA_n = -[(n-1)A_1A_{n-1} + (n-2)(A_2 - 1)A_{n-2} + (n-3)A_3A_{n-3} + \cdots + 2A_{n-2}A_2 + A_{n-1}A_1]$$

$$= (n-2)A_{n-2} - \sum_{m=1}^{n-1} mA_mA_{n-m} \qquad (2\text{-}51)$$

where $A_o = \beta_s \exp(i\frac{1}{3}\pi)$ and $A_oA_1 = 1$. For $n > 3$, the recursion formula for the B_n is

$$nB_n = B_1B_{n-3} + 2B_2B_{n-4} + \cdots + (n-2)B_{n-2}B_1 + (n-1)B_{n-1}B_o$$

$$= \sum_{m=1}^{n-2} mB_mB_{n-m-2} \qquad (2\text{-}52)$$

where $B_o = \alpha_s \exp(i\frac{1}{3}\pi)$, $B_1 = 1$ and $B_2 = 0$.

Since the $A_n(\tau)$ and $B_n(\tau)$ are complex quantities, the calculations based upon Eqs. (2-51) and (2-52) are somewhat more involved than one might suspect. However, one can readily see that each of these quantities must be of the form of a real number times the factor $\exp(i\frac{1}{3}m\pi)$, where m is an integer. This is quite obvious since each of the explicit forms is representable as an integral power of τ multiplied times a polynomial in $z = \tau^3$ in which the coefficients are real. Since $\tau = \alpha_s \exp(i\frac{1}{3}\pi)$ or $\tau = \beta_s \exp(i\frac{1}{3}\pi)$, $\tau^3 = -\alpha_s^3$ or $\tau^3 = -\beta_s^3$, and hence the polynomial is real. If one "plays around" with the explicit forms for a while, it becomes clear that a suitable set of real coefficients can be defined by means of the relations

$$C_n(\beta) = A_n(\tau)\exp(-i\tfrac{1}{3}\pi)\exp(i\tfrac{2}{3}n\pi) \quad , \quad \tau = \beta\exp(i\tfrac{1}{3}\pi) \tag{2-53}$$

$$D_n(\alpha) = B_n(\tau)\exp(-i\tfrac{1}{3}\pi)\exp(-i\tfrac{2}{3}n\pi) \quad , \quad \tau = \alpha\exp(i\tfrac{1}{3}\pi) \tag{2-54}$$

Eqs. (2-21) and (2-22) then take the forms

$$\tau_s(q) = \left[\beta_s + \sum_{n=1}^{\infty} C_n(\beta_s) q^n \exp(-i\tfrac{2}{3}n\pi) \right] \exp(i\tfrac{1}{3}\pi) \tag{2-55}$$

$$\tau_s(q) = \left[\alpha_s + \sum_{n=1}^{\infty} D_n(\alpha_s) q^{-n} \exp(i\tfrac{2}{3}n\pi) \right] \exp(i\tfrac{1}{3}\pi) \tag{2-56}$$

The expressions in the brackets [...] becomes purely real when $q = Q\exp(i\frac{2}{3}\pi)$ or $q = Q\exp(-i\frac{1}{3}\pi)$, where Q is real and positive. In this case the roots $\tau_s(q)$ lies along the ray $\arg(\tau) = \frac{1}{3}\pi$ and we observe that this is illustrated by the root loci depicted in Fig. 2-2. In Table 2-9 we present values for the coefficients C_n for use in Eq. (2-55) for s = 1(1)5. Similar tables of the coefficients D_n for use in Eq. (2-56) are given in Table 2-10.

(Text continues on p. 2-64)

Table 2-9

COEFFICIENTS IN THE EXPANSION OF $\tau_s(q)$ IN SERIES OF POWERS OF q

s = 1

n	C_n		n	C_n	
0	0.10187927916474710890 2E	1	40	-.43450039839476856133 5E	-12
1	0.98155368934566564334 97E	+0	41	-0.59936583720035197224 8E	-12
2	-0.47283779525397403410 4E	+0	42	.25776260002643435788 4E	-12
3	.13440524531392944729 6E	+0	43	.13277983638019863636 3E	-12
4	-.71602938859209580269E	-2	44	-.10744291383610046579 5E	-12
5	-.98774282120637696024 7E	-2	45	-.19142228801387823900 6E	-13
6	.20128847168449974735 6E	-2	46	.37439079134687626594 5E	-13
7	.13462118035028363782 9E	-2	47	-.17666437483341173401 8E	-14
8	-.56668183510136790690 5E	-3	48	-.11337107645703595881 4E	-13
9	-.19194548486765931239 5E	-3	49	.29013724000544148587 5E	-14
10	.15557115743545349859 6E	-3	50	.29519485284101323658 6E	-14
11	.18916689089679143143 8E	-4	51	-.15230115342664931307 0E	-14
12	-.40870251099793881790 8E	-4	52	-.61133877896127375017 7E	-15
13	.24179084285559739402 2E	-5	53	.60615193090988927859 2E	-15
14	.10054412469337743780 2E	-4	54	.66506395408094948795 1E	-16
15	-.24604427443158134182 1E	-5	55	-.20495253271518981323 8E	-15
16	-.22325772962695294382 9E	-5	56	.21678887050098105155 0E	-16
17	.10845740096401966706 7E	-5	57	.60237154078484583664 7E	-16
18	.40970966684901685698 3E	-6	58	-.19511946278463794373 8E	-16
19	-.38009141647458677677 1E	-6	59	-.15019459909324023156 2E	-16
20	-.42322225242223595775 E	-7	60	.92845571586849734978 1E	-17
21	.11617517415717004888 6E	-6	61	.28244522308929290762 6E	-17
22	-.10491804636115500769 6E	-7	62	-.35367534944795050297 1E	-17
23	-.31482430296256068517 3E	-7	63	-.15803605765732402167 0E	-18
24	.92452961606875395493 4E	-8	64	.11573629847955301414 0E	-17
25	.73792846770156102050 2E	-8	65	-.19181766556719008652 6E	-18
26	-.41842339811879769425 8E	-8	66	-.32833642943736497918 3E	-18
27	-.13477462039333872944 0E	-8	67	.12985412991371753757 9E	-18
28	.15124611057458781508 1E	-8	68	.77486418714831781100 8E	-19
29	.98706365158131984016 6E	-10	69	-.57664810893608478424 5E	-19
30	-.47296345270391202255 7E	-9	70	-.12651177866742399956 9E	-19
31	.65725283233185074546 7E	-10	71	.21074349462059793418 5E	-19
32	.12943890327127395662 9E	-9	72	-.37644689028468273486 0E	-21
33	-.46134257672511148689 3E	-10	73	-.66616749277340296364 2E	-20
34	-.29941065860438752794 6E	-10	74	.15187711426620977814 4E	-20
35	.20144823552162073834 2E	-10	75	.18143181729565122691 5E	-20
36	.50372341365009438348 1E	-11	76	-.86768671431 6634096336E	-21
37	-.71968806609140234318 7E	-11	77	-.39931899469301251568 9E	-21
38	-.73940983916856290706 7E	-13	78	.36222565456751322006 9E	-21
39	.22291110964115985238 3E	-11	79	.51994725285181650888 5E	-22
40	-.43450039839476856133 5E	-12	80	-.12723204976377352460 8E	-21

Table 2-9 (Cont'd)

COEFFICIENTS IN THE EXPANSION OF $\tau_s(q)$ IN SERIES OF POWERS OF q

s = 2			s = 3		
n	C_n		n	C_n	
0	.32481975821798365378 8E	1	0	.48200992111787356394 3E	1
1	.30786304548903360888 42E	+0	1	.20746460937584188666 8E	+0
2	-0.14589576585143720362 E	-1	2	-0.44648006459751916 58 E	-2
3	-.30210423227166980947 6E	-1	3	-.14155016044432003300 7E	-1
4	.50763640151679428451 5E	-2	4	.10703302461047620853 4E	-2
5	.49635748316817315445 1E	-2	5	.17028186134496154083 7E	-2
6	-.16300168451047944796 4E	-2	6	-.24112190919635475490 1E	-3
7	-.86750051743351010992 51E	-3	7	-.23724023289804516065 5E	-3
8	.49346434431170064276 9E	-3	8	.52387810540994268604 9E	-4
9	.13413787220397524336 2E	-3	9	.34720586456148142539 7E	-4
10	-.14134597983144194099 0E	-3	10	-.11084729320666862019 1E	-4
11	-.11352364802041992223 8E	-4	11	-.50926297101976206695 8E	-5
12	.38096788303774829557 5E	-4	12	.22934443200343523137 6E	-5
13	-.34181462922891765561 7E	-5	13	.71992799024589129877 9E	-6
14	-.95158669079838683172 1E	-5	14	-.46467583692225332216 22E	-6
15	.25857311472755995240 3E	-5	15	-.92680510376643926992 1E	-7
16	.21289805242624463328 E	-5	16	.92185733199627310209 0E	-7
17	-.10977718809939555909 5E	-5	17	.93997744918046565562 4E	-8
18	-.39007077117699433980 0E	-6	18	-.17883258401771898438 2E	-7
19	.38075054993374097804 2E	-6	19	-.22686619297749298390 2E	-9
20	.38669222760589291073 5E	-7	20	.33834042741925203744 7E	-8
21	-.11596761196786855781 9E	-6	21	-.25398784984426853681 9E	-9
22	.11154549201012280362 8E	-7	22	-.62157569401023839860 4E	-9
23	.31382435243162237173 3E	-7	23	.10437450820500473125 2E	-9
24	-.93616392044720597682 1E	-8	24	.11010360320065704125 0E	-9
25	-.73493345169140300170 6E	-8	25	-.30232657019321717459 4E	-10
26	.42037524229676693554 E	-8	26	-.18581810957867060172 4E	-10
27	.13401164996597343768 8E	-8	27	.76167545013724096019 8E	-11
28	-.15155222680911970800 2E	-8	28	.29222052257793486015 7E	-11
29	-.96925889471678985495 5E	-10	29	-.17702755814782993572 6E	-11
30	.47339165828588738302 1E	-9	30	-.40788806167839688454 9E	-12
31	-.66116692746948640618 4E	-10	31	.38864386269919582222 4E	-12
32	-.12948544598639486453 58E	-9	32	.43624738602138088875 1E	-13
33	.46216299446958825093 9E	-10	33	-.81442705578216297027 7E	-13
34	.29942297972273218868 2E	-10	34	-.82244987486714947417 0E	-15
35	-.20161293566433576504 3E	-10	35	.16352934126658916424 E	-13
36	-.50358612789976318381 1E	-11	36	-.14286831002285407985 E	-14
37	.72000456489099601150 2E	-11	37	-.31434612748914996969 2E	-14
38	.73352357654112349643 5E	-13	38	.59592701806484259270 4E	-15
39	-.22296900968651322241 8E	-11	39	.57520552222222623802 2E	-15
40	.43467669684689957920 7E	-12	40	-.17719622650538895533 7E	-15

Table 2-9 (Cont'd)

COEFFICIENTS IN THE EXPANSION OF $\tau_s(q)$ IN SERIES OF POWERS OF q

n	s = 4 C_n		n	s = 5 C_n	
0	.6163307355639486547 64E	1	0	.7372177255047770177 09E	1
1	.1622505486579361020 33E	+0	1	.1356451378478853765 00E	+0
2	-0.2135642360567450681 E	-2	2	-0.1247908371246331331 E	-0
3	-.8718858881084836369 03E	-2	3	-.6110240121452966726 06E	-2
4	.4024106198428258762 27E	-3	4	.1969567281545797851 04E	-3
5	.8351690615791607382 71E	-3	5	.4926363955840293871 24E	-3
6	-.7154117172979184632 30E	-4	6	-.2937394787911048069 02E	-4
7	-.9404845776049475486 60E	-4	7	-.1694413557027894801 74E	-4
8	.1232508066944196747 55E	-4	8	.4252784253958900345 99E	-5
9	.1135564936006185256 38E	-4	9	.4828986444343769560 19E	-5
10	-.2079429461380929992 31E	-5	10	-.6042107438535025789 62E	-6
11	-.1415520831483895940 45E	-5	11	-.5172559235493433021 18E	-6
12	.3452889380803864287 85E	-6	12	.8468079518451075875 19E	-7
13	.1783101535764832298 11E	-6	13	.5662222483073771682 68E	-7
14	-.5657431681692507930 20E	-7	14	-.1174087860691555187 48E	-7
15	-.2233980237859649832 81E	-7	15	-.6260145124692711725 30E	-8
16	.9158646136902640579 54E	-8	16	.1613063897018287038 52E	-8
17	.2740038333227925815 55E	-8	17	.6929211994711748138 63E	-9
18	-.1465809412640893655 25E	-8	18	-.2198170728134133951 25E	-9
19	-.3221567669461565744 82E	-9	19	-.7618796038025403926 93E	-10
20	.2319548368965501633 20E	-9	20	.2972854857540833475 92E	-10
21	.3498761850779111308 26E	-10	21	.8251671403662615959 29E	-11
22	-.3628149290590101300 47E	-10	22	-.3991269701448401812 72E	-11
23	-.3208160106368888866 67E	-11	23	-.8708534773077954989 49E	-12
24	.5605857491059785532 31E	-11	24	.5319945667309152100 03E	-12
25	.1661746372564530083 39E	-12	25	.8807901367759149338 86E	-13
26	-.8547217250441557604 28E	-12	26	-.7039202569532885351 90E	-13
27	.2385646492056084512 32E	-13	27	-.8279317502824639157 06E	-14
28	.1284037878020003967 11E	-12	28	.9243918209187596221 75E	-14
29	-.1111246255262674368 76E	-13	29	.6727619270616150993 81E	-15
30	-.1896570557354036212 70E	-13	30	-.1204304237074091145 42E	-14
31	.2822629235725341732 24E	-14	31	-.3585591427304927009 89E	-16
32	.2745763601661127777 14E	-14	32	.1555692259593426602 92E	-15
33	-.5988799779211708304 55E	-15	33	-.1948059751686161736 96E	-17
34	-.3878784164286995939 68E	-15	34	-.1991105129560054525 74E	-16
35	.1161122431301609901 10E	-15	35	.1098893161531685599 30E	-17
36	.5309325272565802138 81E	-16	36	.2522420722403347332 27E	-17
37	-.2128350137349513667 95E	-16	37	-.2506916832385049489 22E	-18
38	-.6961437966330455999 63E	-17	38	-.3158809006496259202 02E	-18
39	.3748708667244772137 62E	-17	39	.4626557643375356595 03E	-19
40	.8561361834993754990 07E	-18	40	.3903436297271893691 35E	-19

Table 2-10

COEFFICIENTS IN THE EXPANSION OF $\tau_s(q)$ IN SERIES OF POWERS OF $\frac{1}{q}$

$s = 1$

n	D_n		n	D_n	
0	0.233810741045976703849E	1	40	-.700691631307825915940E	7
1	-0.100000000000000000000E	1	41	-0.172926385120568210974E	8
2	0.000000000000000000000E	0	42	.302402694909812084630E	8
3	.779369136819922346163E	+0	43	.296000071565089620851E	8
4	-.250000000000000000000E	+0	44	-.102975125398930052371E	9
5	-.109334925256937550787E	1	45	-.191061717492799559388E	8
6	.909263992956576070523E	+0	46	.300923950953512690432E	9
7	.164740570689507454422E	1	47	-.133232366387904713760E	9
8	-.264226069370932411401E	1	48	-.764222912713176726357E	9
9	-.213710405546052183249E	1	49	.850994329133701780251E	9
10	.691693414275288923156E	1	50	.162675880891976768802E	10
11	.112297788720997886992E	1	51	-.341788385744768603123E	10
12	-.166366387432300987771E	2	52	-.247043062983368563710E	10
13	.656326185545065316075E	1	53	.112194546138271726615E	11
14	.363549113840041841541E	2	54	-.815317134722237362844E	6
15	-.366549831925517812206E	2	55	-.318350972974930095218E	11
16	-.690160345205570526425E	2	56	.198978249303328144756AE	11
17	.131615045605843940331E	3	57	.780657206146936957945E	11
18	.981386219147608638941E	2	58	-.105461120000807422780E	12
19	-.393408408913159305757E	3	59	-.156329720659350104282E	12
20	-.213671996252882455892E	2	60	.398523735722135716625E	12
21	.103281515436985573212E	4	61	.196308040671876342016E	12
22	-.557005440585398217545E	3	62	-.126091466443172167816E	13
23	-.238101946411483180501E	4	63	.229775799379091896537E	12
24	.289471314674802657161E	4	64	.346149996770127167665E	13
25	.458512006481859752533E	4	65	-.283001749930191764780E	13
26	-.104802124890582268055E	5	66	-.813564326005073585241E	13
27	-.596169025865546235994E	4	67	.132197256989378198077E	14
28	.317985515060398198593E	5	68	.150225118847528990443E	14
29	-.318078127520901200706E	4	69	-.473745720389461439785E	14
30	-.842338171333703320929E	5	70	-.133433575875402257475E	14
31	.599334412880064201119E	5	71	.144494564473025958993E	15
32	.193151282158636859046E	6	72	-.525601236919508639310E	14
33	-.281949289996349905166E	6	73	-.382518041958423898398E	15
34	-.357114653412512609012E	6	74	.392093656248619176020E	15
35	.994137337829276568794E	6	75	.854485756965898248908E	15
36	.373873844573088761079E	6	76	-.166790003306127340751E	16
37	-.297606709730805582319E	7	77	-.141193967004010822819E	16
38	.797089484675489713151E	6	78	.569998253336990313415E	16
39	.776451218363028956443E	7	79	.479966949271702811900E	15
40	-.700691631307825915940E	7	80	-.167565476382404105172E	17

Table 2-10 (Cont'd)

COEFFICIENTS IN THE EXPANSION OF $\tau_s(q)$ IN SERIES OF POWERS OF $\frac{1}{q}$

n	S = 2 D_n		n	S = 3 D_n	
0	0.40879494441309706166E	1	0	0.55205598280955510591 3E	1
1	-0.10000000000000000000E	1	1	-0.10000000000000000000E	1
2	0.00000000000000000000E	0	2	0.00000000000000000000E	0
3	.13626498147103235 3888E	1	3	.18401866093651836863 7E	1
4	-.25000000000000000000E	+0	4	-.25000000000000000000E	+0
5	-.33422661315541423310 0E	1	5	-.60953161631164760522 1E	1
6	.15897581171620441286 9E	1	6	.21468843775927143007 7E	1
7	.95807249818746611815 3E	1	7	.23856826821173090638 4E	2
8	-.80771431512558439665 7E	1	8	-.14730347394198150459 5E	2
9	-.28960632394700688655 6E	2	9	-.10040808457934071911 4E	3
10	.37698212929673685384 2E	2	10	.93089488066151591677 3E	2
11	.87799588209547027215 5E	2	11	.43699417676693186920 3E	3
12	-.16776010502034424762 E	3	12	-.56354575378613821636 E	3
13	-.25596760649324786660 9E	3	13	-.19234822331387866517 9E	4
14	.72153942689197438625 0E	3	14	.33190967984033483649 1E	4
15	.67408112320871534234 1E	3	15	.64220456394637625830 3E	4
16	-.30149423740187542736 3E	4	16	-.19156026667346727872 2E	5
17	-.13536169456412941944 1E	4	17	-.36083404452007356589 1E	5
18	.12252648528028400080 9E	5	18	.10873191889309948424 4E	6
19	.15747671668186029040 9E	3	19	.14797467397614144215 1E	6
20	-.48352355261314256564 4E	5	20	-.60806001575549044 0192E	6
21	.20006478266582259760 9E	5	21	-.55844264531695755255 1E	6
22	.18449600271980195677 6E	6	22	.33525081259921336365 9E	7
23	-.16021114591821377490 9E	6	23	.17586350760320354211 9E	7
24	-.67527140633483645861 8E	6	24	-.18221760084114102297 0E	8
25	.94582100656328777389 8E	6	25	-.28707072019486256444 9E	7
26	.23365860889822990708 2E	7	26	.97563861320080920421 4E	8
27	-.49049438891386544004 9E	7	27	-.19513003891671341399 3E	8
28	-.74256351139623963902 1E	7	28	-.51387383573281713114 1E	9
29	.23541732926286161632 1E	8	29	.28988255472696043906 9E	9
30	.20206656962773089103 8E	8	30	.26566154189056143660 2E	10
31	-.10681916879758598583 1E	9	31	-.25304551104220958585 2E	10
32	-.36083199954860278271 2E	8	32	-.13435237959928836107 6E	11
33	.46242386495158434072 5E	9	33	.18652896164343077413 5E	11
34	-.58135210651448059582 8E	8	34	.66126465733360568438 6E	11
35	-.19149945190416935844 7E	10	35	-.12608526886612937893 4E	12
36	.11152316680964965437 0E	10	36	-.31416970010554998013 6E	12
37	.75687337539762131987 0E	10	37	.80691030633072560724 9E	12
38	-.81687051951502371410 1E	10	38	.14207628813947656005 7E	13
39	-.28319299567073987733 0E	11	39	-.49648672851710567965 4E	13
40	.47636693050378431222 0E	11	40	-.59527643107747323145 1E	13

Table 2-10 (Cont'd)

COEFFICIENTS IN THE EXPANSION OF $\tau_s(q)$ IN SERIES OF POWERS OF $\frac{1}{q}$

n	s = 4 D_n		n	s = 5 D_n	
0	0.67867080900717589987 8E	1	0	0.79441335871208531231 4E	1
1	-0.10000000000000000000 0E	1	1	-0.10000000000000000000 0E	1
2	0.00000000000000000000 0E	0	2	0.00000000000000000000 0E	0
3	.22622360300239196662 6F	1	3	.26480445290402843743 8E	1
4	-.25000000000000000000 0E	+0	4	-.25000000000000000000 0E	+0
5	-.92118813399690925710 2E	1	5	-.12621851690004326655 6E	2
6	.26392753683612396106 4E	1	6	.30893852838803317701 1E	1
7	.44477392581963797345 3E	2	7	.71442625673015338398 1E	2
8	-.22262046571591973713 2E	2	8	-.30502808250843789417 8E	2
9	-.23228352442201209735 0E	3	9	-.43850984002386033737 3E	3
10	.17309728321801953369 9E	3	10	.27772238761129951298 3E	3
11	.12648311946022584830 1E	4	11	.28159704388179158087 1E	4
12	-.12911934807469749795 4E	4	12	-.24273667209309540032 2E	4
13	-.70443566562794051335 5E	4	13	-.18577388704725118927 3E	5
14	.93890885643963765245 9E	4	14	.20699854191806475403 5E	5
15	.39645387268744284345 3E	5	15	.12454270956564370000 8E	6
16	-.67071868651487557769 5E	5	16	-.17359980380390922019 0E	6
17	-.22344495893915186304 2E	6	17	-.84221735539301479890 4E	6
18	.47265241738751729389 6E	6	18	.14380244497858747231 3E	7
19	.12510924856069787478 1E	7	19	.57129613191148106304 8E	7
20	-.32934632124075415021 9E	7	20	-.11795878607678005642 7E	8
21	-.68993757847480297807 2E	7	21	-.38683916852576508903 3E	8
22	.22723090469442822637 6E	8	22	.95968434054035401939 8E	8
23	.37060156025890152736 4F	8	23	.26024287375362030069 7E	9
24	-.15535187839025849561 4E	9	24	-.77516633862212363889 6E	9
25	-.19061209019556454563 3E	9	25	-.17302861683355895769 5E	10
26	.10528202590388115980 7E	10	26	.62201794845390905698 7E	10
27	.90934796372571287042 4E	9	27	.11294073167800880703 2E	11
28	-.70729904149739206252 8E	10	28	-.49603915717611724862 3E	11
29	-.37292133126462144565 8E	10	29	-.71685660630614633499 2E	11
30	.47095083986000163444 4E	11	30	.39320395265180675146 4E	12
31	.97299035351654527136 8E	10	31	.43564768876878347768 9E	12
32	-.31065649608197847179 4E	12	32	-.30983481324309292886 1E	13
33	.33410016397423107414 9E	11	33	-.24617841479479212679 6E	13
34	.20286911897967158776 4E	13	34	.24267319464210282436 0E	14
35	-.86018838152675081037 0E	12	35	.12070518876998135916 5E	14
36	-.13102387981298476635 6E	14	36	-.18889171089186739715 1E	15
37	.98540089774212335366 7E	13	37	-.39689976395566139663 5E	14
38	.83576911862230053407 1E	14	38	.14607471246791980053 8E	16
39	-.92154732766552384100 5E	14	39	-.10664173720124325764 3E	15
40	-.52553165395830652513 0E	15	40	-.11218300096384527988 8E	17

By using Eqs. (2-55) and (2-56) in Eqs. (2-21) and (2-22), respectively, it can be seen that the C_n and D_n satisfy the recursion formulae

$$nC_oC_n = -(n-2)C_{n-2} + \sum_{m=1}^{n-1} mC_mC_{n-m} \qquad (2\text{-}57)$$

where $C_o = \beta_s$ and $C_oC_1 = 1$. In Program 2-1 we show an algorithm which will permit one to compute these coefficients. In order to relate the FORTRAN variables to the mathematical variables, we let N denote the largest value of n for which the C_n are desired, and we let the C_n be denoted in the FORTRAN program by C(I), where n = I. Finally, we denote the value of β_s by the symbol BET. The subroutine to accomplish the setting up of the array C(1), C(2), C(3),...,C(N) has been written to yield double precision results. However, the author wishes to direct the attention of the reader to the manner in which the constants 0 and 1 have been inserted in a DATA statement. This feature is important because the use of 0.0D0 and 1.0D0 has been avoided in the program's executable statements and therefore the only changes that must be made to

Program 2-1

```
      SUBRØUTINE CØEFSQ(BET,N,C)
      DØUBLE PRECISIØN BET, C(N),ZERØ,ØNE,DFLØAT
      DATA ZERØ,ØNE/0.0D0,1.0D0/
      DFLØAT(KK) = DBLE(FLØAT(KK))
      C(1) = ØNE/BET
      C(2) = -C(1)*C(1)/(ØNE + ØNE)
      DØ 1 I = 3,N
      MF = I - 1
      C(I) = DFLØAT(I-2)*C(I-2)
      DØ 2 M = 1,MF
      C(I) = C(I) + DFLØAT(M)*C(M)*C(I-M)
    2 CØNTINUE
      C(I) = - C(I)/DFLOAT(I)
    1 CØNTINUE
      RETURN
      END
```

convert the program to single precision is to delete the DØUBLE PRECISIØN type statements, the arithmetic statement function DFLOAT, and insert "DATA ZERØ,ØNE/0.0,1.0/" in place of the DATA statement that appears in the program.*

In order to check on the numerical stability of the recursion formula in Eq. (2-57), the calculation of the $C_n(\tau_1^o)$ in Table 2-9 was repeated in single precision arithmetic. The results are listed in Table 2-11. The process seems to be "self-correcting" since there are some entries which are good to only four figures in the middle of the table, but the result for n = 80 displays agreement to six figures (after round off is used to write C_{80} = -0.127230E-21).

In order to obtain the recursion formula for the coefficients D_n of Eq. (2-56), we need merely insert Eq. (2-54) in Eq. (2-52) and find that

$$nD_n = -\sum_{m=1}^{n-2} mD_m D_{n-m-2} \qquad (2\text{-}58)$$

where $D_o = \alpha_s$, $D_1 = -1$ and $D_2 = 0$. In Program 2-2 we present an algorithm for the obtaining of the array $D_1, D_2, D_3, \cdots, D_N$. However, before we consider this program, we will make a digression in order to emphasis the importance in practical problems of the case

*It is a good programming practice to avoid the use of constants such as 0.0, 1.0, and 3.1415927 in the "body" of any program in which there exists the possibility that one might need to go to DØUBLE PRECISIØN since one would then have to replace many of the executable statements with new statements containing 0.0D0, 1.0D0, and 3.14159265358979320, respectively. The use of devices such as ZERØ,ØNE,PI in the executable statements and the inserting of the constants by means of a DATA statement permits one to readily convert a program written in double precision to use it in single precision, or vice versa.

of q very large. We will then introduce another array which is important in practical problems, and use Program 2-2 to set up both of these arrays.

In Fig. 2-5 we have presented several curves which show that for vertically polarized waves the cases in which q tends to infinity plays a vital role include virtually all cases of propagation

Table 2-11

THE COEFFICIENTS $C_n(\tau_1^o)$ AS COMPUTED IN SINGLE PRECISION

n	C_n	n	C_n		C_n
0	0.10187930E 01	34	-0.29925379E-10	67	0.12985392E-18
1	0.98155367E 00	35	0.20137136E-10	68	0.77486458E-19
2	-0.47283778E 00	36	0.50409587E-11	69	-0.57664740E-19
3	0.13440523E-00	37	-0.71986632E-11	70	-0.12651211E-19
4	-0.71602707E-02	38	-0.73094466E-13	71	0.21074336E-19
5	-0.98774498E-02	39	0.22287107E-11	72	-0.37643501E-21
6	0.20129035E-02	40	-0.43431269E-12	73	-0.66616721E-20
7	0.13461956E-02	41	-0.59945250E-12	74	0.15187661E-20
8	-0.56666862E-03	42	0.25780213E-12	75	0.18143185E-20
9	-0.19195578E-03	43	0.13276179E-12	76	-0.86768527E-21
10	0.15557887E-03	44	-0.10743466E-12	77	-0.39931940E-21
11	0.18911074E-04	45	-0.19145934E-13	78	0.36227523E-21
12	-0.40866278E-04	46	0.37440696E-13	79	0.51994960E-22
13	0.24151776E-05	47	-0.17673479E-14	80	-0.12723196E-21
14	0.10056239E-04	48	-0.11336792E-13	81	0.10060423E-22
15	-0.24616357E-05	49	0.29012308E-14	82	0.38772412E-22
16	-0.22318140E-05	50	0.29520079E-14	83	-0.11394486E-22
17	0.10840956E-05	51	-0.15230350E-14	84	-0.10073223E-22
18	0.41000339E-06	52	-0.61132877E-15	85	0.57878818E-23
19	-0.39026834E-06	53	0.60614715E-15	86	0.20240678E-23
20	-0.42217450E-07	54	0.66508472E-16	87	-0.22905735E-23
21	0.11611411E-06	55	-0.20495320E-15	88	-0.16954143E-24
22	-0.10456789E-07	56	0.21679091E-16	89	0.77408635E-24
23	-0.31502187E-07	57	0.60237029E-16	90	-0.10850725E-24
24	0.92562826E-08	58	-0.19411866E-16	91	-0.22683739E-24
25	0.73732547E-08	59	-0.15019482E-16	92	0.82855039E-25
26	-0.41809671E-08	60	0.92845527E-17	93	0.55753642E-25
27	-0.13494919E-08	61	0.28244545E-17	94	-0.38520651E-25
28	0.15133817E-08	62	-0.35367505E-17	95	-0.98833368E-26
29	0.98226314E-10	63	-0.15803798E-18	96	0.14534777E-25
30	-0.47271572E-09	64	0.11573625E-17	97	0.12684577E-27
31	0.65598900E-10	65	-0.19181695E-18	98	-0.47271444E-26
32	0.12950258E-09	66	-0.32833645E-18	99	0.95721225E-27
33	-0.46165995E-10			100	0.13275000E-26

over land when the frequency exceeds several megacycles and all cases for propagation over the sea when the frequency exceeds several hundred megacycles. For the case of horizontally polarized waves, the case $q = \infty$ plays an important role throughout the wavelength range from the infra-red to the longest waves for all the values of the conductivity and dielectric constant which occur in nature. These circumstances certainly justify our giving this case special attention. It is noteworthy that Pryce and Domb devoted their entire paper (Ref. 25) to a consideration of the limiting case in which $q = \infty$.

In many cases, the magnitude of q will be so large that one need only used the approximation of Eq. (E-7), namely,

$$\exp(i\xi\tau_s) \approx \exp(i\xi\tau_s^{\infty})\exp(i\xi/q) \tag{2-59}$$

to approximate the functions $W(x,y,y_o,q)$ and $F(x,y,q)$ defined in Eqs. (E-25) and (E-26) in the following manner:

$$W(x,y,y_o,q) \approx \exp(ix/q)W(x,y,y_o,\infty) \tag{2-60}$$

$$F(x,y,q) \approx \exp(ix/q)F(x,y,\infty) \tag{2-61}$$

When Eq. (2-59) is not sufficiently accurate, we can often evaluate the functions for large q by means of the expansions given below.

Since this case of $q \to \infty$ is so important, let us evaluate the coefficients $F_n(\alpha)$ in the expansion

$$\frac{d\tau_s(q)}{d\tau_s^{\infty}} = 1 + \sum_{n=3}^{\infty} F_n(\alpha_s)q^{-n}\exp(i\xi n\pi) \tag{2-62}$$

The reader will have to "look ahead" to the discussion in Section 5 of the Olver relation and the "normalization constant" to grasp some of the reasons why this expansion is so important. However,

we will assert (without a demonstration of the method of proof) that the importance lies in the fact that the following relationship

$$\frac{1}{(1 - \tau_s/q^2)[w_1'(\tau_s)]^2} = \frac{1}{[w_1'(\tau_s^\infty)]^2} \frac{d\tau_s(q)}{d\tau_s^\infty}$$

$$= \frac{1}{4\pi[Ai'(-\alpha_s)]^2} \frac{d\tau_s(q)}{d\alpha_s} \qquad (2\text{-}63)$$

permits us to express the diffraction function $V_{11}(\xi,q)$ in the form

$$V_{11}(\xi,q) = -2\sqrt{-i\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi\tau_s)}{(\tau_s - q^2)[w_1(\tau_s)]^2}$$

$$= 2\sqrt{-i\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi\tau_s)}{[w_1'(\tau_s^\infty)]^2} \frac{d\tau_s(q)}{d\tau_s^\infty} \qquad (2\text{-}64)$$

Therefore, with the aid of the coefficients D_n which are needed in Eq. (2-56) to compute $\tau_s = \tau_s(q)$ and the coefficients F_n which are needed in Eq. (2-62) to compute $d\tau_s(q)/d\tau_s^\infty$, we can evaluate the diffraction function $V_{11}(\xi,q)$ for the case in which q is very large. Since $V_{11}(\xi,q)$ is the diffraction function which is required when both source and receiver are at a very great height (in terms of wavelength) we observe that this function will be very frequently encountered in practice since the case of large values of q occurs in the high frequency range and it is therefore going to generally be the case that the antennas for transmission and reception are at a "great height." We should also make the observation that the diffraction function $V_o(\xi,q)$ can be readily evaluated for the case of $q \to \infty$ by expressing the residue series representation in the form

$$V_o(\xi,q) = -i\sqrt{i\pi\xi}\; q^{-2} \sum \frac{\exp(i\xi\tau_s)}{1 - \tau_s/q^2}$$

$$= -i\,\frac{\sqrt{i\pi\xi}}{q^2} \sum \exp(i\xi\tau_s)\,\frac{d\tau_s(q)}{d(\frac{1}{q})} \qquad (2\text{-}65)$$

Therefore, only the coefficients D_n are required in order to compute $V_o(\xi,q)$ for very large values of q.

By suitably combining the ideas employed to arrive at Eqs. (2-64) and (2-65), we can also evaluate the diffraction function $V_1(\xi,q)$. Since

$$V_1(\xi,q) = 2\sqrt{i\pi} \sum \frac{\exp(i\xi\tau_s)}{(\tau_s - q^2)w_1(\tau_s)} \qquad (2\text{-}66)$$

we need only observe that

$$\frac{1}{[(1 - \tau_s/q^2)w_1'(\tau_s)]^2} = \frac{1}{[w_1'(\tau_s^\infty)]^2}\left[\frac{d\tau_s(q)}{d\tau_s^\infty}\,\frac{d\tau_s(q)}{d(\frac{1}{q})}\right] \qquad (2\text{-}67)$$

to see that we can express $V_1(\xi,q)$ in the form

$$V_1(\xi,q) = -\frac{2\sqrt{i\pi}}{q^2} \sum \frac{\exp(i\xi\tau_s)}{w_1'(\tau_s^\infty)} \left[\frac{d\tau_s(q)}{d\tau_s^\infty}\,\frac{d\tau_s(q)}{d(\frac{1}{q})}\right]^{\frac{1}{2}} \qquad (2\text{-}68)$$

These results reveal that the knowledge of the coefficients D_n and F_n permit one to evaluate the three types of diffraction functions. Let us therefore return the to the development of the background theory for Program 2-2 in which these coefficients are to be computed.

Since $F_n(\alpha) = dD_n(\alpha)/d\alpha$, Eq. (2-58) leads to

$$F_n(\alpha) = \frac{dD_n(\alpha)}{d\alpha} = -\sum_{m=1}^{n-2} m[F_m D_{n-m-2} + D_m F_{n-m-2}]$$

$$= -\frac{(n-1)}{n}\sum_{m=1}^{n-2} F_m D_{n-m-2} \qquad (2\text{-}69)$$

where $F_o = 1$, $F_1 = 0$ and $F_2 = 0$. Program 2-2 has been used to set up an array for each of these coefficients. In writing this program we have done what is so often required in translating a

Program 2-2

```
      SUBRØUTINE LARGEQ(ALP,N,D,F)
      DØUBLE PRECISIØN  ALP,D(N),F(N),DFLØAT,ZERØ,ØNE,AN,AN1
      DATA ZERØ,ØNE/0.0D0,1.0D0/
      DFLØAT(KK) = DBLE(FLØAT(KK))
      D(1) = ALP
      D(2) = -ØNE
      D(3) = ZERØ
      F(1) = ØNE
      F(2) = ZERØ
      F(3) = ZERØ
      DØ 1 I = 4,N
      AN = DFLØAT(I-1)
      K  = I-3
      D(I) = ZERØ
      F(I) = ZERØ
      DØ 2 J = 1,K
      L    = K-J+1
      AN1  = DFLØAT(J)
      D(I) = D(I) + AN1*D(J+1)*D(L)
      F(I) = F(I) + AN1*(D(J+1)*F(L) + D(L)*F(J+1))
    2 CØNTINUE
      D(I) = - D(I)/AN
      F(I) = - F(I)/AN
    1 CØNTINUE
      RETURN
      END
```

mathematical expression into FORTRAN instructions - we have dimensioned the arrays D(N) and F(N), but since we need a place to put D_o and F_o, we have used the following convention

$$\begin{aligned} D_o &= D(1) & F_o &= F(1) \\ D_1 &= D(2) & F_1 &= F(2) \\ D_2 &= D(3) & F_2 &= F(3) \\ &\vdots & &\vdots \\ D_{N-1} &= D(N) & F_{N-1} &= F(N) \end{aligned}$$

The quantity denoted by ALP in the subroutine's argument list is the Airy function root α_s which is required in order to determine the coefficients for the root $\tau_s(q)$. In Table 2-12 we give the computer output which results from running the subroutine contained in Program 2-2. The case given in the table is that for s = 1. These tables were actually computed with the subroutine given in Program 2-3 rather than the one which is designated as Program 2-2. We take this occasion to present the instructions contained in Program 2-3 in order to permit the reader to have an example of a program which employs the NPREC subroutines which have been discussed in the Preface. There is a one-to-one correspondence between the instructions in Program 2-2 and those in Program 2-3, but the use of the NPREC subroutines requires that one put in a number of additional instructions which enable one to employ the computer software which permits the working with 21 decimal digits. The reader who is unfamiliar with NPREC will notice that in order to store the numbers one needs three conventional 36-bit "words," it is necessary to set us an array such as D(3,51) in order to obtain 51 values of D. Also, since this is a FORTRAN II program, the use of adjustatble dimensions in the arrays is not possible and therefore the integer 51 appears explicitly because we set out to obtain the coefficients for n = 0, 1, 2,...,50.

Table 2-12

THE COEFFICIENTS $D_n(\alpha_1)$ AND $F_n(\alpha_1)$

n	$D_n(\alpha_1)$	$F_n(\alpha_1)$
0	.233810741045976700C00E 1	.1000000000000000000000E 1
1	-.1C0000C0000000000000E 1	+C.C
2	+0.0	+0.0
3	.779369136819922333333E +0	.333333333333333333333E +0
4	-.250000000000000000000E +0	-C.C
5	-.109334925256937547187E 1	-.935242964183906799999E +0
6	.909263992956576055554E +0	.388888888888888888888E +0
7	.164740570689507445404E 1	.234289125555C5EEC45E255EE 1
8	-.264226069370932405702E 1	-.226017049677777476666E 1
9	-.213710405546052163332E 1	-.517464491527850323975E 1
10	.691693414275288888169E 1	.9C9C4180713625217804CE 1
11	.112297788720997851922E 1	.911112745519882451207E 1
12	-.166366387432300976017E 2	-.305386585496655844333E 2
13	.656326185545065344842E 1	-.747305883153744299052E 1
14	.363549113840041806875E 2	.900680865684224769793E 2
15	-.366549831925517799523E 2	-.329562820431761948952E 2
16	-.690160345205570436150E 2	-.234546326422511212139E 3
17	.131615045605843931255E 3	.235E1E2445194121175236E 3
18	.981386219147608437748E 2	.522730919468299964612E 3
19	-.393408408913159267568E 3	-.992323025066159018309E 3
20	-.213671996252882116077E 2	-.E82E4E3CC3111E3ECS554E 3
21	.103281515436985560184E 4	.338534769300161144809E 4
22	-.557005440585398234408E 3	.437853476401823925701E 3
23	-.238101946411483142016E 4	-.5555EC1143ECC66323676E 4
24	.289471314674802638095E 4	.495493922692861108129E 4
25	.458512006481859653460E 4	.257420606271768706578E 5
26	-.104802124890582256371E 5	-.303645112186488398690E 5
27	-.596169025865546022592E 4	-.554454619032728532684E 5
28	.317985515060398151420E 5	.122585863334508983208E 6
29	-.318078127520901523539E 4	.838693002088812401608E 5
30	-.842338171333703163827E 5	-.4082444C712226EEE4773221E 6
31	.599334412880064196210E 5	.128148424303529659982E 5
32	.193151282158636813687E 6	.117864891622043676654E 7
33	-.281949289996349875198E 6	-.77E911222724SE5E3CE5CE 6
34	-.357114653412512495799E 6	-.294166033372288326956E 7
35	.994137337829276410111E 6	.412402065801852456725E 7
36	.373873844573088530982E 6	.597E3C3E647246423S376E 7
37	-.297606709730805521484E 7	-.158099112391683853018E 8
38	.797089484675490000935E 6	-.747523525814095398903E 7
39	.776451218363028760465E 7	.5C92936883479065886 41E 8
40	-.700691631307825876177E 7	-.1034384930383755548 54E 8

Program 2-3

```
      SUBRØUTINE LARGEQ(ALP,D,F)
      DIMENSIØN ALP(3),D(3,51),C(3,51),AN(3),AN1(3)
      CØMMØN AN,AN1
      CALL SP2NP(0.0,D(1,3))
      CALL SP2NP(-1.0 ,D(1,2))
      CALL NPREC
      D(1,1) = ALP
      CALL NPØUT
      CALL SP2NP(0.0,F(1,3))
      CALL SP2NP(0.0,F(2,3))
      CALL SP2NP(1.0,F(1,1))
      DØ 1 I = 4,51
      AN = FLØATF(I-1)
      CALL SP2NP(AN,AN)
      K  = I - 3
      CALL SP2NP(0.0,D(1,I))
      CALL SP2NP(0.0,F(1,I))
      DØ 2 J = 1,K
      L = K - J + 1
      AN1 = FLØATF(J)
      CALL SP2NP(AN1,AN1)
      CALL NPREC
      D(1,I) = D(1,I) + AN1*D(1,J+1)*D(1,L)
      F(1,I) = F(1,I) + AN1*(D(1,J+1)*F(1,L) + D(1,L)*F(1,J+1))
      CALL NPØUT
      CALL NP2SP(AN1,AN1)
    2 CØNTINUE
      CALL NPREC
      D(1,I) = - D(1,I)/AN
      F(1,I) = - F(1,I)/AN
      CALL NPØUT
      CALL NP2SP(AN,AN)
    1 CØNTINUE
      RETURN
      END
```

Before leaving the subject of these classical expansions, we will present the numerical values for the coefficients for further terms in the explicit representations for $A_n'(\tau)$ and $B_n'(\tau)$. The data which we have obtained in presented in Block 2-1. This is a FORTRAN IV BLØCK DATA subprogram and the data are placed in a labeled COMMØN which we have designated as BREMM. The data consists of the arrays BREM(18,9) and BR(20,3). The computations which led to the obtaining of these data were undertaken because

```
      BLOCK DATA
      COMMON/BREMM/BREM(18,9),BR(20,3)
      DOUBLE PRECISION BREM,BR
      DATA BREM(1,1), BREM(2,1), BREM(3,1), BREM(3,2), BREM(4,1),          BREM1  D
     A   BREM(4,2), (BREM(5,I),I=1,3), (BREM(6,I),I=1,3), (BREM(7,I),      BREM1  D
     B   I=1,4), (BREM(8,I),I=1,4),(BREM(9,I),I=1,5), (BREM(10,I),         BREM1  D
     C   I=1,5)     /                                                     BREM1  D
     D -0.10000000000000000000D+01, +0.15000000000000000000D+01,          BREM1  1
     E -0.25000000000000000000D+01, -0.66666666666666666667D+00,          BREM1  2
     F +0.43750000000000000000D+01, +0.23333333333333333333D+01,          BREM1  3
     G -0.78750000000000000000D+01, -0.63000000000000000000D+01,          BREM1  4
     H -0.60000000000000000000D+00, +0.14437500000000000000D+02,          BREM1  5
     I +0.16399999999999999999D+02, +0.32222222222222222222D+01,          BREM1  6
     J -0.26812500000000000000D+02, -0.35750000000000000000D+02,          BREM1  7
     K -0.11822222222222222222D+02, -0.57142857142857142857D+00,          BREM1  8
     L +0.50273437500000000000D+02, +0.80437500000000000000D+02,          BREM1  9
     M +0.36675000000000000000D+02, +0.41571428571428571428D+01,          BREM1 10
     N -0.94960937500000000000D+02, -0.17726041666666666667D+03,          BREM1 11
     O -0.10341527777777777778D+03, -0.19274779541446208112 8D+02,        BREM1 12
     P -0.55555555555555555555D+00, +0.18042578125000000000D+03,          BREM1 13
     Q +0.38490833333333333333D+03, +0.27416277777777777777D+03,          BREM1 14
     R +0.72360934744268077601 0D+02, +0.51311111111111111111D+01/        BREM1 15
      DATA (BREM(11,I),I=1,6),(BREM(12,I),I=1,6),(BREM(13,I),I=1,7),       BREM1  D
     A   (BREM(14,I),I=1,7), (BREM(15,I),I=1,8)     /                     BREM1  D
     B -0.34444921875000000000D+03, -0.82667812500000000000D+03,          BREM1 16
     C -0.69615000000000000000D+03, -0.23968454545454545454 5D+03,        BREM1 17
     D -0.28827272727272727272D+02, -0.54545454545454545454 5D+00,        BREM1 18
     E +0.66019433593750000000 0D+03, +0.17605182291666666666 6D+04,      BREM1 19
     F +0.17122895833333333333D+04, +0.73036283389450056115 7D+03,        BREM1 20
     G +0.12713674897119341563 6D+03, +0.61389129389129389129 4D+01,      BREM1 21
     H -0.12696044921875000000 0D+04, -0.37241731770833333333 3D+04,      BREM1 22
     I -0.41100067708333333333 0D+04, -0.20965056341189674522 6D+04,      BREM1 23
     J -0.48385435839880284324 1D+03, -0.40628634328634328634 3D+02,      BREM1 24
     K -0.53846153846153846153 8D+00, +0.24485229492187500000 0D+04,      BREM1 25
     L +0.78352734375000000000 0D+04, +0.96763734375000000000 0D+04,      BREM1 26
     M +0.57536891883116883116 9D+04, +0.16658590909090909090 9D+04,      BREM1 27
     N +0.20616411588411588411 6D+03, +0.71765377479663193948 6D+01,      BREM1 28
     O -0.47338110351562500000 0D+04, -0.16410544921875000000 0D+05,      BREM1 29
     P -0.22427325555555555555 5D+05, -0.15247142561928811928 8D+05,      BREM1 30
     Q -0.53323218255571588903 5D+04, -0.88712486360278952870 1D+03,      BREM1 31
     R -0.54813098746051127003 1D+02, -0.53333333333333333333 3D+00/      BREM1 32
```

```
 DATA (BREM(16,I),I=1,8), (BREM(17,I),I=1,9),(BREM(18,I),I=1,9)/    BREM1  D
A +0.91717588806152343750OD+04, +0.34241233154296875000D+05,       BREM1 33
B +0.51311770887586805555D+05, +0.39286165424520502643 8D+05,      BREM1 34
C +0.16147079150022045854 8D+05, +0.34069692750980158386 8D+04,    BREM1 35
D +0.31505766485718866670 8D+03, +0.82408147408147408147 1D+01,    BREM1 36
E -0.17804002532958984375 0D+05, -0.71216010131835937500 0D+05,    BRE71 37
F -0.11612624194335937500 0D+06, -0.98923729017857142852 5D+05,    BREM1 38
G -0.46803984281250000000 0D+05, -0.12034291181612505141 5D+05,    BREM1 39
H -0.15139770680160175958 2D+02, -0.71503395428101310453 7D+02,    BREM1 40
I -0.52941176470588235294 1D+00, +0.34618893814086914062 5D+05,    BREM1 41
J +0.14770728027343750000 0D+06, +0.26038642154947916666 7D+06,    BREM1 42
K +0.24435742340133101850 5D+06, +0.13095257575147890945 8D+06,    BREM1 43
L +0.39860161242696926246 4D+05, +0.64208609705147770989 6D+04,    BREM1 44
M +0.45985941223305057727 0D+03, +0.93291779389818605504 6D+01/    BREM1 45
 DATA BR(3,1),BR(5,1),BR(6,1),BR(7,1),BR(8,1),BR(9,1),BR(9,2),      BREM2  D
A   BR(10,1),BR(11,1),BR(11,2),BR(12,1),BR(12,2),BR(13,1),BR(13,2),BREM2  D
B   BR(14,1),BR(14,2),BR(15,1),BR(15,2),BR(15,3),BR(16,1),BR(16,2)  BREM2  D
C   ,(BR(17,I),I=1,3),(BR(18,I),I=1,3),(BR(19,I),I=1,3),(BR(20,I),  BREM2  D
D I=1,3)/0.33333333333333333333D0,0.4D0,0.38888888888888888888D0,BREM2  1
E +0.42857142857142857142 9D+00, +0.96666666666666666667 0D+00,    BREM2  2
F +0.50617283950617283950 6D+00, +0.44444444444444444444 4D+00,    BREM2  3
G +0.16628571428571428571 4D+01, +0.19131313131313131313 1D+01,    BREM2  4
H +0.45454545454545454545 4D+00, +0.69231334509112286889 9D+00,    BREM2  5
I +0.24433862433862433862 4D+01, +0.45323076923076923076 9D+01,    BREM2  6
J +0.46153846153846153846 1D+00, +0.35138883162692686502 1D+01,    BREM2  7
K +0.32887033601319315605 0D+01, +0.97418626298255927885 3D+00,    BREM2  8
L +0.86194426807760141093 4D+01, +0.46666666666666666667 0D+00,    BREM2  9
M +0.10604600399600399600 4D+02, +0.41863136863136863136 8D+01,    BREM2 10
N +0.62078274631962773886 0D+01, +0.14391435766225682192 0D+02,    BREM2 11
O +0.47058823529411764705 8D+00, +0.13963268045778333844 1D+01,    BREM2 12
P +0.24753261410742892224 3D+02, +0.51276180609513942847 1D+01,    BREM2 13
Q +0.22762617157278811414 1D+02, +0.22036799290934629280 4D+02,    BREM2 14
R +0.47368421052631578947 3D+00, +0.10709012800723040374 4D+02,    BREM2 15
S +0.49347279188494874769 3D+02, +0.61063710146063087239 5D+01/    BREM2 16
 END
```

the author was interested in obtaining an asymptotic expansion for the function $V_2(0,q)$ which describes the "foward scattered field" which has attracted so much attention in the literature. This function can be defined by the relationship

$$V_2(0,q) = \lim_{\xi \to 0} \left[V_{11}(\xi,q) - \frac{\sqrt{I}}{2\sqrt{\pi}\,\xi} \right] \tag{2-70}$$

An expansion in powers of q can be found "operationally" from the divergent series representation

$$V_2(0,q) = -2\sqrt{-i\pi} \sum_{s=1}^{\infty} \frac{1}{\tau_s [w_1(\tau_s^o)]^2} \frac{d\tau_s(q)}{d\tau_s^o} \tag{2-71}$$

and an expansion in inverse powers of q comes from the divergent series representation

$$V_2(0,q) = 2\sqrt{i\pi} \sum_{s=1}^{\infty} \frac{1}{[w_1'(\tau_s^\infty)]} \frac{d\tau_s^\infty(q)}{d\tau_s^\infty} \tag{2-72}$$

Therefore, the quantities which were sought in the computer program and which have been prepared for re-entry into the computer with the aid of Block 2-1 were the derivatives $A_n(\tau_s^o)$ and $B_n(\tau_s^\infty)$ which appear in the expressions

$$\frac{d\tau_s(q)}{d\tau_s^o} = 1 + A_1'(\tau_s^o)q + A_2'(\tau_s^o)q^2 + \cdots \tag{2-73}$$

$$\frac{d\tau_s(q)}{d\tau_s^\infty} = 1 + B_1'(\tau_s^\infty)q^{-1} + B_2'(\tau_s^\infty)q^{-2} + \cdots \tag{2-74}$$

where the explicit forms for the first few coefficients are

$$A_1'(\tau) = -\frac{1}{\tau^2} \quad , \qquad B_1'(\tau) = B_2'(\tau) = 0$$

$$A_2'(\tau) = \frac{3}{2\tau^4} \quad , \qquad B_3'(\tau) = \frac{1}{3} \quad , \quad B_4'(\tau) = 0$$

$$A_3'(\tau) = -\frac{5}{2\tau^6} - \frac{2}{3\tau^3} \quad , \qquad B_5'(\tau) = \frac{2}{5}\tau$$

$$A_4'(\tau) = \frac{35}{8\tau^8} + \frac{7}{3\tau^5} \quad , \qquad B_6'(\tau) = \frac{7}{18}$$

The array BREM(18,9) in Block 2-1 contains the numerical values of the coefficients E_n^m in the expansion

$$A_n'(\tau) = \frac{1}{\tau^{2n}} \left[\sum_{m=1}^{K} E_n^m \tau^{3m} \right] \quad , \quad K = [(n-1)/2] \qquad (2\text{-}75)$$

where the [...] denote that K is the integral part of the expression enclosed in brackets. The data go as far as n = 18. The relation between the notation in Eq. (2-75) and that in Block 2-1 is E_N^M = BREM(N,M).

In Table 2-13 we give values for $C_n(\beta) = A_n(\tau)\exp[i\frac{1}{3}(2n-1)]$ and $E_n(\beta) = A_n'(\tau)\exp(i\frac{2}{3}n\pi)$ for the case in which $\beta = \beta_1$ and $\tau = \tau_1^o = \beta_1 \exp(i\frac{1}{6}\pi)$.

The numbers which appear in the array BR(20,3) contain the non-zero coefficients which appear in the expansion

$$B_n'(\tau) = \sum_{m=1}^{K} F_n^m \tau^{m-1} \quad , \quad K = [(n-1)/2] \qquad (2\text{-}76)$$

In Table 2-14 we give values of the F_n^m for $n \leq 38$. The array BR(20,3) can be used to set up an array F(20,9) which is related to the coefficients F_n^m by means of the relation F_N^M = F(N,M). This array can be set up by the following set of FORTRAN statements:

(Text continues on p. 2-84)

Table 2-13

THE COEFFICIENTS $C_n(\beta_1)$ AND $E_n(\beta_1)$

n	$C_n(\beta_1)$	$E_n(\beta_1)$
0	.10187929716474710890 2E 1	.10000000000000000000E 1
1	.98155368934565643349 7E +0	-.96344764506806941597 1E +0
2	-.47283779525397403410 5E +0	.13923470471808129930 4E 1
3	.13440524531392944729 7E +0	-.16053054125344253173 0E 1
4	-.71602938885920958033 1E -2	.16436381013601944822 8E 1
5	-.98774282120637696019 7E -2	-.15639534210030e465911E 1
6	.20128847168449974731 1E -2	.14043300102476047666 7E 1
7	.13462118035028363786 9E -2	-.11963904081477613925 0E 1
8	-.56668183510136790723 0E -3	.97200460070773450190 9E +0
9	-.19194546486765931214 3E -3	-.75771755754966854524 5E +0
10	.15557115743545349840 6E -3	.56974005571320268559 6E +0
11	.18916689089679143283 0E -4	-.41467814609907902685 4E +0
12	-.40870251099793881889 1E -4	.29286454687132957931 6E +0
13	.24179084285597400745 E -5	-.20114822081866261170 6E +0
14	.10054412469337743735 3E -4	.13466140710953970261 6E +0
15	-.24604427443158133887 6E -5	-.88045331403569040094 6E -1
16	-.22325772762695294571 5E -5	.56306184217195336401 9E -1
17	.10845740096401966824 6E -5	-.35262793224434505018 0E -1
18	.40970966684901684975 0E -6	.21652329914413193894 1E -1
19	-.38009141647458677241 0E -6	-.13050841963263201666 0E -1
20	-.42322222524222362163 5E -7	.77297532412165772684 1E -2
21	.11617517415717005039 2E -6	-.45022764394089775786 8E -2
22	-.10491804636115501631 8E -7	.25807341603517489946 9E -2
23	-.31482430296256068030 8E -7	-.14568860416957410666 5E -2
24	.92452961606875392783 0E -8	.81060131724229592635 1E -3
25	.73792846770156103538 8E -8	-.44478953910721467867 1E -3
26	-.41842339811879770231 1E -8	.24060560662108797766 8E -3
27	-.13477462039333872514 2E -8	-.12668969964730067211 3E -3
28	.15124611057456761281 2E -8	.67924894048143916885 1E -4
29	.98706365158131995873 5E-10	-.35428739307444012608 0E -4
30	-.47296345270391202867 1E -9	.18267522927603778885 0E -4
31	.65725283233185077656 8E-10	-.93133978987356338091 2E -5
32	.12943890327127395506 2E -9	.46967614687308792551 9E -5
33	-.46134257672511147905 4E-10	-.23440789228599980176 5E -5
34	-.29941065860438753183 0E-10	.11582455486567746928 4E -5
35	.20144823552162074024 0E-10	-.56667166275756340520 9E -6
36	.50372341365009437432 9E-11	.27453892031472087236 8E -6
37	-.71968806609140233879 4E-11	-.13176663923238432826 9E -6
38	-.73940983916856311752 9E-13	.62689633747060563585 7E -7
39	.22291110964115985337 8E-11	-.29569778631045041304 5E -7
40	-.43450039839476856594 5E-12	.13823902955464913194 5E -7

Table 2-14

COEFFICIENTS FOR POLYNOMIAL REPRESENTATION FOR $F_n(\tau)$

m	F_n^m
	n = 3
1	.33333333333333333333E +0
	n = 5
1	+0.0
2	.40000000000000000000E +0
	n = 6
1	.38888888888888888888E +0
2	+0.0
	n = 7
1	+0.0
2	+0.0
3	.42857142857142857142E +0
	n = 8
1	+0.0
2	.96666666666666666666E +0
3	+0.0
	n = 9
1	.50617283950617283950E +0
2	+0.0
3	+0.0
4	.44444444444444444444E +0
	n = 10
1	+0.0
2	+0.0
3	.16628571428571428571E 1
4	+0.0
	n = 11
1	+0.0
2	.19131313131313131313E 1
3	+0.0
4	+0.0
5	.45454545454545454545E +0

m	F_n^m
	n = 12
1	.69231334509112286889 9E +0
2	+0.0
3	+0.0
4	.24433862433862433862 4E 1
5	+0.0
	n = 13
1	+0.0
2	+0.0
3	.45323076923076923076 9E 1
4	+0.0
5	+0.0
6	.46153846153846153846 1E +0
	n = 14
1	+0.0
2	.35138883162692686502 1E 1
3	+0.0
4	+0.0
5	.32887033601319315605 0E 1
6	+0.0
	n = 15
1	.97418626298255927885 3E +0
2	+0.0
3	+0.0
4	.86194426607760141093 4E 1
5	+0.0
6	+0.0
7	.46666666666666666666 6E +0
	n = 16
1	+0.0
2	+0.0
3	.10604600399600399600 3E 2
4	+0.0
5	+0.0
6	.41863136863136863136 8E 1
7	+0.0

Table 2-14 (Cont'd)

m	F_n^m
	n = 17
1	+0.0
2	.62078274631962773886OE 1
3	+0.0
4	+0.0
5	.14391435766225682192OE 2
6	+0.0
7	+0.0
8	.47058823529411764705 8E +0
	n = 18
1	.13963268045778333844 1E 1
2	+0.0
3	+0.0
4	.24753261410742892224 3E 2
5	+0.0
6	+0.0
7	.51276160609513942847 1E 1
8	+0.0
	n = 19
1	+0.0
2	+0.0
3	.22762617157278811414 1E 2
4	+0.0
5	+0.0
6	.22036799290934629280 4E 2
7	+0.0
8	+0.0
9	.47368421052631578947 3E +0
	n = 20
1	+0.0
2	.10709012800723040374 4E 2
3	+0.0
4	+0.0
5	.49347279188494874769 3E 2
6	+0.0
7	+0.0
8	.61063710146063087239 5E 1
9	+0.0

m	F_n^m
	n = 21
1	.20274541706387119449 4E 1
2	+0.0
3	+0.0
4	.62976153561769082051 1E 2
5	+0.0
6	+0.0
7	.31722460490714458968 3E 2
8	+0.0
9	+0.0
10	.47619047619047619047 6E +0
	n = 22
1	+0.0
2	+0.0
3	.46215836994616496847 8E 2
4	+0.0
5	+0.0
6	.88328698564512966828 7E 2
7	+0.0
8	+0.0
9	.71178476181853609379 5E 1
10	+0.0
	n = 23
1	+0.0
2	.18179563827781342377 0E 2
3	+0.0
4	+0.0
5	.14595480003533956902 6E 3
6	+0.0
7	+0.0
8	.43598397463553960935 3E 2
9	+0.0
10	+0.0
11	.47826086956521739130 4E +0

Table 2-14 (Cont'd)

m	F_n^m	F_n^m
	n = 24	n = 27
1	.29721955453724352443 2E 1	1 .43894641905860423882 7E 1
2	+0.0	2 +0.0
3	+0.0	3 +0.0
4	.14784974246976871366 3E 3	4 .32775596625936247427 3E 3
5	+0.0	5 +0.0
6	+0.0	6 +0.0
7	.14615634302642722384 6E 3	7 .55925210221413858286 8E 3
8	+0.0	8 +0.0
9	+0.0	9 +0.0
10	.81583573518558038668 8E 1	10 .74455120828592138079 6E 2
11	+0.0	11 +0.0
	n = 25	12 +0.0
1	+0.0	13 .48148148148148148148 1E +0
2	+0.0	n = 28
3	.90251283172271150237 1E 2	1 +0.0
4	+0.0	2 +0.0
5	+0.0	3 .17124298381901058361 4E 3
6	.29904171501835304336 1E 3	4 +0.0
7	+0.0	5 +0.0
8	+0.0	6 .90144077773070778140 2E 3
9	.57800927734241553548 9E 2	7 +0.0
10	+0.0	8 +0.0
11	+0.0	9 .33857790613371638217 4E 3
12	.47999999999999999999 9E +0	10 +0.0
	n = 26	11 +0.0
1	+0.0	12 .10315164280750113490 5E 2
2	.30506844158278786159 0E 2	13 +0.0
3	+0.0	n = 29
4	+0.0	1 +0.0
5	.39058332019856206733 2E 3	2 .50748887559993113176 0E 2
6	+0.0	3 +0.0
7	+0.0	4 +0.0
8	.22777316477408251957 5E 3	5 .97210916814928243547 5E 3
9	+0.0	6 +0.0
10	+0.0	7 +0.0
11	.92249428966770410585 1E 1	8 .97467266869810339474 4E 3
12	+0.0	9 +0.0

Table 2-14 (Cont'd)

m	F_n^m
	n = 29 (Cont'd)
10	+0.0
11	.936766189615653190600E 2
12	+0.0
13	+0.0
14	.482758620689655172413E +0
	n = 30
1	.652069314549479024577E 1
2	+0.0
3	+0.0
4	.695945579296443016777E 3
5	+0.0
6	+0.0
7	.188022396839860045014E 4
8	+0.0
9	+0.0
10	.484400309450673263990E 3
11	+0.0
12	+0.0
13	.114270663855553701652E 2
14	+0.0
	n = 31
1	+0.0
2	+0.0
3	.317794011464592426888E 3
4	+0.0
5	+0.0
6	.249550314037575970858E 4
7	+0.0
8	+0.0
9	.160582473370178115902E 4
10	+0.0
11	+0.0
12	.115573044314335074603E 3
13	+0.0
14	+0.0
15	.483870967741935483870E +0

	F_n^m
	n = 32
1	+0.0
2	.838496463337888808773E 2
3	+0.0
4	+0.0
5	.228979320141244052940E 4
6	+0.0
7	+0.0
8	.362400322674403857380E 4
9	+0.0
10	+0.0
11	.671479258008439025267E 3
12	+0.0
13	+0.0
14	.125588863013868467797E 2
15	+0.0
	n = 33
1	.973305410627465997017E 1
2	+0.0
3	+0.0
4	.142898572373963308229E 4
5	+0.0
6	+0.0
7	.574360013033138447610E 4
8	+0.0
9	+0.0
10	.252699166255536099042E 4
11	+0.0
12	+0.0
13	.140245109198466331867E 3
14	+0.0
15	+0.0
16	.484848484848484848484E +0
	n = 34
1	+0.0
2	+0.0
3	.579477224434114964707E 3
4	+0.0
5	+0.0
6	.647172918046845199628E 4
7	+0.0

Table 2-14 (Cont'd)

m	F_n^m
	n = 34 (Cont'd)
8	+0.0
9	.655539933695388262864E 4
10	+0.0
11	+0.0
12	.906443355298408425055E 3
13	+0.0
14	+0.0
15	.137091867046603180395E 2
16	+0.0
	n = 35
1	+0.0
2	.137786191682490519142E 3
3	+0.0
4	+0.0
5	.516500505178988111748E 4
6	+0.0
7	+0.0
8	.121316535158478055037E 5
9	+0.0
10	+0.0
11	.382751190307577062323E 4
12	+0.0
13	+0.0
14	.167787507355019997021E 3
15	+0.0
16	+0.0
17	.485714285714285714284E +0
	n = 36
1	.145855972182003225723E 2
2	+0.0
3	+0.0
4	.285628962836130224192E 4
5	+0.0
6	+0.0
7	.162922813170870526888E 5
8	+0.0
9	+0.0
10	.112546659995619454984E 5
11	+0.0
12	+0.0
13	.119629354980511123509E 4

m	F_n^m
	n = 36 (Cont'd)
14	+0.0
15	+0.0
16	.148767067820837517100E 2
17	+0.0
	n = 37
1	+0.0
2	+0.0
3	.104162580780469409211E 4
4	+0.0
5	+0.0
6	.159372661320496727412E 5
7	+0.0
8	+0.0
9	.239040540638447963500E 5
10	+0.0
11	+0.0
12	.561303917260750868718E 4
13	+0.0
14	+0.0
15	.198289640531145198050E 3
16	+0.0
17	+0.0
18	.486486486486486486485E +0
	n = 38
1	+0.0
2	.225406614380351157893E 3
3	+0.0
4	+0.0
5	.112501909112857603875E 5
6	+0.0
7	+0.0
8	.374247923442398553549E 5
9	+0.0
10	+0.0
11	.164955702084799466403E 5
12	+0.0
13	+0.0
14	.154838749050109722940E 4
15	+0.0
16	+0.0
17	.160603453183037816598E 2
18	+0.0

```
      CØMMØN /BREMM/BREM(18,9),BR(20,3)
      DØUBLE PRECISIØN F(20,9),BREM,BR
      DØ 1 I = 1,20
      DØ 2 J = 1,9
      F(I,J) = 0.0D0
    1 CØNTINUE
    2 CØNTINUE
      F(3,1) = BR(3,1)
      F(5,2) = BR(5,1)
      DØ 3 K = 2,6
      I0     = 3*K
      I1     = I0 + 1
      I2     = I0 + 2
      F(I0,1)= BR(I0,1)
      F(I1,3)= BR(I1,1)
      F(I2,2)= BR(I2,1)
    3 CØNTINUE
      F(9,3) = BR(9,2)
      F(11,5)= BR(11,2)
      DØ 4 K = 4,6
      I0     = 3*K
      I1     = I0 + 1
      I2     = I0 + 2
      DØ 5 J = 2,3
      J0     = 3*J - 2
      J1     = J0 + 2
      J2     = J0 + 1
      F(I0,J0) = BR(I0,J)
      F(I1,J1) = BR(I1,J)
      F(I2,J2) = BR(I2,J)
    5 CØNTINUE
    4 CONTINUE
```

As an illustration of the usefulness of the results given in the relations which have been labeled Eq. (2-75) and Eq. (2-76), let us consider Eq. (2-72) in further detail. We can show that the function $V_2(\xi,\infty)$ is an entire function of ξ which can be expanded in a Taylor series of the form

$$V_2(\xi,\infty) = 2\sqrt{i\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi\tau_s^{\infty})}{[w_1'(\tau_s^{\infty})]^2} = \sum_{n=0}^{\infty} a_n \frac{\xi^n}{n!} \qquad (2\text{-}77)$$

Let us now observe that

$$\frac{d^r V_2(\xi,\infty)}{d\xi^r} = 2\sqrt{i\pi} \sum_{s=1}^{\infty} \frac{(i\tau_s^\infty)^r \exp(i\xi\tau_s^\infty)}{[w_1'(\tau_s^\infty)]^2} = \sum_{n=r}^{\infty} a_n \frac{\xi^{n-r}}{(n-r)!} = \sum_{n=0}^{\infty} a_{n+r} \frac{\xi^n}{n!}$$

Therefore, we can express a function of the form

$$F(\xi) = 2\sqrt{i\pi} \sum_{s=1}^{\infty} [c_o + c_1\tau_s^\infty + c_2(\tau_s^\infty)^2 + \cdots + c_m(\tau_s^\infty)^m] \frac{\exp(i\xi\tau_s^\infty)}{[w_1'(\tau_s^\infty)]^2}$$

in terms of the function $V_2(\xi,\infty)$ according to the rule

$$F(\xi) = \left[c_o + c_1 \frac{d}{d(i\xi)} + c_2 \frac{d^2}{d(i\xi)^2} + \cdots + c_m \frac{d^m}{d(i\xi)^m}\right] V_2(\xi,\infty)$$

If we now use this rule in the representation which results from using Eq. (2-76) in Eq. (2-74), we find that Eq. (2-72) leads to

$$V_2(0,q) = V_2(0,\infty) + \sum_{n=1}^{\infty} A_n q^{-n} \qquad (2\text{-}78)$$

where

$$A_n = \sum_{m=1}^{K} (-i)^{m-1} F_n^m a_{m-1} \qquad (2\text{-}79)$$

The coefficients F_n^m are those which were defined in Eq. (2-76) and the coefficients a_j are those defined by Eq. (2-77).

The form of the result in Eq. (2-79) provides an excellent example of a situation which occurs frequently in numerical analysis. Since we know a recursion formula which will permit us to compute the quantities $B_n'(\tau) = F_n(\alpha)\exp(i\frac{2}{3}n\pi)$ without having to take recourse to the representation in Eq. (2-76) it is tempting to dismiss this representation in the form of a polynomial in τ as being "useless." However, we see in Eq. (2-79) that the coefficients in the polynomial representation can play an important role in analytical work.

2.5 Programs for Step-by-Step Integration

In this discussion we want to present some methods by which one can compute the root t(q) of

$$w_1'(t) - q\, w_1(t) = 0 \tag{2-80}$$

by integrating the differential equation

$$\frac{dt}{dq} = \frac{1}{t - q^2} \tag{2-81}$$

with the initial conditions

$$t_o = t(q_o)$$

We can use the differential equation to compute

$$\left[\frac{d^n t}{dq^n}\right]_{q=q_o}$$

and use these expressions in the Taylor series to obtain for t(q) a representation of the form

$$t(q) = t_o + \sum_{n=1}^{\infty} A_n (q - q_o)^n \tag{2-82}$$

The explicit form of the first few A_n are

$$A_1 = \frac{1}{t_o - q^2}$$

$$A_2 = -\frac{1}{2}\left[\frac{1}{\left(t_o - q_o^2\right)^3} - \frac{2q_o}{\left(t_o - q_o^2\right)^2}\right]$$

$$A_3 = -\frac{1}{2}\frac{1}{(t_o - q_o^2)^5} - \frac{5}{3}\frac{q_o}{(t_o - q_o^2)^4} + \frac{4}{3}\frac{q_o^2}{(t_o - q_o^2)^3} + \frac{1}{3}\frac{1}{(t_o - q_o^2)^2}$$

$$A_4 = -\frac{5}{8}\frac{1}{(t_o - q_o^2)^7} + \frac{35}{12}\frac{q_o}{(t_o - q_o^2)^6} - \frac{13}{3}\frac{q_o^2}{(t_o - q_o^2)^5} + 2\frac{q_o^3}{(t - q_o^2)^4}$$

$$- \frac{7}{12}\frac{1}{(t_o - q_o^2)^4} + \frac{q_o}{(t_o - q_o^2)^3}$$

$$A_5 = \frac{7}{8}\frac{1}{(t_o - q_o^2)^9} - \frac{105}{20}\frac{q_o}{(t_o - q_o^2)^8} + \frac{34\, q_o^2}{3(t_o - q_o^2)^7} + \left(\frac{21}{20} - \frac{154}{15}\, q_o^3\right)\frac{1}{(t_o - q_o^2)^6}$$

$$\left(\frac{16}{5}\, q_o^4 - \frac{49}{15}\right)\frac{q_o}{(t_o - q_o^2)^5} + \frac{12}{5}\frac{q_o^2}{(t_o - q_o^2)^4} + \frac{1}{5(t_o - q_o^2)^3}$$

$$A_6 = -\frac{21}{16}\frac{1}{(t_o - q_o^2)^{11}} + \frac{157}{24}\frac{q_o}{(t_o - q_o^2)^{10}} - \frac{364}{18}\frac{q_o^2}{(t_o - q_o^2)^9}$$

$$+ \left(\frac{2134}{90}\, q_o^3 - \frac{231}{20}\right)\frac{1}{(t_o - q_o^2)^8} + \left(\frac{709}{90}\, q_o - \frac{1128}{90}\, q_o^4\right)\frac{1}{t_o - q_o^{2\,7}}$$

$$+ \left(\frac{16}{3}\, q_o^5 - \frac{1006}{90}\right)\frac{1}{(t_o - q_o^2)^6} + \left(-\frac{58}{90} + \frac{16}{3}\, q_o^3\right)\frac{1}{(t_o - q_o^2)^5} + \frac{q_o}{(t_o - q_o^2)^4}$$

Since t_o and q_o are complex quantities, the programming of these coefficients would be quite tedious. However, we can develop a much simpler method for the computation of the A_n if we insert Eq. (2-82) in Eq. (2-81) and obtain a recursion formula for the coefficients.

We observe that

$$\frac{dt}{dq} = A_1 + \sum_{n=1}^{\infty} (n + 1)A_{n+1}(q - q_o)^n$$

and

$$t - q^2 = t - [q_o + (q - q_o)]^2$$

$$= t_o - q_o^2 + (A_1 - 2q^o)(q - q_o) + (A_2 - 1)(q - q_o)^2$$

$$+ \sum_{n=3}^{\infty} A_n(q - q_o)^n$$

The differential equation

$$(t - q^2)\frac{dt}{dq} = 1$$

then leads to the relationships

$$(t_o - q_o^2)(1A_1) = 1$$

$$(t_o - q_o^2)(2A_2) + (A_1 - 2q_o)(1A_1) = 0$$

$$(t_o - q_o^2)(3A_3) + (A_1 - 2q_o)(2A_2) + (A_2 - 1)(1A_1) = 0$$

$$(t_o - q_o^2)(4A_4) + (A_1 - 2q_o)(3A_3) + (A_2 - 1)(2A_2) + A_3(1A_1) = 0$$

The general term in the series can be evaluated from the relationship

$$(t_o - q_o^2)(nA_n) = -[(A_1 - 2q_o)(n-1)A_{n-1} + (A_2 - 1)(n-2)A_{n-2}]$$

$$- \sum_{m=3}^{n-1} (n - m)A_m A_{(n-m)} \qquad (2\text{-}83)$$

It would be rather straight foward to construct a computer program to evaluate the A_n. However, since the terms which appear in Eq. (2-82) are of the form

$$B_n = A_n(q - q_o)^n \tag{2-84}$$

it would advantageous to have the program evaluate B_n directly and to stop the summation when a sufficient number of terms have been summed. This is much better computer practice than to first construct the A_n and then go back and sum the polynomial which results when Eq. (2-82) is truncated.

Although it is wasteful of storage space in the computer, a simplication is obtained in programming this set of relationships if we define an auxiliary set of coefficients C_n by means of the following statements

$$U = (q - q_o)/(t_o - q_o^2) \tag{2-85}$$

$$C_1 = B_1 - 2q_oU \tag{2-86a}$$

$$C_2 = B_2 - U^2 \tag{2-86b}$$

$$C_n = B_n \quad , \quad n \geq 3 \tag{2-86c}$$

The recursion relationships then take the form

$$(t_o - q_o^2)(1B_1) = 0 \tag{2-89a}$$

$$(t_o - q_o^2)(2B_2) = - C_1B_1 \tag{2-89b}$$

$$(t_o - q_o^2)(3B_3) = - (2C_1B_2 + C_2B_1) \tag{2-89c}$$

or, in general

$$(t_o - q_o^2)(nB_n) = - \sum_{j=1}^{n-1} (n-j)C_jB_{n-j} \tag{2-89d}$$

Program 2-4

```
      SUBRØUTINE CMRTSM(QO,TO,Q,T,EPSI,CØNVRG)
      LØGICAL CØNVRG
      CØMPLEX QO,TO,Q,T,B(50),C(50),U,F,TEST
C------LØGICAL VARIABLE CØNVRG = .FALSE. IF SUM ØF LAST THREE
C------TERMS FAILS TØ BE LESS THAN EPSI*CABS(T).
      CØNVRG = .TRUE.
      U      = Q - QO
      F      = 1.0/(TO - QO*QO)
      B(1)   = U*F
      C(1)   = B(1) - 2.0*QO*U
      T      = TO + B(1)
      TEST   = B(1)
      B(2)   = -(F/FLØAT(2))*C(1)*B(1)
      C(2)   = B(2) - U*U
      T      = T + B(2)
      TEST   = TEST + B(2)
      B(3)   = -(FLØAT(3))*(2.0*C(1)*B(2) + C(2)*B(1))
      C(3)   = B(3)
      T      = T + B(3)
      TEST   = TEST + B(3)
      DØ 1 I = 4,50
      B(I)   = 0.0
      INDEX  = I - 1
      DØ 2 J = 1,INDEX
      B(I)   = B(I) + FLØAT(I-J)*C(J)*B(I-J)
    2 CØNTINUE
      B(I)   = - (F/FLØAT(I))*B(I)
      C(I)   = B(I)
      T      = T + B(I)
      TEST   = TEST + B(I) - B(I-3)
      TMAG   = EPSI*CABS(T)
      TESTMG = CABS(TEST)
      IF(TESTMG.LT.TMAG) GØ TØ 10
    1 CØNTINUE
      CØNVRG = .FALSE.
   10 RETURN
      END
```

In Program 2-4 we present a FORTRAN subroutine which will evaluate Eq. (2-82) by employing the recursion relation given in Eq. (2-89). The relation between the mathematical variables and the FORTRAN variables are

$$t_o = TO \quad , \quad t = T$$

$$q_o = QO \quad , \quad q = Q$$

The user will have to supply the value of EPSI which is used to control the exit from the DØ-loop. The author generally employed

EPSI = 1.0E-8

However, for some applications in which less accuracy is required a smaller value of EPSI will cut down on the amount of work which the computer is asked to perform. The LØGICAL variable CØNVRG is employed to permit the user to know whether the series converged when fewer than 50 terms were summed. The criteron for convergence to the accuracy specified by EPSI is based upon whether the last three terms combine to have an absolute value which is less than EPSI times the absolute value of T.

When q_o and q are very large, the representation in Eq. (2-82) becomes quite useless. When this is the case, one can replace Eq. (2-82) with the alternative form

$$t(q) = t_o + \sum_{n=1}^{\infty} B_n (q^{-1} - q_o^{-1})^n \qquad (2\text{-}90)$$

It will be convenient to define

$$x_o = \frac{1}{q_o} \quad , \quad x = \frac{1}{q}$$

and observe that Eq. (2-81) takes the form

$$(1 - x^2 t)\frac{dt}{dx} = 1 \qquad (2\text{-}91)$$

Let us express Eq. (2-90) in the form

$$t(x) = t_o + \sum_{n=1}^{\infty} B_n (x - x_o)^n \qquad (2\text{-}92)$$

If we insert Eq. (2-92) in Eq. (2-91) we can show that the recursion formula for the B_n is

$$(1-x_o^2 t_o)(n+1)B_{n+1} = \sum_{r=1}^{n} rB_r(B_{n-r-1} + 2x_o B_{n-r} + x_o^2 B_{n-r-1}) \quad (2\text{-}93)$$

The initial values are

$$B_{-1} = 0 \quad , \quad B_o = t_o \quad , \quad B_1 = \frac{1}{1 - x_o^2 t_o}$$

$$B_2 = \frac{2t_o x_o + x_o^2 - 2t_o^2 x_o^3}{2(1 - x_o^2 t_o)^3}$$

$$B_3 = \frac{2t_o + 4x_o + 2t_o^2 x_o^2 + 2t_o x_o^3 + (3 - 10t_o^3)x_o^4 - 6t_o^2 x_o^5 + 6t_o^4 x_o^6}{t_o(1 - x_o^2 t_o)^5}$$

For $q_o \to \infty$, we have $x_o = 0$. For this limiting case, the recursion formula in Eq. (2-93) is identical with Eq. (2-52).

Since the terms which occur in Eq. (2-92) are of the form

$$D_n = B_n(x - x_o)^n$$

it would be advantageous to program a recursion formula for the D_n and compute only as many of the D_n as are required to obtain the desired accuracy for $t(x)$. Let us define

$$U = x - x_o$$

$$A_1 = x_o^2 D_1 + 2x_o U t_o$$

$$A_2 = x_o^2 D_2 + 2x_o U D_1 + U^2 t_o^2$$

$$A_3 = x_o^2 D_3 + 2x_o U D_2 + U^2 D_1$$

and, in general

$$A_n = x_o^2 D_n + 2x_o U D_{n-1} + U^2 D_{n-2} \quad , \quad n > 3$$

We can then express Eq. (2-93) in the form

$$(1 - x_o^2 t_o)(n+1)D_{n+1} = \sum_{r=1}^{n} rD_r A_{n-r+1} \qquad (2\text{-}94)$$

The introduction of the A_n has greatly simplified the appearance of the equation that defines the recursion process by which the D_n are to be generated. Although the introduction of the A_n means that we must set up arrays in the computer storage for both the D_n and the A_n, the price paid for the using of computer storage space for the A_n is more than compensated for in the saving of the operations which would be repeated in each step of the recursion if we were to program the form in Eq. (2-93).

In Program 2-5 we present a FORTRAN subroutine which permits us to evaluate the sum in Eq. (2-92). The FORTRAN variables are related to the mathematical variables in the following manner:

$$x_o = XO \quad , \quad x = X$$

$$t_o = TO \quad , \quad t = T$$

The quantity EPSI plays the same role as the corresponding criterion in Program 2-4. The value

$$EPSI = 1.0E\text{-}8$$

can be employed in most applications of the program.

In Programs 2-6 and 2-7 we present double precision versions of Programs 2-4 and 2-5, respectively.

Program 2-6 provides the FORTRAN statements for

SUBRØUTINE RØØTSM(QOR,QOI,TOR,TOI,QR,QI,TR,TI,EPSI,CØNVRG)

The FORTRAN variables in the argument list are related to the

(Text continues on p. 2-99)

Program 2-5

```
      SUBRØUTINE CMRTLG(XO,TO,X,T,EPSI,CØNVRG)
      LØGICAL CONVRG
      CØMPLEX XO,TO,X,T,U,U2,C2XOU,XO2,F,D(50),A(50),TEST
C------LØGICAL VARIABLE CØNVRG = .FALSE. IF SUM ØF LAST THREE
C------TERMS FAILS TØ BE LESS THAN EPSI*CABS(T).
      CØNVRG = .TRUE.
      U      = X - XO
      U2     = U*U
      C2XOU  = 2.0*XO*U
      XO2    = XO*XO
      F      = 1.0/(1.0 - XO2*TO)
      T      = TO
      D(1)   = U*F
      T      = T + D(1)
      TEST   = D(1)
      A(1)   = C2XOU*TO + XO2*D(1)
      D(2)   = A(1)*D(1)*F/FLØAT(2)
      T      = T + D(2)
      TEST   = TEST + D(2)
      A(2)   = TO*U2 + C2XOU*D(1) + XO2*D(2)
      D(3)   = (2.0*A(1)*D(2) + A(2)*D(1))*F/FLØAT(3)
      T      = T + D(3)
      TEST   = TEST + D(3)
      A(3)   = U2*D(1) + C2XOU*D(2) + XO2*D(3)
      DØ 1 I = 4,50
      INDEX  = I - 1
      D(I)   = 0.0
      DØ 2 J = 1,INDEX
      D(I)   = D(I) + FLØAT(I-J)*A(J)*D(I-J)
    2 CØNTINUE
      D(I)   = D(I)*F/FLØAT(I)
      A(I)   = U2*D(I-2) + C2XOU*D(I-1) +XO2*D(I)
      T      = T + D(I)
      TEST   = TEST + D(I) - D(I-3)
      TMAG   = EPSI*CABS(T)
      TESTMG = CABS(TEST)
      IF(TESTMG.LT.TMAG) GØ TØ 10
    1 CØNTINUE
      CØNVRG = .FALSE.
   10 RETURN
      END
```

```
      SUBROUTINE ROOTSM(Q0R,Q0I,T0R,T0I,QR,QI,TR,TI,EPSI,CONVRG)
      LOGICAL CONVRG
      DOUBLE PRECISION  Q0R,Q0I,T0R,T0I,QR,QI,TR,TI,EPSI
      DOUBLE PRECISION  DFLOAT,DSQRT,BR(50),BI(50),CR(50),CI(50),UR,UI,
     A        FR,FI,TESTR,TESTI,TEST,TESTMG
     B
      DFLOAT(K) = DBLE(FLOAT(K))
      CONVRG = .TRUE.
      UR     = QR - Q0R
      UI     = QI - Q0I
      TR     = T0R - Q0R*Q0R + Q0I*Q0I
      TI     = T0I - 2.0D0*Q0R*Q0I
      FI     = TR*TR + TI*TI
      FR     = TR/FI
      FI     =-TI/FI
C-----(FR,FI) IS THE RECIPROCAL OF (T0 - Q0*Q0)
      BR(1)  = UR*FR - UI*FI
      BI(1)  = UR*FI + UI*FR
      CR(1)  = BR(1) - 2.0D0*(Q0R*UR - Q0I*UI)
      CI(1)  = BI(1) - 2.0D0*(Q0R*UI + Q0I*UR)
      TR     = T0R + BR(1)
      TI     = T0I + BI(1)
      TESTR  = BR(1)
      TESTI  = BI(1)
      CR(2)  = CR(1)*BR(1) - CI(1)*BI(1)
      CI(2)  = CR(1)*BI(1) + CI(1)*BR(1)
      BR(2)  =-(FR*CR(2) - FI*CI(2))/DFLOAT(2)
      BI(2)  =-(FR*CI(2) + FI*CR(2))/DFLOAT(2)
      CR(2)  = BR(2) - UR*UR + UI*UI
      CI(2)  = BI(2) - 2.0D0*UR*UI
      TR     = TR + BR(2)
      TI     = TI + BI(2)
      TESTR  = TESTR + BR(2)
      TESTI  = TESTI + BI(2)
```

```
      CR(3)  = (DFLOAT(2))*(CR(1)*BR(2) - CI(1)*BI(2)) + CR(2)*BR(1)
     A         - CI(2)*BI(1)
      CI(3)  = (DFLOAT(2))*(CR(1)*BI(2) + CI(1)*BR(2)) + CR(2)*BI(1)
     A         + CI(2)*BR(1)
      BR(3)  =-(FR*CR(3) - FI*CI(3))/DFLOAT(3)
      BI(3)  =-(FR*CI(3) + FI*CR(3))/DFLOAT(3)
      CR(3)  = BR(3)
      CI(3)  = BI(3)
      TR     = TR + BR(3)
      TI     = TI + BI(3)
      TESTR  = TESTR + BR(3)
      TESTI  = TESTI + BI(3)
      DO 1 I = 4,50
      CR(I)  = 0.0D0
      CI(I)  = 0.0D0
      INDEX  = I - 1
      DO 2 J = 1,INDEX
      CR(I)  = CR(I) + DFLOAT(I-J)*(CR(J)*BR(I-J) - CI(J)*BI(I-J))
      CI(I)  = CI(I) + DFLOAT(I-J)*(CR(J)*BI(I-J) + CI(J)*BR(I-J))
    2 CONTINUE
      BR(I)  =-(FR*CR(I) - FI*CI(I))/DFLOAT(I)
      BI(I)  =-(FR*CI(I) + FI*CR(I))/DFLOAT(I)
      CR(I)  = BR(I)
      CI(I)  = BI(I)
      TR     = TR + BR(I)
      TI     = TI + BI(I)
      TESTR  = TESTR + BR(I) - BR(I-3)
      TESTI  = TESTI + BI(I) - BI(I-3)
      TEST   = DSQRT(TESTR**2 + TESTI**2)
      TESTMG = DSQRT(TR**2 + TI**2)*EPSI
      IF(TEST.LT.TESTMG) GO TO 10
    1 CONTINUE
      CONVRG = .FALSE.
   10 RETURN
      END
```

```
      SUBROUTINE ROOTLG(X0R,X0I,T0R,T0I,XR,XI,TR,TI,EPSI,CONVRG)
      LOGICAL CONVRG
      DOUBLE PRECISION  X0R,X0I,T0R,T0I,XR,XI,TR,TI,EPSI
      DOUBLE PRECISION  DR(50),DI(50),AR(50),AI(50),TMAG,TESTR,TESTI,
     A                  DFLOAT,UR,UI,U2R,U2I,X02R,X02I,FR,FI,U2X0R,U2X0I
     B                 ,DSQRT,TEST,TEPSI
      DFLOAT(K) = DBLE(FLOAT(K))
      CONVRG = .TRUE.
      UR     = XR - X0R
      UI     = XI - X0I
      U2R    = UR*UR - UI*UI
      U2I    = 2.0D0*UR*UI
      U2X0R  = 2.0D0*(UR*X0R - UI*X0I)
      U2X0I  = 2.0D0*(UR*X0I + UI*X0R)
      X02R   = X0R*X0R - X0I*X0I
      X02I   = 2.0D0*X0R*X0I
      TESTR  = 1.0D0 - X02R*T0R + X02I*T0I
      TESTI  =       - X02R*T0I - X02I*T0R
      TMAG   = TESTR*TESTR + TESTI*TESTI
      FR     = TESTR/TMAG
      FI     =-TESTI/TMAG
      TR     = T0R
      TI     = T0I
      DR(1)  = UR*FR - UI*FI
      DI(1)  = UR*FI + UI*FR
      TR     = TR + DR(1)
      TI     = TI + DI(1)
      TESTR  = DR(1)
      TESTI  = DI(1)
      AR(1)  = U2X0R*T0R - U2X0I*T0I + X02R*DR(1) - X02I*DI(1)
      AI(1)  = U2X0R*T0I + U2X0I*T0R + X02R*DI(1) + X02I*DR(1)
      AR(2)  =(AR(1)*DR(1) - AI(1)*DI(1))/DFLOAT(2)
      AI(2)  =(AR(1)*DI(1) + AI(1)*DR(1))/DFLOAT(2)
      DR(2)  = AR(2)*FR - AI(2)*FI
      DI(2)  = AR(2)*FI + AI(2)*FR
      TR     = TR + DR(2)
      TI     = TI + DI(2)
      TESTR  = TESTR + DR(2)
```

```
      TESTI  = TESTI + DI(2)
      AR(2)  = U2R*T0R - U2I*T0I + U2X0R*DR(1) - U2X0I*DI(1)
     A                           + X02R*DR(2) - X02I*DI(2)
      AI(2)  = U2R*T0I + U2I*T0R + U2X0R*DI(1) + U2X0I*DR(1)
     A                           + X02R*DI(2) - X02I*DR(2)
      AR(3)  = (DFLOAT(2)*(AR(1)*DR(2) - AI(1)*DI(2)) + AR(2)*DR(1)
     A        - AI(2)*DI(1))/DFLOAT(3)
      AI(3)  = (DFLOAT(2)*(AR(1)*DI(2) + AI(1)*DR(2)) + AR(2)*DI(1)
     A        + AI(2)*DR(1))/DFLOAT(3)
      DR(3)  = AR(3)*FR - AI(3)*FI
      DI(3)  = AR(3)*FI + AI(3)*FR
      TR     = TR + DR(3)
      TI     = TI + DI(3)
      TESTR  = TESTR + DR(3)
      TESTI  = TESTI + DI(3)
      DO 1 I = 4,50
      INDEX  = I-1
      AR(I)  = 0.0D0
      AI(I)  = 0.0D0
      DO 2 J = 1,INDEX
      AR(I)  = AR(I) + DFLOAT(I-J)*(AR(J)*DR(I-J) - AI(J)*DI(I-J))
      AI(I)  = AI(I) + DFLOAT(I-J)*(AR(J)*DI(I-J) + AI(J)*DR(I-J))
    2 CONTINUE
      DR(I)  = (AR(I)*FR - AI(I)*FI)/DFLOAT(I)
      DI(I)  = (AR(I)*FI + AI(I)*FR)/DFLOAT(I)
      TR     = TR + DR(I)
      TI     = TI + DI(I)
      TESTR  = TESTR + DR(I) - DR(I-3)
      TESTI  = TESTI + DI(I) - DI(I-3)
      TEPSI  = DSQRT(TR**2 + TI**2)*DABS(EPSI)
      TEST   = DSQRT(TESTR**2 + TESTI**2)
      IF(TEST.LT.TEPSI) GO TO 10
      AR(I)  = X02R*DR(I) - X02I*DI(I) + U2X0R*DR(I-1) - U2X0I*DI(I-1)
     A        +U2R*DR(I-2) - U2I*DI(I-2)
      AI(I)  = X02R*DI(I) + X02I*DR(I) + U2X0R*DI(I-1) + U2X0I*DR(I-1)
     A        +U2R*DI(I-2) + U2I*DR(I-2)
    1 CONTINUE
      CONVRG = .FALSE.
   10 RETURN
      END
```

mathematical variables in the following manner:

$$q_o = QOR + iQOI \quad , \quad q = QR + iQI$$

$$t_o = TOR + iTOI \quad , \quad t = TR + iTI$$

The value to be assigned to EPSI will depend upon the accuracy which the user wishes to obtain. CØNVRG is a logical variable which tells whether or not the series converged to the desired accuracy by the time it has summed 50 terms. The user will have to supply the non-executable statements

```
LØGICAL CØNVRG
DØUBLE PRECISIØN QOR,QOI,TOR,TOI,QR,QI,TR,TI,EPSI
```

in the calling program.

The statements in Program 2-7 constitute the subroutine entitled

```
SUBRØUTINE RØØTLG(XOR,XOI,TOR,TOI,XR,XI,TR,TI,EPSI,CØNVRG)
```

The calling program will have to contain the statements

```
LØGICAL CØNVRG
DØUBLE PRECISIØN XOR,XOI,TOR,TOI,XR,XI,TR,TI,EPSI
```

The FORTRAN variables are related to the mathematical variables in the following manner:

$$x_o = XOR + iXOI \quad , \quad x = XR + iXI$$

$$t_o = TOR + iTOI \quad , \quad t = TR + iTI$$

Programs 2-6 and 2-7 were employed to generate the tables which are presented in Appendix D. In Program 2-8 we present a subroutine which we employed to generate tables such as the one in Table D-1a. The values in Table D-1a were obtained with a CALL statement of the form

```
CALL PRØPSM(1.018'929'164'471D0,45.0D0,0.05D0,41,TR,TI)
```

The reader will recognize the first argument to be the root β_1

defined by $Ai'(-\beta_1) = 0$.

It should be noted that Programs (2-4 through (2-7) can be used for step-by-step integration of the Riccati equation $(t-q^2)\frac{dt}{dq} = 1$ and that the calculations made in this manner (by employing a program such as in Program 2-8) do not have the boundaries of regions of applicability such as illustrated in Fig. 2-3.

Program 2-8

```
      SUBROUTINE PROPSM(T,ARG,QSTEP,N,TR,TI)
      LOGICAL CONVRG
      DOUBLE PRECISION T,ARG,QSTEP,TR(N),TI(N)
      DOUBLE PRECISION PI0180,COS60,SIN60,DFLOAT,Q0R,Q0I,EPSI,
     A   RAD,T0R,T0I,QR,QI,TREAL,TIMAG,DCOS,DSIN
      DATA PI0180,COS60,SIN60/0.17453292519943   D-1, 0.5D0,
     A                        0.86602540378443865D0 /
      DFLOAT(K) = DBLE(FLOAT(K))
      TR(1)  = T*COS60
      TI(1)  = T*SIN60
      Q0R    = 0.0D0
      Q0I    = 0.0D0
      EPSI   = 1.0D-17
      RAD    = PI0180*ARG
      DO 1 I = 2,N
      INDEX  = I
      T0R    = TR(I-1)
      T0I    = TI(I-1)
      QI     = QSTEP*DFLOAT(I-1)
      QR     = QI*DCOS(RAD)
      QI     = QI*DSIN(RAD)
      CALL ROOTSM(Q0R,Q0I,T0R,T0I,QR,QI,TREAL,TIMAG,EPSI,CONVRG)
      TR(I)  = TREAL
      TI(I)  = TIMAG
      Q0R    = QR
      Q0I    = QI
      IF(.NOT.CONVRG) GO TO 2
    1 CONTINUE
      RETURN
    2 WRITE(6,150) INDEX,T,ARG,QSTEP
      DO 9 I = INDEX,N
      TR(I)  = 0.0D0
      TI(I)  = 0.0D0
    9 CONTINUE
      RETURN
  150 FORMAT(46H0  SUBROUTINE PROPSM COULD NOT HANDLE INDEX = ,I4/
     A 11H  WITH T = ,D22.16,8H  ARG = ,D22.16,10H  QSTEP = ,D22.16)
      END
```

REFERENCES FOR SECTION 2

1. V. A. Fock, Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965. (This is a collection of 17 papers published by Fock between 1945 and 1963. The appendix is devoted to a discussion of the Airy function and a table is given for u(t) and v(t) for real values of t.)

2. J. R. Wait, "Electromagnetic Surface Waves," Advances in Radio Research, J. A. Saxton, ed., London, Academic Press, 1964, pp. 157-215. (This survey paper will provide the reader with an excellent introduction to the subject matter and the bibliography can be used to lead the reader to many classic papers, as well as to some of the numerous papers published by Wait during the past decade.)

3. Lockheed Missiles and Space Division, General Research in Diffraction Theory, Vol. 1, by N. A. Logan, LMSD-288087, Sunnyvale, Calif., Dec., 1959

4. N. A. Logan and K. S. Yee, "A Mathematical Model for Diffraction by Convex Surfaces," Electromagnetic Waves, R. E. Langer, Ed., Madison, Wisc., Univ. of Wisconsin Press, 1962

5. J. C. P. Miller, The Airy Integral, British Association Mathematical Tables, Part-Volume B, Cambridge, Cambridge University Press, 1946

6. B. van der Pol and H. Bremmer, "Diffraction of Electromagnetic Waves from an Electrical Point Source Round a Finitely Conducting Sphere with Applications of Radio-Telegraphy and the Theory of the Rainbow," Part I: Phil. Mag., Vol. 24, 1937, pp. 141-176; Part II: Phil. Mag., Vol. 24, 1937, pp. 825-864; Part III: Phil. Mag. , Vol. 25, 1938, pp. 817-837

7. Harvard University Computation Laboratory, Tables of the Modified Hankel Functions of Order One-Third and of Their Derivatives, Annals of the Computation Laboratory of Harvard University, Vol. II, Cambridge, Mass., Harvard University Press, 1945. (The "Introduction" of this book was written by W. E. Furry and H. A. Arnold.)

8. H. Bremmer, Terrestrial Radio Waves: Theory of Propagation, New York, Elsevier, 1949

9. C.C.I.R. Resolution No. 11, Atlas of Ground-Wave Propagation Curves for Frequencies between 30 Mc/s and 300 Mc/s," Geneva, Union Internationale des Télécommunications, 1955. (This Atlas was prepared by the Mathematical Centre, Amsterdam, under the supervision of A. van Wyngaarden and B. van der Pol.)

10. H. Jeffreys, "The Effect on Love Waves of Heterogeneity in the Lower Layer," Monthly Notices Roy. Astro. Soc., Geophys. Suppl., Vol. 2, 1928, pp. 101-111

11. Air Force Cambridge Research Center, The Zeros and Maxima of the Airy Function and Its First Derivative to 25 Significant Figures, by M. E. Sherry, Report AFCRC-TR-59-135, ASTIA Document No. AD 214568, Bedford, Mass., April, 1959

12. R. S. Elliott, "Spherical Surface Wave Antennas," IRE Trans. Antennas and Propagation, Vol. AP-4, 1956, pp. 422-428

13. K. A. Norton, "The Calculation of Ground-Wave Intensity over a Finitely Conducting Spherical Earth," Proc. IRE, Vol. 5, Dec., 1941, pp. 623-639

14. H. H. Howe, "Note on the Solution of Riccati's Differential Equation," J. Research NBS, Vol., 64B, Apr.-June, 1960, pp. 95-98

15. J. R. Johler, L. C. Walters, and C. M. Lilley, Low- and Very-Low Radiofrequency Tables of Ground Wave Parameters for the Spherical Earth Theory: The Roots of Riccati's Differential Equation, NBS Technical Note No. 7 , Washington, Office of Technical Services, U. S. Dept. Commerce, Feb., 1959

16. E. M. Gyunninen, G. I. Makarov, V. V. Novikov, and S. T. Rybachek, "Propagation of Electromagnetic Pulses and Their Harmonics Over the Earth's Surface," Problems of Wave-Diffraction and Propagation, Vol. III, E. M. Gyunninen, ed., Leningrad, Publishing House of Leningrad State University, 1964. (A translation can be purchased from the Joint Publications Research Service, OTS, U. S. Dept. Commerce, Report No. JPRS:26,487,TT:64-41745, 22 Set., 1964.)

17. M. G. Belkina, Tables to Calculate the Electromagnetic Field in the Shadow Region for Various Soils, Moscow, Soviet Radio Press, 1949. (A translation of the introductory material was made by M. D. Friedman and made available to the scientific public by the Air Force Cambridge Research Center in November, 1956. The ASTIA Document No. is AD 110298.)

18. I. H. Gerks, "Use of a High-Speed Computer for Ground-Wave Calculations," IRE Trans. Antennas and Propagation, Vol. AP-10, May, 1962, pp. 292-299.

19. C. Domb, "Tables of Functions Occurring in the Diffraction of Electromagnetic Waves by the Earth," Advances in Physics, Quarterly Supplement of Phil. Mag., Vol. 5, 1953, pp. 96-102

20. N. S. Kapany, J. J. Burke, and K. Frame, "Diffraction by Apertures of Wavelength Dimensions," Applied Optics, Vol. 4, Oct., 1965, pp. 1229-1238. (Consult at later issue of this journal for an errata which is currently "in press.")

21. W. Streifer and R. D. Kodis, "On the Solution of a Transcendental Equation Arising in the Theory of Scattering by a Dielectric Cylinder," Quart. Appl. Math., Vol. 21, Jan., 1964, pp. 285-298

22. F. W. J. Olver, "The Asymptotic Expansion of Bessel Functions of Large Order," Trans. Roy. Soc. (London), Vol. 247A, 1954, pp. 328-368

23. The Radio Research Laboratories, Atlas of Radio Wave Propagation Curves for Frequencies Between 30 and 10,000 Mc/s, Tokyo, Ministry of Postal Services, 1957

24. M. H. L. Pryce, "The Diffraction of Radio Waves by the Curvature of the Earth," Advances in Physics, Quarterly Supplement of Phil. Mag., Vol. 5, 1953, pp. 96-102

25. C. Domb and M. H. L. Pryce, "The Calculation of Field Strengths Over a Spherical Earth," J.I.E.E., Vol. 94, Part III, 1947, pp. 325-339

Section 3

EXPANSIONS IN THE LIGHTED REGION OF A CIRCULAR CYLINDER

Consider the problem which arises when one wants to compute the field in the lighted region when the plane wave $\exp(-ikx - i\omega t)$ illuminates a perfectly conducting cylinder of radius a. If the radius of the cylinder is much greater than the wavelength, and if the field is observed for points outside the shadow region, the field can be described by the optical approximations

$$u(\rho,\varphi) \sim \exp(-ikx - i\omega t) \pm \sqrt{\frac{a\cos\alpha}{2D + a\cos\alpha}}\ \exp\left[ik(D - a\cos\alpha) - i\omega t\right] \qquad (3\text{-}1)$$

where the + sign is used when the magnetic vector is parallel to the axis of the cylinder and $U(\rho,\varphi)$ represents the magnetic field (the so-called Neuman problem since the derivative of the wave function must vanish at the surface) and the - sign is used when the electric vector is parallel to the axis of the cylinder and $U(\rho,\varphi)$ represents the electric field (the so-called Dirichlet problem since the wave function must vanish at the surface). The geometry is illustrated in Fig. 3-1. The angle α is the angle of




Fig. 3-1 Geometry of the Reflection Problem

incidence of the wave at the point of reflection, and the distance D is the distance of the observation point (ρ,φ) from the point of reflection.

In the following discussion we will present several extensions of the optical approximation which are based upon the diffraction function

$$V_1(\xi,\zeta,q) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t) \left[v(t-\zeta) - \frac{v'(t) - q\, v(t)}{w_1'(t) - q\, w_1(t)}\, w_1(t-\zeta) \right] dt \quad (3\text{-}2)$$

which has been employed by Fock (Ref. 1), Belkina and Vainshtein (Ref. 2), Azriliant and Belkina (Ref. 3), Fedorov (Ref. 4), Wait (Ref. 5), Goodrich (Ref. 6), Logan and Yee (Ref. 7), and many other authors. Fock showed that for $\sqrt{\zeta} - \xi > 0$ that $V_1(\xi,\zeta,q)$ can be expressed in the form

$$V_1(\xi,\zeta,q) = \exp\left[i\left(\xi\zeta - \frac{\xi^3}{3}\right)\right] + P(\xi,\zeta,q) \quad (3\text{-}3)$$

$$P(\xi,\zeta,q) = -\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(i\xi t)\, \frac{v'(t) - q\, v(t)}{w_1'(t) - q\, w_1(t)}\, w_1(t-\zeta)\, dt \quad (3\text{-}4)$$

and that $P(\xi,\zeta,q)$ can be shown to have the asymptotic property

$$P(\xi,\zeta,q) \xrightarrow[p>>1]{} -\frac{q - ip}{q + ip}\sqrt{\frac{p}{\sigma}}\, \exp\left[i\, \frac{1}{12}\left(\sigma^3 + 3\sigma^2 p - 3\sigma p^2 + 3p^3\right)\right] \quad (3\text{-}5)$$

where

$$p = \frac{1}{3}\left[\sqrt{\xi^2 + 3\zeta} - 2\xi\right], \qquad \sigma = \sqrt{\xi^2 + 3\zeta} \quad (3\text{-}6)$$

Let us express $P(\xi,\zeta,q)$ in the form of a product of a slowly varying complex amplitude and the phase factor given appearing in Eq. (3-5).

$$P(\xi,\zeta,q) = A(p,\sigma,q)\exp\left[i\frac{1}{12}(\sigma^3 + 3\sigma^2 p - 3\sigma p^2 + 3p^3)\right] \tag{3-7}$$

Since Fock has shown that $P(\xi,\zeta,q)$ is a solution of

$$\frac{\partial^2 P}{\partial \zeta^2} + i\frac{\partial P}{\partial \xi} + \zeta P = 0 \tag{3-8}$$

we can show that $A(p,\sigma,q)$ is a solution of

$$\frac{1}{4\sigma^2}\frac{\partial^2 A}{\partial p^2} + \frac{9}{4\sigma^2}\frac{\partial^2 A}{\partial \sigma^2} + \frac{3}{2\sigma^2}\frac{\partial^2 A}{\partial \sigma \partial p} - \frac{3}{4\sigma^3}\frac{\partial A}{\partial p} + \left(2i - \frac{9}{4\sigma^3}\right)\frac{\partial A}{\partial \sigma} + \frac{i}{\sigma}A = 0 \tag{3-9}$$

In order to see how $A(p,\sigma,q)$ can be used to describe the reflected wave from the circular cylinder, we can start from a form of the Laplace operator which was used by Keller, Lewis, and Seckler (Ref. 8) to study a slightly different aspect of this problem. The form which we need is

$$\nabla^2 = \left(1 + \frac{u}{s^2}\right)\frac{\partial^2}{\partial s^2} + \left(\frac{1}{s} - \frac{w}{s^2} - \frac{u}{s^3}\right)\frac{\partial}{\partial s} - \frac{2v}{s^2}\frac{\partial^2}{\partial s \partial \beta} + \frac{1}{s^2}\frac{\partial^2}{\partial \beta^2} + \frac{v}{s^3}\frac{\partial}{\partial \beta} \tag{3-10}$$

where

$$s = D + \frac{a}{2}\cos\alpha \qquad\qquad \beta = \pi - 2\alpha$$

and

$$u = \frac{9}{16}a^2\cos^2\beta/2 \qquad v = -\frac{3}{4}a\cos\beta/2 \qquad w = \frac{3}{8}a\sin\beta/2$$

If we make a change of scale from α to σ and from $\sin(\beta/2)$ to p

$$p = \left(\frac{ka}{2}\right)^{1/3}\sin\beta/2 \qquad\qquad \sigma = 2\left(\frac{ka}{2}\right)^{1/3}\frac{s}{a} \tag{3-11}$$

the wave equation takes the form

$$\left(\nabla^2 + k^2\right)\Phi = \left[\left(\frac{ka}{2}\right)^{4/3}L_0 + \left(\frac{ka}{2}\right)^{2/3}L_1 + \left(\frac{ka}{2}\right)^{0}L_2\right]\Phi \tag{3-12}$$

where L_0 is the differential operator

$$L_0 = \frac{9}{\sigma^2}\frac{\partial^2}{\partial\sigma^2} + \frac{1}{\sigma^2}\frac{\partial^2}{\partial p^2} + \frac{6}{\sigma^2}\frac{\partial^2}{\partial\sigma\partial p} - \frac{3}{\sigma^3}\frac{\partial}{\partial p} + \left(8i - \frac{9}{\sigma^3}\right)\frac{\partial}{\partial\sigma} + \frac{4i}{\sigma} \qquad (3\text{-}13)$$

and L_1 and L_2 are similar differential operators which involve only σ and ρ. If we assume that $\Phi(\rho,\varphi)$ can be expressed in the form of an asymptotic series

$$\Phi(\rho,\sigma) = \left(\sum_{n=0}^{\infty} \Phi_n(p,\sigma)\ (ka/2)^{-\frac{2}{3}n}\right) \exp[ik(D - a\cos\alpha)] \qquad (3\text{-}14)$$

we then find that

$$L_0\Phi_0 = 0$$

so that Φ_0 satisfies the same partial differential equation as does $A(p,\sigma,q)$. An examination of the boundary conditions reveals that we can identify Φ_0 with the slowly-varying function $A(p,\sigma,q)$.

$$\Phi_0(p,\sigma) = A(p,\sigma,q) \qquad (3\text{-}15)$$

Therefore, we can express $U(\rho,\varphi)$ in the form

$$u(\rho,\varphi) \sim \exp(-ikx - i\omega t) + A(p,\sigma,q)\exp\left[ik(D - a\cos\alpha) - i\omega t\right] \qquad (3\text{-}16)$$

provided the higher-order terms in $(ka/2)^{-\frac{2}{3}}$ are neglectable. Since

$$A(p,\sigma,q) \xrightarrow[p>>1]{} -\frac{q-ip}{q+ip}\sqrt{\frac{p}{\sigma}} = -\frac{q-ip}{q+ip}\sqrt{\frac{a\cos\alpha}{2D + a\cos\alpha}} \qquad (3\text{-}17)$$

we can express the approximation for the case when the magnetic vector is parallel to the axis of the cylinder in the form

$$H_z(\rho,\varphi) \sim \exp(-ikx - i\omega t) + A(p,\sigma,0)\exp\left[ik(D - a\cos\alpha) - i\omega t\right] \qquad (3\text{-}18)$$

and the case when the electric vector is parallel to the axis in the form

$$E_z(\rho,\varphi) \sim \exp(-ikx - i\omega t) + A(p,\sigma,\infty)\exp\left[ik(D - a\cos\alpha) - \right] \qquad (3\text{-}19)$$

If we assume that $A(p,\sigma,q)$ possesses an asymptotic expansion of the form

$$A(p,\sigma,q) = \sqrt{\frac{p}{\sigma}} \sum_{n=0}^{\infty} \frac{A_n(p,q)}{\sigma^n} \tag{3-20}$$

then the differential equation Eq. (3-9) leads to the recursion formulae

$$A_{n+1}(p,q) = -\frac{1}{2(n+1)} \left\{ \left[\frac{1}{4} \frac{d^2 A_n}{dp^2} + \frac{1}{4p} \frac{dA_n}{dp} - \frac{1}{16p^2} A_n \right] - \left[\frac{3}{2} n - \frac{dA_{n-1}}{dp} + \frac{3n}{4p} A_{n-1} \right] + \left[\frac{45}{16} + \frac{9}{4} (n-2)(n+1) A_{n-2} \right] \right\} \tag{3-21}$$

for which the initial conditions are

$$A_0(p,q) = \sqrt{2/p}\ V_{11}(-2p,q) \exp(-i\tfrac{2}{3}p^3),\ A_m = 0,\ m < 0 \tag{3-22}$$

Logan (Ref. 9) has shown that $V_{11}(\xi,0)$ and $V_{11}(\xi,\infty)$ can be expanded in asymptotic series of the form

$$V_{11}(\xi,0) \underset{\xi \to -\infty}{\longrightarrow} -\frac{\sqrt{-\xi}}{2} \exp\left(-i\frac{\xi^3}{12}\right) \left\{ 1 + i\frac{2}{\xi^3} - \frac{28}{\xi^6} - i\frac{896}{\xi^9} + \frac{43120}{\xi^{12}} + i\frac{2754752}{\xi^{15}} + \cdots \right\} \tag{3-23}$$

$$V_{11}(\xi,\infty) \underset{\xi \to -\infty}{\longrightarrow} \frac{\sqrt{-\xi}}{2} \exp\left(-i\frac{\xi^3}{12}\right) \left\{ 1 - i\frac{2}{\xi^3} + \frac{20}{\xi^6} + i\frac{560}{\xi^9} - \frac{25520}{\xi^{12}} - i\frac{1601600}{\xi^{15}} + \cdots \right\} \tag{3-24}$$

In Tables 3-1 and 3-2 we list some of the leading terms in these asymptotic expansions. The author has tables which contain further terms in these series, but until appropriate summation techniques for summing these types of expansions can be found their usefulness is limited.

Table 3-1

LEADING TERMS IN THE ASYMPTOTIC EXPANSION OF $A_n(p,0)$

$$A_0(p,0) = -1 + \frac{1}{4}\frac{1}{p^3} + \frac{7}{16}\frac{1}{p^6} - i\frac{7}{4}\frac{1}{p^9} + \cdots$$

$$A_1(p,0) = -i\frac{1}{32}\frac{1}{p^2} + \frac{35}{128}\frac{1}{p^5} - i\frac{1001}{512}\frac{1}{p^8} + \cdots$$

$$A_2(p,0) = -i\frac{3}{16}\frac{1}{p} + \frac{465}{2048}\frac{1}{p^4} - i\frac{10857}{8192}\frac{1}{p^7} + \cdots$$

$$A_3(p,0) = +i\frac{15}{32}\frac{1}{p^0} + \frac{45}{512}\frac{1}{p^3} - i\frac{63525}{65536}\frac{1}{p^6} + \cdots$$

$$A_4(p,0) = -\frac{105}{1024}\frac{1}{p^2} - i\frac{28665}{32768}\frac{1}{p^5} + \cdots$$

$$A_5(p,0) = -\frac{315}{512}\frac{1}{p} - i\frac{45675}{65536}\frac{1}{p^4} + \cdots$$

$$A_6(p,0) = +\frac{3465}{2048}\frac{1}{p^0} - i\frac{3465}{16384}\frac{1}{p^3} + \cdots$$

$$A_7(p,0) = +i\frac{45045}{65536}\frac{1}{p^2} + \cdots$$

$$A_8(p,0) = +i\frac{135135}{32768}\frac{1}{p} + \cdots$$

$$A_9(p,0) = -i\frac{765765}{65536}\frac{1}{p^0} + \cdots$$

Table 3-2

LEADING TERMS IN THE ASYMPTOTIC EXPANSION OF $A_n(p,\infty)$

$$A_0(p,\infty) = 1 + i\frac{1}{4}\frac{1}{p^3} + \frac{5}{16}\frac{1}{p^6} - i\frac{35}{32}\frac{1}{p^9} + \ldots$$

$$A_1(p,\infty) = + i\frac{1}{32}\frac{1}{p^2} + \frac{35}{128}\frac{1}{p^5} - i\frac{715}{512}\frac{1}{p^8} + \ldots$$

$$A_2(p,\infty) = + i\frac{3}{16}\frac{1}{p} + \frac{495}{2048}\frac{1}{p^4} - i\frac{8745}{8192}\frac{1}{p^7} + \ldots$$

$$A_3(p,\infty) = - i\frac{15}{32}\frac{1}{p^0} + \frac{75}{512}\frac{1}{p^3} - i\frac{60315}{65536}\frac{1}{p^6} + \ldots$$

$$A_4(p,\infty) = + \frac{105}{1024}\frac{1}{p^2} - i\frac{30135}{32768}\frac{1}{p^5} + \ldots$$

$$A_5(p,\infty) = + \frac{315}{512}\frac{1}{p} - i\frac{55125}{65536}\frac{1}{p^4} + \ldots$$

$$A_6(p,\infty) = - \frac{3465}{2048}\frac{1}{p^0} - i\frac{10395}{16384}\frac{1}{p^3} + \ldots$$

$$A_7(p,\infty) = - i\frac{45045}{65536}\frac{1}{p^2} + \ldots$$

$$A_8(p,\infty) = - i\frac{135135}{32768}\frac{1}{p} + \ldots$$

$$A_9(p,\infty) = + i\frac{765765}{65536}\frac{1}{p^0} + \ldots$$

Our present state of knowledge of the type of series which are displayed in Tables 3-1 and 3-2 is so limited that we do not know how to "sum" the series when $p \to 0$. This type of expansion can only be used when p is very large and positive, which corresponds to the physical situation of being well above the horizon in the lighted region, and the radius of curvature of the cylinder being very large in comparison with the wavelength. The author has had some limited success with obtaining numerical results from these series by employing a special form of the Euler transformation, namely,

$$S = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \ldots$$
$$= (1 - x)^{-1} \left(a_0 + \sum_{k=1}^{\infty} [x/(1 - x)]^k \, \Delta^k a_0 \right) \qquad (3\text{-}25)$$

where

$$\Delta a_0 = a_1 - a_0$$
$$\Delta^2 a_0 = \Delta a_1 - \Delta a_0$$
$$\ldots\ldots\ldots$$

Since the series are characterized by the appearance of factors such as signs +, i, -1, -i, etc, the author has found it most useful to let $\exp(i\pi/2)$ be a factor of the variable x which appears in Eq. (3-25). Although the author has been able to use methods of this type to extrapolate values from the asymptotic series to much smaller value than can be obtained by a straightfoward use of the asymptotic series, the method does not appear useful enough at the present state of its development to warrant a further discussion at this time.

In order to obtain practical results for smaller values of p, it is much simpler to reconsider Eq. (3-22) and to realize that the function $V_{11}(-2p,q)$ offers a means of obtaining approximate values of $A(p,\sigma,q)$ provided σ is very large. Since $V_{11}(-2p,q)$ can be

numerically integrated by employing an integral representation such as that in Eq. (1-50), a more attractive approach to the computation of $A(p,\sigma,q)$ would appear to be to use Eq. (3-21) to find the $A_n(p,q)$ by means of numerical integration. Work along these lines has been begun by the author but the progress to date has been too limited to be able to report at this time upon the usefulness of this method. Furthermore, since Eq. (3-20) is an asmptotic series in inverse powers of σ, the usefulness of this representation (for small values of p) may be so limited that these studies may be more "academic" than "practical."

However, there are some very practical aspects of Eq. (3-16) which should be discussed at this point. When Eq. (3-22) is inserted into Eq. (3-20) and Eq. (3-16), there remains the limitation that p must be positive. It is relatively easy to eliminate this restriction and obtain a representation which holds down to $p = 0$ (i.e., on the horizon) and even to points slightly below the horizon. In order to achieve this result, we need to return to Eq. (3-7) in which the function $A(p,\sigma,q)$ was defined in terms of the function $P(\xi,\zeta,q)$. From Eq. (3-6), we find that the inverse relations are

$$2\xi = \sigma - 3p \quad , \qquad 4\zeta = \sigma^2 + 2\sigma p - 3p^2 \tag{3-26}$$

We can then show that for large positive values of σ,

$$\frac{2}{3}\zeta^{3/2} \longrightarrow \frac{1}{12}(\sigma^3 + 3\sigma^2 p - 3\sigma p^2 + 3p^3) - \frac{2}{3}p^3 \tag{3-27}$$

We now want to use an asymptotic property of Eq. (3-2) in order to show that in the vicinity of the "horizon" (defined as $\xi - \sqrt{\zeta} = 0$) that one can obtain an extension of Eq. (3-16) which shows that the field in this region is very closely related to the classical Fresnel integral representation for the field behind an opaque screen. The asymptotic formulae which contain this result are

described by the relations

$$V_1(\xi,\zeta,q) - \exp[i(\xi\zeta - \tfrac{1}{3}\xi^3)] \xrightarrow[\substack{\zeta \to \infty \\ \xi < 0}]{} F(\zeta)[-\mu K(-\tau) + V_2(x,q)] \quad (3\text{-}28)$$

$$V_1(\xi,\zeta,q) \xrightarrow[\substack{\zeta \to \infty \\ \xi > 0}]{} F(\zeta)[\mu K(\tau) + V_2(x,q)] \quad (3\text{-}29)$$

where the parameters μ, x, and τ are defined by

$$\mu = \zeta^{1/4} \quad , \quad x = \xi - \sqrt{\zeta} \quad , \quad \tau = \mu x$$

and $F(\zeta)$ denotes the factor

$$F(\zeta) = [\zeta^{-1/4}] \exp(i\tfrac{2}{3}\zeta^{3/2})$$

and $K(\tau)$ denotes a special form of the familiar Fresnel integrals

$$K(\tau) = \exp[-i(\tau^2 + \pi/4)] \frac{1}{\sqrt{\pi}} \int_\tau^\infty \exp(is^2)ds \quad (3\text{-}30)$$

which has the properties

$$K(0) = \frac{1}{2}$$

$$K(\tau) \xrightarrow[\tau \to \infty]{} \frac{\exp(i\pi/4)}{2\sqrt{\pi}\,\tau}$$

The function $V_2(x,q)$ has already been defined by Eq. (1-51) in which it is related to a function $V_{11}(x,q)$ which was defined in Eq. (1-50). From these former results we see that we can express $V_2(x,q)$ in the form

$$V_2(x,q) = -\frac{\exp(i\pi/4)}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(ixt) \frac{v'(t) - q\,v(t)}{w_1'(t) - q\,w_1(t)}\,dt - \frac{\exp(i\pi/4)}{2\sqrt{\pi}\,x} \quad (3\text{-}31)$$

However, this is not a convenient form for numerical evaluation since the portion taken along the negative t-axis gives rise to a singular function of x which is exactly cancelled off by the factor involving the reciprocal of x. Since $V_2(x,q)$ is actually

an entire function of x, it can actually be developed in a Taylor series about x = 0 if one starts from a proper representation. One of the most convenient representations for use with numerical integration techniques is obtained by making use of the property of the Airy functions which assert that

$$w_1(t\exp(i\tfrac{2}{3}\pi)) = \exp(i\tfrac{1}{3}\pi)w_2(t)$$

$$v(t\exp(i\tfrac{2}{3}\pi)) = \frac{1}{2}\exp(-i\pi/6)w_1(t)$$

where, for the sake of the present discussion, we can assume that t is real and positive. These properties permit us to deform the section of the contour in Eq. (3-31) that runs from $-\infty$ to 0 to a contour that runs from $\infty\exp(i\frac{2}{3}\pi)$ to 0 and arrive at the representation

$$V_2(x,q) = -\frac{\exp(i\pi/4)}{\sqrt{\pi}}\left[\int_0^\infty \exp(ixt)\frac{v'(t) - qv(t)}{w_1'(t) - qw_1(t)}\,dt + \exp(-i\tfrac{1}{3}\pi)\int_0^\infty \exp\left[-xt\frac{(\sqrt{3}+i)}{2}\right]\frac{v'(t) - q^*v(t)}{w_2'(t) - q^*w_2(t)}\,dt\right] \quad (3\text{-}32)$$

where

$$q^* = q\exp(i\tfrac{2}{3}\pi)$$

and $w_1(t)$ and $w_2(t)$ are related to the real-valued functions (i.e., real-valued when t is real) u(t) and v(t) by means of the relations

$$w_1(t) = u(t) + i\,v(t)$$

$$w_2(t) = u(t) - i\,v(t)$$

Let us now collect some of the properties of Eq. (3-16) which can be deduced from the results which we have been discussing.

We observe that for points "high" above the horizon (i.e., for $p > 0$ and $\sigma \to \infty$) that Eq. (3-22) leads to

$$U(\rho,\varphi) - \exp(-ikx - i\omega t) \approx$$

$$\sqrt{2/\sigma}\; V_{11}(-2p,q)\exp(-i\tfrac{2}{3}p^3)\exp[ik(D - a\cos\alpha) - i\omega t] \qquad (3\text{-}33)$$

If we then let $p \to \infty$, we arrive at the asymptotic approximation

$$u(\rho,\varphi) \sim \exp(-ikx - i\omega t) - \frac{q-ip}{q+ip}\sqrt{\frac{p}{\sigma}}\exp\left[ik(D-a\cos\alpha) - i\omega t\right] \qquad (3\text{-}34)$$

which is the extension to the impedance boundary condition

$$\frac{\partial U}{\partial \rho} = ikZU, \quad \text{for } \rho = a$$

$$q = -i(ka/2)^{\frac{1}{3}}Z$$

of the special problems which were expressed in Eq. (3-1) since the introduction of the definitions of p and σ change the appearance of (3-34) to the following form

$$u(\rho,\varphi) \sim \exp(-ikx - i\omega t) - \frac{q-ip}{q+ip}\sqrt{\frac{a\cos\alpha}{2D + a\cos\alpha}}\exp\left[ik(D-a\cos\alpha) - i\omega t\right] \qquad (3\text{-}35)$$

However, the analytical "tools" which we have developed by the introduction of the functions $V_{11}(x,q)$ and $V_2(x,q)$ permit us to use either Eq. (3-28) or Eq. (3-29) to arrive at a result which holds <u>on the horizon</u>, i.e., for $p = 0$, for the case when $\sigma \to \infty$. This result is of the form

$$U(\rho,\varphi) \xrightarrow[p\to 0]{} \left\{\frac{1}{2} + \left(\frac{ka}{2}\right)^{1/3}\sqrt{\frac{2}{kD}}\; V_2(0,q)\right\}\exp(-ikx - i\omega t) \qquad (3\text{-}36)$$

The extensions to the optical results which were made in the paper by Keller, Lewis, and Seckler (Ref. 8) belong to the class of results of which the expansions in Tables 3-1 and 3-2 are examples. However, these representations (which are effectively in inverse

integral powers of (ka) become useless for $p \to 0$. These new results have the considerable advantage of being useful for $p \to 0$ as well as $p \to \infty$.

The results which we have obtained provide us with an interesting means of obtaining the fields at the surface of the cylinder. If we consider the meanings of p and σ, we see that the fields on the surface of the cylinder are obtained by letting $\sigma \to p$. From Eq. (3-16) we see that

$$\begin{aligned} U(a,\varphi) &= [1 + A(p,p,q)] \exp(-ika\cos\varphi - i\omega t) \\ &= V_1(-p,q) \exp[-i\tfrac{1}{3}p^3 - ika\cos\varphi - i\omega t] \end{aligned} \tag{3-37}$$

where $V_1(z,q)$ (which was already introduced in Eq. (1-31) is defined by means of either a Fourier integral

$$V_1(z,q) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \exp(izt) \frac{1}{w_1'(t) - q w_1(t)} dt \tag{3-38}$$

or a residue series

$$V_1(z,q) = i2\sqrt{\pi} \sum_{s=1}^{\infty} \frac{\exp(i z t_s)}{(t_s - q^2) w_1(t_s)} = -i2\sqrt{\pi}\, \frac{1}{q} \sum_{s=1}^{\infty} \frac{\exp(i z t_s)}{\left(1 - \frac{t_s}{q^2}\right) w_1'(t_s)} \tag{3-39}$$

where the roots t_s are solutions of the transcendental equation

$$w_1'(t_s) - q w_1(t_s) = 0 .$$

Let us employ these results for the case of the electromagnetic scattering problems which we considered in Eqs. (3-18) and (3-19). When the magnetic field of the incident plane wave is parallel to the axis of the cylinder, we obtain the magnetic field at the surface of the cylinder by considering the case $\sigma \to p$. For this case we obtain

$$H_z(a,\varphi) = \left[1 + A(p,p,0)\right] \exp(-ika\cos\varphi - i\omega t)$$
$$= G(-p) \exp\left[-ika\cos\varphi - i\omega t\right] \qquad (3\text{-}40)$$

where G(-p) is the current distribution function

$$G(-p) = \exp\left(-i\frac{p^3}{3}\right)\frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty}\frac{\exp(-ipt)}{w_1'(t)}\,dt \qquad (3\text{-}41)$$

which has the asymptotic property that as $p \to \infty$

$$G(-p) \approx 2\left\{1 - i\frac{1}{4p^3} - \frac{1}{p^6} - i\frac{469}{64}\frac{1}{p^9} + \frac{5005}{64}\frac{1}{p^{12}} + i\frac{1122121}{1024}\frac{1}{p^{15}} + \ldots\right\} \qquad (3\text{-}42)$$

Since the tangential component of the electric field vanishes on the surface of the cylinder, we observe that $V_1(-p, 0, \infty) = 0$ and hence Eq. (3-3) will not directly give us the field distribution upon the cylinder. However, if we use the Maxwell equation

$$\sqrt{\frac{\mu}{\epsilon}}\,H_\varphi = i\frac{\partial E_z}{\partial(kp)}$$

we find that Eq. (3-19) leads to

$$\left[\sqrt{\frac{\mu}{\epsilon}}H_\varphi\right]_{\rho=a} = i\left(\frac{2}{ka}\right)^{1/3}\left[-i2p + \frac{1}{p}\frac{\partial A}{\partial\sigma}\right]_{\sigma=p}\exp(-ika\cos\varphi - i\omega t)$$

$$= i\left(\frac{2}{ka}\right)^{1/3} F(-p)\exp(-ika\cos\varphi - i\omega t) \qquad (3\text{-}43)$$

where F(-p) is the current distribution function

$$F(-p) = \exp(-i\frac{p^3}{3})\frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty}\frac{\exp(-ipt)}{w_1(t)}\,dt$$

which has the asymptotic property that as $p \to \infty$

$$F(-p) = -i2p + \frac{1}{2p^2} - \sum_{n=1}^{\infty}\frac{n\,A_n(p,\infty)}{p^{n+1}}$$
$$\approx -i2p\left\{1 + \frac{1}{4p^3} + \frac{1}{2p^6} - i\frac{175}{64}\frac{1}{p^9} - \frac{395}{16}\frac{1}{p^{12}} + i\frac{318175}{1024}\frac{1}{p^{15}} + \ldots\right\} \qquad (3\text{-}44)$$

The expansions in inverse integral powers of p which have been given in Tables 3-1 and 3-2, as well as the expansions for G(-p) and F(-p) which have been given in Eqs. (3-42) and (3-44), respectively, become useless when p tends to zero. The physical interpretation of p in terms of the cosine of the angle of incidence reveals that all the expansions of this type become useless for grazing incidence. However, we must bear in mind that even in some cases where the expansions in inverse powers of p appear to be useful, the terms of order $(ka/2)^{-\frac{2}{3}}$ which we have neglected in Eq. (3-14) may become important. We have shown that by means of the functions such as $V_2(x,q)$ and $V_1(x,q)$ that the restriction to non-grazing angles of incidence can be readily relaxed by evaluating these integrals by numerical means. However, it is very difficult to relax the restrictions to large values of ka and much research is need in this direction. It is interesting to observe that the above results reveal that for p tending to zero, i.e., for grazing incidence, it is no longer satisfactory to decompose the fields into the unperturbed field and the reflected field. Since the obstacle tends to block off the incident field, a Fresnel integral representation such as given in Eqs. (3-28) and (3-29) becomes necessary. The author's experience with these types of expansions had led him to conjecture that one cannot expect to be able to analytically continue the expansions in inverse powers of (ka) from the lighted region into the shadow region. However, in Section 2 the author has discussed an example in which it is shown how one can start with the residue series which is valid in the shadow region and analytically extend these series into the lighted region provided one introduces the diffraction functions as analytical "tools" to be employed to cross the horizon. The situation that exists in the vicinity of the horizon appears to be be a type of "Stokes phenomenon." Two types of series are required in the lighted region (namely, the incident and the reflected waves), whereas

in the shadow region one works with the total field. It is the experience of the present author that the "connection formulae" can be found when one wishes to pass from the shadow region into the lighted region, but that the connection formulae for the reverse direction present a much higher order of difficulty. Since we are actually working with a partial differential equation (the wave equation) the phenomenon to which we refer is actually a generalization of the "Stokes phenomenon" since the latter has arisen as a topic in the discussion of the asymptotic solution of ordinary differential equations.

In the course of studying these expansions in inverse integral power of (ka), the author has extended the computations of Keller, Lewis, and Seckler (Ref. 8) so as to obtain further terms in the asymptotic expansions. In fact, some of the further terms in the series employed by Keller and his associates are identical with the terms which have been displayed in Tables 3-1 and 3-2. The additional terms which have been evaluated are, at the moment, more or less of purely academic interest. However, if means of "summing" these asymptotic series can be discovered, then the additional terms may prove to be of considerable interest. Therefore, the author is taking the opportunity presented by the occasion of preparing this report to include these tables. The tables were generated on the IBM 7090 by employing a FORTRAN II program which was augmented by the use of the NPREC subroutines which the author has discussed in the Preface. In order to make it easier for the reader to utilize these results, we will introduce the notation which appears in Ref. 8. In Fig. 3-2 we reproduce the illustration (Fig. 10) of Ref. 8 which provides the geometric meanings for the symbols which are employed.

A plane wave exp(ikx) is incident from the left on a circular cylinder whose equation is

$$x = - b_o \cos \varphi = - b_o \sin \beta/2 \ , \ y = b_o \sin \varphi = b_o \cos \beta/2 \ .$$




Fig. 3-2 The Geometry as Defined by Keller, Lewis, and Seckler (Ref. 8)

These authors use the scalar wave equation

$$(\nabla^2 + k^2)U(x,y) = 0$$

to determine a recursion relation to permit them to determine the $V_n(x,y)$ which occur in the expansion

$$U(x,y) \sim \exp[ik\psi(x,y)]\sum_{n=0}^{\infty} \frac{V_n(x,y)}{(ik)^n}$$

where $\psi(x,y)$ is the phase function which satisfies the eiconal equation of geometrical optics

$$(\nabla\psi)^2 = 1 \ .$$

These authors found it convenient to employ a set of coordinates (s,β), where s is the distance measured along a ray from the caustic, and β is the angle between a ray and the positive x-axis.

The origin of $\underline{s}$ is not shown in the figure. The caustic is a distance

$$s_o = (b_o/2) \cos \theta = (b_o/2) \sin (\beta/2)$$

measured back along the dashed line which is the extension to the interior of the cylinder of the reflected ray (i.e., the dashed line which is used to define the angle β).

In their Eq. (200), these authors express the field in the form

$$U(x,y) = U(s,\beta) \approx \exp(ikx) + D(s,\beta)F(s,\beta) \qquad (3\text{-}45)$$

where

$$D(s,\beta) = \frac{1}{2}\sqrt{\frac{b_o}{2s}\sin\frac{\beta}{2}}\ \exp\left\{ik\left(s - \frac{3}{2}b_o \sin\frac{\beta}{2}\right)\right\} \qquad (3\text{-}46)$$

and

$$F(s,\beta) = \sum_{n=0}^{\infty}\sum_{j=0}^{3n}\sum_{t=0}^{n} a_{jtn}\left(i16b_o k \sin\frac{\beta}{2}\right)^{-n}\left(\frac{b_o}{2s}\right)^{j}\left(\sin\frac{\beta}{2}\right)^{j-2t} \qquad (3\text{-}47)$$

Recursion relationships are given for the a_{jtn}, but we will refer the reader to Ref. 8 if he wishes to examine the intricate nature of these formulae. Our aim in this report is to present to the reader the explicit values for the a_{jtn} which we have obtained for the cases $j = 0(1)6$ for the boundary condition $U = 0$ (the so-called Dirichlet problem) and the boundary condition $\partial U/\partial \nu = 0$ (the so-called Neuman problem). We have chosen to employ only two subscripts, and hence we have made the definitions

$$a_{0jt} = A_{jt}\ , \qquad a_{4jt} = E_{jt}$$
$$a_{1jt} = B_{jt}\ , \qquad a_{5jt} = F_{jt}$$
$$a_{2jt} = C_{jt}\ , \qquad a_{6jt} = G_{jt}$$
$$a_{3jt} = D_{jt}\ ,$$

For the Dirichlet problem, the leading term (i.e., the coefficient which corresponds to that occuring in geometrical optics) is

$$a_{000} = A_{00} = -2 \qquad (3\text{-}48)$$

In Table 3-3 we give the coefficients B_{jt} and C_{jt} for the Dirichlet problem. Tables 3-4 through 3-7 contain the other terms in the series for this problem which have been evaluated.

Table 3-3

COEFFICIENTS B_{jt} AND C_{jt} FOR THE DIRICHLET PROBLEM

B_{jt}			C_{jt}			
j \ t	0	1	j \ t	0	1	2
0	-6	16	0	15	-528	640
1	-6	2	1	-18	-354	560
2	-18	12	2	-87	-438	495
3	30	-30	3	-540	60	300
			4	945	-1260	210
			5	3150	-4410	1260
			6	-3465	6930	-3465

Table 3-4

COEFFICIENTS D_{jt} FOR THE DIRICHLET PROBLEM

j \ t	0	1	2	3
0	-105	23112	-90240	71680
1	-75	23985	-108144	91520
2	-540	18888	-81291	69960
3	-3060	6780	-62415	60315
4	-25830	27720	-69615	60270
5	43470	-34650	-73395	55125
6	346500	-471240	79695	41580
7	-180180	540540	-405405	45045
8	-945945	2162160	-1486485	270270
9	765765	-2297295	2297295	-765765

Table 3-5

COEFFICIENTS E_{jt} FOR THE DIRICHLET PROBLEM

j \ t	0	1	2	3	4
0	1181.25	-1200984	10507584.0	-22124544	13066240.0
1	2205.0	-1783299	17177688.0	-38025600	23152640.0
2	1822.5	-1338327	13367695.5	-30515976	18980160.0
3	-14895.0	-1077645	11616765.0	-26670435	16605000.0
4	-144821.25	-1532265	10799617.5	-23531445	14665901.25
5	-1638630.0	92610	7833420.0	-19866420	13229370.0
6	2671515.0	-3069990	5318775.0	-18724860	12969495.0
7	37567530.0	-54144090	24054030.0	-21351330	13243230.0
8	2331078.75	27297270	-22567545.0	-19729710	12499987.5
9	-204459255.0	487792305	-339999660.0	45945900	10720710.0
10	21824302.5	-189143955	327364537.5	-174594420	14549535.0
11	392837445.0	-1265809545	1440403965.0	-654729075	87297210.0
12	-250979478.75	1003917915	-1505876872.5	1003917915	-250979478.75

Table 3-6a

COEFFICIENTS F_{jt} FOR THE DIRICHLET PROBLEM

j \ t	0	1	2
0	-18191.25	72049590.0	-1189662912.0
1	-53156.25	140589546.75	-2493[illegible]76.0
2	-33941.25	115874419.5	-222401[illegible]4.5
3	-37406.25	113075966.25	-21091[illegible]207.5
4	-865856.25	112610925.0	-1878[illegible]2775.0
5	-9485201.25	54580128.75	-1559640285.0
6	-130756106.25	163009192.5	-1571646037.5
7	204222768.75	-31114833.75	-1675876702.5
8	4466718506.25	-638972340.0	1162431270.0
9	2666747131.25	-363546933.75	-2988015030.0
10	-36435673023.75	91072814332.5	-690884669[illegible]47.5
11	-24170415018.75	30892300188.75	133346[illegible]8827.5
12	132266185301.25	-45477481549[illegible]5.	549143099505.[illegible]
13	18823460906.25	18823460906.25	-175685635125.[illegible]
14	-207058069968.75	865879201687.5	-139293610700[illegible].5
15	109176073256.25	-545880366281.25	109176073256[illegible].5

(Table is concluded on the next page.)

Table 3-6b

COEFFICIENTS F_{jt} FOR THE DIRICHLET PROBLEM

j \ t	3	4	5
0	4837567488.0	-6980886528.0	3280076800.0
1	10577896128.0	-15687469056.0	7513088000.0
2	9886569924.0	-15115684416.0	7394503680.0
3	9127981957.5	-13695202920.0	6609998400.0
4	7930033650.0	-11799834716.25	5680423350.0
5	6843403035.0	-10290725786.25	4995269923.75
6	6266154510.0	-9287917773.75	4547717212.5
7	5553170122.5	-8369484873.75	4240074588.75
8	4176752580.0	-7528843722.5	4117563450.0
9	3903509530.0	-7777492222.5	4225108387.5
10	18856197360.0	-8795193907.5	4379410035.0
11	-17263023277.5	-7049249707.5	4255736987.5
12	-259010822070.0	28360681098.75	4015671660.0
13	225881530875.0	-94117304531.25	62744866968.75
14	1054113810750.0	-357645757216.75	376469218 12.5
15	-1091760732562.5	545680366281.25	-109176073256.25

Table 3-7a

COEFFICIENTS G_{jt} FOR THE DIRICHLET PROBLEM

j \ t	0	1	2
0	354729.375	-4899226590.0	137443572568.0
1	1400726.25	-11889983283.75	353103599970.0
2	1273269.375	-11381431736.25	369123784232.625
3	2082150.0	-11878860960.0	370580264955.0
4	2035687.5	-11008263420.0	339630798150.0
5	-52421985.0	-9304907535.0	311613956955.0
6	-758372422.5	-13367793285.0	301540337752.5
7	-12515933430.0	-2479456980.0	2527469094l5.0
8	187038664l31.25	-11009178180.0	1818922505l0.0
9	59123758l437.5	-873421221172.5	567608756715.0
10	72187154l641.25	-58303624l267.5	1964951075l8.75
11	-6435463023990.0	1673697029004l.0	-12982622828175.0
12	-9653172711682.5	183877605311l0.0	-667203846309l.0
13	326524301853l5.0	-120696031330875.0	15608413783l625.0
14	36705748767187.5	-100667868926625.0	760844289830l2.5
15	-9520153587945l.0	43146384150870l.0	-748947862537875.0
16	-32152353573965.625	8122699850265l.0	0.0
17	131993872566806.25	-6802761124596l3.75	1421472473796375.0
18	-592280197415l5.625	3553681174490l3.75	-888420296122734.375

(Table continues on next page.)

Table 3-7b

COEFFICIENTS G_{jt} FOR THE DIRICHLET PROBLEM

j \ t	3	4	5
0	-937260303360.0	244084104806l4.0	-267930694336O.0
1	-2498278112832.0	6671675630592.0	-74575226880O0.0
2	-2753248927788.0	761297522294l4.0	-8721456457600.0
3	-270144613881l2.5	7360090004100.0	-6345820571200.0
4	-2457927602392.5	6644016395060.625	-7477774083990.0
5	-2232804008452.5	5949249780835.25	-6620485137503.75
6	-2028862629832.5	5285640393061.875	-5827940150051.25
7	-1772818610062.5	4648813932262.5	-5153677211182.5
8	-1585806725002.5	4178670406595.625	-4664542970587.5
9	-1619620925422.5	3855431313483.75	-4306063901391.25
10	-1452705596842.5	3381430148971.875	-4001998734483.75
11	2066346785002.5	270669364l655.0	-3846096829575.0
12	-304036540557l.5	3217180448356.875	-4080926324475.0
13	-812295082974l7.5	1579288370034.3.75	-4536454078406.25
14	3802339103062.5	-1478582854l859.375	-3068224127716.75
15	6081107280373l2.5	-217260385779937.5	1987004533263l.5
16	-1692229135471l7.5	1776840592245l6.875	-6092024887698l.5
17	-1523006221924687.5	863036859090650.25	-233527620695118.75
18	11845603948303l2.5	-8884202961227l4.375	3553681184490l3.75

(Table concluded on next page.)

Table 3-7c

COEFFICIENTS G_{jt} FOR THE DIRICHLET PROBLEM

j \ t	6
0	1045037056000.0
1	294878904 3200.0
2	3512107315200.0
3	3335791488000.0
4	296984 4006000.0
5	260717134 0410.0
6	22885304 98629.375
7	2040565008982.5
8	187417059204 3.75
9	1772671312912.5
10	1726104090335.625
11	174913054 8165.0
12	184068349 7152.5
13	193254 1986375.0
14	192940474 2890.625
15	196516931 8612.5
16	338445827094 3.75
17	203067496256 62.5
18	-5922801974 1515.625

The reader will observe that some of the entries in these tables contain as many as 18 significant figures. The author chose to stop the computer program with the coefficients G_{jt} since to continue to any higher-order terms would not permit the exact determination of the coefficients. The author is well aware of the fact that these coefficients already contain more significant figures than can be expected to be of significance in a practical

problem. However, the author feels that a scientist should be permitted to occasionally enjoy the "luxury" of obtaining exact results which have an esthetic value even if their practical value is questionable. However, in the particular case of the coefficients which have been presented above, it may be that the effective use of summation techniques in order to sum these series will actually require these exact results for the coefficients. Let us now turn to the second case of the Keller, Lewis, and Seckler recursion formula which was run on the IBM 7090.

For the Neuman problem, the leading term (i.e., the coefficient which corresponds to that occuring in geometrical optics) is

$$a_{000} = A_{00} = 2 \tag{3-49}$$

In Table 3-8 we give the coefficients B_{jt} and C_{jt} for the Neuman problem. Tables 3-9 through 3-12 contain the other terms in the series for this problem which have been evaluated.

Table 3-8

COEFFICIENTS B_{jt} AND C_{jt} FOR THE NEUMAN PROBLEM

B_{jt}			C_{jt}			
j \ t	0	1	j \ t	0	1	2
0	6	16	0	-15	-528	896
1	6	-2	1	18	-318	560
2	18	-12	2	87	-426	465
3	-30	30	3	540	-540	180
			4	-945	1260	-210
			5	-3150	4410	-1260
			6	3465	-6930	3465

Table 3-9

COEFFICIENTS D_{jt} FOR THE NEUMAN PROBLEM

j \ t	0	1	2	3
0	105	23112	-126336	114688
1	75	23535	-138096	128128
2	540	18600	-96789	86856
3	3060	5700	-65265	63525
4	25830	-2520	-71505	57330
5	-43470	105210	-97965	45675
6	-346500	526680	-190575	13860
7	180180	-540540	405405	-45045
8	945945	2162160	1486485	-270270
9	-765765	2297295	-2297295	765765

Table 3-10

COEFFICIENTS E_{jt} FOR THE NEUMAN PROBLEM

t \ j	0	1	2	3	4
0	-1181.25	-1200984	14474688.0	-34750464	22077440.0
1	-2205.0	-1775949	23133528.0	-57321600	37044224.0
2	-1822.5	-1327977	17333680.5	-43190280	28034496.0
3	14895.0	-1054515	14487555.0	-35566125	227411320.0
4	144821.25	-1535415	13210942.5	-29766555	18487218.75
5	1638630.0	-1755810	9070740.0	-23149980	14878710.0
6	-2671515.0	7283430	3329865.0	3329865	1297425.0
7	-37567530.0	57026970	-6756750.0	-24054030	11981970.0
8	-2331078.75	-38108070	65810745.0	-34324290	9121612.5
9	204459255.0	-500044545	37675380.0	-82702620	1531530.0
10	-21824302.5	189143955	-327364537.5	174594420	-14549535.0
11	-392837445.0	1265809545	-1440403965.0	654729075	-87297210.0
12	250979478.75	-1003917915	1505876872.5	-1003917915	250979478.75

Table 3-11a

COEFFICIENTS F_{jt} FOR THE NEUMAN PROBLEM

j \ t	0	1	2
0	18191.25	72049590.0	-1605837888.0
1	53156.25	140440709.25	-3370646760.0
2	33941.25	115532644.5	-2964851887.5
3	37406.25	112536753.75	-2779312432.5
4	865856.25	112393575.0	-2439376905.0
5	9485201.25	52560191.25	-1970244675.0
6	130756106.25	13 8967.5	-1954005322.5
7	-204222768.75	900303153.75	-2301056257.5
8	-4466718506.25	7104857760.0	-2896483590.0
9	-2868747131.25	-180509954 6.25	7435578150.0
10	36435673023.75	-94215513892.5	72114770227.5
11	241704 15018.75	-287971671 48.75	-2660382 4747.5
12	-132266185301.25	4587904 87155.0	-5652057861 45.0
13	-18823460906.25	-18823460906.25	175685635125.0
14	207058069968.75	-865879201687.5	1392936107062.5
15	-109176073056.25	545880366281.25	-1091760732562.5

(Table is concluded on the next page.)

Table 3-11b

COEFFICIENTS F_{jt} FOR THE NEUMAN PROBLEM

j \ t	3	4	5
0	7404675072.0	-11469545472.0	5641732096.0
1	16096961088.0	-25456091136.0	12694528000.0
2	14666724804.0	-23700901056.0	11992866816.0
3	13221905962.5	-20759495400.0	10283179200.0
4	11142795510.0	-17135697363.75	8380599990.0
5	9251998245.0	-14213492133.75	6925025756.25
6	8210393730.0	-12216371186.25	5883263347.5
7	7069519957.5	-10402388246.25	5035591811.25
8	4729184460.0	-8781950677.5	4408914510.0
9	1750538790.0	-8542491457.5	4081910332.5
10	-733296560.0	-10759381132.5	3768329565.0
11	4499752797.5	-15997213732.5	2728037812.5
12	283104852030.0	-4423367738.75	0.0
13	-225881530875.0	94117304531.25	-62744869 68.75
14	-10541138 10750.0	35764575 7218.75	-37646921812.5
15	1091760732562.5	-5458803662 81.25	109176073256.25

Table 3-12a

COEFFICIENTS G_{jt} FOR THE NEUMAN PROBLEM

j \ t	0	1	2
0	-354729.375	-4899226590.0	181732327632.0
1	-1400726.25	-11886381416.25	471713468130.0
2	-1273269.375	-11369792643.75	493165448463.375
3	-2082150.0	-11861529660.0	493784995905.0
4	-2035687.5	-10987567920.0	449568654570.0
5	52421985.0	-9249823485.0	409455822825.0
6	758372422.5	-13433718375.0	394506450607.5
7	12515933430.0	-16615118520.0	327045483765.0
8	-187038664431.25	88663695120.0	20276898420.0
9	-591237584437.5	943451961952.5	1065554339 85.0
10	-721871541641.25	82823227987.5	16629354654 41.25
11	6435463023990.0	-1757659485582 0.0	143837430486 75.0
12	9653172711682.5	-1759667321412 0.0	257605336989 0.0
13	-32652430185375.0	12280425895237 5.0	-16240882069 9125.0
14	-3670574876718 7.5	10036669355212 5.0	-7186797374 0062.5
15	952015358794 50.0	-43321065868080 0.0	75768194839 8375.0
16	32152353573965.625	-8122699850265 0.0	0.0
17	-1319938725668 06.25	6802761124596 93.75	-14214724737 96375.0
18	59228019741515.625	-35536811844909 3.75	88842029612273 4.375

(Table continues on next page.)

Table 3-12b

COEFFICIENTS G_{jt} FOR THE NEUMAN PROBLEM

j \ t	3	4	5
0	-1395776937984.0	388911539404 8.0	-44610705162 24.0
1	-375452864 4288.0	106990457978 88.0	-12464072957952.0
2	-4120084137132.0	12105282445536.0	-1440244 8162816.0
3	-400651065 9787.5	11534155406340.0	-13521037575360.0
4	-3595959838327.5	102083298446 19.375	-11817369920790.0
5	-3217416555307.5	8938503496203.75	-10164946510136.25
6	-287690457804 7.5	7739219430298.125	-8645108453808.75
7	-2456455358857.5	6590920773937.5	-7334426516917.5
8	-2133839285077.5	5722828821084.375	-6345237370972.5
9	-2183792928817.5	5128131594836.25	-5588995390728.75
10	-2391514342717.5	4311898373908.125	-4901269118996.25
11	-3503258861602.5	2864614297545.0	-4410473292225.0
12	+7296977365177.5	9591180780 43.125	-4613002819425.0
13	8404047845 9437.5	-7359973214343.75	-6004684029093.75
14	-176564063300 62.5	3406105250985 9.375	-9279966226781.25
15	-625578899758312.5	2347285575009 37.5	-2860413119313 7.5
16	16922291354 7187.5	-17768405922454 6.875	6092024887698 7.5
17	1523006221924687.5	-863036859090656.25	2335276206951 18.75
18	-1184560394830312.5	8884202961227 34.375	-3553681184490 93.75

(Table concluded on next page.)

Table 3-12c

COEFFICIENTS G_{jt} FOR THE NEUMAN PROBLEM

j \ t	6
0	1794850684928.0
1	5071917154304.0
2	5951569096704.0
3	5525039063040.0
4	4776418491600.0
5	4051514138490.0
6	3408147408290.625
7	2885500155037.5
8	2496129902516.25
9	2201422370647.5
10	1969012214544.375
11	1811548053315.0
12	1725232936927.5
13	1581170716125.0
14	1082349002109.375
15	-218352146512.5
16	-3384458270943.75
17	-20306749625662.5
18	5922801974151 5.625

The reader may be interested to know that the "run time" for each of these problems on the IBM 7090 was 0.05 hour. Since the "cost" of computer time on such a computer is generally between $300 and $500 per hour, we see that the obtaining of these tables "cost" less than $50 in actual computer expenses. Needless to say, the true cost was far greater because of the time required to program the recursion formulae and the debugging of the programs. However, when one stops for a moment to realize the tremendous amount of

labour that would have been required to have obtained only the first few terms in these series on a desk-type manual computer, one can begin to see vividly that the electronic computer has opened an era in which recursion formulae such as those derived a decade ago by Keller, Lewis, and Seckler may take on an entirely new importance. Granted that the series in their present forms are of limited usefulness because of their asymptotic nature, the fact that a summation scheme may be devised which will extend the value of these series to smaller value of ka makes these series stand out as a promising means of obtaining practical results for the scattering properties of smooth obstacles. The studies made by the present author have only involved the case of the circular cylinder and the two cases of the Dirichlet and the Neuman problem. However, in their classic paper, Keller, Lewis, and Seckler (Ref. 8) consider as their Example 24 the "Diffraction of a Place Wave by a Circular Cylinder ($\partial U/\partial \upsilon = ikZU$)." Time under the present research program was not available for the completion of a program based upon this case of the impedance boundary condition which would have led to the results of the Dirichlet and the Neuman problems for the special cases of $Z = \infty$ and $Z = 0$, respectively.

Although the author has devoted these efforts to the circular cylinder problem, it should be emphasized that the classic paper (Ref. 8) which was used to obtain Tables 3-3 through 3-12 also showed how one could apply similar methods to other obstacles. Even the case of a sphere (which was briefly considered as Examples 19 and 20 in Ref. 8) offers "rich rewards" to anyone who attempts to follow up the problem of working out the expansion in inverse powers of ka and then to attempt to find means of summing the series. It should also be borne in mind that with the coming of age of the computer that some of the actual algebra which is required when one follows the methods of Ref. 8 can be mechanized.

The present author has prepared several computer programs which are valuable for carrying out of tedious algebraic manipulations with polynomials. Although these have proven to be of considerable advantage, time has not permitted the application of these methods to attempting to extend the work on the finding of expansions in inverse powers of $k = 2\pi/\lambda$ for points in the lighted region. However, as an indication of what may lie just ahead in the field of carrying out tedious algebraic manipulations by means of electronic computers, the author wishes to call attention to a paper by Brown (Ref. 9) which describes the ALPAK system which has been under development at the Bell Telephone Laboratories. The present author readily admits that his own experience has not led him to be as optimistic as to lead him to agree with Brown's "...rule of thumb that one man-hour equals one 7090-second." However, once the particular algebra of a specific problem has been so organized as to be performed by the computer, then this rule of thumb is certainly very close to being consistent with the author's experience. The author believes that within the next decade that someone will rework the theory of Ref. 8 by employing routines on a digital computer to carry out the tedious algebra. The results will certainly lead to an enhancement of our understanding of the "breakdown" of geometrical optics as the wavelength increases and becomes comparable with the radii of curvature of the scattering obstacle.

REFERENCES FOR SECTION 3

1. V. A. Fock, Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965. (See, in particular, the paper "The Field of a Plane Wave Near the Surface of a Conducting Body," pp. 90-111. This paper originally appeared (in English) in J. Physics USSR, Vol. 10, 1946, pp. 399-409.)

2. M. G. Belkina and L. A. Vainshtein, "Radiation Characteristics of Spherical Surface Antennas," Diffraction of Electromagnetic Waves by Certain Bodies of Revolution, Moscow, Soviet Radio Press, 1957. The author has an unpublished translation of this important monograph which was made available to him in early 1958 by K. M. Siegel who was then the head of the Radiation Laboratory, University of Michigan.

3. P. A. Azriliant and M. G. Belkina, Numerical Results of the Theory of Diffraction of Radiowaves Around the Earth's Surface, Moscow, Soviet Radio Press, 1957. The author has a translation of the text which preceeds these tables. The translation was made by M. D. Friedman but for financial reasons the translation has not been prepared in a form which would permit its distribution in report form.

4. A. A. Federov, "Asymptotic Solution of the Problem of Diffraction of a Plane Electromagnetic Wave by a Perfectly Conducting Sphere," Zhur. Tech. Fiz., Vol. 28, 1958, pp. 1451-1462.

5. J. R. Wait, Electromagnetic Radiation from Cylindrical Structures, New York, Pergamon Press, 1958.

6. R. F. Goodrich, "Fock Theory-An Appraisal and Exposition," IRE Trans. on Antennas and Propagation, Vol. AP-7, 1959, pp. S26-S38.

7. N. A. Logan and K. S. Yee, "A Mathematical Model for the Theory of Diffraction by Convex Surfaces," Electromagnetic Waves: Proceedings of a Symposium Conducted by the Mathematics Research Center, United States Army, at the University of Wisconsin, April 10-12, 1961, R. E. Langer, ed., Madison, Wisc., University of Wisconsin Press, 1962

8. J. B. Keller, R. M. Lewis, and B. D. Seckler, "Asymptotic Solution of Some Diffraction Problems," Comm. Pure Appl. Math., Vol. 9, 1956, pp. 207-265.

9. W. S. Brown, "The ALPAK System for Nonnumerical Algebra on a Digital Computer — I: Polynomials in Several Variables and Truncated Power Series with Polynomial Coefficients," B.S.T.J., Vol. 42, Sept., 1963, pp. 2081-2119.

Section 4

ASYMPTOTIC EXPANSIONS FOR THE HEIGHT GAIN FUNCTION FOR THE CASE OF NORMAL REFRACTION

4.1 Introduction

This section deals with a number of aspects of the problem of obtaining and of employing asymptotic representations for the class of second-order differential equations which arise naturally when one attempts to solve the wave equation by the method of separation of variables. The choice of the material to be included is based upon the author's experience as to what results he has found useful (or interesting) in his work on the theory of the diffraction of waves by convex surfaces. The author has chosen to refer to the type of problem to be considered by the use of the term "height gain function" since the variable in the differential equation is generally related in some way to the height (or the distance) of the observation point from the convex surface. The restriction to the case of "normal refraction" is included in the title because all the asymptotic results to be discussed in this section arise from the consideration of differential equations which have only one "turning point." Therefore, we have excluded all problems in which the index of refraction above the convex (or the plane) surface varies in such a manner as to give rise to the phenomena of superrefraction, tropospheric waveguide modes, or similar complications of an already complex problem.

4.2 Problems Leading to the Height Gain Function

The height gain function $F(y,\lambda_s)$ with which we shall be concerned is the solution of the differential equation

$$\frac{d^2F}{dy^2} + [-\lambda_s + k^2 f(y)]\, F = 0 \qquad (4\text{-}1)$$

for the case of "normal refraction" which we shall define by the requirements that $f(y) > 0$ and $f'(y) > 0$ for $y > 0$. The eigenvalues λ_s are to be defined by the impedance boundary condition

$$\left(\frac{dF}{dy} + ikZF \right)_{y=0} = 0 \qquad (4\text{-}2)$$

The function $F(y,\lambda_s)$ occurs in many problems which involve the propagation of waves over convex surfaces or over plane surfaces above which the index of refraction is a function only of the height above the surface.

Let us cite several examples. The solution of the problem of propagation of axially symmetric waves in a horizontally stratified atmosphere above a plane surface is governed by the equation

$$\frac{\partial^2 G}{\partial x^2} + \frac{1}{x}\frac{\partial G}{\partial x} + \frac{\partial^2 G}{\partial y^2} + k^2 n^2(y) G = 0 \,. \qquad (4\text{-}3)$$

The solution is of the form

$$G(x,y) = \sum_{s=1}^{\infty} \frac{i}{4} H_0^{(1)}(\mu_s x)\Phi_s(y)\Phi_s(y_0) \qquad (4\text{-}4)$$

where $\Phi(y)$ is a solution of

$$\frac{d^2\Phi}{dy^2} + [-\mu_s{}^2 + k^2 n^2(y)]\Phi = 0 \qquad (4\text{-}5)$$

which appears in the standard form of Eq. (4-1) with $\lambda_s = \mu_s{}^2$ and $f(y) = n^2(y)$.

As a second example, let us consider the problem of the propagation of two-dimensional waves over the surface of a circular cylinder

when the index of refraction is a function only of the distance from the center of the cylinder.

$$\frac{\partial^2 G}{\partial \rho^2} + \frac{1}{\rho}\frac{\partial G}{\partial \rho} + \frac{1}{\rho^2}\frac{\partial^2 G}{\partial \varphi^2} + k^2 n^2(\rho) G = 0 \qquad (4\text{-}6)$$

The solution is of the form

$$G(\rho,\varphi) = \sum_{s=1}^{\infty} \frac{i}{2} \frac{\exp(i\nu_s \varphi)}{\nu_s} \Phi_s(\rho)\Phi_s(\rho_0) \qquad (4\text{-}7)$$

where $\Phi(\rho)$ is a solution of

$$\frac{d^2\Phi}{d\rho^2} + \frac{1}{\rho}\frac{d\Phi}{d\rho} + \left[-\frac{\nu_s^2}{\rho^2} + k^2 n^2(\rho)\right]\Phi = 0 \qquad (4\text{-}8)$$

Eq. (4-8) does not immediately take on the appearance of the standard form which we have taken to be that of Eq. (4-1). However, if we make the transformations

$$\rho = a \exp(y/a) \ , \quad y = a \log(\rho/a)$$

and define

$$\Phi(\rho) = F(y)$$

we will find that $F(y)$ satisfies Eq. (4-1) provided we make the definitions

$$\lambda_s = (\nu_s/a)^2 \ , \quad f(y) = [\rho n(\rho)/a]^2$$

Another good example is the problem of the propagation of axially symmetric waves over the surface of a sphere when the index of refraction is a function of the radial distance r alone.

$$\frac{\partial^2 G}{\partial r^2} + \frac{2}{r}\frac{\partial G}{\partial r} + \frac{1}{r^2 \sin\theta}\frac{\partial}{\partial \theta}\left(\sin\theta \frac{\partial G}{\partial \theta}\right) + k^2 n^2(r) G = 0 \qquad (4\text{-}9)$$

The solution is of the form

$$G(r,\theta) = \sum_{s=1}^{\infty} \frac{P_{\nu_s - 1/2}(-\cos\theta)}{4\cos(\nu_s \pi)} \Psi_s(r)\Psi_s(r_0) \qquad (4\text{-}10)$$

where $\Psi_s(r)$ is a solution of

$$\frac{d^2\Psi}{dr^2} + \frac{2}{r}\frac{d\Psi}{dr} + \left[- \frac{\nu_s^2 - 1/4}{r^2} + k^2 n^2(r) \right]\Psi = 0 \qquad (4\text{-}11)$$

Eq. (4-11) can be brought into the standard form of Eq. (4-1) by making the transformations

$$r = a \exp(y/a) \ , \quad y = a \log(r/a)$$

and the definition

$$\Psi(r) = \frac{1}{\sqrt{r}} F(y)$$

along with the identification of λ_s and $f(y)$ as being

$$\lambda_s = (\nu_s/a)^2 \ , \quad f(y) = [rn(r)/a]^2$$

There are several other classical problems which we could cite which would lead to an illustration of the fact that Eq. (4-1) is of paramount importance in the study of wave propagation problems. However, in order to focus more sharply on the subject matter of this Section, let us cite one more example which can serve as a model for most of the above problems. We will consider the height gain function $F(y,\lambda_s)$ as it arises in the solution of the separable wave equation

$$\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} + k^2 \left[f(y) + g(x) \right] U = - \delta(x - x_o)\, \delta(y - y_o) \qquad (4\text{-}12)$$

Barring certain pathological cases which we need not consider here, it is possible to obtain solutions of Eq. (4-12) in the form of a series of normal modes of the form

$$U(x, y; x_o, y_o) = \sum_{s=1}^{\infty} \frac{\Psi^+(x_>, \lambda_s)\, \Psi^-(x_<, \lambda_s)}{W_s} \quad \frac{F(y, \lambda_s)\, F(y_o, \lambda_s)}{N_s} \qquad (4\text{-}13)$$

where $\Psi(x,\lambda)$ and $F(y,\lambda)$ are solutions of the ordinary differential equations

$$\frac{d^2\Psi}{dx^2} + \left[\lambda + k^2\, g(x)\right]\, \Psi = 0 \qquad (4\text{-}14)$$

$$\frac{d^2F}{dy^2} + \left[-\lambda + k^2\, f(y)\right] F = 0 \qquad (4\text{-}15)$$

If we assume an $\exp(-i\omega t)$ time dependance, we must require that the "distance" functions $\Psi^{\pm}(x,\lambda_s)$ have the properties

$$\Psi^{\pm}(x, \lambda_s) \xrightarrow[k\to\infty]{} \frac{\text{constant}}{\sqrt[4]{k^2 g(x) + \lambda_s}}\, \exp\left[\pm i \int_0^x \sqrt{k^2 g(u) + \lambda_s}\; du\right] \qquad (4\text{-}16)$$

and the "height gain" functions $F(y,\lambda_s)$ have the property

$$F(y, \lambda_s) \xrightarrow[k\to\infty]{} \frac{\text{constant}}{\sqrt[4]{k^2 f(y) - \lambda_s}}\, \exp\left[i \int_0^y \sqrt{k^2 f(v) - \lambda_s}\; dv\right] \qquad (4\text{-}17)$$

The properties which have been prescribed in Eqs. (4-16) and (4-17) are asymptotic properties which are based upon the so-called WKB approximation for which there exists a very large literature. In order to complete the description of Eq. (4-13) we need to set down the fact that the eigenvalues λ_s are defined by the homogeneous boundary condition

$$\left[\frac{\partial F(y, \lambda_s)}{\partial y} + i\, k\, Z\, F(y, \lambda_s)\right]_{y=0} = 0 \qquad (4\text{-}18)$$

The Wronskian W_s is defined by

$$W_s = W(\lambda_s) = \Psi^+ \frac{d\Psi^-}{dx} - \Psi^- \frac{d\Psi^+}{dx} \qquad (4\text{-}19)$$

and the normalization constant is defined by

$$N_s = N(\lambda_s) = \int_0^\infty \left[F(y, \lambda_s)\right]^2 dy \qquad (4\text{-}20)$$

Although a long list of references could be cited for the benefit of the reader who is not familiar with results such as those which are summarized in Eqs. (4-12) through (4-20), the author wishes to acknowledge the fact that the summary given above follows that presented by Friedlander (Ref. 1) more closely than that of any other source with which the author is familiar. Although the monograph by Friedlander (Ref. 2) is generally more accessible to the general scientific public, the author highly commends the original research report to a reader who wishes to acquaint himself with this subject matter.

4.3 Survey of Some Previous Work on the Height Gain Function

Perhaps the most extensive papers on the height gain function and related problems are the two papers (Refs. 3 and 4) by Bremmer. There is a marked similiarity between many of the results obtained by Bremmer and results which will appear below. However, tho work reported in this report has been undertaken independently of that of Bremmer and a detailed comparison of the asymptotic expansions have not been undertaken by the present author.*

*Although many of the results in this report have been made available to Bremmer (and to his collaborators at the National Bureau of Standards at Boulder, Colorado) the present author has not been informed of any discrepancies between the results herein reported and those reported in the papers published by Bremmer. Since these two pieces of work have been carried out independently and follow somewhat different "lines of attack," it is hoped that in the near future such a comparison will be made.

The cylindrical (Bessel) functions $Z_\upsilon(kx)$ are solutions of

$$x^2 \frac{d^2Z}{dx^2} + x \frac{dZ}{dx} + (k^2x^2 - \upsilon^2)Z = 0 .$$

Since the transformation $y = a \log(x/a)$ leads to the operator

$$x^2 \frac{d^2}{dx^2} + x \frac{d}{dx} = a^2 \frac{d^2}{dy^2}$$

the cylindrical functions are the solutions

$$F(y) = Z_\upsilon[(ka)\exp(y/a)]$$

of the equation

$$\frac{d^2F}{dy^2} + [-(\upsilon/a)^2 + k^2\exp(2y/a)]F = 0$$

which is of the form of Eq. (4-1) with $\lambda = (\upsilon/a)^2$ and $f(y) = \exp(2y/a)$. Therefore, it is not surprising that much of the literature that is appropriate to the study of the asymptotic behavior of the height gain function for $k \to \infty$ has had its origin in the interest of the author in developing asymptotic formulae for the cylindrical functions. For a summary of work related to the asymptotic expansions for Bessel functions (and a good guide to the literature), the reader is advised to consult Olver's preface to a table of zeros of Bessel functions which was prepared by the Royal Society Mathematical Tables Committee (Ref. 5). Erdélyi (Ref. 6) has also recently written a survey of the literature related to the study of Eq. (4-1). However, these surveys fail to include some results by Pekeris (Ref. 7), Fock (Ref. 8), Imai (Refs. 9 and 10), and Friedlander (Ref. 1). In many respects, the research reported in Refs. 7 through 10 is more directly applicable to the propagation problems than the more general results which can be found in the other sources.

It will be convenient to start our discussion with a paper by

Pekeris (Ref. 7) in which he defined

$$\lambda_s = - k^2 \mu_s$$

and sought solutions of

$$\frac{d^2F}{dy^2} + k^2 \left[\mu_s + f(y) \right] F = 0 \qquad (4\text{-}21)$$

He showed that one could express $F(y, -k^2 \mu_s)$ in the form

$$F(y, -k^2 \mu_s) \sim \frac{1}{\sqrt[4]{k^2 \left[\mu_s + f(y) \right]}} \left[u^{1/2} H_{1/3}^{(2)}(u) + \frac{3A}{2k^{4/3}} u^{5/6} H_{-2/3}^{(2)}(u) \right.$$

$$\left. - \frac{B}{2k^2} u^{3/2} H_{4/3}^{(2)}(u) + 0(k^{-8/3}) + \cdots \right] \qquad (4\text{-}22)$$

where

$$u(y) = k \int_{y_1}^{y} \sqrt{\mu_s + f(h)} \, dh \qquad (4\text{-}23)$$

and y_1 is the turning point defined by

$$\mu_s + f(y_1) = 0 \qquad (4\text{-}24)$$

The constants in Eq. (4-22) are defined by

$$A = \frac{(1.5)^{4/3}}{315 (\dot{f})^{2/3}} \left[9 \left(\frac{\ddot{f}}{\dot{f}} \right)^2 - 10 \left(\frac{\dddot{f}}{\dot{f}} \right) \right]$$

$$B = \frac{1}{450 \, \dot{f}} \left[- 25 \left(\frac{\ddddot{f}}{\dot{f}} \right) + 70 \left(\frac{\ddot{f} \, \dddot{f}}{\dot{f}^2} \right) - 42 \left(\frac{\ddot{f}}{\dot{f}} \right)^3 \right]$$

where

$$\dot{f} = \frac{df(y_1)}{dy} ,$$

$$\ddot{f} = \frac{d^2 f(y_1)}{dy^2} , \quad \text{etc.}$$

Pekeris showed that he could also express Eq. (4-22) in the form

$$F(y, -k^2\mu_s) \sim \frac{1}{\sqrt[4]{k^2\left[\mu_s + f(y)\right]}}\, u^{1/6}\left(u^{2/3} + A/k^{4/3}\right)^{\frac{1}{2}} H_{1/3}^{(2)}\left[\left(u^{2/3} + A/k^{4/3}\right)^{\frac{3}{2}}\right]$$

or

$$F(y, -k^2\mu_s) \approx \frac{1}{\sqrt[4]{k^2\left[\mu_s + f(y)\right]}}\, u^{1/6} h_2\left[\zeta + \left(\frac{3A^{3/2}}{2k^2}\right)^{2/3}\right] \qquad (4\text{-}25)$$

where

$$\zeta = \left(\frac{3u}{2}\right)^{2/3} \qquad (4\text{-}26)$$

The function $h_2(\chi)$ in Eq. (4-25) is defined by

$$h_2(\chi) \equiv \left(\frac{2}{3}\right)^{1/3} \chi^{\frac{1}{2}} H_{1/3}^{(2)}\left(\frac{2}{3}\chi^{3/2}\right) = z^{1/3} H_{1/3}^{(2)}(z) \quad , \quad z \equiv \left(\frac{3\chi}{2}\right)^{2/3} \qquad (4\text{-}27)$$

These are the Airy functions which were tabulated by the Harvard University Computation Laboratory (Ref. 11).

Imai (Ref. 9) rediscovered some of the results which had been derived by Pekeris. In a later paper, Imai (Ref. 10) found an extension of Eq. (4-25) which can be obtained by employing a result that was used by Cherry (Ref. 12). From Cherry's Lemma 1, we can show that $F(y,\lambda_s)$ can be expressed in the form

$$F(y, \lambda_s) = \left(\frac{d\eta}{dy}\right)^{\frac{1}{2}} h_2(\eta) \qquad (4\text{-}28)$$

where $\eta(y)$ is a solution of the equation

$$\eta(\dot{\eta})^2 - \frac{3}{4}\left(\frac{\ddot{\eta}}{\dot{\eta}}\right)^2 + \frac{1}{2}\frac{\dddot{\eta}}{\dot{\eta}} = -\lambda_s + k^2 f(y) \qquad (4\text{-}29)$$

where

$$\dot{\eta} = \frac{dn}{dy}\,,\ \ddot{\eta} = \frac{d^2n}{dy^2}\,,\ \dddot{\eta} = \frac{d^3n}{dy^3}$$

Eq. (4-25) is an example of Cherry's Lemma. Thus, let

$$\eta = \zeta + \left(\frac{3}{2}\right)^{2/3} \frac{A}{k^{4/3}} + O\left(k^{-6/3}\right) \qquad (4\text{-}30)$$

so that

$$\frac{d\eta}{dy} \sim \frac{d\xi}{dy} = \left(\frac{2}{3}\right)^{1/3} u^{-1/3} \frac{du}{dv} = \left(\frac{2}{3}\right)^{1/3} u^{-1/3} \sqrt{k^2 \left[\mu_s + f(y)\right]} \qquad (4\text{-}31)$$

Therefore, Eq. (4-25) can be expressed in the form

$$F\left(y_1, -k^2 \mu_s\right) \sim \left(\frac{2}{3}\right)^{1/6} \left(\frac{d\eta}{dy}\right)^{-1/2} h_2(\eta) \qquad (4\text{-}32)$$

where η is defined in Eq. (4-30). The term B/k^2 in Eq. (4-22) can be used to extend Eq. (4-30) to include a term in B/k^2. This step was apparently not carried out until the publication of Imai's 1956 paper (Ref. 10).

In order to appreciate Imai's results, we use Eq. (4-27) to express Eq. (4-28) in the form

$$F(y, \lambda_s) \sim \left(\frac{d\eta}{dy}\right)^{-1/2} Z^{1/3} H_{1/3}^{(2)}(Z) \quad , \quad Z = \frac{2}{3} \eta^{3/2} \qquad (4\text{-}33)$$

From Eq. (20) of Ref. 10, we find that Z has been approximated by

$$Z = \sqrt{k^2 + \lambda_o} \left[\xi + \frac{1}{5} \alpha_1 \kappa^{-2} \xi^2\right]^{3/2} \qquad (4\text{-}34)$$

where

$$\xi = [u(y)/k]^{2/3} + \frac{\alpha}{\kappa^2}$$

$$u(y) = k \int_{y_1}^{y} \sqrt{\mu_s + f(h)}\, dh$$

$$\kappa^2 = k^2 + \alpha_o$$

The constants α, α_0, and α_1 will be defined below.

We have taken some liberties with Imai's notation in that we have substituted some of the notation employed by Pokoris. Actually, Imai (Ref. 10) studied the equation

$$\frac{d^2\Phi}{dx} + a^2 P(x) \Phi = 0 \qquad (4\text{-}35)$$

under the assumption that

$$P(x) = a_1 x \left(1 + b_1 x + b_2 x^2 + b_3 x^3 + \cdots\right) \tag{4-36}$$

If we let

$$a^2 = k^2$$

$$\chi = y - y_1$$

$$a_1 = f'(y_1) = \left[\frac{df}{dy}\right]_{y = y_1} = \dot{f}$$

$$a_1 b_n = \frac{1}{n!} \left[\frac{d^{n+1} f}{dy^{n+1}}\right]_{y = y_1}$$

we can compare the papers of Imai and Pekeris. The constants α, α_0, and α_1 used by Imai are defined by the relations

$$\alpha = -\frac{3}{35}\left(\frac{3}{2} a_1\right)^{-2/3}\left(5b_2 - 3b_1^2\right) , \tag{4-37}$$

$$\alpha_o = -\frac{4}{75}\frac{1}{a_1}\left(25b_3 - 35b_1 b_2 + 14b_1^3\right) , \tag{4-38}$$

$$\alpha_1 = -\frac{27}{26950}\left(\frac{3}{2} a_1\right)^{-4/3}\left(6125b_4 - 4350b_2^2 - 9800b_1 b_3 + 12080b_1^2 b_2 - 3624b_1^4\right) . \tag{4-39}$$

We observe that the A and B of Eq. (4-22) are essentially the same quantities as the α and α_0 defined by Eqs. (4-37) and (4-38), respectively.

Let us use the definition of Z in Eq. (4-33) to write

$$\eta = \left(\frac{3}{2} Z\right)^{2/3} \sim \left(\frac{3}{2}\right)^{2/3} \sqrt[3]{k^2 + \alpha_o} \left[\xi + \frac{1}{5}\alpha_1 \kappa^{-2} \xi^2\right]$$
$$\sim \left(\frac{3k}{2}\right)^{2/3}\left[\xi + \frac{1}{3}\frac{\alpha_o}{k^2}\xi + \frac{1}{5}\alpha_1 k^{-2}\xi^2\right] \tag{4-39}$$

From the definition of ξ which occured in Eq. (4-34), we find that

$$\eta \sim \left(\frac{3k}{2}\right)^{2/3}\left\{\left[\frac{u(y)}{k}\right]^{2/3} + \frac{\alpha}{k^2} + \frac{1}{3}\,\frac{\alpha_o}{k^2}\left[\frac{u(y)}{k}\right]^{2/3} + \frac{1}{5}\,\alpha_1 k^{-2}\left[\frac{u(y)}{k}\right]^{4/3}\right\} \quad (4\text{-}40)$$

$$= \left(\frac{3}{2}\right)^{2/3}\left\{u^{2/3}(y) + \frac{\alpha}{k^{4/3}} + \frac{1}{3}\,\frac{\alpha_o}{k^{6/3}}\,u^{2/3}(y) + \frac{1}{5}\,\frac{\alpha_1}{k^{8/3}}\,u^{4/3}(y)\right\} \quad (4\text{-}41)$$

If we recall the definition of ζ given in Eq. (4-26), we can write

$$\eta = \zeta + \left(\frac{3}{2}\right)^{2/3}\frac{\alpha}{k^{4/3}} + \frac{1}{3}\,\frac{\alpha_o}{k^2}\,\zeta + \frac{1}{5}\left(\frac{2}{3}\right)^{2/3}\frac{\alpha_1}{k^{8/3}}\,\zeta^2 + O\left(k^{-10/3}\right) \quad (4\text{-}42)$$

When this result is compared with Eq. (4-30), we see that $\alpha = A$. The simplications we have made in Imai's results have led us to recognize that we have available two additional terms in Eq. (4-30). Imai was apparently not familiar with the work of Pekeris and Cherry so that the use of Eq. (4-42) in Eq. (4-28) to obtain a useful representation for $F(y,\lambda_s)$ is a new result. Further terms in Eq. (4-42) can be generated with the help of Eq. (4-29).

4.4 The Research Reported by Fock

In a classic paper on radiowave propagation, Fock (Ref. 8) studied the equation

$$\frac{d^2G}{du^2} + \left[-t + u + \beta_o u^2\right]G = 0 \quad (4\text{-}43)$$

He showed that

$$G(u,\,t) = w(t-u) - \frac{\beta_o}{15}\left[(3u+2t)\,w(t-u) + (3u^2+4ut+8t^2)\,w'(t-u)\right] + O\left(\beta_o^2\right) \quad (4\text{-}44)$$

where $w(z) = w_1(z)$ is Fock's Airy function

$$w(z) = \sqrt{\frac{\pi}{3}}\exp\left(i\frac{2\pi}{3}\right)(-z)^{1/2}\,H_{1/3}^{(1)}\left[\frac{2}{3}(-z)^{3/2}\right] \quad (4\text{-}45)$$

Fock also used the form

$$G(u,\,t) \sim \left(-\frac{dX}{du}\right)^{-\frac{1}{2}}\,w(X) \quad (4\text{-}46)$$

in which

$$X \sim (t - u) - \frac{\beta_o}{15} (3u^2 + 4tu + 8t^2) + O\left(\beta_o^2\right) \tag{4-47}$$

From Fock's paper, it is not clear whether Fock realized the importance of the form of Eq. (4-46) which is essentially the same as Cherry's form for the solution which we gave in Eq. (4-28). The relations between the two types of Airy functions are

$$w_2(-\eta) = \frac{\sqrt{\pi}}{\sqrt[6]{12}} \exp\left(-i\frac{2\pi}{3}\right) h_2(\eta) \tag{4-48a}$$

$$w_1(-\eta) = \frac{\sqrt{\pi}}{\sqrt[6]{12}} \exp\left(i\frac{2\pi}{3}\right) h_1(\eta) \tag{4-48b}$$

The expressions given in Eq. (4-28) and Eq. (4-46) are special cases of a more general result. Let F(y) and Z(x) be solutions of the equations

$$\frac{d^2F}{dy^2} + P(y)\, F = 0 \qquad \frac{d^2Z}{dx^2} + Q(x)\, Z = 0 \tag{4-49}$$

We can then show that

$$F(y) = \left(\frac{d\eta}{dy}\right)^{-\frac{1}{2}} Z(\eta) \tag{4-50}$$

provided $\eta(y)$ is a solution of the non-linear equation

$$(\dot{\eta})^2 Q(\eta) - \frac{3}{4}\left(\frac{\dot{\eta}}{\eta}\right)^2 + \frac{1}{2}\frac{\ddot{\eta}}{\eta} = P(y) \tag{4-51}$$

Fock arrived at Eq. (4-47) by using the Langer (Ref. 13) asymptotic estimate (the first term in the Pekeris-Imai expansions)

$$\frac{2}{3}(-X)^{\frac{3}{2}} \sim \int_{u_1}^{u} \sqrt{v - t + \beta_o v^2}\; dv \tag{4-52}$$

where u_1 is the turning point which is defined by

$$u_1 - t + \beta_o u_1^2 = 0 \tag{4-53}$$

Fock did not indicate how one could obtain higher-order corrections of Eq. (4-47) or Eq. (4-52).

Eq. (4-47) has the advantage that it does not involve the turning point. This is a very desirable feature when one sets out to solve for the eigenvalues t_s which are defined by

$$\left[\frac{\partial G}{\partial u} + qG\right]_{u=0} = 0 \tag{4-54}$$

For example, Fock showed that

$$t_s(q) = \tau_s + \frac{\beta_o}{15}\left[8\tau_s^2 - \frac{3 + 4\tau_s q}{\tau_s - q^2}\right] + O\left(\beta_o^2\right) \tag{4-55}$$

where τ_s is a solution of

$$w_1'(\tau_s) - qw_1(\tau_s) = 0 \tag{4-56}$$

In order to see the advantages of Eq. (4-55), let us take $q = \infty$ and write

$$t_s(\infty) = \tau_s^\infty + \frac{8}{15}\beta_o\left[\tau_s^\infty\right]^2 + O\left(\beta_o^2\right) \tag{4-57}$$

where

$$w_1(\tau_s^\infty) = 0 \tag{4-58}$$

Let us now compare this with the procedure which involves the use of the turning point. Let us seek the roots λ_s defined by

$$F(0,\lambda_s) = 0 \tag{4-59}$$

Let us express $F(y,\lambda_s)$ in the form

$$F(y, \lambda_s) = \left(\frac{d\eta}{dy}\right)^{-1/2} w(-\eta) \tag{4-60}$$

where η is given by Eq. (4-42). We find that the roots of $F(0,\lambda_s) = 0$ are to be obtained from the relationship

$$-\tau_s^\infty = \zeta_o + \left(\frac{3}{2}\right)^{2/3}\frac{\alpha}{k^{4/3}} + \frac{1}{3}\frac{\alpha_o}{k^2}\zeta_o + \frac{1}{5}\left(\frac{2}{3}\right)^{2/3}\frac{\alpha_1}{k^{8/3}}\zeta_o^2 + \cdots \tag{4-61}$$

where ζ_0 is defined by

$$\frac{2}{3}\zeta_o^{3/2} = u(0) = k\int_{y_1}^{0}\sqrt{\mu_s + f(h)}\,dh = \int_{y_1}^{0}\sqrt{k^2 f(h) - \lambda_s}\,dh \qquad (4\text{-}62)$$

and the turning point y_1 is defined by

$$k^2 f(y_1) - \lambda_s = 0$$

Since λ_s appears explicitly in the factor $\sqrt{k^2 f(h) - \lambda_s}$ in the integrand of Eq. (4-62), and implicitly in the integration limit y_1, the inversion of Eq. (4-61) to solve for λ_s requires some skillful analysis. A method for achieving this inversion was described by Pekeris (Ref. 7). A comparison of Fock's procedure with that used by Pekeris shows very clearly that Fock's method is more direct and hence it would be advantageous to extend Fock's analysis to include higher-order terms.

4.5 The Contributions of Friedlander

Friedlander (Ref. 1) considered the problem of determining the behavior, as $s \to \infty$, of the eigenvalues λ of the differential equation

$$\frac{d^2y}{dx^2} + \left[\lambda - s^2 g(x)\right] y = 0 \qquad (4\text{-}63)$$

subject to the conditions that $y \to 0$ as $x \to \infty$ and either $y = 0$ or $dy/dx = 0$ at $x = 0$. He assumed that $g(x)$ was characterized by the conditions $g(0) > 0$ and $g'(0) \geq 0$ and assumed that $g(x)$ could be represented in the form of a Taylor series

$$g(x) = g_o + g_1 x + g_2 x^2 + g_3 x^3 + \cdots \quad , \quad g_m = \frac{1}{m!}\,g^{(m)}(0) \qquad (4\text{-}64)$$

Friedlander assumed that λ could be expressed in the form of an asymptotic series in inverse integral powers of $s^{\frac{2}{3}}$

$$\lambda = s^2 \sum_{m=0}^{\infty} c_m s^{-\frac{2m}{3}} \quad , \quad c_o = g_o \qquad (4\text{-}65)$$

and that $y(x)$ could be expressed in the form of a similar series

$$y(x) = \sum_{m=0}^{\infty} Y_m(\eta)\, s^{-\frac{2m}{3}} \quad , \quad \eta = s^{2/3} \tag{4-66}$$

Friedlander showed that the coefficients $Y_m(\eta)$ are determined by a system of recurrence relations

$$\frac{d^2Y_o}{d\eta^2} + (c_1 - g_1\eta)\, Y_o = 0 \tag{4-67a}$$

$$\frac{d^2Y_1}{d\eta^2} + (c_1 - g_1\eta)\, Y_1 = (g_2\eta^2 - c_2)\, Y_o \tag{4-67b}$$

$$\frac{d^2Y_2}{d\eta^2} + (c_1 - g_1\eta)\, Y_2 = (g_3\eta^3 - c_3)\, Y_o + (g_2\eta^2 - c_2)\, Y_1 \tag{4-67c}$$

$$\frac{d^2Y_m}{d\eta^2} + (c_1 - g_1\eta)\, Y_m = \sum_{j=1}^{m} \left(g_{j+1}\eta^{j+1} - c_{j+1}\right) Y_{m-j} \tag{4-67d}$$

Friedlander showed that

$$Y_o(\eta) = L_o\, Ai\,(g_1^{1/3}\eta - \alpha) \quad , \quad \alpha = c_1\, g_1^{-2/3} \tag{4-68}$$

where L_0 is a constant. He also showed that

$$Y_1(\eta) = \frac{\pi L_o}{g_1^{1/3}} \int_0^{\eta} \left[Ai\left(g_1^{1/3}\xi - \alpha\right) Bi\left(g_1^{1/3}\eta - \alpha\right) - Ai\left(g_1^{1/3}\eta - \alpha\right) Bi\left(g_1^{1/3}\xi - \alpha\right) \right] \left[g_2\xi^2 - c_2\right] Ai\left(g_1^{1/3}\xi - \alpha\right) d\xi \tag{4-69}$$

and gave a similar result for $Y_m(\eta)$. However, Friedlander did not evaluate these integrals and hence $Y_0(\eta)$ is the only $Y_m(\eta)$ which is given in its final form.

If we put $s = \pm ik$ in Eq. (4-63), and replace x by y, g(x) by f(x), and y(x) by $F(y,\lambda)$ we find that the function being studied (for

the pulse problem by Friedlander) is essentially the height gain function which was being studied by Fock.

$$\left[\mathrm{Ai}\left(s^{2/3} g_1^{1/3} y - \alpha\right)\right]_{s = \mp i k} = \frac{\exp\left(\mp i \frac{\pi}{6}\right)}{2\sqrt{\pi}} w_{1,2}\left(\alpha e^{\pm i\frac{\pi}{3}} - k^{2/3} g_1^{1/3} y\right) \quad (4\text{-}70)$$

Therefore, we can use Friedlander's theory as a starting point to obtain an extension of Fock's results. Let us return to the differential equation for $F(y, \lambda_s)$

$$\frac{d^2F}{dy^2} + \left[-\lambda_s + k^2 f(y)\right] F = 0 \quad (4\text{-}71)$$

and express $f(y)$ in the form

$$f(y) = f_o + f_1 + f_2 y^2 + f_3 y^3 + \cdots = f_o + f_1\left[y + h_2 y^2 + h_3 y^3 + \cdots\right] \quad (4\text{-}72)$$

where

$$h_n = \frac{f_n}{f_1} = \frac{1}{n! f_1}\left[\frac{d^n f(y)}{dy^n}\right]_{y=0} \quad (4\text{-}73)$$

Let us define t_s by means of

$$\lambda_s = k^2 f_o + k^{4/3} f_1^{2/3} t_s \quad (4\text{-}74)$$

and u by means of

$$u = k^{2/3} f_1^{1/3} y \quad (4\text{-}75)$$

We then find that if we define the parameter β to be

$$\beta = k^{-2/3} f_1^{-1/3} \quad (4\text{-}76)$$

that Eq. (4-71) [which is identical with Eq. (4-1)] takes the form

$$\frac{d^2G}{du^2} + \left[-t_s + u + \beta h_2 u^2 + \beta^2 h_3 u^3 + \beta^3 h_4 u^4 + \cdots\right] G = 0 \quad (4\text{-}77)$$

where

$$G(u, t_s) = F(y, \lambda_s) \quad (4\text{-}78)$$

A comparison of Eq. (4-78) with Eq. (4-43) reveals that if we let $\beta_0 = \beta h_2$ and neglect the terms in β^n for $n > 3$ in Eq. (4-77) that we have Fock's approximate differential equation.

Let us follow Friedlander [cf. Eq. (4-66)] and seek a representation for G(u,t) in the form

$$G(u, t) = \sum_{n=0}^{\infty} G_n (u, t) \beta^n \tag{4-79}$$

We then find that the $G_n(u,t)$ satisfy the equations

$$\left[\frac{d^2}{du^2} + (-t + u)\right] G_o = 0$$

$$\left[\frac{d^2}{du^2} + (-t + u)\right] G_1 = - h_2 u^2 G_o$$

$$\left[\frac{d^2}{du^2} + (-t + u)\right] G_2 = - h_2 u^2 G_1 - h_3 u^3 G_o$$

$$\left[\frac{d^2}{du^2} + (-t + u)\right] G_n = - \sum_{j=2}^{n+1} h_j u^j G_{n+1-j}$$

This set of recursive formulae is similar to Eq. (4-67). If we define

$$t = t(\tau) = \tau + \sum_{n=1}^{\infty} D_n(\tau) \beta^n = \sum_{n=0}^{\infty} D_n(\tau) \beta^n , \quad D_o = \tau \tag{4-80}$$

and

$$H (u, t) = G (u, t(\tau)) = \sum_{n=0}^{\infty} H_n (u, \tau) \beta^n \tag{4-81}$$

we find that

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] H_o = 0 \tag{4-82a}$$

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] H_1 = \left[- h_2 u^2 + D_1\right] H_o \tag{4-82b}$$

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] H_2 = \left[-h_2 u^2 + D_1\right] H_1 + \left[-h_3 u^3 + D_2\right] H_o \qquad (4\text{-}82c)$$

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] H_n = \sum_{j=2}^{n+1} \left[-h_j u^j + D_{j-1}\right] H_{n+1-j} \qquad (4\text{-}82d)$$

which is essentially the same as Eq. (4-67). Therefore, the theory given in Friedlander's paper for results such as those given in Eq. (4-68) and Eq. (4-69) can be used to determine the $H_n(u,\tau)$.

4.6 Determination of the $H_n(u,\tau)$

From the differential equation satisfied by the Airy function

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] w(\tau - u) = 0 \qquad (4\text{-}83)$$

we can show that

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] u^n w(\tau - u) = n(n-1)\, u^{n-2} w(\tau - u) - 2n\, u^{n-1} w'(\tau - u) \qquad (4\text{-}84)$$

and

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] u^n w'(\tau - u) = \left[(2n+1)\, u^n - 2\tau n u^{n-1}\right] w(\tau - u) + n(n-1)\, u^{n-2} w'(\tau - u) \qquad (4\text{-}85)$$

These results can be used to construct solution of the inhomogeneous equations

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] A_n(u, \tau) = u^n w(\tau - u) \qquad (4\text{-}86)$$

$$\left[\frac{d^2}{du^2} + (-\tau + u)\right] B_n(u, \tau) = u^n w'(\tau - u) \qquad (4\text{-}87)$$

If $A_n(u,\tau)$ and $B_n(u,\tau)$ are known, we can solve Eq. (4-82). However, it may be more convenient to define

$$x = \tau - u \quad , \quad u = \tau + x \qquad (4\text{-}88)$$

and replace Eq. (4-82) by

$$\left[\frac{d^2}{dx^2} - x\right] H_n = \sum_{j=2}^{n+1} \left[-h_j (\tau + x)^j + D_{j-1}\right] H_{n+1-j} \qquad (4\text{-}89)$$

In Table 4-1 we give some solutions of

$$\frac{d^2y}{dx^2} - x\,y = f(x)$$

which can be used to construct solutions of Eq. (4-89). For example, the solution of Eq. (4-82a) is

$$H_0(u,\tau) = c_0 w(\tau - u) \qquad (4\text{-}90a)$$

where c_0 is a constant. We can then show that the solution of Eq. (4-82b) is

$$H_1(u,\tau) = c_1 w(\tau - u) + c_0 \left\{ -\frac{1}{5} h_2 u\, w(\tau - u) - \left[\frac{1}{5} h_2 u^2 + \frac{4}{15} h_2 \tau u + \frac{8}{15} \tau^2 h_2 - D_1\right] w'(\tau - u) \right\} \qquad (4\text{-}90b)$$

where c_1 is a constant.

Before determining $H_2(u,\tau)$ it would be desirable to determine the constants c_0 and c_1 (or, at least determine the ratio c_1/c_0). Since the function $H(u,\tau)$ satisfies the homogeneous boundary condition derived from Eq. (4-54)

$$\left[\frac{\partial H}{\partial u} + qH\right]_{u=0} = 0 \qquad (4\text{-}91)$$

we must require that $H_n(u,\tau_s)$ satisfy the equations

$$\left[\frac{\partial H_o}{\partial u} + q\,H_o\right]_{u=0} = 0 \qquad (4\text{-}92a)$$

$$\left[\frac{\partial H_1}{\partial u} + q\,H_1\right]_{u=0} = 0 \qquad (4\text{-}92b)$$

We observe that Eq. (4-90a) and Eq. (4-92a) lead to the equation

$$w'(\tau_s) - q\,w(\tau_s) = 0 \qquad (4\text{-}93)$$

Table 4-1

PARTICULAR SOLUTIONS OF $\frac{d^2y}{dx^2} - xy = f(x)$

f(x)	y(x)
0	$w(x)$
$w(x)$	$w'(x)$
$w'(x)$	$\frac{1}{2} x\, w(x)$
$x\, w(x)$	$\frac{1}{3} x\, w'(x)$
$x\, w'(x)$	$\frac{1}{4} x^2 w(x) - \frac{1}{2} w'(x)$
$x^2 w(x)$	$-\frac{1}{5} x\, w(x) + \frac{1}{5} x^2 w'(x)$
$x^2 w'(x)$	$\left[\frac{1}{6} x^3\right] w(x) - x \frac{1}{3} w'(x)$
$x^3 w(x)$	$\left[-\frac{3}{14} x^2\right] w(x) + \left[\frac{3}{7} + x^3 \frac{1}{7}\right] w'(x)$
$x^3 w'(x)$	$\left[\frac{3}{10} x + \frac{1}{8} x^4\right] w(x) - \frac{3}{10} x^2 w'(x)$
$x^4 w(x)$	$\left[-\frac{2}{9} x^3\right] w(x) + \left[\frac{4}{9} x + \frac{1}{9} x^4\right] w'(x)$
$x^4 w'(x)$	$\left[\frac{3}{7} x^2 + \frac{1}{10} x^5\right] w(x) + \left[-\frac{6}{7} - \frac{2}{7} x^3\right] w'(x)$
$x^5 w(x)$	$\left[-\frac{6}{11} x - \frac{5}{22} x^4\right] w(x) + \left[\frac{6}{11} x^2 + \frac{1}{11} x^5\right] w'(x)$
$x^5 w'(x)$	$\left[\frac{5}{9} x^3 + \frac{1}{12} x^6\right] w(x) + \left[-\frac{10}{9} x - \frac{5}{18} x^4\right] w'(x)$
$x^6 w(x)$	$\left[-\frac{90}{91} x^2 - \frac{3}{13} x^5\right] w(x) + \left[\frac{180}{91} + \frac{60}{91} x^3 + \frac{1}{13} x^6\right] w'(x)$
$x^6 w'(x)$	$\left[\frac{18}{11} x + \frac{15}{22} x^4 + \frac{1}{14} x^7\right] w(x) + \left[-\frac{18}{11} x^2 - \frac{3}{11} x^5\right] w'(x)$
$x^7 w(x)$	$\left[-\frac{14}{9} x^3 - \frac{7}{30} x^6\right] w(x) + \left[\frac{28}{9} x + \frac{7}{9} x^4 + \frac{1}{15} x^7\right] w'(x)$
$x^7 w'(x)$	$\left[\frac{45}{13} x^2 + \frac{21}{26} x^5 + \frac{1}{16} x^8\right] w(x) + \left[-\frac{90}{3} - \frac{30}{13} x^3 - \frac{7}{26} x^6\right] w'(x)$
$x^8 w(x)$	$\left[-\frac{1008}{187} x - \frac{420}{187} x^4 - \frac{4}{17} x^7\right] w(x) + \left[\frac{1008}{187} x^2 + \frac{168}{187} x^5 + \frac{1}{17} x^8\right] w'(x)$
$x^8 w'(x)$	$\left[\frac{56}{9} x^3 + \frac{14}{15} x^6 + \frac{1}{18} x^9\right] w(x) + \left[-\frac{112}{9} x - \frac{28}{9} x^4 - \frac{4}{15} x^7\right] w'(x)$

which determines τ_s uniquely. If we observe that Eq. (4-90b) and Eq. (4-92b) lead to

$$\left[\frac{\partial H_1(u,\tau)}{\partial u} + q\,H_1(u,\tau)\right]_{u=0} = -c_1\left[w'(\tau) - q\,w(\tau)\right] + c_o\left\{\left[-\frac{1}{5}h_2 w(\tau)\right.\right.$$

$$\left.\left. - \frac{4}{15}h_2\tau w'(\tau) + \frac{8}{15}h_2\tau^3 w(\tau) - D_1\tau\,w(\tau)\right] + q\left[-\frac{8}{15}h_2\tau^2 + D_1\right]w'(\tau)\right\} = 0 \quad (4\text{-}94)$$

from which it is seen that the value of D_1 is given by

$$(\tau - q^2)\,D_1 = h_2\left[-\frac{1}{5} - \frac{4}{15}\tau q + \frac{8}{15}\tau^2(\tau - q^2)\right] \quad (4\text{-}95)$$

Therefore, we observe that the boundary condition given by Eq. (4-91) determines the constants D_n but leaves the values of c_n undetermined.

The c_n can be determined by requiring that the eigenfunctions $F(y,\lambda_s)$ have the property

$$\int_0^\infty F^2(y,\lambda_s)\,dy = 1 \quad (4\text{-}96)$$

This leads to

$$\int_0^\infty H^2(u,\tau_s)\,du = k^{2/3} f_1^{1/3} = \beta^{-1} \quad (4\text{-}97)$$

If we use the property

$$H^2 = \left[\sum_{n=0}^{\infty} H_n\beta^n\right]^2 = H_o^2 + 2H_o H_1\beta + (2H_o H_2 + H_1^2)\beta^2 + \cdots \quad (4\text{-}98)$$

and require that

$$\int_0^\infty \left[H_o(u,\tau_s)\right]^2 du = c_o^2 \int_0^\infty w^2(\tau_s - u)\,du = k^{2/3} f_1^{1/3} \quad (4\text{-}99)$$

we observe that Eq. (4-97) is satisfied provided

$$\int_0^\infty H_o(u,\tau_s)\,H_1(u,\tau_s)\,du = 0 \quad (4\text{-}100a)$$

$$\int_0^\infty \left[2H_o(u,\tau_s)\,H_2(u,\tau_s) + H_1^2(u,\tau_s)\right]du = 0 \quad (4\text{-}100b)$$

along with similar equations involving H_3, H_4, etc., This procedure has the advantage that the ratio c_n/c_0 is independent of β.

We can determine c_0 from Eq. (4-99) by observing that

$$\int_0^\infty w^2(\tau_s - u)\, du = \tau_s \left[w^2(\tau_s)\right] - \left[w'(\tau_s)\right]^2 \tag{4-101}$$

since

$$\frac{d}{du}\left\{(\tau - u)\left[w(\tau - u)\right]^2 - \left[w'(\tau - u)\right]^2\right\} = -\left[w(\tau - u)\right]^2$$

Therefore, we find that

$$c_o^{\,2} = \frac{k^{2/3} f_1^{\,1/3}}{\tau_s\left[w(\tau_s)\right]^2 - \left[w'(\tau_s)\right]^2} = \frac{k^{2/3} f_1^{\,1/3}}{\left[\tau_s - q^2\right]\left[w(\tau_s)\right]^2} \tag{4-102}$$

In order to determine c_1 we must evaluate the integrals that occur in the equation

$$\begin{aligned}\frac{c_1}{c_o}\int_0^\infty w^2(\tau - u)\, du &= \frac{c_1}{c_o}\left\{\tau\left[w(\tau)\right]^2 - \left[w'(\tau)\right]^2\right\} = -D_1\int_0^\infty w(\tau - u)w'(\tau - u)\, du \\ &\quad + \frac{h_2}{15}\int_0^\infty \left[3uw(\tau - u) + (3u^2 + 4u\tau + 8\tau^2)w'(\tau - u)\right]w(\tau - u)\, du \\ &= -\frac{D_1}{2}\left[w(\tau)\right]^2 + \frac{h_2}{15}\left[-3\int_{-\infty}^{\tau}(\zeta + \tau)\, w^2(\zeta)\, d\zeta\right. \\ &\quad \left. + \int_{-\infty}^{\tau}(15\tau^2 - 10\zeta\tau + 3\zeta^2)\, w(\zeta)\, w'(\zeta)\, d\zeta\right]\end{aligned} \tag{4-103}$$

We can show that

$$3\int_{-\infty}^{\tau}(\zeta + \tau)\, w^2(\zeta)\, d\zeta = \left\{4\tau^2\left[w(\tau)\right]^2 - 4\tau\left[w'(\tau)\right]^2 + w(\tau)\, w'(\tau)\right\}$$

$$\int_{-\infty}^{\tau}(15\tau^2 - 10\zeta\tau + 3\zeta^2)\, w(\zeta)\, w'(\zeta)\, d\zeta = \left\{8\tau^2\left[w(\tau)\right]^2 - 4\tau\left[w'(\tau)\right]^2 - w(\tau)\, w'(\tau)\right\}$$

Since τ_s is a solution of $w'(\tau) = q\, w(\tau)$, we find that the integrals in Eq. (4-103) lead to the result

$$\frac{c_1}{c_o}(\tau - q^2) = -\frac{D_1}{2} + \frac{h_2}{15}\left|4\tau^2 - 2q\right| \qquad (4\text{-}104)$$

Because of the labour involved in the determination of the constants c_n which will occur in the terms $c_n w(\tau - u)$ which are to be added to the solutions of the functions $H_n(u,\tau)$ which satisfy Eq. (4-82), it would be most convenient to follow Friedlander's suggestion and take $c_0 = 1$ and $c_n = 0$ for $n \geq 1$. We should then replace Eq. (4-81) by

$$H(u, t) = G(u, t(\tau)) = c(\beta,\tau)\left[w(\tau - u) + \sum_{n=1}^{\infty} H_n(u, \tau)\,\beta^n\right] \qquad (4\text{-}105)$$

and determine $C(\beta,\tau)$ from Eq. (4-97). From Eq. (4-90b) we know that for this case

$$H_1(u,\tau) = -\frac{1}{5}h_2 uw(\tau - u) - \left[\frac{1}{5}h_2u^2 + \frac{4}{15}h_2\tau u + \frac{8}{15}h_2\tau^2 - D_1\right]w'(\tau-u) \qquad (4\text{-}106)$$

From Eq. (4-105) and Eq. (4-97), we find that $C(\beta,\tau)$ is determined by the relation

$$\frac{1}{\beta} = C^2(\beta, \tau)\, w^2(\tau)\left\{(\tau - q^2) + \left[D_1 - \frac{4}{15}h_2\,(2\tau^2 - q)\right]\beta + O(\beta^2)\right\} \qquad (4\text{-}107)$$

We can also seek a solution in the form of Eq. (4-105) in which $H_1(u,\tau)$ is of the form

$$H_1(u, \tau) = (c_1 - \frac{1}{5}h_2u)\,w(\tau-u) - \left[\frac{1}{5}h_2u^2 + \frac{4}{15}h_2\tau u + \frac{8}{15}h_2\tau^2 - D_1\right]w'(\tau-u) \qquad (4\text{-}108)$$

where c_1 is a constant independent of β. For example, Eq. (4-44), which was taken from Fock's paper, is of the form of Eq. (4-108) if we take

$$c_1 = -\frac{2}{15}h_2\tau \qquad (4\text{-}109)$$

Let us now turn to the problem of the determination of $H_2(u,\tau)$. We have discussed the role played by the choice of c_1 in the determination of $H_2(u,\tau)$. For example, a term $c_1 w(\tau - y)$ in $H_1(u,\tau)$

leads to a term in $H_2(u,\tau)$ which is of the form

$$c_1\left\{-\frac{1}{5}h_2 u\, w(\tau-u) - \left[\frac{1}{5}h_2 u^2 + \frac{4}{15}h_2\tau u + \frac{8}{15}h_2\tau^2 - D_1\right] w'(\tau-u)\right\} \qquad (4\text{-}110)$$

If we take c_1 to be defined by Eq. (4-109), we find that Eq. (4-82c), with the aid of the entries in Table 4-1, leads to

$$\begin{aligned} H_2(u,\tau) = c_2\, w(\tau-u) + \Big[& -\frac{1}{50}h_2{}^2 u^5 - \frac{1}{30}h_2{}^2\tau u^4 - \frac{4}{45}h_2{}^2\tau^2 u^3 + \frac{1}{5}h_2 D_1 u^3 \\ & + \frac{1}{15}h_2 D_1 \tau u^2 + \left(-\frac{3}{14}h_3 + \frac{9}{70}h_2{}^2\right)u^2 + \frac{4}{15}h_2 D_1\tau^2 u \\ & \left(-\frac{6}{35}h_3 + \frac{22}{105}h_2{}^2\right)\tau u - \frac{1}{2}D_1{}^2 u\Big] w(\tau-u) + \Big[-\frac{1}{7}h_3 \\ & - \frac{3}{35}h_2{}^2\Big)u^3 - \left(\frac{6}{35}h_3 - \frac{22}{105}h_2{}^2\right)\tau u^2 - \left(\frac{8}{35}h_3 \\ & - \frac{144}{315}h_2{}^2\right)\tau^2 u - \frac{7}{15}h_2 D_1 u - \left(\frac{6}{35}h_3 - \frac{288}{315}h_2{}^2\right)\tau^3 \\ & - \frac{6}{5}h_2 D_1\tau + \left(\frac{3}{7}h_3 - \frac{9}{35}h_2{}^2 + D_2\right)\Big] w'(\tau-u) \end{aligned} \qquad (4\text{-}111)$$

In the next section we will show than an extension of Fock's results summarized in Eq. (4-46) and Eq. (4-47) suggests that a useful value for the arbitrary constant c_2 is

$$c_2 = \frac{32}{225}h_2{}^2\tau^5 - \frac{8}{15}h_2 D_1\tau^3 + \left(-\frac{4}{35}h_3 + \frac{58}{315}h_2{}^2\right)\tau^2 + \frac{1}{2}D_1{}^2\tau - \frac{2}{15}h_2 D_1 \qquad (4\text{-}112)$$

The obtaining of $H_3(u,\tau)$, $H_4(u,\tau)$, etc., by the above methods is exceedingly tedious although it is a very straight-foward process if one uses Table 4-1 along with Eq. (4-89).

4.7 Determination of $\eta(u,t)$

Let us follow Eq. (4-50) and express the solution of Eq. (4-77) in the form

$$G(u,t) = \left(\frac{d\eta}{du}\right)^{-\frac{1}{2}} w_{1,2}(-\eta) = \frac{\sqrt{\pi}}{\sqrt[6]{12}}\exp\left(\pm i\frac{2\pi}{3}\right)\left(\frac{d\eta}{du}\right)^{-\frac{1}{2}} h_{1,2}(\eta) \qquad (4\text{-}113)$$

where $w(z)$ and $h(\eta)$ are solutions of the Airy differential equations

$$\frac{d^2w}{dz^2} - z\,w = 0 \quad , \qquad \frac{d^2h}{d\eta^2} + \eta\, h = 0$$

According to Eq. (4-51), the function $\eta(u,t)$ is a solution of

$$\eta\,(\dot{\eta})^2 - \frac{3}{4}\left(\frac{\dot{\eta}}{\eta}\right)^2 + \frac{1}{2}\left(\frac{\ddot{\eta}}{\eta}\right) = u - t_s + \sum_{n=2}^{\infty} h_n\, u^n\, \beta^{n-1} \qquad (4\text{-}114)$$

where the dot denotes differentiation with respect to u. We observe that Eq. (4-114) is essentially the same equation as Eq. (4-29) except that we have used Eq. (4-74) to define t_s and Eq. (4-75) to define u.

Although Eq. (4-114) is non-linear, and of the third order, it is simpler to work with than the linear, second order equation which defines G(u,t). The simplicity comes from the fact that $\eta(u,t)$ can be expressed in the form

$$\eta(u,\,t) = (u - t) + \sum_{n=1}^{\infty} p_n\,(u,\,t)\,\beta^n \qquad (4\text{-}115)$$

where $p_n(u,t)$ is a sum of homogeneous polynomials in u and t for which the degree of the polynomials differ by three. Thus

$$p_1\,(u,\,t) = a_1\,u^2 + b_1\,u\,t + c_1\,t^2 \qquad (4\text{-}116a)$$

$$p_2\,(u,\,t) = a_2\,u^3 + b_2\,u^2\,t + c_2\,u\,t^2 + d_2\,t^3 + \alpha_2 \qquad (4\text{-}116b)$$

$$p_3\,(u,\,t) = a_3\,u^4 + b_3\,u^3\,t + c_3\,u^2\,t^2 + d_3\,u\,t^3 + e_3\,t^4 + \alpha_3\,u + \beta_3\,t \qquad (4\text{-}116c)$$

If we put Eq. (4-116) in Eq. (4-115), and then use Eq. (4-114), we can determine the constants $a_n, b_n, \ldots, \alpha_n, \beta_n, \ldots$, etc. The first several terms in the quantities in Eq. (4-114) are

$$\eta\left[\frac{d\eta}{du}\right]^2 = (u - t) + \beta\left[5a_1\,u^2 + (3b_1 - 4a_1)\,ut + (c_1 - 2b_1)\,t^2\right] + \cdots \qquad (4\text{-}117a)$$

$$-\frac{3}{4}\left[\frac{d\eta}{du}\right]^{-2}\left[\frac{d^2\eta}{du^2}\right]^2 = 4a_1^{\,2}\,\beta^2 + \cdots \qquad (4\text{-}117b)$$

$$\frac{1}{2}\left[\frac{d\eta}{du}\right]^{-1}\left[\frac{d^3\eta}{du^3}\right] = 6a_2\,\beta^2 + \cdots \qquad (4\text{-}117c)$$

In Eq. (4-118) we give the resulting expression for $\eta(u,t)$ for terms as high as those in β^4.

$$\eta(u,t) = [u - t]$$

$$+\beta\left[\frac{1}{5}h_2 u^2 + \frac{4}{15}h_2 t u + \frac{8}{15}h_2 t^2\right]$$

$$+\beta^2\left[\left(\frac{1}{7}h_3 - \frac{8}{175}h_2^2\right)u^3 + \left(\frac{6}{35}h_3 - \frac{68}{525}h_2^2\right)tu^2 + \left(\frac{8}{35}h_3 - \frac{496}{1575}h_2^2\right)t^2u + \left(\frac{16}{35}h_3 - \frac{1328}{1575}h_2^2\right)t^3 - \left(\frac{3}{7}h_3 - \frac{9}{35}h_2^2\right)\right]$$

$$+\beta^3\left[\left(\frac{1}{9}h_4 - \frac{22}{315}h_2 h_3 + \frac{148}{7875}h_2^3\right)u^4 + \left(\frac{8}{63}h_4 - \frac{3212}{17325}h_2 h_3 + \frac{608}{7875}h_2^3\right)tu^3 + \left(\frac{16}{105}h_4 - \frac{2288}{5775}h_2 h_3\right.\right.$$
$$\left.+\frac{1808}{7875}h_2^3\right)t^2u^2 + \left(\frac{64}{315}h_4 - \frac{13376}{17325}h_2 h_3 + \frac{43712}{70875}h_2^3\right)t^3u + \left(\frac{128}{315}h_4 - \frac{33088}{17325}h_2 h_3 + \frac{128512}{70875}h_2^3\right)t^4$$
$$\left.-\left(\frac{4}{9}h_4 - \frac{28}{45}h_2 h_3 + \frac{56}{225}h_2^3\right)u - \left(\frac{80}{63}h_4 - \frac{1628}{693}h_2 h_3 + \frac{1768}{1575}h_2^3\right)t\right]$$

$$+\beta^4\left[\left(\frac{1}{11}h_5 - \frac{28}{495}h_2 h_4 - \frac{15}{539}h_3^2 + \frac{5464}{121275}h_2^2 h_3 - \frac{29584}{3031875}h_2^4\right)u^5 + \left(\frac{10}{99}h_5 - \frac{1504}{10395}h_2 h_4 - \frac{1707}{24255}h_3^2\right.\right.$$
$$+\frac{321392}{1819125}h_2^2 h_3 - \frac{93584}{1819125}h_2^4\Big)tu^4 + \left(\frac{80}{693}h_5 - \frac{2176}{7425}h_2 h_4 - \frac{16656}{121275}h_3^2 + \frac{888592}{1819125}h_2^2 h_3\right.$$
$$-\frac{327808}{1819125}h_2^4\Big)t^2u^3 + \left(\frac{32}{231}h_5 - \frac{8768}{17325}h_2 h_4 - \frac{3488}{13475}h_3^2 + \frac{708704}{606375}h_2^2 h_3 - \frac{2914624}{5457375}h_2^4\right)t^3u^2$$
$$+\left(\frac{128}{693}h_5 - \frac{46336}{51975}h_2 h_4 - \frac{18176}{40425}h_3^2 + \frac{4738432}{1819125}h_2^2 h_3 - \frac{24237568}{16372125}h_2^4\right)t^4u + \left(\frac{256}{693}h_5 - \frac{109568}{51975}h_2 h_4\right.$$
$$-\frac{42688}{40425}h_3^2 + \frac{38593536}{5457375}h_2^2 h_3 - \frac{371579392}{81860625}h_2^4\Big)t^5 - \left(\frac{6}{11}h_5 - \frac{388}{495}h_2 h_4 - \frac{1044}{2695}h_3^2 + \frac{115372}{121275}h_2^2 h_3\right.$$
$$-\frac{165512}{606375}h_2^4\Big)u^2 - \left(\frac{112}{99}h_5 - \frac{3568}{1485}h_2 h_4 - \frac{8828}{8085}h_3^2 + \frac{1330096}{363825}h_2^2 h_3 - \frac{2327168}{1819125}h_2^4\right)tu$$
$$\left.-\left(\frac{1808}{693}h_5 - \frac{71552}{10395}h_2 h_4 - \frac{24076}{8085}h_3^2 + \frac{12920064}{1091475}h_2^2 h_3 - \frac{25097808}{5457375}h_2^4\right)t^2\right] + \cdots \qquad (4\text{-}118)$$

LOCKHEED MISSILES & SPACE COMPANY

In many applications it is more convenient to let t be defined by Eq. (4-80). If we recall the definition of $H(u,\tau)$ in Eq. (4-81) we can write

$$H(u,\tau) = G\left[u, t(\tau)\right] = \left(\frac{d\gamma}{du}\right)^{-\frac{1}{2}} w_{1,2}(-\gamma) \qquad (4\text{-}119)$$

where γ is defined by

$$\begin{aligned}
\gamma(u,\tau) = \eta\left[u, t(\tau)\right] &= (u-\tau) + \beta\left[\tfrac{1}{5} h_2 u^2 + \tfrac{4}{15} h_2 \tau u + \tfrac{8}{15} h_2 \tau^2 - D_1\right] \\
&+ \beta^2\Big[(\tfrac{1}{7} h_3 - \tfrac{8}{175} h_2^2) u^3 + \tfrac{6}{35} h_3 - \tfrac{68}{525} h_2^2) \tau u^2 + (\tfrac{8}{35} h_3 \\
&\quad - \tfrac{496}{1575} h_2^2) \tau^2 u + \tfrac{4}{15} h_2 D_1 u + (\tfrac{16}{35} h_3 \\
&\quad - \tfrac{1328}{1575} h_2^2) \tau^3 + \tfrac{16}{15} h_2 D_1 \tau - (\tfrac{3}{7} h_3 - \tfrac{9}{35} h_2^2 + D_2)\Big] \\
&+ \beta^3\Big[(\tfrac{1}{9} h_4 - \tfrac{22}{315} h_2 h_3 + \tfrac{148}{7875} h_2^3) u^4 + (\tfrac{8}{63} h_4 \\
&\quad - \tfrac{3212}{17325} h_2 h_3 + \tfrac{608}{7875} h_2^3) \tau u^3 + (\tfrac{16}{105} h_4 \\
&\quad - \tfrac{2288}{5775} h_2 h_3 + \tfrac{1808}{7875} h_2^3) \tau^2 u^2 + (\tfrac{6}{35} h_3 \\
&\quad - \tfrac{68}{525} h_2^2) D_1 u^2 + (\tfrac{64}{315} h_4 - \tfrac{13376}{17325} h_2 h_3 \\
&\quad + \tfrac{43712}{70875} h_2^3) \tau^3 u - (\tfrac{4}{9} h_4 - \tfrac{28}{45} h_2 h_3 + \tfrac{56}{225} h_2^3) u \\
&\quad + (\tfrac{16}{35} h_3 \tau D_1 + \tfrac{4}{15} h_2 D_2 - \tfrac{992}{1575} h_2^2 \tau D_1) u \\
&\quad + (\tfrac{128}{315} h_4 - \tfrac{33088}{17325} h_2 h_3 + \tfrac{128512}{70875} h_2^3) \tau^4 \\
&\quad + (\tfrac{48}{35} h_3 - \tfrac{1328}{525} h_2^2) \tau^2 D_1 - (\tfrac{80}{63} h_4 \\
&\quad - \tfrac{1628}{693} h_2 h_3 + \tfrac{1768}{1575} h_2^3 - \tfrac{16}{15} h_2 D_2) \tau \\
&\quad + (\tfrac{8}{15} h_2 D_1^2 - D_3)\Big] + O(\beta^4)
\end{aligned} \qquad (4\text{-}120)$$

Let us turn now to the construction of the derivative of $F(y,\lambda)$ or $G(u,t)$. We begin by observing that

$$\frac{\partial F(y,\lambda)}{\partial y} = k^{2/3} f_1^{1/3} \frac{\partial G(u,t)}{\partial u} = \frac{1}{\beta} \frac{\partial G(u,t)}{\partial u} = \frac{1}{\beta} \frac{\partial H(u,\tau)}{\partial u} \tag{4-121}$$

From Eq. (4-113), we can write

$$\frac{\partial G}{\partial u} = \sqrt{\frac{d\eta}{du}} \left\{ w'_{1,2}(-\eta) + \frac{1}{2} \left[\frac{d\eta}{du}\right]^{-2} \frac{d^2\eta}{du^2} w_{1,2}(-\eta) \right\} \tag{4-122a}$$

or

$$\frac{\partial G}{\partial u} = \frac{\sqrt{\pi}}{\sqrt[6]{12}} \exp(\pm i \tfrac{2\pi}{3}) \sqrt{\frac{d\eta}{du}} \left\{ h'_{1,2}(\eta) - \frac{1}{2} \frac{\eta}{(\eta)^2} h_{1,2}(\eta) \right\} \tag{4-122b}$$

Let us now represent the expression in curly braces in Eq. (4-122b) in the form

$$h'(\eta) - \frac{1}{2}\frac{\eta}{[\eta]^2} h(\eta) = \left[1 + \alpha_1 \beta + \alpha_2 \beta^2 + \alpha_3 \beta^3 + \cdots\right] h'(\eta + \gamma_1\beta + \gamma_2\beta^2 + \gamma_3 \beta^3 + \cdots) \tag{4-123}$$

Let us first consider a more general result. Let us solve for the A_n and B_n which are defined in Eq. (4-124)

$$h'(\eta) - \Delta h(\eta) = [\,1 + \Delta A_1 + \Delta^2 A_2 + \Delta^3 A_3 + \cdots] h'[\eta + \Delta B_1 + \Delta^2 B_2 + \Delta^3 B_3 + \cdots] \tag{4-124}$$

In order to obtain this result, we observe that

$$\begin{aligned}
h^{(2)}(\eta) &= -\eta\, h(\eta) \\
h^{(3)}(\eta) &= -\eta\, h'(\eta) - h(\eta) \\
h^{(4)}(\eta) &= \eta^2 h(\eta) - 2h'(\eta) \\
h^{(5)}(\eta) &= 4\eta\, h(\eta) + \eta^2 h'(\eta) \\
h^{(6)}(\eta) &= (4 - \eta^3) h(\eta) + 6\eta\, h'(\eta) \\
h^{(7)}(\eta) &= (10 - \eta^3) h'(\eta) - 9\eta^2 h(\eta) \\
h^{(n)}(\eta) &= \frac{d}{d\eta^n} h(\eta)
\end{aligned} \tag{4-125}$$

and use the Taylor series expansion

$$
\begin{aligned}
&h'(\eta + \Delta B_1 + \Delta^2 B_2 + \Delta^3 B_3 + \Delta^4 B_4 + \cdots) \\
&= h'(\eta) - \left[\Delta B_1 + \Delta^2 B_2 + \Delta^3 B_3 + \Delta^4 B_4 + \cdots\right] \eta\, h(\eta) \\
&\quad - \frac{1}{2}\left[\Delta^2 B_1^{\,2} + 2\Delta^3 B_1 B_2 + (2B_1 B_3 + B_2^{\,2})\,\Delta^4 + \cdots\right]\left[\eta h'(\eta) + h(\eta)\right] \\
&\quad + \frac{1}{6}\left[\Delta^3 B_1^{\,3} + 3\Delta^4 B_1^{\,2} B_2 + \cdots\right]\left[\eta^2 h(\eta) - 2h'(\eta)\right] \\
&\quad + \frac{1}{24}\left[\Delta^4 B_1^{\,4} + \cdots\right]\left[4\eta\, h(\eta) + \eta^2 h'(\eta)\right] + \cdots
\end{aligned}
\tag{4-126}
$$

in Eq. (4-124). By equating the coefficients of Δ on each side of Eq. (4-124) we find that

$$
\begin{aligned}
A_1 &= 0 & B_1 &= \frac{1}{\eta} \\
A_2 &= \frac{1}{2\eta} & B_2 &= -\frac{1}{2\eta^3} \\
A_3 &= -\frac{1}{6\eta^3} & B_3 &= \frac{1}{2}\frac{1}{\eta^5} - \frac{1}{3}\frac{1}{\eta^2} \\
A_4 &= \frac{9}{8}\frac{1}{\eta^5} - \frac{1}{8}\frac{1}{\eta^2} & B_4 &= -\frac{5}{8}\frac{1}{\eta^7} + \frac{5}{12}\frac{1}{\eta^4}
\end{aligned}
\tag{4-127}
$$

In order to use Eq. (4-123), we define

$$
\begin{aligned}
2\Delta = \left(\frac{d\eta}{du}\right)^{-2}\frac{d^2\eta}{du^2} &= \left\{\frac{2}{5}h_2\beta + \beta^2\left[+\frac{12}{35}h_3 t - \frac{248}{525}h_2^{\,2} t\right]\right. \\
&\quad + \beta^3\left[\left(\frac{4}{3}h_4 - \frac{88}{105}h_2 h_3 + \frac{1960}{2625}h_2^{\,3}\right)u^2 + \left(\frac{16}{21}h_4\right.\right. \\
&\qquad \left. - \frac{6424}{5775}h_2 h_3 + \frac{3360}{2625}h_2^{\,3}\right) t\, u + \left(\frac{32}{105}h_4\right. \\
&\qquad \left.\left. - \frac{4576}{5775}h_2 h_3 + \frac{7360}{7875}h_2^{\,3}\right) t^2\right] \\
&\quad + \beta^4\left[\left(\frac{20}{11}h_5 - \frac{1264}{495}h_2 h_4 + \frac{217696}{121275}h_2^{\,2} h_3 - \frac{300}{539}h_3^{\,2}\right.\right. \\
&\qquad \left. - \frac{2657920}{3031875}h_2^{\,4}\right) u^3 + \left(\frac{40}{33}h_5 - \frac{11648}{3465}h_2 h_4 - \frac{6828}{8085}h_3^{\,2}\right. \\
&\qquad \left. + \frac{2366032}{606375}h_2^{\,2} h_3 - \frac{7118240}{3031875}h_2^{\,4}\right) t\, u^2 + \left(\frac{160}{231}h_5\right. \\
&\qquad - \frac{137856}{51975}h_2 h_4 - \frac{33312}{40425}h_3^{\,2} + \frac{2905696}{606375}h_2^{\,2} h_3 \\
&\qquad \left. - \frac{9343744}{3031875}h_2^{\,4}\right) t^2 u + \left(\frac{64}{231}h_5 - \frac{23168}{17325}h_2 h_4\right.
\end{aligned}
$$

(eq. concluded on following page)

$$- \frac{6976}{13475} h_3^2 + \frac{2048192}{606375} h_2^2 h_3 - \frac{61579520}{27286875} h_2^4) t^3$$
$$+ (- \frac{12}{11} h_5 + \frac{952}{495} h_2 h_4 + \frac{2088}{2695} h_3^2 - \frac{291112}{121275} h_2^2 h_3$$
$$+ \frac{451760}{606375} h_2^4)\Big] + O(\beta^5)\Big\} \qquad (4\text{-}128)$$

From the definition of $\eta(u,t)$ in Eq. (4-118), we can obtain a representation for the reciprocals of η which occur in Eq. (4-127). We first define

$$z = u - t \qquad (4\text{-}129)$$

and then find that

$$\frac{1}{\eta} = \frac{1}{z}\Big\{1 - \frac{\beta}{z}\Big[\frac{1}{5} h_2 u^2 + \frac{4}{15} h_2 t u + \frac{8}{15} h_2 t^2\Big]$$
$$+ \beta^2\Big[-\frac{1}{z}\Big[(\frac{1}{7} h_3 - \frac{8}{175} h_2^2) u^3 + (\frac{6}{35} h_3 - \frac{68}{525} h_2^2) tu^2 + (\frac{8}{35} h_3$$
$$- \frac{496}{1575} h_2^2) t^2 u + (\frac{16}{35} h_3 - \frac{1328}{1575} h_2^2) t^3 - (\frac{3}{7} h_3 - \frac{9}{35} h_2^2)\Big]$$
$$+ \frac{1}{z^2}\Big[\frac{1}{25} h_2^2 u^4 + \frac{8}{75} h_2^2 tu^3 + \frac{64}{225} h_2^2 t^2 u^2 + \frac{64}{225} h_2^2 t^3 u + \frac{64}{225} h_2^2 t^4\Big]\Big]$$
$$+ \beta^3\Big[-\frac{1}{z}\Big[(\frac{1}{9} h_4 - \frac{22}{315} h_2 h_3 + \frac{148}{7875} h_2^3) u^4 + (\frac{8}{63} h_4 - \frac{3212}{17325} h_2 h_3$$
$$+ \frac{608}{7875} h_2^3) tu^3 + (\frac{16}{105} h_4 - \frac{2288}{5775} h_2 h_3 + \frac{1808}{7875} h_2^3) t^2 u^2$$
$$+ (\frac{64}{315} h_4 - \frac{13376}{17325} h_2 h_3 + \frac{43712}{70875} h_2^3) t^3 u + (\frac{128}{315} h_4 - \frac{33088}{17325} h_2 h_3$$
$$+ \frac{128512}{70875} h_2^3) t^4 - (\frac{4}{9} h_4 - \frac{28}{45} h_2 h_3 + \frac{56}{225} h_2^3) u - (\frac{80}{63} h_4 - \frac{1628}{693} h_2 h_3$$
$$+ \frac{1768}{1575} h_2^3) t\Big] + \frac{1}{z^2}\Big[(\frac{2}{35} h_2 h_3 - \frac{16}{875} h_2^3) u^5 + (\frac{8}{105} h_2 h_3$$
$$- \frac{64}{2625} h_2^3) tu^4 + (\frac{1136}{2625} h_2 h_3 - \frac{1920}{7875} h_2^3) t^2 u^3 + (\frac{1472}{2625} h_2 h_3$$
$$- \frac{15200}{23625} h_2^3) t^3 u^2 - (\frac{6}{35} h_2 h_3 - \frac{18}{175} h_2^3) u^2 + (\frac{256}{525} h_2 h_3 - \frac{18560}{23625} h_2^3) t^4 u$$
$$- (\frac{24}{105} h_2 h_3 - \frac{72}{525} h_2^3) tu + (\frac{256}{525} h_2 h_3 - \frac{21248}{23625} h_2^3) t^5 - (\frac{48}{105} h_2 h_3$$
$$- \frac{144}{525} h_2^3) t^2\Big] - \frac{1}{z^3}\Big[\frac{1}{125} h_2^3 u^6 + \frac{4}{125} h_2^3 t u^5 + \frac{120}{1125} h_2^3 t^2 u^4$$
$$+ \frac{640}{3375} h_2^3 t^3 u^3 + \frac{960}{3375} h_2^3 t^4 u^2 + \frac{768}{3375} h_2^3 t^5 u + \frac{512}{3375} h_2^3 t^6\Big]\Big] + O(\beta^4)\Big\} \qquad (4\text{-}130)$$

We can now use Eq. (4-129) and Eq. (4-130) to compute the argumant

$$\zeta(u,t) = \eta + \Delta B_1 + \Delta^2 B_2 + \Delta^3 B_3 + \cdots$$

$$= \eta + \gamma_1\beta + \gamma_2\beta^2 + \gamma_3\beta^3 + \cdots \qquad (4\text{-}131)$$

in Eq. (4-123). The execution of these tedious algebraic manipulations has been carried out only for the special case u = 0.

$$\zeta(0,t) = -t + \beta\left[\frac{8}{15}h_2 t^2 - \frac{1}{5t}h_2\right]$$
$$+ \beta^2\left[\left(\frac{16}{35}t^3 - \frac{3}{5}\right)h_3 + \left(-\frac{1328}{1575}t^3 + \frac{406}{1050} + \frac{1}{50}\frac{1}{t^3}\right)h_2^2\right]$$
$$+ \beta^3\left[\left(\frac{128}{315}t^4 - \frac{448}{315}t\right)h_4 + \left(-\frac{33088}{17325}t^4 + \frac{47564}{17325}t + \frac{3}{25}\frac{1}{t^2}\right)h_2 h_3\right.$$
$$\left. + \left(\frac{128512}{70875}t^4 - \frac{10648}{7875}t - \frac{182}{2625}\frac{1}{t^2} - \frac{1}{250}\frac{1}{t^5}\right)h_2^3\right] \qquad (4\text{-}132)$$

Since the use of Eq. (4-123) involves a considerable amount of algebraic manipulation of such complex expressions as Eq. (4-130), it may be desirable in future work to seek a representation for the derivative of G(u,t) which is of the form

$$\frac{dG(u,t)}{du} = A(u,t)\,h'_{1,2}(\zeta) \qquad (4\text{-}133)$$

where ζ is defined by an expression similar to Eq. (4-114) and the amplitude function A(u,t) involves the derivative of ζ with respect to u. Let us indicate how this can be done by considering the derivative of $F(y,\lambda)$ with respect to y. Let us define $K(y,\lambda)$ by means of

$$\frac{dF(y,\lambda)}{dy} = \sqrt{k^2 f(y) - \lambda}\;\; K(y,\lambda) \qquad (4\text{-}134)$$

Since

$$\left\{\frac{d^2}{dy^2} + \frac{k^2 f'(y)}{\lambda - k^2 f(y)}\frac{d}{dy} + \left[-\lambda + k^2 f(y)\right]\right\}\frac{dF(y,\lambda)}{dy} = 0$$

we can show that

$$\frac{d^2K}{dy^2} + \left\{\left[-\lambda + k^2 f(y)\right] - \frac{3}{4}\left[\frac{k^2 f'(y)}{\lambda - k^2 f(y)}\right]^2 - \frac{1}{2}\frac{k^2 f''(y)}{[\lambda - k^2 f(y)]}\right\}K = 0 \qquad (4\text{-}135)$$

For an "approximately identical" differential equation, we take

$$\frac{d^2y}{dx^2} + \left[x - \frac{3}{4}\frac{1}{x^2}\right] y = 0 \qquad (4\text{-}136)$$

which has the solution

$$y(x) = \frac{1}{\sqrt{x}}\, h'(x)$$

where h(x) is a solution of Airy's equation

$$\frac{d^2h}{dx^2} + x\,h = 0$$

We can then define $\xi(y,\lambda)$ by means of

$$K(y,\lambda) = [\partial\xi(y,\lambda)/\partial y]^{-1/2}\, y[\xi(y,\lambda)] \qquad (4\text{-}137)$$

and show that

$$\left(\xi - \frac{3}{4}\frac{1}{\xi^2}\right)(\dot{\xi})^2 - \frac{3}{4}\left(\frac{\ddot{\xi}}{\dot{\xi}}\right)^2 + \frac{1}{2}\frac{\dddot{\xi}}{\dot{\xi}} =$$

$$\left\{\left[-\lambda + k^2 f(y)\right] - \frac{3}{4}\left[\frac{k^2 f'(y)}{k^2 f(y) - \lambda}\right]^2 + \frac{1}{2}\frac{k^2 f''(y)}{k^2 f(y) - \lambda}\right\} \qquad (4\text{-}138)$$

where the dots denote differentiation with respect to y. We can then represent the derivative of $F(y,\lambda)$ in the form

$$\frac{dF(y,\lambda)}{dy} = \sqrt{\frac{k^2 f(y) - \lambda}{\xi}}\left(\frac{1}{k^{2/3} f_1^{1/3}}\frac{d\xi}{dy}\right)^{-1/2} h'_{1,2}(\xi) \qquad (4\text{-}139)$$

If we define t as in Eq. (4-74) and u as in Eq. (4-75), and use the property $F(y,\lambda) = G(u,t)$, we can show that

$$\frac{dG(u,t)}{du} = \sqrt{\frac{u - t + \beta h_2 u^2 + \beta^2 h_3 u^3 + \cdots}{\zeta}}\left(\frac{d\zeta}{du}\right)^{-1/2} h'_{1,2}(\zeta) \qquad (4\text{-}140)$$

where

$$\zeta = \zeta(u,t) = \xi(y,\lambda)$$

From Eq. (4-138), we find that ζ is a solution of non-linear differential equation

$$\left(\zeta - \frac{3}{4}\ \frac{1}{\zeta^2}\right)(\dot{\zeta})^2 - \frac{3}{4}\left(\frac{\ddot{\zeta}}{\dot{\zeta}}\right)^2 + \frac{1}{2}\ \frac{\dddot{\zeta}}{\dot{\zeta}} = \left\{ \left[- t+u + \beta h_2 u^2 + \beta^2 h_3 u^3 + \cdots \right] - \frac{3}{4}\left[\frac{1 + 2\beta h_2 u + 3\beta^2 h_3 u^2 + \cdots}{u - t + \beta h_2 u^2 + \beta^2 h_3 u^3 + \cdots} \right]^2 + \frac{1}{2}\left[\frac{2\beta h_2 + 6\beta^2 h_3 u + 12\beta^3 h_4 u^2 + \cdots}{u - t + \beta h_2 u^2 + \beta^2 h_3 u^3 + \cdots} \right] \right\} \qquad (4\text{-}141)$$

where the dots denote differentiation with respect to u.

We can solve for $\zeta(u,t)$ in much the same manner as we solve for $\eta(u,t)$ by using Eq. (4-114). However, we cannot express $\zeta(u,t)$ in form

$$\zeta(u,\ t) = u - t + \sum_{n=1}^{\infty} p_n\ (u,\ t)\ \beta^n$$

where $p_n(u,t)$ is the sum of homogeneous polynomials in u and t. Our studies of the behavior of of the coefficients of β^n in the expansion of $\zeta(u,t)$ led us to see that if we defined

$$z = u - t \qquad (4\text{-}142)$$

that we could express $\zeta(u,t)$ in the form

$$\zeta(u,\ t) = \zeta\ [z + t, t] = z + \sum_{n=1}^{\infty} q_n\ (z,\ t)\ \beta^n \qquad (4\text{-}143)$$

where $q_n(z,t)$ is the sum of homogeneous polynomials in z and t which are of the form

$$q_1(z,\ t) = a_1\ z^2 + b_1\ zt + c_1\ t^2 + \alpha_1\ z^{-1} + \beta_1\ t^{-1} \qquad (4\text{-}144a)$$

$$q_2(z,\ t) = a_2\ z^3 + b_2\ z^2\ t + c_2\ zt^2 + d_2\ t^3 + \alpha_2 + A_2\ z^{-3} + B_2\ z^{-2}\ t^{-1} + C_2\ z^{-1}\ t^{-2} + D_2\ t^{-3} \qquad (4\text{-}144b)$$

The coefficient $q_n(z,t)$ begins with a homogeneous polynomial of degree (n + 1), followed by homogeneous polynomials of degree (n - 2), (n - 2), (n - 5), (n - 8), ..., (-2n + 1).

If we reconsider Eq. (4-123), we find that $\zeta(u,t)$ is defined in terms of $\eta(u,t)$ by means of Eq. (4-131). Therefore, Eq. (4-132) is an example of the expansion defined by Eq. (4-143).

Let us conclude this discussion by observing that the expansion for γ in Eq. (4-120) can be used in Eq. (4-119) to obtain an expansion of the form of Eq. (4-81). Thus, if we define Ω by means of the relation

$$\gamma = u - \tau - \Omega \tag{4-145}$$

we find from Eq. (4-120) that

$$\Omega = \Omega(u, t) = \sum_{n=1}^{\infty} r_n(u, \tau)\, \beta^n \tag{4-146}$$

where $r_n(u,\tau)$ for $n \leq 3$ can be found in Eq. (4-120). Then we can write

$$w(-\gamma) = w(\tau - u) + \Omega\, w'(\tau-u) + \frac{1}{2}\Omega^2 (\tau - u)\, w(\tau - u) + \frac{1}{6}\Omega^3 [w(\tau - u) + (\tau - u)\, w'(\tau - u)] + \cdots \tag{4-147}$$

We then find that

$$w(-\gamma) = w(\tau - u) + \beta r_1(u, \tau)\, w'(\tau - u) + \beta^2 \left[\frac{1}{2}(\tau - u)\, r_1{}^2(u, \tau)\, w(\tau - u) + r_2(u, \tau)\, w'(\tau - u)\right] + \beta^3 \left\{ \left[(\tau - u)\, r_1(u, \tau)\, r_2(u, \tau) + \frac{1}{6} r_1{}^3(u, \tau)\right] w(\tau - u) + \left[r_3(u, \tau) + \frac{1}{6}(\tau - u)\, r_1{}^3(u, \tau)\right] w'(\tau - u) \right\} + O(\beta^4)$$

$$\left(\frac{d\gamma}{du}\right)^{-\frac{1}{2}} = \left[1 + \sum_{n=1}^{\infty} [dr_n(u,\tau)/du]\, \beta^n\right]^{-1/2} = 1 - \frac{\beta}{2}\frac{dr_1(u, \tau)}{du} - \frac{\beta^2}{2}\left\{\frac{dr_2(u, \tau)}{du} - \frac{3}{4}\left[\frac{dr_1(u, \tau)}{du}\right]^2\right\} - \frac{\beta^3}{2}\left\{\frac{dr_3(u, \tau)}{du} - \frac{3}{2}\frac{dr_1(u, \tau)}{du}\frac{dr_2(u, \tau)}{du} + \frac{5}{8}\left[\frac{dr_1(u, \tau)}{du}\right]^3\right\} + O(\beta^4)$$

We can use these results to show that

$$H(u,\tau) = G\left[u,\, t(\tau)\right] = \left(\frac{d\gamma}{du}\right)^{-1/2} w(-\gamma) = w(\tau - u)\left\{1 - \beta\left[\frac{1}{5} h_2 u + \frac{2}{15} h_2 \tau\right]\right.$$

$$+ \beta^2 \left[-\frac{1}{50} h_2^2 u^5 - \frac{1}{30} h_2^2 \tau u^4 - \frac{4}{45} h_2^2 \tau^2 u^3 + \frac{1}{5} h_2 D_1 u^3\right.$$

$$+ \frac{1}{15} h_2 D_1 \tau u^2 + \left(-\frac{3}{14} h_3 + \frac{9}{70} h_2^2\right) u^2 + \frac{4}{15} h_2 D_1 \tau^2 u$$

$$+ \left(-\frac{6}{35} h_3 + \frac{22}{105} h_2^2\right) \tau u - \frac{1}{2} D_1^2 u + \frac{32}{225} h_2^2 \tau^5 - \frac{8}{15} h_2 D_1 \tau^3$$

$$\left. + \left(-\frac{4}{35} h_3 + \frac{58}{315} h_2^2\right) \tau^2 + \frac{1}{2} D_1^2 \tau - \frac{2}{15} h_2 D_1\right]$$

$$+ \beta^3 \left[\left(-\frac{1}{35} h_2 h_3 + \frac{31}{2625} h_2^3\right) u^6 + \left(-\frac{23}{525} h_2 h_3 + \frac{173}{5250} h_2^3\right) \tau u^5\right.$$

$$+ \left(-\frac{2}{21} h_2 h_3 + \frac{139}{1575} h_2^3\right) \tau^2 u^4 + \left(\frac{1}{7} h_3 - \frac{5}{42} h_2^2\right) D_1 u^4$$

$$+ \left(-\frac{8}{105} h_2 h_3 + \frac{2552}{14175} h_2^3\right) \tau^3 u^3$$

$$+ \left(\frac{1}{35} h_3 - \frac{19}{63} h_2^2\right) D_1 \tau u^3 + \left(-\frac{2}{9} h_4 + \frac{223}{630} h_2 h_3\right.$$

$$\left. - \frac{473}{3150} h_2^3 + \frac{1}{5} h_2 D_2\right) u^3 + \frac{16}{675} h_2^3 \tau^4 u^2$$

$$+ \left(\frac{2}{35} h_3 - \frac{78}{315} h_2^2\right) D_1 \tau^2 u^2 + \left(-\frac{4}{21} h_4 + \frac{572}{1155} h_2 h_3\right.$$

$$\left. + \frac{1}{15} h_2 D_2 - \frac{146}{525} h_2^3\right) \tau u^2 + \frac{4}{15} h_2 D_1^2 u^2$$

$$- \frac{32}{3375} h_2^3 \tau^5 u + \left(\frac{8}{35} h_3 - \frac{16}{35} h_2^2\right) D_1 \tau^3 u$$

$$+ \left(-\frac{16}{105} h_4 + \frac{748}{1155} h_2 h_3 - \frac{742}{1575} h_2^3 + \frac{4}{15} h_2 D_2\right) \tau^2 u$$

$$+ \frac{19}{30} h_2 D_1^2 \tau u + \left(-\frac{3}{5} h_3 + \frac{7}{15} h_2^2 - D_2\right) D_1 u$$

$$+ \left(\frac{128}{525} h_2 h_3 - \frac{35008}{70875} h_2^3\right) \tau^6 + \left(-\frac{16}{35} h_3 + \frac{512}{315} h_2^2\right) D_1 \tau^4$$

$$+ \left(-\frac{32}{315} h_4 + \frac{704}{3465} h_2 h_3 + \frac{3404}{14175} h_2^3 - \frac{8}{15} h_2 D_2\right) \tau^3$$

$$- \frac{7}{5} h_2 D_1^2 \tau^2 + \left(\frac{1}{5} h_3 + \frac{1}{9} h_2^2 + D_2\right) D_1 \tau$$

$$\left. \left. + \left(\frac{2}{9} h_4 - \frac{14}{45} h_2 h_3 + \frac{28}{225} h_2^3 - \frac{2}{15} h_2 D_2 + \frac{1}{6} D_1^3\right)\right] + O(\beta^4)\right\}$$

(Eq. concluded on following page)

$$
\begin{aligned}
+ w'(\tau - u)\Bigg\{ & \beta\left[-\frac{1}{5} h_2 u^2 - \frac{4}{15} h_2 \tau u - \frac{8}{15} h_2 \tau^2 + D_1\right] \\
& + \beta^2\left[-\left(\frac{1}{7} h_3 - \frac{3}{35} h_2^2\right)u^3 - \left(\frac{6}{35} h_3 - \frac{22}{105} h_2^2\right)\tau u^2 - \left(\frac{8}{35} h_3 - \frac{144}{315} h_2^2\right)\tau^2 u\right. \\
& \quad \left. - \frac{7}{15} h_2 D_1 u - \left(\frac{16}{35} h_3 - \frac{288}{315} h_2^2\right)\tau^3 - \frac{6}{5} h_2 D_1 \tau + \left(\frac{3}{7} h_3 - \frac{9}{35} h_2^2 + D_2\right)\right] \\
& + \beta^3\left[\frac{1}{750} h_2^3 u^7 + \frac{1}{250} h_2^3 \tau u^6 + \frac{14}{1125} h_2^3 \tau^2 u^5 - \frac{1}{50} h_2^2 D_1 u^5\right. \\
& \quad + \frac{28}{2025} h_2^3 \tau^3 u^4 - \frac{1}{30} h_2^2 D_1 \tau u^4 + \left(-\frac{1}{9} h_4 + \frac{89}{630} h_2 h_3\right. \\
& \quad \left. - \frac{169}{3150} h_2^3\right) u^4 + \frac{32}{2025} h_2^3 \tau^4 u^3 - \frac{4}{45} h_2^2 D_1 \tau^2 u^3 + \left(-\frac{8}{63} h_4\right. \\
& \quad \left. + \frac{104}{315} h_2 h_3 - \frac{292}{1575} h_2^3\right) \tau u^3 + \frac{1}{10} h_2 D_1^2 u^3 - \frac{32}{3375} h_2^3 \tau^5 u^2 \\
& \quad + \left(-\frac{16}{105} h_4 + \frac{68}{105} h_2 h_3 - \frac{106}{225} h_2^3\right) \tau^2 u^2 + \frac{1}{30} h_2 D_1^2 \tau u^2 \\
& \quad + \left(-\frac{27}{70} h_3 + \frac{71}{210} h_2^2\right) D_1 u^2 - \frac{128}{10125} h_2^3 \tau^6 u + \left(-\frac{64}{315} h_4\right. \\
& \quad \left. + \frac{704}{693} h_2 h_3 - \frac{14008}{14175} h_2^3\right) \tau^3 u + \frac{2}{15} h_2 D_1^2 \tau^2 u + \left(-\frac{22}{35} h_3\right. \\
& \quad \left. + \frac{118}{105} h_2^2\right) D_1 \tau u + \left(\frac{4}{9} h_4 - \frac{223}{315} h_2 h_3 + \frac{473}{1575} h_2^3 - \frac{7}{15} h_2 D_2 - \frac{1}{6} D_1^3\right) u \\
& \quad - \frac{256}{10125} h_2^3 \tau^7 + \frac{32}{225} h_2^2 D_1 \tau^5 + \left(-\frac{128}{315} h_4 + \frac{1408}{693} h_2 h_3\right. \\
& \quad \left. - \frac{28688}{14175} h_2^3\right) \tau^4 - \frac{4}{15} h_2 D_1^2 \tau^3 + \left(-\frac{52}{35} h_3 + \frac{922}{315} h_2^2\right) D_1 \tau^2 \\
& \quad + \left(\frac{80}{63} h_4 - \frac{8338}{3465} h_2 h_3 + \frac{1822}{1575} h_2^3 - \frac{6}{5} h_2 D_2 + \frac{1}{6} D_1^3\right) \tau \\
& \quad \left. + \left(-\frac{2}{3} h_2 D_1^2 + D_3\right)\right] + O(\beta^4)\Bigg\}
\end{aligned}
\tag{4-148}
$$

If we differentiate with respect to u, we find that

$$\frac{\partial H(u,\tau)}{\partial u} = w'(\tau - u)\Big\{ -1 - \beta\left[\tfrac{1}{5}h_2 + \tfrac{2}{15}h_2\tau\right] + \beta^2\left[\tfrac{1}{50}h_2^2u^5 + \tfrac{1}{30}h_2^2\tau u^4 + \tfrac{4}{45}h_2^2\tau^2u^3\right.$$
$$-\tfrac{1}{5}h_2D_1u^3 - \tfrac{1}{15}h_2D_1\tau u^2 + (-\tfrac{3}{14}h_3 + \tfrac{9}{70}h_2^2)u^2 - \tfrac{4}{15}h_2D_1\tau^2u$$
$$+ (-\tfrac{6}{35}h_3 + \tfrac{22}{105}h_2^2)\tau u + \tfrac{1}{2}D_1^2u - \tfrac{32}{225}h_2^2\tau^5 + \tfrac{8}{15}h_2D_1\tau^3$$
$$\left. + (-\tfrac{4}{35}h_3 + \tfrac{86}{315}h_2^2)\tau^2 - \tfrac{1}{2}D_1^2\tau - \tfrac{1}{3}h_2D_1\right]$$
$$+ \beta^3\left[(\tfrac{1}{35}h_2h_3 - \tfrac{13}{5250}h_2^3)u^6 + (\tfrac{23}{525}h_2h_3 - \tfrac{47}{5250}h_2^3)\tau u^5\right.$$
$$+ (\tfrac{2}{21}h_2h_3 - \tfrac{41}{1575}h_2^3)\tau^2u^4$$
$$+ (-\tfrac{1}{7}h_3 + \tfrac{2}{105}h_2^2)D_1u^4 + (\tfrac{8}{105}h_2h_3 - \tfrac{1768}{14175}h_2^3)\tau^3u^3$$
$$+ (-\tfrac{1}{35}h_3 + \tfrac{53}{315}h_2^2)D_1\tau u^3 + (-\tfrac{2}{9}h_4 + \tfrac{133}{630}h_2h_3 - \tfrac{203}{3150}h_2^3$$
$$- \tfrac{1}{5}h_2D_2)u^3 + \tfrac{16}{675}h_2^3\tau^4u^2 + (-\tfrac{2}{35}h_3 - \tfrac{2}{105}h_2^2)D_1\tau^2u^2$$
$$+ (-\tfrac{4}{21}h_4 + \tfrac{572}{1155}h_2h_3 - \tfrac{146}{525}h_2^3 - \tfrac{1}{15}h_2D_2)\tau u^2 + \tfrac{1}{30}h_2D_1^2u^2$$
$$- \tfrac{32}{3375}h_2^3\tau^5u + (-\tfrac{8}{35}h_3 + \tfrac{16}{35}h_2^2)D_1\tau^3u$$
$$+ (-\tfrac{16}{105}h_4 + \tfrac{748}{1155}h_2h_3 - \tfrac{742}{1575}h_2^3 - \tfrac{4}{15}h_2D_2)\tau^2u - \tfrac{17}{30}h_2D_1^2\tau u$$
$$+ (-\tfrac{6}{35}h_3 + \tfrac{22}{105}h_2^2 + D_2)D_1u + (-\tfrac{128}{525}h_2h_3 + \tfrac{34112}{70875}h_2^3)\tau^6$$
$$+ (\tfrac{16}{35}h_3 - \tfrac{512}{315}h_2^2)D_1\tau^4 + (-\tfrac{32}{315}h_4 + \tfrac{2816}{3465}h_2h_3$$
$$- \tfrac{10604}{14175}h_2^3 + \tfrac{8}{15}h_2D_2)\tau^3 + \tfrac{23}{15}h_2D_1^2\tau^2 + (-\tfrac{29}{35}h_3$$
$$+ \tfrac{319}{315}h_2^2 - D_2)D_1\tau + (\tfrac{2}{9}h_4 - \tfrac{25}{63}h_2h_3 + \tfrac{277}{1575}h_2^3$$
$$\left.\left. - \tfrac{1}{3}h_2D_2 - \tfrac{1}{3}D_1^3)\right] + O(\beta^4)\right\}$$

(Eq. concluded on following page)

$$
\begin{aligned}
&+ w(\tau - u)\Big\{\beta\Big[-\tfrac{1}{5}h_2u^3 - \tfrac{1}{15}h_2\tau u^2 - \tfrac{4}{15}h_2\tau^2u + D_1u + \tfrac{8}{15}h_2\tau^3 - \tfrac{1}{5}h_2 - D_1\tau\Big] \\
&+ \beta^2\Big[u^4(-\tfrac{1}{7}h_3 - \tfrac{1}{70}h_2^2) + \tau u^3(-\tfrac{1}{35}h_3 - \tfrac{1}{105}h_2^2) \\
&\quad + \tau^2u^2(-\tfrac{2}{35}h_3 - \tfrac{2}{105}h_2^2) + u^2(\tfrac{2}{15}h_2D_1) + \tau^3u(-\tfrac{8}{35}h_3 + \tfrac{144}{315}h_2^2) \\
&\quad + \tau u(-\tfrac{3}{5}h_2D_1) + D_2u + \tau^4(\tfrac{16}{35}h_3 - \tfrac{288}{315}h_2^2) + \tau^2(\tfrac{22}{15}h_2D_1) \\
&\quad + \tau(-\tfrac{3}{5}h_3 + \tfrac{7}{15}h_2^2 - D_2) - \tfrac{1}{2}D_1^2\Big] \\
&+ \beta^3\Big[\tfrac{1}{750}h_2^3u^8 + \tfrac{1}{375}h_2^3\tau u^7 + \tfrac{19}{2250}h_2^3\tau^2u^6 - \tfrac{1}{50}h_2^2D_1u^6 + \tfrac{14}{10125}h_2^3\tau^3u^5 \\
&\quad - \tfrac{1}{75}h_2^2D_1\tau u^5 + (-\tfrac{1}{9}h_4 - \tfrac{19}{630}h_2h_3 + \tfrac{271}{15750}h_2^3)u^5 \\
&\quad + \tfrac{4}{2025}h_2^3\tau^4u^4 - \tfrac{1}{18}h_2^2D_1\tau^2u^4 + (-\tfrac{1}{63}h_4 - \tfrac{19}{630}h_2h_3 + \tfrac{52}{1575}h_2^3)\tau u^4 \\
&\quad + \tfrac{1}{10}h_2D_1^2u^4 - \tfrac{256}{10125}h_2^3\tau^5u^3 + \tfrac{4}{45}h_2^2D_1\tau^3u^3 + (-\tfrac{8}{315}h_4 - \tfrac{4}{63}h_2h_3 \\
&\quad + \tfrac{106}{1575}h_2^3)\tau^2u^3 - \tfrac{1}{15}h_2D_1^2\tau u^3 + (\tfrac{13}{70}h_3 - \tfrac{29}{210}h_2^2)D_1u^3 \\
&\quad - \tfrac{32}{10125}h_2^3\tau^6u^2 + (-\tfrac{16}{315}h_4 + \tfrac{484}{3465}h_2h_3 + \tfrac{326}{14175}h_2^3)\tau^3u^2 \\
&\quad + \tfrac{1}{10}h_2D_1^2\tau^2u^2 + (-\tfrac{11}{70}h_3 - \tfrac{5}{42}h_2^2)D_1\tau u^2 + (-\tfrac{2}{9}h_4 + \tfrac{223}{630}h_2h_3 - \tfrac{1}{6}D_1^3 \\
&\quad - \tfrac{473}{3150}h_2^3 + \tfrac{2}{15}h_2D_2)u^2 - \tfrac{128}{10125}h_2^3\tau^7u + \tfrac{32}{225}h_2^2D_1\tau^5u \\
&\quad + (-\tfrac{64}{315}h_4 + \tfrac{704}{693}h_2h_3 - \tfrac{14008}{14175}h_2^3)\tau^4u - \tfrac{2}{5}h_2D_1^2\tau^3u \\
&\quad + (-\tfrac{26}{35}h_3 + \tfrac{412}{315}h_2^2)D_1\tau^2u + (\tfrac{4}{9}h_4 - \tfrac{2453}{3465}h_2h_3 + \tfrac{473}{1575}h_2^3 - \tfrac{3}{5}h_2D_2 \\
&\quad + \tfrac{1}{3}D_1^3)\tau u + (-\tfrac{2}{15}h_2D_1^2 + D_3)u + \tfrac{256}{10125}h_2^3\tau^8 - \tfrac{32}{225}h_2^2D_1\tau^6 \\
&\quad + (+\tfrac{128}{315}h_4 - \tfrac{1408}{693}h_2h_3 + \tfrac{142768}{70875}h_2^3)\tau^5 + \tfrac{4}{15}h_2D_1^2\tau^4 \\
&\quad + (\tfrac{12}{7}h_3 - \tfrac{1066}{315}h_2^2)D_1\tau^3 + (-\tfrac{64}{45}h_4 + \tfrac{962}{315}h_2h_3 - \tfrac{2564}{1575}h_2^3 \\
&\quad + \tfrac{22}{15}h_2D_2 - \tfrac{1}{6}D_1^3)\tau^2 + (\tfrac{13}{10}h_2D_1^2 - D_3)\tau + (-\tfrac{3}{5}h_3 + \tfrac{7}{15}h_2^2 \\
&\quad - D_2)D_1\Big] + O(\beta^4)\Big\}
\end{aligned}
\tag{4-149}
$$

A useful check on the coefficients in Eqs. (4-148) and (4-149) can be obtained from the Wronskian relation

$$W(H_1, H_2) = H_1 \frac{dH_2}{du} - H_2 \frac{dH_1}{du} = -2i \tag{4-150}$$

This property is readily derived from the Wronskian for the Airy functions

$$W(w_1, w_2) = w_1(z) \frac{dw_2(z)}{dz} - w_2(z) \frac{dw_1(z)}{dz} = 2i \tag{4-151}$$

by using the representations

$$H_{1,2}(u, \tau) = \left(\frac{d\eta}{du}\right)^{-1/2} w_{1,2}(-\eta) \tag{4-152a}$$

$$\frac{dH_{1,2}(u, \tau)}{du} = -\left(\frac{d\eta}{du}\right)^{-1/2} w'_{1,2}(-\eta) - \frac{1}{2} \frac{\frac{d^2\eta}{du^2}}{\left[\frac{d\eta}{du}\right]^2} w_{1,2}(-\eta) \tag{4-152b}$$

Let us express Eq. (4-148) in the form

$$H_{1,2} = w_{1,2}(\tau - u) + \sum_{n=1}^{\infty} \left[A_n(u, \tau)\, w_{1,2}(\tau - u) + B_n(u, \tau)\, w'_{1,2}(\tau - u)\right] \beta^n \tag{4-153a}$$

and use Eq. (4-149) in the form

$$\frac{dH_{1,2}(u, \tau)}{du} = -w'_{1,2}(\tau - u) + \sum_{n=1}^{\infty} \left[C_n(u, \tau)\, w_{1,2}(\tau - u) + D_n(u, \tau)\, w'_{1,2}(\tau - u)\right] \beta^n \tag{4-153b}$$

We can show that the coefficients A_n, B_n, C_n, and D_n have the property

$$\left(\sum_{n=1}^{\infty} A_n(u, \tau)\, \beta^n\right)\left(\sum_{n=1}^{\infty} D_n(u, \tau)\, \beta^n\right) = \left(\sum_{n=1}^{\infty} B_n(u, \tau)\, \beta^n\right)\left(\sum_{n=1}^{\infty} C_n(u, \tau)\, \beta^n\right) \tag{4-154}$$

or $A_1 D_1 = B_1 C_1$ and

$$\sum_{j=1}^{n+1} A_j D_{n-j} = \sum_{j=1}^{n+1} B_j C_{n-j}, \quad n \geq 2 \tag{4-155}$$

4.8 Application to Bessel Functions

An interest check on the accuracy of the expansion given in Eq. (4-118) can be obtained by making use of certain results for the Hankel functions which were obtained by Schobe (Ref. 14). The result which we need is

$$H_\nu^{(1)}(kx) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{kx}\right)^{1/3}\left\{\sum_{m=0}^{\infty}(-1)^m p_m(\xi)\left(\frac{2}{kx}\right)^{\frac{2m}{3}}\right\} w_1[T(\xi,kx)] \qquad (4\text{-}156)$$

where the argument of the Airy function is

$$T(\xi,kx) = \sum_{n=0}^{\infty}(-1)^n q_n(\xi)\left(\frac{2}{kx}\right)^{\frac{2n}{3}}$$

where

$$\xi = \left(\frac{2}{kx}\right)^{2/3}(\nu - kx)$$

and

$$p_0(t) = 1 \qquad q_0(t) = t$$

$$p_1(t) = \frac{1}{75}t \qquad q_1(t) = \frac{1}{60}t^2$$

$$p_2(t) = \frac{13}{1260}t^2 \qquad q_2(t) = \frac{2}{1575}t^3 + \frac{1}{140}$$

$$p_3(t) = \frac{109}{56780}t^3 + \frac{1}{900} \qquad q_3(t) = \frac{41}{283500}t^4 + \frac{4}{1575}t$$

$$p_4(t) = \frac{203743}{523908000}t^4 + \frac{6761}{7276500}t \qquad q_4(t) = \frac{6553}{327442500}t^5 + \frac{1409}{1819125}t^2$$

In order to use Schobe's expansion, we must first observe that the Hankel function $H_\nu^{(1)}(kx)$ is a solution of

$$\frac{d^2H_\nu^{(1)}(kx)}{dx^2} + \frac{1}{x}\frac{dH_\nu^{(1)}(kx)}{dx} + \left[k^2 - \frac{\nu^2}{x^2}\right]H_\nu^{(1)}(kx) = 0 \qquad (4\text{-}157)$$

Therefore, we can show that

$$H_\nu^{(1)}(kx) = F_1[a \log(x/a), (\nu/a)^2] \qquad (4\text{-}158)$$

where $F_1(y,\lambda)$ is a solution of

$$\frac{d^2F_1(y,\lambda)}{dy^2} - \left[-\lambda + k^2\exp\left(2\frac{y}{a}\right)\right]F_1(y,\lambda) = 0 \qquad (4\text{-}159)$$

We start our comparison Schobe's expansion and our Eq. (4-118) by observing that $f(y) = \exp(2y/a)$ and therefore the result that corresponds to Eq. (4-72) is

$$f(y) = f_0 + f_1[y + h_2 y^2 + h_3 y^3 + \cdots]$$

$$= 1 + \frac{2}{a}\left[y + \frac{1}{a}\, y^2 + \frac{2}{3a^2}\, y^3 + \frac{1}{3a^3}\, y^4 + \cdots\right] \qquad (4\text{-}160)$$

Therefore, for this case we find that

$$f_0 = 1 \quad , \quad f_1 = \frac{2}{a} \qquad (4\text{-}161a)$$

and

$$h_2 = \frac{1}{a} \quad , \quad h_3 = \frac{2}{3a^2} \quad , \quad h_4 = \frac{1}{3a^3} \quad , \quad \text{etc.} \qquad (4\text{-}161b)$$

From Eq. (4-74) we find that t is defined by

$$\lambda = \frac{\nu^2}{a^2} = k^2 + k^{4/3} \left(\frac{2}{a}\right)^{2/3} t \qquad (4\text{-}162a)$$

or

$$t = \frac{\nu^2 - (ka)^2}{4\,(ka/2)^{4/3}} \qquad (4\text{-}162b)$$

From Eq. (4-75) we find that

$$u = k^{\frac{2}{3}} f_1^{\frac{1}{3}} y = k^{\frac{2}{3}} (2/a)^{\frac{1}{3}} a \log(x/a) = 2\,(ka/2)^{\frac{2}{3}} \log(x/a) \qquad (4\text{-}163)$$

From Eq. (4-76) we find that

$$\beta = \frac{1}{k^{2/3} f_1^{1/3}} = \frac{1}{k^{2/3}} \left(\frac{a}{2}\right)^{1/3} = \frac{a}{2} \left(\frac{2}{ka}\right)^{2/3} \qquad (4\text{-}164)$$

Therefore, we can express the Hankel function in the form

$$H_\nu^{(1)}(kx) \sim (\text{constant})\, F_1\left(a \log \frac{x}{a} \, , \, \frac{\nu^2}{a^2}\right) = (\text{constant})\, G_1(u, t) \qquad (4\text{-}165)$$

where $G_1(u,t)$ is defined by

$$G_1(u, t) = F_1(y, \lambda) = F_1\left(a \log \frac{x}{a} \, , \, \frac{\nu^2}{a^2}\right) \qquad (4\text{-}166)$$

where u is defined by Eq. (4-163), t is defined by Eq. (4-162b), and β is defined by Eq. (4-164).

From Eq. (4-113), we find that $G_1(u,t) = \sqrt{du/d\eta}\; w_1(-\eta)$ where η is defined by Eq. (4-118). Therefore, we can express Eq. (4-165) in the form

$$H_\nu^{(1)}(kx) = -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}\left(\frac{d\eta}{du}\right)^{-1/2} w_1(-\eta) \tag{4-167}$$

where the choice of the constant of the form

$$(\text{constant}) = -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}$$

follows from the result

$$H_\nu^{(1)}(kx) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3} w_1(t-u) \tag{4-168}$$

which is valid when $kx \gg 1$ and $\nu \approx kx \approx ka$.

If we compare Eq. (4-167) with Eq. (4-156), we see that Schobe's expansion leads to

$$\eta = -\sum_{n=0}^{\infty} (-1)^n q_n(\xi)\left(\frac{2}{kx}\right)^{\frac{2n}{3}} \tag{4-169}$$

In order to compare Eq. (4-169) with Eq. (4-118), we need to express the right side of Eq. (4-169) in terms of u, t, and β. As a preliminary to this step, we need to express ξ and $(2/kx)$ in terms of u, t, and β. Since

$$x = a\exp(y/a) = a\exp[\beta u/a]$$

we find that

$$\left(\frac{2}{kx}\right)^{\lambda} = \left(\frac{2}{ka}\right)^{\lambda}\exp\left(-\frac{\lambda\beta u}{a}\right) = \left(\frac{2\beta}{a}\right)^{\frac{3\lambda}{2}}\left\{1-\beta\frac{\lambda u}{a}+\frac{1}{2}\beta^2\frac{\lambda^2u^2}{a^2}+\cdots\right\} \tag{4-170}$$

The relationship of ξ to u, t, and β is somewhat more complex. From Eq. (4-156) we find that

$$\nu^2 = (kx)^2 + 4\left(\frac{kx}{2}\right)^{4/3}\xi + \left(\frac{kx}{2}\right)^{2/3}\xi^2 \tag{4-171}$$

whereas from Eq. (4-162a), we have

$$\nu^2 = (ka)^2 + 4\left(\frac{ka}{2}\right)^{4/3} t \tag{4-172}$$

If we use Eq. (4-170), we find that

$$(ka)^2 \exp\left(\frac{2\beta u}{a}\right)\left[1+\frac{1}{2}\left(\frac{2}{ka}\right)^{2/3}\xi \exp\left(-\frac{2\beta u}{3a}\right)\right]^2 = (ka)^2\left[1+\left(\frac{2}{ka}\right)^{2/3} t\right]$$

or

$$\exp\left(\frac{\beta u}{a}\right)\left[1+\frac{1}{2}\left(\frac{2}{ka}\right)^{2/3}\xi \exp\left(-\frac{2\beta u}{3a}\right)\right] = \sqrt{1+\left(\frac{2}{ka}\right)^{2/3} t}$$

Therefore

$$\xi = \xi(u, t) = \frac{a}{\beta} \exp\left(-\frac{\beta u}{3a}\right)\left[\sqrt{1+\frac{2\beta}{a}t} - \exp\left(\frac{\beta u}{a}\right)\right]$$

$$= (t-u) + \sum_{n=1}^{\infty} E_n(u, t)\left(\frac{\beta}{a}\right)^n \qquad (4\text{-}173)$$

where the coefficients $E_n(u,t)$ are defined by

$$E_1(u, t) = -\left[\frac{1}{2}t^2 + \frac{1}{3}tu + \frac{1}{6}u^2\right]$$

$$E_2(u, t) = -\left[\frac{1}{2}t^3 + \frac{1}{6}t^2u + \frac{1}{18}tu^2 - \frac{1}{18}u^3\right]$$

$$E_3(u, t) = \left[-\frac{5}{8}t^4 - \frac{1}{6}t^3u - \frac{1}{36}t^2u^2 - \frac{1}{162}tu^3 - \frac{5}{648}u^4\right]$$

$$E_4(u, t) = \left[\frac{7}{8}t^5 + \frac{5}{24}t^4u + \frac{1}{36}t^3u^2 + \frac{1}{324}t^2u^3 + \frac{1}{1944}tu^4 - \frac{11}{9720}u^5\right]$$

By the use of Eq. (4-170) and Eq. (4-173) in Eq. (4-168), we have verified the correctness of all the coefficients in Eq. (4-118)

The Correctness of the coefficients in the expansion of $\zeta(0,t)$ given in Eq. (4-132) can be verified in a similar manner if we use an expansion for the derivative which is in the form of Eq. (4-140), namely,

$$H_\nu^{(1)'}(kx) \sim \frac{1}{\sqrt{\pi}}\left(\frac{2}{kx}\right)^{2/3}\left\{1+\sum_{n=1}^{\infty} B_n(\xi)\left(\frac{2}{kx}\right)^{\frac{2n}{3}}\right\} w_1'(-\xi) \qquad (4\text{-}174a)$$

$$-\zeta = \xi + \sum_{n=1}^{\infty} C_n(\xi)\left(\frac{2}{kx}\right)^{\frac{2n}{3}} \qquad (4\text{-}174b)$$

The coefficient $C_1(\xi)$ was obtained by Scott (Ref. 15). A search

of the literature failed to turn up further terms. Therefore, we have extended Schobe's paper by showing that

$$B_1(t) = \frac{1}{15}\, t \qquad , \qquad C_1(t) = \frac{6 - t^3}{60\, t}$$

$$B_2(t) = -\frac{37}{6300}\, t^2 - \frac{1}{200}\, \frac{1}{t} \qquad , \qquad C_2(t) = \frac{2}{1575}\, t^3 - \frac{39}{1800} - \frac{1}{200}\, \frac{1}{t^3}$$

$$B_3(t) = \frac{239}{283500}\, t^3 + \frac{943}{252000} + \frac{1}{6000}\, \frac{1}{t^3} \ ,$$

$$C_3(t) = -\frac{41}{283500}\, t^4 + \frac{1447}{252000}\, t + \frac{29}{12000}\, \frac{1}{t^2} + \frac{1}{2000}\, \frac{1}{t^5}$$

$$B_4(t) = -\frac{53863}{374220000}\, t^4 - \frac{43993}{23760000}\, t - \frac{79}{720000}\, \frac{1}{t^2} - \frac{1}{8000}\, \frac{1}{t^5}$$

$$C_4(t) = \frac{6553}{327442500}\, t^5 - \frac{260737}{99792000}\, t^2 - \frac{1679}{1108800}\, \frac{1}{t} - \frac{7}{20000}\, \frac{1}{t^4} - \frac{5}{80000}\, \frac{1}{t^7}$$

Since we have not carried out the algebraic details for $\zeta(u,t)$ defined in Eq. (4-131) except for the case $u = 0$, we are limited to considering only the expansion for $\zeta(0,t)$ which was given in Eq. (4-132). For the case of the derivative of the Hankel function as defined by Eq. (4-174), we would need the results

$$\left(\frac{2}{kx}\right) = \left(\frac{2\beta}{a}\right)^{\frac{3\lambda}{2}}$$

$$\xi = \frac{a}{\beta}\sqrt{1 + \frac{2\beta}{a}t} - 1$$

$$= t - \frac{1}{2}t^2\left(\frac{\beta}{a}\right) + \frac{1}{2}t^3\left(\frac{\beta}{a}\right)^2 - \frac{5}{8}t^4\left(\frac{\beta}{a}\right)^3 + \frac{7}{8}t^5\left(\frac{\beta}{a}\right)^4 + \ldots$$

along with the values of h_n which were defined in Eq. (4-161b).

Additional confidence in the correctness of the coefficients in Eq. (4-148) and Eq. (4-149) has been acquired by using Eq. (4-170) and Eq. (4-173) in Schobe's expansions for $H_\nu^{(1)}(kx)$ and $H_\nu^{(1)\prime}(kx)$ which are of the form

$$H_\nu^{(1)}(kx) \approx -\frac{i}{\sqrt{\pi}}\left(\frac{2}{kx}\right)^{1/3} w_1(\xi)\left\{1 - \left(\frac{1}{15}\xi\right)\left(\frac{2}{kx}\right)^{2/3} + \left(\frac{1}{7200}\xi^5 + \frac{13}{1260}\xi^2\right)\left(\frac{2}{kx}\right)^{4/3}\right.$$
$$- \left(\frac{283}{9072000}\xi^6 + \frac{463}{226800}\xi^3 + \frac{1}{900}\right)\left(\frac{2}{kx}\right)^{6/3}$$
$$+ \left(\frac{1}{311040000}\xi^{10} + \frac{599}{95256000}\xi^7 + \frac{26897}{59875200}\xi^4\right.$$
$$\left.\left. + \frac{7939}{8316000}\xi\right)\left(\frac{2}{kx}\right)^{8/3} + \cdots\right\}$$
$$-\frac{i}{\sqrt{\pi}}\left(\frac{2}{kx}\right)^{1/3} w_1'(\xi)\left\{0 - \left(\frac{1}{60}\xi^2\right)\left(\frac{2}{kx}\right)^{2/3} + \left(\frac{1}{420}\xi^3 + \frac{1}{140}\right)\left(\frac{2}{kx}\right)^{4/3}\right.$$
$$- \left(\frac{1}{1296000}\xi^7 + \frac{13}{32400}\xi^4 + \frac{19}{6300}\xi\right)\left(\frac{2}{kx}\right)^{6/3}$$
$$+ \left(\frac{17}{72576000}\xi^8 + \frac{2521}{33264000}\xi^5 + \frac{359}{346500}\xi^2\right)\left(\frac{2}{kx}\right)^{8/3}$$
$$\left. + \cdots\right\} \qquad (4\text{-}175)$$

and

$$H_\nu^{(1)'}(kx) \approx \frac{i}{\sqrt{\pi}}\left(\frac{2}{kx}\right)^{2/3} w_1'(\xi)\left\{1 + \left(\frac{2}{kx}\right)^{2/3}\frac{1}{15}\xi + \left(\frac{2}{kx}\right)^{4/3}\left(\frac{1}{7200}\xi^5 - \frac{19}{2520}\xi^2\right)\right.$$
$$- \left(\frac{2}{kx}\right)^{6/3}\left(\frac{61}{4536000}\xi^6 - \frac{283}{226800}\xi^3 - \frac{23}{12600}\right)$$
$$+ \left(\frac{2}{kx}\right)^{8/3}\left(\frac{\xi^{10}}{311040000} + \frac{443\xi^7}{381024000} - \frac{1483\xi^4}{5987520}\right.$$
$$\left.\left. - \frac{9809\xi}{8316000}\right) + \cdots\right\}$$
$$+\frac{i}{\sqrt{\pi}}\left(\frac{2}{kx}\right)^{2/3} w_1(\xi)\left\{0 - \left(\frac{2}{kx}\right)^{2/3}\left(\frac{\xi^3}{60} - \frac{1}{10}\right) + \left(\frac{2}{kx}\right)^{4/3}\left(\frac{1}{3360}\xi^4 - \frac{1}{60}\xi\right)\right.$$
$$- \left(\frac{2}{kx}\right)^{6/3}\left(\frac{1}{1296000}\xi^8 - \frac{181}{4536000}\xi^5 - \frac{103}{25200}\xi^2\right)$$
$$+ \left(\frac{2}{kx}\right)^{8/3}\left(\frac{\xi^9}{725760} - \frac{13\xi^6}{887040} - \frac{3821\xi^3}{178200} - \frac{947}{2772000}\right)$$
$$\left. + \cdots\right\} \qquad (4\text{-}176)$$

In order to emphasize the generality of the new expansions

described by Eq. (4-148) and Eq. (4-149), let us observe that the terms up to and including $(kx/2)^{-2}$ in Schobe's results described by Eq. (4-175) and Eq. (4-176) are obtained from the new results by using the following definitions

$$D_n = 0 \quad \text{for all } n$$

$$u = 0$$

$$\beta = \frac{x}{2}\left(\frac{2}{kx}\right)^{2/3}$$

$$t = \xi + \frac{1}{4}\left(\frac{2}{kx}\right)^{2/3}\xi^2$$

and

$$h_2 = \frac{1}{x} \quad , \quad h_3 = \frac{2}{3x^2} \quad , \quad h_4 = \frac{1}{3x^3} \quad , \text{ etc.}$$

4.9 Numerical Aspects of Employing These Expansions

If a numerical anlyst were faced with the problem of evaluating the function $F(y,\lambda)$ which satisfies Eq. (4-1), he would probably find in Ref. 5 (and in the references which have been cited by Olver) a number of methods which he would prefer when they were compared with the expansions which we have expounded upon in the "forty-plus" pages of this Section. The author will readily agree that he himself tends to favor the basing of future work on more computer work and less algebraic work. However, the expansions in this Section are the outgrowths of earlier studies by Fock, Pekeris, and Friedlander. The expansions are also very similar to those in the recent paper by Bremmer (Ref. 2). The author feels that the type of expansions which have been the subject of this section have not outlived their usefulness. Even as the more erudite methods become developed into computer programs which are "de-bugged" and become operational, these expansions will continue to provide some valuable checks and guidelines.

REFERENCES FOR SECTION 4

1. Institute of Mathematical Sciences, New York University, Propagation of a Pulse in an Inhomogeneous Medium, by F. G. Friedlander, Report EM - 76, Contract AF 19(122)-42, New York, 1955

2. F. G. Friedlander, Sound Pulses, Cambridge, Cambridge University Press, 1958. (Parts of Chapter 6, "Some Other Diffraction Problems," have been obtained by making a condensation of Ref. 1.)

3. H. Bremmer, " On the Theory of Wave Propagation Through a Concentrically Stratified Troposphere with a Smooth Profile. Part I. Discussion of the Extended W.K.B. Approximation," J. Research NBS, Vol. 64D, Sept. - Oct., 1960, pp. 467-482

4. H. Bremmer, "On the Theory of Wave Propagation Through a Concentrically Stratified Troposphere with a Smooth Profile. Part II. Expansion of the Rigorous Solution," J. Research NBS, Vol. 66D, Jan.-Feb., 1962, pp. 31-52

5. F. W. J. Olver, "Definitions, Formulae, and Methods," Bessel Functions, Part III, Zeros and Associated Values, Royal Society Mathematical Tables, Vol. 7, Cambridge, Cambridge University Press, 1960

6. A. Erdélyi, "Asymptotic Solutions of Differential Equations with Transition Points or Singularities," J. of Math. Physics, Vol. 1, Jan.-Feb., 1960, pp. 16-26

7. C. L. Pekeris, "Asymptotic Solutions for the Normal Modes in the Theory of Microwave Propagation," J. Appl. Physics, Vol. 17, 1946, pp. 1108-1124

8. V. A. Fock, "Propagation of the Direct Wave Around the Earth With Due Account for Diffraction and Propagation," Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965. (This classic paper was first published in Bulletin de l'Académie des Sciences de l'URSS, ser. phys., Vol. 12, No. 2, 1948, pp. 81-97.)

9. I. Imai, "Asymptotic Solutions of Ordinary Linear Differential Equations of the Second Order," Phys. Rev., Vol. 80, 1950, p. 1112

10. I. Imai, "A Refinement of the WKB Method and Its Application to the Electromagnetic Wave Theory," IRE Trans. on Antennas and Propagation, Vol. AP-4, July, 1956, pp. 233-239

11. Harvard University Computation Laboratory, Tables of the Modified Hankel Functions of Order One-Third and of Their Derivatives, Annals of the Computation Laboratory of Harvard University, Vol. II, Cambridge, Mass., Harvard University Press, 1945

12. T. M. Cherry, "Uniform Asymptotic Formulas for Functions with Transition Points," Trans. Am. Math. Soc., Vol. 68, 1950, pp. 224-257

13. R. E. Langer, "On Asymptotic Solutions of Ordinary Differential Equations with Application to Bessel Functions of Large Order," Trans. Am. Math. Soc., Vol. 33, 1931, pp. 23-64

14. W. Schobe, "Eine an die Nicholsonformel Anschliessende Asymptotische Entwicklung für Zylinderfunktionen," Acta Mathematica, Vol. 92, 1954, pp. 265-307

15. Atomic Energy Research Establishment, An Asymptotic Series for the Radar Scattering Cross Section of a Spherical Target," Report T/M 30, Harwell, England, Sept., 1949

Section 5

ASYMPTOTIC EXPANSIONS FOR THE PROPAGATION CONSTANTS AND THE NORMALIZATION INTEGRAL

5.1 The Eigenvalue Problem

The eigenvalues λ_s or t_s are defined in the radiowave propagation problems by the homogeneous boundary conditions (sometimes referred to as "impedance boundary conditions") which has been given above as Eq. (4-54), namely

$$\left[\frac{dG(u, t_s)}{du} + q\, G(u, t_s)\right]_{u=0} = 0 \tag{5-1a}$$

We can also write

$$\left[\frac{\partial H(u, \tau_s)}{\partial u} + q\, H(u, \tau_s)\right]_{u=0} = 0 \tag{5-1b}$$

and

$$\left[\frac{\partial F(y, \lambda_s)}{\partial y} + k^{2/3} f_1^{1/3} q\, F(y, \lambda_s)\right]_{y=0} = 0 \tag{5-1c}$$

We have found the relations between λ_s, t_s, and τ_s to be

$$\lambda_s = k^2 f_o + k^{4/3} f_1^{2/3} t_s = k^2 f_o + k^{4/3} f_1^{2/3}\left[\tau_s + \sum_{n=1}^{\infty} D_n(\tau_s, q)\, \beta^n\right] \tag{5-2}$$

where

$$\beta = k^{-\frac{2}{3}} f_1^{-\frac{1}{3}}$$

Eqs. (4-148) and (4-149) are ideally suited for the determination of the coefficients $D_n(\tau_s, q)$ for n = 1, 2, and 3.

The case $q = \infty$ is quite easy. Since

$$H(0, \tau_s^{\infty}) = 0 \tag{5-3a}$$

we find that τ_s^∞ is defined by

$$w(\tau_s^\infty) = 0 \tag{5-3b}$$

From Eq. (4-148), we find that in order to satisfy Eq. (5-3a) we must have

$$D_1 = \frac{8}{15} h_2 \tau^2 \tag{5-4a}$$

$$D_2 = (\frac{16}{35} h_3 - \frac{48}{175} h_2^2)\, \tau^3 - (\frac{3}{7} h_3 - \frac{9}{35} h_2^2) \tag{5-4b}$$

$$D_3 = (\frac{128}{315} h_4 - \frac{11968}{17325} h_2 h_3 + \frac{22912}{70875} h_2^3)\, \tau^4 - (\frac{80}{63} h_4 - \frac{6556}{3465} h_2 h_3 + \frac{1336}{1575} h_2^3)\, \tau \tag{5-4c}$$

$$D_4 = (\frac{256}{693} h_5 - \frac{41984}{51975} h_2 h_4 - \frac{17344}{40425} h_3^2 + \frac{7521792}{5457375} h_2^2 h_3 - \frac{41575168}{81860625} h_2^4)\, \tau^5 + (-\frac{1808}{693} h_5 + \frac{50432}{10395} h_2 h_4 + \frac{19324}{8085} h_3^2 - \frac{8047680}{1091475} h_2^2 h_3 + \frac{14141280}{5457375} h_2^4)\, \tau^2 \tag{5-4d}$$

where all the τ are to be assumed to be τ_s^∞. The roots of

$$w_{1,2}(\tau_s^\infty) = 0 \tag{5-5a}$$

are well known (See Section 2). They can be expressed in the form

$$\tau_s^\infty = a_s \exp(\pm \tfrac{1}{3}\pi) \tag{5-5b}$$

where a_s denotes the roots of the Airy function

$$Ai(-a_s) = 0$$

In Table 2-1 we have listed some 15 decimal values which were computed for the author by G. F. Miller and P. H. Haines at the National Physical Laboratory (Teddington, England) in 1958. Sherry (Ref. 1) has obtained 25 decimal values for these constants.

For the case q = 0, we must have

$$\frac{\partial H(0, \tau_s^0)}{\partial u} = 0 \tag{5-6a}$$

This leads to

$$w'(\tau_s^0) = 0 \tag{5-6b}$$

as the condition which defines τ_s^0. From Eq. (4-149), we find that in order to satisfy Eq. (5-6) we must have

$$D_1 = (\frac{8}{15}\tau^2 - \frac{1}{5\tau})\, h_2 \tag{5-7a}$$

$$D_2 = (\frac{16}{35} h_3 - \frac{48}{175} h_2^2)\, \tau^3 - (\frac{3}{5} h_3 - \frac{7}{25} h_2^2) - \frac{1}{50} h_2^2 \frac{1}{\tau^3} \tag{5-7b}$$

$$D_3 = (\frac{128}{315} h_4 - \frac{11968}{17325} h_2 h_3 + \frac{22912}{70875} h_2^3)\, \tau^4 + (-\frac{448}{315} h_4 + \frac{33308}{17325} h_2 h_3 - \frac{12176}{15750} h_2^3)\, \tau - \frac{1}{375} h_2^3 \frac{1}{\tau^2} - \frac{1}{250} h_2^3 \frac{1}{\tau^5} \tag{5-7c}$$

where all τ are to be taken to be τ_s^0. The roots of

$$w_{1,2}'(\tau_s^0) = 0 \tag{5-8a}$$

are well known. They can be expressed in the form

$$\tau_s^0 = \beta_s \exp(\pm\tfrac{1}{3}\pi) \tag{5-8b}$$

where β_s denotes the roots of the derivative of the Airy function

$$Ai'(-\beta_s) = 0$$

In Table 2-2 we list the 15 decimal values by Miller and Haines. Values which are given to 25 decimals have been obtained by Sherry (Ref. 1).

For the case of arbitrary q, we can use Eq. (4-153a) and Eq. (4-153b) to express Eq. (5-1b) in the form

$$-w'(\tau_s) + \sum_{n=1}^{\infty} \left[C_n(0, \tau_s)\, w(\tau_s) + D_n(0, \tau_s)\, w'(\tau_s)\right] \beta^n + q\, w(\tau_s) + q \sum_{n=1}^{\infty} \left[A_n(0, \tau_s)\, w(\tau_s) + B_n(0, \tau_s) w'(\tau_s)\right] \beta^n = 0 \tag{5-9}$$

Since the coefficient of each power of β must vanish, we find that τ_s must be defined by

$$w'(\tau_s) - q\, w(\tau_s) = 0 \tag{5-10}$$

We can use Eq. (5-10) to express the coefficient of β^n in Eq. (5-9) in the form

$$q\left[D_n(0, \tau_s) + q\, B_n(0, \tau_s)\right] + \left[C_n(0, \tau_s) + q\, B_n(0, \tau_s)\right] = 0 \tag{5-11}$$

For n = 1, Eq. (5-11) takes the form

$$q\left\{-\frac{2}{15}h_2\tau + q\left[-\frac{8}{15}h_2\tau^2 + D_1\right]\right\} + \left\{\left[\frac{8}{15}h_2\tau^3 - \frac{1}{5}h_2 - D_1\tau\right] + q\left[-\frac{2}{15}h_2\tau\right]\right\} = 0 \tag{5-12}$$

We can solve Eq. (5-12) for D_1 and find that

$$D_1 = \frac{h_2}{15}\left[8\tau^2 - \frac{3 + 4\tau q}{\tau - q^2}\right] \tag{5-13}$$

For n = 2, Eq. (5-11) takes the form

$$\begin{aligned} q\Bigg\{&\left[-\frac{32}{225}h_2{}^2\tau^5 + \frac{8}{15}h_2D_1\tau^3 + (-\frac{4}{35}h_3 + \frac{86}{315}h_2{}^2)\,\tau^2 - \frac{1}{2}D_1{}^2\tau - \frac{1}{3}h_2D_1\right] \\ &+ q\left[-(\frac{16}{35}h_3 - \frac{288}{315}h_2{}^2)\,\tau^3 - \frac{6}{5}h_2D_1\tau + (\frac{3}{7}h_3 - \frac{9}{35}h_2{}^2 + D_2)\right]\Bigg\} \\ &+ \Bigg\{\left[\tau^4(\frac{16}{35}h_3 - \frac{288}{315}h_2{}^2) + \tau^2(\frac{22}{15}h_2D_1) + \tau(-\frac{3}{5}h_3 + \frac{7}{15}h_2{}^2 - D_2) - \frac{1}{2}D_1{}^2\right] \\ &+ q\left[\frac{32}{225}h_2{}^2\tau^5 - \frac{8}{15}h_2D_1\tau^3 + (-\frac{4}{35}h_3 + \frac{58}{315}h_2{}^2)\,\tau^2 + \frac{1}{2}D_1{}^2\tau - \frac{2}{15}h_2D_1\right]\Bigg\} = 0 \end{aligned} \tag{5-14}$$

If we combine the similar terms in Eq. (5-14) we find that D_2 is given by

$$\begin{aligned} (\tau - q^2)\,D_2 = {}&(\tau - q^2)\,\tau^3(\frac{16}{35}h_3 - \frac{288}{315}h_2{}^2) + \frac{22}{15}h_2D_1\tau^2 + (-\frac{3}{5}h_3 + \frac{7}{15}h_2{}^2)\,\tau \\ &- \frac{1}{2}D_1{}^2 + q\left[(-\frac{8}{35}h_3 + \frac{16}{35}h_2{}^2)\,\tau^2 - \frac{7}{15}h_2D_1\right] + q^2\left[-\frac{6}{5}h_2D_1\tau\right. \\ &\left. + (\frac{3}{7}h_3 - \frac{9}{35}h_2{}^2)\right] \end{aligned} \tag{5-15}$$

If we use Eq. (5-13), we can eliminate D_1 from Eq. (5-15) to obtain the form

$$(\tau - q^2)^2 D_2 = (\tau - q^2)\tau^3 \left(\frac{16}{35} h_3 - \frac{288}{315} h_2^2\right) + \frac{16}{25} h_2^2 \tau^4 - \frac{14}{225} h_2^2 \tau^2 \left(\frac{3+4\tau q}{\tau - q^2}\right)$$
$$+ \tau \left(-\frac{3}{5} h_3 + \frac{7}{15} h_2^2\right) - \frac{1}{450} h_2^2 \left(\frac{3 + 4\tau q}{\tau - q^2}\right)^2$$
$$+ q \left[\left(-\frac{8}{35} h_3 + \frac{328}{1575} h_2^2\right)\tau^2 + \frac{7}{225} h_2^2 \left(\frac{3 + 4\tau q}{\tau - q^2}\right)\right]$$
$$+ q^2 \left[-\frac{16}{25} h_2^2 \tau^3 + \left(\frac{3}{7} h_3 - \frac{9}{35} h_2^2\right) + \frac{2}{25} h_2^2 \tau \left(\frac{3+4\tau q}{\tau + q^2}\right)\right] \qquad (5\text{-}16)$$

For n = 3, Eq. (5-11) takes the form

$$q\left\{\left[\left(-\frac{128}{525} h_2 h_3 + \frac{34112}{70875} h_2^3\right)\tau^6 + \left(\frac{16}{35} h_3 - \frac{512}{315} h_2^2\right) D_1 \tau^4 + \left(-\frac{32}{315} h_4\right.\right.\right.$$
$$\left. + \frac{2816}{3465} h_2 h_3 - \frac{10604}{14175} h_2^3 + \frac{8}{15} h_2 D_2\right)\tau^3 + \frac{23}{15} h_2 D_1^2 \tau^2 + \left(-\frac{29}{35} h_3\right.$$
$$\left.\left.\left. + \frac{319}{315} h_2^2 - D_2\right) D_1 \tau + \left(\frac{2}{9} h_4 - \frac{25}{63} h_2 h_3 + \frac{277}{1575} h_2^3 - \frac{1}{3} h_2 D_2 - \frac{1}{3} D_1^3\right)\right]\right\}$$
$$+q\left[-\frac{256}{10125} h_2^3 \tau^7 + \frac{32}{225} h_2^2 D_1 \tau^5 + \left(-\frac{128}{315} h_4 + \frac{1408}{693} h_2 h_3 - \frac{28688}{14175} h_2^3\right)\tau^4\right.$$
$$- \frac{4}{15} h_2 D_1^2 \tau^3 + \left(-\frac{52}{35} h_3 + \frac{922}{315} h_2^2\right) D_1 \tau^2 + \left(\frac{80}{63} h_4 - \frac{8338}{3465} h_2 h_3\right.$$
$$\left.\left. + \frac{1822}{1575} h_2^3 - \frac{6}{5} h_2 D_2 + \frac{1}{6} D_1^3\right)\tau + \left(-\frac{2}{3} h_2 D_1^2 + D_3\right)\right]\Bigg\}$$
$$+\left\{\left[\frac{256}{10125} h_2^3 \tau^8 - \frac{32}{225} h_2^2 D_1 \tau^6 + \left(+\frac{128}{315} h_4 - \frac{1408}{693} h_2 h_3 + \frac{142768}{70875} h_2^3\right)\tau^5\right.\right.$$
$$+ \frac{4}{15} h_2 D_1^2 \tau^4 + \left(\frac{12}{7} h_3 - \frac{1066}{315} h_2^2\right) D_1 \tau^3 + \left(+\frac{962}{315} h_2 h_3 - \frac{2564}{1575} h_2^3 - \frac{64}{45} h_4\right.$$
$$\left.\left. + \frac{22}{15} h_2 D_2 - \frac{1}{6} D_1^3\right)\tau^2 + \left(\frac{13}{10} h_2 D_1^2 - D_3\right)\tau + \left(-\frac{3}{5} h_3 + \frac{7}{15} h_2^2 - D_2\right) D_1\right]$$
$$+q\left[\left(\frac{128}{525} h_2 h_3 - \frac{35008}{70875} h_2^3\right)\tau^6 + \left(-\frac{16}{35} h_3 + \frac{512}{315} h_2^2\right) D_1 \tau^4 + \left(-\frac{32}{315} h_4\right.\right.$$
$$+ \frac{704}{3465} h_2 h_3 - \frac{3404}{14175} h_2^3 - \frac{8}{15} h_2 D_2\Big)\tau^3 - \frac{7}{5} h_2 D_1^2 \tau^2 + \left(\frac{1}{5} h_3 + \frac{1}{9} h_2^2\right.$$
$$\left.\left.\left. + D_2\right) D_1 \tau + \left(\frac{2}{9} h_4 - \frac{14}{45} h_2 h_3 + \frac{28}{225} h_2^3 - \frac{2}{15} h_2 D_2 + \frac{1}{6} D_1^3\right)\right]\Bigg\} = 0 \qquad (5\text{-}17)$$

If we solve for D_3, and combine similar terms, we find that

$$(\tau - q^2)\, D_3 = \frac{256}{10125} h_2^3 \tau^8 - \frac{32}{225} h_2^2 D_1 \tau^6 + \left(\frac{128}{315} h_4 - \frac{1408}{693} h_2 h_3 + \frac{142768}{70875} h_2^3\right) \tau^5$$
$$+ \frac{4}{15} h_2 D_1^2 \tau^4 + \left(\frac{12}{7} h_3 - \frac{1066}{315} h_2^2\right) D_1 \tau^3 + \left(-\frac{64}{45} h_4 + \frac{962}{315} h_2 h_3\right.$$
$$\left. - \frac{2564}{1575} h_2^3 + \frac{22}{15} h_2 D_2 - \frac{1}{6} D_1^3\right) \tau^2 + \frac{13}{10} h_2 D_1^2 \tau + \left(-\frac{3}{5} h_3 + \frac{7}{15} h_2^2 - D_2\right) D_1$$
$$+ q \left[-\frac{128}{10125} h_2^3 \tau^6 + \left(-\frac{64}{315} h_4 + \frac{704}{693} h_2 h_3 - \frac{14008}{14175} h_2^3\right) \tau^3 + \frac{2}{15} h_2 D_1^2 \tau^2 \right.$$
$$\left. + \left(-\frac{22}{35} h_3 + \frac{354}{1575} h_2^2\right) \tau + \left(\frac{4}{9} h_4 - \frac{223}{315} h_2 h_3 + \frac{473}{1575} h_2^3 - \frac{7}{15} h_2 D_2 - \frac{1}{6} D_1^3\right) \right]$$
$$+ q^2 \left[-\frac{256}{10125} h_2^3 \tau^7 + \frac{32}{225} h_2^2 D_1 \tau^5 + \left(-\frac{128}{315} h_4 + \frac{1408}{693} h_2 h_3 - \frac{28688}{14175} h_2^3\right) \tau^4 \right.$$
$$- \frac{4}{15} h_2 D_1^2 \tau^3 + \left(-\frac{52}{35} h_3 + \frac{922}{315} h_2^2\right) D_1 \tau^2 + \left(\frac{80}{63} h_4 - \frac{8338}{3465} h_2 h_3\right.$$
$$\left. \left. + \frac{1822}{1575} h_2^3 - \frac{6}{5} h_2 D_2 + \frac{1}{6} D_1^3\right) \tau - \frac{2}{3} h_2 D_1^2 \right] \qquad (5\text{-}18)$$

Because of the complexity of Eq. (5-18) in its present form, we have not used Eq. (5-13) and Eq. (5-16) to eliminate D_1 and D_2. The algebra has become so tedious that we have not attempted to obtain D_4 except for the $q = \infty$ (See. Eq. (5-4d).

5.2 Another Expansion for λ_s

The expansions for $\tau_s(q)$ given in Eq. (2-21) and Eq. (2-22) can be used in Eq. (5-13), Eq. (5-16), and Eq. (5-18) to obtain representations for the solutions of Eq. (5-1c) in the form

$$\lambda_s = k^2 f_o + k^{4/3} f_1^{2/3} \left[\tau_s^o + \sum_{n=1}^{\infty} P_n(\tau_s^o, \frac{q}{\beta})\, \beta^n \right] \qquad (5\text{-}19)$$

or

$$\lambda_s = k^2 f_o + k^{4/3} f_1^{2/3} \left[\tau_s^\infty + \sum_{n=1}^{\infty} Q_n(\tau_s^\infty, \frac{1}{q\beta})\, \beta^n \right] \qquad (5\text{-}20)$$

where the coefficients $P_n(\tau_s^o, \frac{q}{\beta})$ and $Q_n(\tau_s^\infty, \frac{1}{q\beta})$ are of the form

$$P_n(\tau_s^o, \frac{q}{\beta}) = D_n(\tau_s^o, 0) + \sum_{m=1}^{\infty} a_m^n (\tau_s^o) (\frac{q}{\beta})^m \qquad (5\text{-}21)$$

$$Q_n(\tau_s^\infty, \frac{1}{q\beta}) = D_n(\tau_s^\infty, \infty) + \sum_{m=1}^{\infty} b_m^n (\tau_s^\infty) \frac{1}{(\beta q)^m} \qquad (5\text{-}22)$$

where $D_n(\tau_s^o, 0)$ is given by Eq. (5-7) and $D_n(\tau_s^\infty, \infty)$ is given by Eq. (5-4).

It is perhaps more convenient to determine $P_n(\tau_s^o, \frac{q}{\beta})$ and $Q_n(\tau_s^\infty, \frac{1}{\beta q})$ by expressing the boundary condition in the form

$$\frac{\partial H(0, \tau)}{\partial u} + \beta Q H(0, \tau) = 0 \quad , \quad q = \beta Q \qquad (5\text{-}23)$$

when q is small, and in the form

$$H(0, \tau) + \beta P \frac{\partial H(0, \tau)}{\partial u} = 0 \quad , \quad \frac{1}{q} = \beta P \qquad (5\text{-}24)$$

when q is large.

For the case of small q, Eq. (5-23) leads to

$$w'(\tau_s^o) = 0 \qquad (5\text{-}25a)$$

$$c_1(0, \tau_s^o) + Q = 0 \qquad (5\text{-}25b)$$

$$c_2(0, \tau_s^o) + QA_1(0, \tau_s^o) = 0 \qquad (5\text{-}25c)$$

$$c_{n+1}(0, \tau_s^o) + QA_n(0, \tau_s^o) = 0 \ , \ n \geq 1 \qquad (5\text{-}25d)$$

where the $C_n(u,\tau)$ and $A_n(u,\tau)$ are defined in Eq. (4-153). From Eq. (5-25) we find that

$$P_1(\tau, Q) = (\frac{8}{15}\tau^2 - \frac{1}{5\tau}) h_2 + \frac{Q}{\tau} \qquad (5\text{-}26a)$$

$$P_2(\tau, Q) = (\frac{16}{35} h_3 - \frac{48}{175} h_2^2) \tau^3 - (\frac{3}{5} h_3 - \frac{7}{25} h_2^2) - \frac{1}{50} h_2^2 \frac{1}{\tau^3} + \frac{4}{5} h_2 Q + \frac{1}{5} h_2 \frac{Q}{\tau^3} - \frac{1}{2} \frac{Q^2}{\tau^3} \qquad (5\text{-}26b)$$

$$P_3(\tau, Q) = \left(\frac{128}{315} h_4 - \frac{11968}{17325} h_2 h_3 + \frac{22912}{70875} h_2^3\right) \tau^4 + \left(-\frac{448}{315} h_4 + \frac{33308}{17325} h_2 h_3 - \frac{12176}{15750} h_2^3\right) \tau - \frac{1}{375} h_2^3 \frac{1}{\tau^2} - \frac{1}{250} h_2^3 \frac{1}{\tau^5}$$
$$+ Q\left[-\frac{272}{315} h_2^2 \tau + \frac{8}{7} h_3 \tau + \frac{1}{25} h_2^2 \frac{1}{\tau^2} + \frac{3}{50} h_2^2 \frac{1}{\tau^5}\right]$$
$$+ Q^2\left[-\frac{1}{5} h_2 \frac{1}{\tau^2} - \frac{3}{10} h_2 \frac{1}{\tau^5}\right] + Q^3\left[\frac{1}{3} \frac{1}{\tau^2} + \frac{1}{2} \frac{1}{\tau^5}\right] \qquad (5\text{-}26c)$$

For the case of large q, Eq. (5-24) leads to

$$w(\tau_s^\infty) = 0 \qquad (5\text{-}27a)$$
$$B_1(0, \tau_s^\infty) - P = 0 \qquad (5\text{-}27b)$$
$$B_2(0, \tau_s^\infty) + P\, D_1(0, \tau_s^\infty) = 0 \qquad (5\text{-}27c)$$
$$B_{n+1}(0, \tau_s^\infty) + P\, D_n(0, \tau_s^\infty) = 0 \quad , \quad n \geq 1 \qquad (5\text{-}27d)$$

where the $B_n(u,\tau)$ and $D_n(u,\tau)$ are defined by Eq. (4-153).* From Eq. (5-27), we find that

$$Q_1(\tau, P) = \frac{8}{15} h_2 \tau^2 + P \qquad (5\text{-}28a)$$
$$Q_2(\tau, P) = \left(\frac{16}{35} h_3 - \frac{48}{175} h_2^2\right) \tau^3 - \left(\frac{3}{7} h_3 - \frac{9}{35} h_2^2\right) + \frac{4}{3} h_2 \tau P \qquad (5\text{-}28b)$$
$$Q_3(\tau, P) = \left(\frac{128}{315} h_4 - \frac{11968}{17325} h_2 h_3 + \frac{22912}{70875} h_2^3\right) \tau^4 - \left(\frac{80}{63} h_4 - \frac{6556}{3465} h_2 h_3 + \frac{1336}{1575} h_2^3\right) \tau + \left(\frac{8}{5} h_3 - \frac{32}{45} h_2^2\right) \tau^2 P + h_2 P^2 + \frac{\tau}{3} P^3 \qquad (5\text{-}28c)$$

Some unpublished work of the present author can be used as a numerical check on the coefficients in Eq. (5-26). It has been shown that the roots ν_s of the Hankel function combination

$$H_{\nu_s}^{(1,2)'}(ka) + \left(\frac{2}{ka}\right) Z\, H_{\nu_s}^{(1,2)}(ka) = 0 \qquad (5\text{-}29)$$

---

*The reader is cautioned to avoid confusing the $D_n(\tau,q)$ defined by Eq. (5-2) with the $D_n(u,\tau)$ defined by Eq. (4-153). The context in which these functions are used should provide the means to distinguish between them.

can be expressed in the form

$$\nu_s = ka + \left(\frac{ka}{2}\right)^{1/3}\Bigg\{ \tau + \left(\frac{2}{ka}\right)^{2/3}\left[\frac{1}{60}\tau^2 - \frac{1}{10}\frac{1}{\tau} + \frac{Z}{\tau}\right]$$
$$+ \left(\frac{2}{ka}\right)^{4/3}\left[\left(-\frac{1}{1400}\tau^3 + \frac{1}{50} - \frac{1}{200}\frac{1}{\tau^3}\right) + Z\left(\frac{1}{10}\frac{1}{\tau^3} - \frac{1}{10}\right) - Z^2\left(\frac{1}{2}\frac{1}{\tau^3}\right)\right]$$
$$+ \left(\frac{2}{ka}\right)^{6/3}\left[\left(\frac{281}{4536000}\tau^4 - \frac{611}{126000}\tau - \frac{1}{3000}\frac{1}{\tau^2} - \frac{1}{2000}\frac{1}{\tau^5}\right)\right.$$
$$\left. + Z\left(\frac{41}{2520}\tau + \frac{1}{100}\frac{1}{\tau^2} + \frac{3}{200}\frac{1}{\tau^5}\right) - Z^2\left(\frac{1}{10}\frac{1}{\tau^2} + \frac{3}{20}\frac{1}{\tau^5}\right)\right.$$
$$\left. + Z^3\left(\frac{1}{3}\frac{1}{\tau^2} + \frac{1}{2}\frac{1}{\tau^5}\right)\right]$$
$$+ \left(\frac{2}{ka}\right)^{8/3}\left[\left(-\frac{73769}{10478160000}\tau^5 + \frac{56299}{24948000}\tau^2 + \frac{1679}{1108800}\frac{1}{\tau}\right.\right.$$
$$\left. - \frac{7}{120000}\frac{1}{\tau^4} - \frac{1}{16000}\frac{1}{\tau^7}\right) + Z\left(-\frac{97}{32400}\tau^2 - \frac{803}{42000}\frac{1}{\tau}\right.$$
$$\left. + \frac{7}{3000}\frac{1}{\tau^4} + \frac{1}{400}\frac{1}{\tau^7}\right) + Z^2\left(\frac{2017}{25200}\frac{1}{\tau} - \frac{7}{200}\frac{1}{\tau^4} - \frac{3}{80}\frac{1}{\tau^7}\right)$$
$$\left. + Z^3\left(-\frac{11}{90}\frac{1}{\tau} + \frac{7}{30}\frac{1}{\tau^4} + \frac{1}{4}\frac{1}{\tau^7}\right) + Z^4\left(-\frac{7}{12}\frac{1}{\tau^4} - \frac{5}{8}\frac{1}{\tau^7}\right)\right]$$
$$\left. + \cdots\cdot\right\} \qquad (5\text{-}30)$$

where

$$\tau = \tau_s^o = \beta_s \exp(\pm i\pi/3) \qquad (5\text{-}31)$$

where the + sign is used with $H_\nu^{(1)}(ka)$ and the - sign with $H_\nu^{(2)}(ka)$. If we express ν_s^2 in the form

$$\left(\frac{\nu_s}{ka}\right)^2 = 1 + \left(\frac{2}{ka}\right)^{2/3}\left[\tau_s^o + \sum_{n=1}^{\infty} E_n(\tau_s^o, Z)\left(\frac{2}{ka}\right)^{2n/3}\right] \qquad (5\text{-}32)$$

we find that

$$E_1(\tau, Z) = \frac{4}{15}\tau^2 - \frac{1}{10}\frac{1}{\tau} + \frac{Z}{\tau} \qquad (5\text{-}33a)$$

$$E_2(\tau, Z) = \frac{4}{525}\tau^3 - \frac{3}{100} - \frac{1}{200}\frac{1}{\tau^3} + Z\left(\frac{1}{10}\frac{1}{\tau^3} + \frac{2}{5}\right) - \frac{Z^2}{2\tau^3} \qquad (5\text{-}33b)$$

$$E_3(\tau, Z) = -\frac{16}{70875}\tau^4 + \frac{34}{7875}\tau - \frac{1}{3000}\frac{1}{\tau^2} - \frac{1}{2000}\frac{1}{\tau^5} + Z\left(-\frac{8}{315}\tau + \frac{1}{100}\frac{1}{\tau^2} + \frac{3}{200}\frac{1}{\tau^5}\right) - Z^2\left(\frac{1}{10}\frac{1}{\tau^2} + \frac{3}{20}\frac{1}{\tau^5}\right) + Z^3\left(\frac{1}{3}\frac{1}{\tau^2} + \frac{1}{2}\frac{1}{\tau^5}\right) \quad (5\text{-}33c)$$

$$E_4(\tau, Z) = \frac{1472}{81860625}\tau^5 + \frac{859}{24948000}\tau^2 + \frac{53}{173250}\frac{1}{\tau} - \frac{21}{360000}\frac{1}{\tau^4} - \frac{1}{16000}\frac{1}{\tau^7} + Z\left(\frac{8}{2025}\tau^2 + \frac{3}{1750}\frac{1}{\tau} + \frac{7}{3000}\frac{1}{\tau^4} + \frac{1}{400}\frac{1}{\tau^7}\right) + Z^2\left(-\frac{38}{1575}\frac{1}{\tau} - \frac{7}{200}\frac{1}{\tau^4} - \frac{3}{80}\frac{1}{\tau^7}\right) + Z^3\left(\frac{2}{45}\frac{1}{\tau} + \frac{7}{30}\frac{1}{\tau^4} + \frac{1}{4}\frac{1}{\tau^7}\right) + Z^4\left(-\frac{7}{12}\frac{1}{\tau^4} - \frac{5}{8}\frac{1}{\tau^7}\right) \quad (5\text{-}33d)$$

Let us recall [from Eq. (4-161b) and Eq. (4-164)] that for the Hankel functions

$$\beta = \left(\frac{a}{2}\right)\left(\frac{2}{ka}\right)^{2/3}, \quad h_2 = \frac{1}{a}, \quad h_3 = \frac{2}{3a^2}, \quad h_4 = \frac{1}{3a^3} \quad .$$

If we let

$$Q = \left(\frac{2}{a}\right) Z$$

we find that we can check the numerical coefficients in Eq. (5-26) by a comparison with Eq. (5-33) since

$$\left(\frac{a}{2}\right)^n P_n(\tau, Q) = E_n\left(\tau, \frac{a}{2}Q\right) \quad (5\text{-}34)$$

We have also shown that the roots ν_s defined by

$$H_{\nu_s}^{(1,2)}(ka) + \left(\frac{2}{ka}\right)^{1/3}\Gamma\, H_{\nu_s}^{(1,2)}(ka) = 0 \quad (5\text{-}35)$$

can be represented by an asymptotic expansion of the form

$$\nu_s = ka + \left(\frac{ka}{2}\right)^{1/3}\left\{\tau + \left(\frac{2}{ka}\right)^{2/3}\left[\frac{1}{60}\tau^2 + \Gamma\right] + \left(\frac{2}{ka}\right)^{4/3}\left[-\frac{1}{1400}\tau^3 - \frac{1}{140} + \frac{1}{6}\tau\Gamma\right] + \left(\frac{2}{ka}\right)^{6/3}\left[\frac{281}{4536000}\tau^4 + \frac{29}{12600}\tau - \frac{1}{360}\tau^2\Gamma + \frac{1}{4}\Gamma^2 + \frac{1}{3}\tau\Gamma^3\right] + \cdots\right\} \quad (5\text{-}36)$$

where

$$\tau = t_s^{\infty} = \alpha_s \exp(\pm i\pi/3) \quad (5\text{-}37)$$

where the + sign is used with $H_\nu^{(1)}(ka)$. If we express ν_s^2 in the form

$$\left(\frac{\nu_s}{ka}\right)^2 = 1 + \left(\frac{2}{ka}\right)^{2/3}\left[\tau_s^\infty + \sum_{n=1}^{\infty} F_n(\tau_s^\infty, \Gamma)\left(\frac{2}{ka}\right)^{2n/3}\right] \tag{5-38}$$

we find that

$$F_1(\tau, \Gamma) = \frac{4}{15}\tau^2 + \Gamma \tag{5-39a}$$

$$F_2(\tau, \Gamma) = \frac{4}{525}\tau^3 - \frac{1}{140} + \frac{2}{3}\tau\,\Gamma \tag{5-39b}$$

$$F_3(\tau, \Gamma) = -\frac{16}{70875}\tau^4 - \frac{2}{1575}\tau + \frac{4}{45}\tau^2\,\Gamma + \frac{1}{4}\Gamma^2 + \frac{1}{3}\tau\,\Gamma^3 \tag{5-39c}$$

If we set

$$P = \left(\frac{2}{a}\right)\Gamma$$

we can use the relation

$$\left(\frac{a}{2}\right)^n Q_n(\tau, P) = F_n\left(\tau, \frac{a}{2}P\right) \tag{5-40}$$

to obtain a method of verifying the accuracy of the coefficients in Eq. (5-28).

5.3 The Turning Points

In the studies by Pekeris (Ref. 2), Imai (Refs. 3 and 4), Bremmer (Ref. 5) and other authors, one needs the values of the turning point y_1 defined by Eq. (4-24), namely

$$\lambda_s + k^2 f(y_1) = k^2[\mu_s + f(y_1)] = 0 \tag{5-41}$$

If we recall the steps which we took in Eqs. (4-74) through (4-77), we see that we can define

$$y_1 = \beta u_1 = [k^{\frac{2}{3}} f_1^{\frac{1}{3}}]^{-1} u_1 \tag{5-42}$$

where u_1 is defined by

$$-t_s + u_1 + \beta h_2 u_1^2 + \beta^2 h_3 u_1^3 + \beta^3 h_4 u_1^4 + \ldots = 0$$

If we invert this expression, we find that

$$u_1 = t_s - \beta h_2 t_s^2 - \beta^2 (h_3 - 2h_2^2) t_s^3 - \beta^3 (h_4 - 5h_2h_3 + 5h_2^3) t_s^4$$
$$- \beta^4 (h_5 - 6h_2h_4 - 3h_3^2 + 21h_2^2h_3 - 14h_2^4) t_s^5 + O(\beta^5) \qquad (5\text{-}43)$$

From Eq. (5-2) and Eq. (5-13), we obtain the result

$$t_s = \tau_s + \beta h_2 \frac{1}{15}\left[8\tau_s^2 - \frac{3 + 4q\tau_s}{\tau_s - q^2}\right] + O(\beta^2) \qquad (5\text{-}44)$$

Therefore, we find that

$$u_1 = \beta\tau_s - \beta^2 h_2 \frac{1}{15}\left[7\tau_s^2 + \frac{3 + 4q\tau_s}{\tau_s - q^2}\right] + O(\beta^3) \qquad (5\text{-}45)$$

For $q = 0$, we can use the results in Eq. (5-7) to show that

$$u_1 = \beta\left\{\tau - \beta\left[\frac{7}{15}h_2\tau^2 + \frac{1}{5\tau}h_2\right] - \beta^2\left[\left(\frac{19}{35}h_3 - \frac{346}{525}h_2^2\right)\tau^3 + \left(\frac{3}{5}h_3 - \frac{17}{25}h_2^2\right)\right.\right.$$
$$\left. + \frac{1}{50}h_2^2\right)\frac{1}{\tau^3}\Big] - \beta^3\left[\left(\frac{187}{315}h_4 - \frac{31097}{17325}h_2h_3 + \frac{85943}{70875}h_2^3\right)\tau^4\right.$$
$$\left. + \left(\frac{448}{315}h_4 - \frac{64493}{17325}h_2h_3 + \frac{36536}{15750}h_2^3\right)\tau + \frac{17}{750}h_2^3\frac{1}{\tau^2} + \frac{1}{250}h_2^3\frac{1}{\tau^5}\right]$$
$$\left. + O(\beta^4)\right\} \qquad (5\text{-}46)$$

where all τ are to be taken to be τ_s^0.

For $q = \infty$, we can use the results in Eq. (5-4) to show that

$$u_1 = \beta\left\{\tau - \frac{7}{15}h_2\tau^2\beta - \beta^2\left[\left(\frac{19}{35}h_3 - \frac{346}{525}h_2^2\right)\tau^3 + \left(\frac{3}{7}h_3 - \frac{9}{35}h_2^2\right)\right]\right.$$
$$- \beta^3\left[\left(\frac{187}{315}h_4 - \frac{31097}{17325}h_2h_3 + \frac{85943}{70875}h_2^3\right)\tau^4 + \left(\frac{80}{63}h_4 - \frac{9526}{3465}h_2h_3\right.\right.$$
$$\left.\left. + \frac{2146}{1575}h_2^3\right)\tau\right]$$
$$- \beta^4\left[\left(\frac{437}{693}h_5 - \frac{116746}{51975}h_2h_4 + \frac{194059}{40425}h_3^2 - \frac{200019867}{5457375}h_2^2h_3\right.\right.$$
$$\left. + \frac{2084790958}{81860625}h_2^4\right)\tau^5 + \left(\frac{1808}{693}h_5 - \frac{76832}{10395}h_2h_4 - \frac{29719}{8085}h_3^2\right.$$
$$\left.\left. + \frac{15327645}{1091475}h_2^2h_3 - \frac{30322830}{5457375}h_2^4\right)\tau^2\right] + O(\beta^5)\Big\} \qquad (5\text{-}47)$$

where all τ are to be taken to be τ_s^∞.

5.4 The Propagation Constants

In many applications we need representations for $\sqrt{\lambda_s}$. For example, in the case of the Hankel functions $H_\nu^{(1)}(ka)$ we defined

$$\lambda = (\nu/a)^2 \tag{5-48}$$

in Eq. (4-162a). In order to treat these problems it will be convenient to define

$$\lambda = \kappa^2 \tag{5-49}$$

We will call κ the propagation constant. It will also be convenient to define

$$f_0 = n_0^2 \tag{5-50}$$

and call n_o the index of refraction at $y = 0$.

We can then express Eq. (5-2) in the form

$$\kappa_s^2 = (kn_o)^2 + k^{4/3} f_1^{2/3} \tau_s + k^{4/3} f_1^{2/3} \sum_{n=1}^{\infty} D_n(\tau_s, q)\, \beta^n \tag{5-51}$$

If we assume that κ_s can be expressed in the form

$$\kappa_s = kn_o + \frac{1}{2} \frac{k^{1/3} f_1^{2/3}}{n_o} \left[\tau_s + \sum_{n=1}^{\infty} N_n(\tau_s, q)\, \beta^n\right] \tag{5-52}$$

we can show that

$$\kappa_s^2 = (kn_o)^2 + k^{4/3} f_1^{2/3} \tau_s + k^{4/3} f_1^{2/3} \left\{ \left[N_1 + \frac{f_1}{4f_o} \tau^2\right]\beta + \left[N_2 + \frac{f_1}{4f_o} 2\tau N_1\right]\beta^2 + \left[N_3 + \frac{f_1}{4f_o}(2\tau N_2 + N_1^2)\right]\beta^3 + \cdots \right\} \tag{5-53}$$

Therefore, we find that

$$N_1(\tau, q) = D_1(\tau, q) - \frac{f_1}{4f_o} \tau^2 \tag{5-54a}$$

$$N_2(\tau, q) = D_2(\tau, q) - \frac{f_1}{4f_o} 2\tau N_1(\tau, q) \tag{5-54b}$$

$$N_3(\tau, q) = D_3(\tau, q) - \frac{f_1}{4f_o} \left[2\tau N_2(\tau, q) + N_1^2(\tau, q)\right] \tag{5-54c}$$

The expansions for ν_s which are given in Eq. (5-30) and Eq. (5-36) can be used to verify the correctness of the coefficients in Eq. (5-54).

5.5 The Characteristic Wave Numbers

Up to this point, we have always considered $\underline{k}$ to be known and we have sought to determine the eigenvalues λ_s. However, in pulse diffraction problems we encounter the case in which λ is known and we must find the characteristic wave numbers k_s for which either

$$F(0,\lambda) = 0 \tag{5-55a}$$

or

$$[dF(y,\lambda)/dy]_{y=0} = 0 \tag{5-55b}$$

The more general problem

$$[dF/dy + i\, k_s\, Z\, F]_{y=0} = 0 \tag{5-55c}$$

will not be considered because the impedance Z is usually a function of k_s.

We can determine k_s for the Dirichlet problem defined by Eq. (5-55a) by using Eq. (5-2) with D_n defined by Eq. (5-4). Let us consider Eq. (5-2) as the definition of a relation between λ and k which makes $F(y,\lambda)$ vanish for $y = 0$. We can then define the characteristic wave numbers k_s by means of Eq. (5-2) which we write in the form

$$\lambda = k_s^{\,2} f_0 + k_s^{\,4/3} f_1^{\,2/3} \tau_s + D_1(\tau_s)\, k_s^{\,2/3} f_1^{\,1/3} + \sum_{n=2}^{\infty} D_n(\tau_s)\, k_s^{\frac{4-2n}{3}} f_1^{\frac{2-n}{3}} \tag{5-56}$$

If we seek a representation for k_s of the form

$$f_0 k_s^{\,2} = \lambda - f_1^{\,2/3} \tau_s \left(\frac{\lambda}{f_0}\right)^{2/3} + \sum_{n=1}^{\infty} F_n(\tau_s)\, f_1^{\frac{2-n}{3}} \left(\frac{\lambda}{f_0}\right)^{\frac{4-2n}{3}} \tag{5-57}$$

we can determine $F_n(\tau_s)$ by inserting Eq. (5-57) in Eq. (5-56) and

equate the coefficients of λ on each side of the equation.

The above suggestion leads to an expansion for k_s^2 whereas in practice we generally want k_s. However, if we express k_s in the form

$$k_s = \sqrt{\frac{\lambda}{f_o}} - \frac{1}{2f_o} f_1^{2/3} \tau_s \left(\frac{\lambda}{f_o}\right)^{1/6} + \sum_{n=1}^{\infty} M_n(\tau_s) f_1^{\frac{2-n}{3}} \left(\frac{\lambda}{f_o}\right)^{\frac{1-2n}{6}} \tag{5-58}$$

and use this in Eq. (5-56) we can relate the $M_n(\tau_s)$ to the $D_n(\tau_s)$.

For the special case of the k_s defined by the Hankel function combination

$$H_\nu^{(1,2)'}(k_s a) - \left(\frac{i}{ka}\right) Z H_\nu^{(1,2)}(k_s a) = 0 \tag{5-59}$$

we have shown that

$$k_s a = \nu - \left(\frac{\nu}{2}\right)^{1/3} \tau + \left(\frac{2}{\nu}\right)^{1/3} R_1(\tau, Z) + \left(\frac{2}{\nu}\right)^{3/3} R_2(\tau, Z) + \left(\frac{2}{\nu}\right)^{5/3} R_3(\tau, Z) + \left(\frac{2}{\nu}\right)^{7/3} R_4(\tau, Z) \tag{5-60}$$

where

$$R_1(\tau, Z) = +\frac{3\tau^2}{20} + \frac{1}{10}\frac{1}{\tau} - \frac{Z}{\tau} \tag{5-61a}$$

$$R_2(\tau, Z) = +\frac{\tau^3}{1400} - \frac{1}{50} + \frac{1}{200}\frac{1}{\tau^3} - Z\left(\frac{1}{10}\frac{1}{\tau^3} - \frac{1}{10}\right) + \frac{1}{2}\frac{Z^2}{\tau^3} \tag{5-61b}$$

$$R_3(\tau, Z) = -\frac{479}{504000}\tau^4 - \frac{509}{126000}\tau + \frac{1}{3000}\frac{1}{\tau^2} + \frac{1}{2000}\frac{1}{\tau^5} - Z\left(-\frac{11}{280}\tau + \frac{1}{100}\frac{1}{\tau^2} + \frac{3}{200}\frac{1}{\tau^5}\right) - Z^2\left(-\frac{1}{10}\frac{1}{\tau^2} - \frac{3}{20}\frac{1}{\tau^5}\right) - Z^3\left(\frac{1}{3}\frac{1}{\tau^2} + \frac{1}{2}\frac{1}{\tau^5}\right) \tag{5-61c}$$

$$R_4(\tau, Z) = -\frac{20231}{129360000}\tau^5 - \frac{20879}{24948000}\tau^2 - \frac{2006}{11088000}\frac{1}{\tau} + \frac{7}{120000}\frac{1}{\tau^4} + \frac{5}{80000}\frac{1}{\tau^7} - Z\left(-\frac{5}{226800}\tau^2 + \frac{819}{126000}\tau^2 + \frac{391}{126000}\frac{1}{\tau} + \frac{28}{12000}\frac{1}{\tau^4} + \frac{1}{400}\frac{1}{\tau^7}\right) - Z^2\left(-\frac{783}{25200} - \frac{21}{600}\frac{1}{\tau^4} - \frac{111}{400}\frac{1}{\tau^7}\right) - Z^3\left(\frac{1}{10}\frac{1}{\tau} + \frac{1}{10}\frac{1}{\tau^4} + \frac{1}{4} + \frac{1}{\tau^7}\right) + Z^4\left(\frac{7}{12}\frac{1}{\tau^4} + \frac{5}{8}\frac{1}{\tau^7}\right) \tag{5-61d}$$

In these results, τ is defined as in Eq. (5-31).

The coefficients $R_1(\tau,0)$, $R_2(\tau,0)$, and $R_3(\tau,0)$ have been previously given by Olver [See Eq. 9.6 of Ref. 6]. The coefficients $R_1(\tau,-1/4)$, $R_2(\tau,-1/4)$, and $R_3(\tau,-1/4)$ have been previously obtained by Miller (G.F.) and Olver [See Eq. (1.39) of Ref. 7].

For the case $Z = \infty$ when k_s is defined by

$$H_\nu^{(1,2)}(k_s a) = 0 \tag{5-62}$$

we have shown that k_s can be represented by an expansion of the form of Eq. (5-60) in which τ is defined by Eq. (5-37) and

$$B_1(\tau, \infty) = \frac{3}{20}\tau^2 \tag{5-63a}$$

$$B_2(\tau, \infty) = \frac{1}{1400}\tau^3 + \frac{1}{140} \tag{5-63b}$$

$$B_3(\tau, \infty) = -\frac{479}{504000}\tau^4 + \frac{1}{12600}\tau \tag{5-63c}$$

$$B_4(\tau, \infty) = -\frac{20231}{129360000}\tau^5 - \frac{551}{1293600} \tag{5-63d}$$

These coefficients were previosuly obtained by Olver (Ref. 6).

These expansions for k_s can be used to extend the treatment of pulse diffraction by a sphere which was recently the subject of a paper by Weston (Ref. 8). Weston derived the coefficients $B_1(\tau_s^\infty,\infty)$ and $B_1(\tau_s^0,1/4)$. The cases $Z = \infty$ and $Z = 1/4$ were of interest to Weston because he sought to obtain results for pulse diffraction problems by expressing the results in the form

$$U(t) = \frac{c}{2\pi}\int_{-\infty}^{\infty} A(k)\, u(k)\, \exp(-i\,k\,c\,t)\, dk \tag{5-64}$$

where c denotes the velocity of propagation, A(k) denotes the frequency spectrum, and u(k) denotes the solution of the boundary value problem for an $\exp(-ikct) = \exp(-i\omega t)$ time dependance. For the case of the plane wave

$$\underset{\sim}{E}_i = \underset{\sim}{\hat{x}}\, \exp[-ik(z + ct)] \tag{5-65}$$

illuminating a perfectly conducting sphere of radius a, the

secondary field is given by

$$\underset{\approx}{E}_s(k) = \frac{i}{k} \sum_{n=1}^{\infty} \frac{2n+1}{n(n+1)} (-i)^n \left\{ \frac{\psi_n'(ka)}{\zeta_n^{(1)'}(ka)} \text{ curl curl } \left[\underset{\approx}{r}\, h^{(1)}(kr)\, P_n^1(\cos\theta) \cos\phi \right] \right.$$

$$\left. + \frac{\psi_n(ka)}{\zeta_n^{(1)}(ka)} \text{ curl } \left[\underset{\approx}{r}\, h_n^{(1)}(kr)\, P_n^{\,1}(\cos\theta) \sin\phi \right] \right\} \qquad (5\text{-}66)$$

where the Riccati Bessel functions are defined by

$$\psi_n(x) = x\, j_n(x) = \sqrt{\frac{\pi x}{2}}\; J_{n+\frac{1}{2}}(x) \qquad (5\text{-}67a)$$

$$\zeta_n^{(1)}(x) = x\, h_n(x) = \sqrt{\frac{\pi x}{2}}\; H^{(1)}_{n+\frac{1}{2}}(x) \qquad (5\text{-}67b)$$

When Eq. (5-66) is used in Eq. (5-64), one finds that U(t) can be evaluated in the form of a series of residues derived from the zeros k_s of the functions

$$\zeta^{(1)}_{\nu-1/2}(k_s a) = \sqrt{\frac{\pi k_s a}{2}}\; H_\nu^{(1)}(k_s a) = 0 \qquad (5\text{-}68a)$$

$$\zeta^{(1)'}_{\nu-1/2}(k_s a) = \sqrt{\frac{\pi k_s a}{2}} \left[H_\nu^{(1)'}(k_s a) + \frac{1}{2k_s a}\, H^{(1)}_{\nu_s}(k_s a) \right] = 0 \qquad (5\text{-}68b)$$

The roots k_s of Eq. (5-68a) depend upon the coefficients $B_n(\tau_s^\infty, \infty)$ given in Eq. (5-63) The roots k_s of Eq. (5-68b) depend upon the coefficients $B_n(\tau_s^0, 1/4)$ which can be derived from Eq. (5-61). The explicit form for the solution of Eq. (5-68b) is

$$k_s a = \nu - \left(\frac{\nu}{2}\right)^{1/3} \tau + \left(\frac{2}{\nu}\right)^{1/3} \left(\frac{3}{20}\tau^2 - \frac{3}{20}\frac{1}{\tau}\right) + \left(\frac{2}{\nu}\right)^{3/3} \left(\frac{\tau^3}{1400} + \frac{1}{200} + \frac{9}{800}\frac{1}{\tau^3}\right)$$

$$+ \left(\frac{2}{\nu}\right)^{5/3} \left(-\frac{479}{504000}\tau^4 + \frac{1457}{252000}\tau - \frac{9}{8000}\frac{1}{\tau^2} - \frac{27}{16000}\frac{1}{\tau^5}\right)$$

$$+ \left(\frac{2}{\nu}\right)^{7/3} \left(-\frac{20231}{129360000}\tau^5 + \frac{3287}{49896000}\tau^2 - \frac{12801}{22176000}\frac{1}{\tau} + \frac{4567}{1920000}\frac{1}{\tau^4}\right.$$

$$\left. + \frac{19605}{1280000}\frac{1}{\tau^7}\right) + \cdots \qquad (5\text{-}69)$$

In acoustic diffraction problems involving a rigid sphere of radius

a, we find that the secondary field is of the form

$$u_s(k) = -\sum_{n=0}^{\infty} (2n+1)(-i)^n \frac{j_n'(ka)}{h_n^{(1)'}(ka)} h_n^{(1)}(kr) P_n(\cos\theta) \qquad (5\text{-}70)$$

where the spherical Bessel functions are defined in Eq. (5-67). If we seek to study a pulse diffraction problem for this case we find that k_s is defined by

$$h_n^{(1)'}(k_s a) = \sqrt{\frac{\pi}{k_s a}} \left[H_\nu^{(1)'}(k_s a) - \frac{1}{2k_s a} H_\nu^{(1)}(k_s a) \right] = 0 \qquad (5\text{-}71)$$

We observe that this case requires the coefficients $B_n(\tau_s^0, -1/4)$ which can be derived from Eq. (5-61). The explicit form for the solution for this case is

$$\begin{aligned} k_s a = \nu &- \left(\frac{\nu}{2}\right)^{1/3} \tau + \left(\frac{2}{\nu}\right)^{1/3} \left(\frac{3}{20}\tau^2 + \frac{7}{20}\frac{1}{\tau}\right) + \left(\frac{2}{\nu}\right)^{3/3} \left(\frac{\tau^2}{1400} - \frac{9}{200} + \frac{49}{800}\frac{1}{\tau^3}\right) \\ &+ \left(\frac{2}{\nu}\right)^{5/3} \left(-\frac{479}{504000}\tau^4 - \frac{499}{36000}\tau + \frac{343}{24000}\tau^2 + \frac{343}{16000}\frac{1}{\tau^5}\right) \\ &+ \left(\frac{2}{\nu}\right)^{7/3} \left(-\frac{20231}{129360000}\tau^5 - \frac{86803}{49896000}\tau^2 + \frac{90907}{22176000}\frac{1}{\tau}\right. \\ &\left. + \frac{12807}{1920000}\frac{1}{\tau^4}\right) + \cdots \end{aligned} \qquad (5\text{-}72)$$

Similar asymptotic expansions for the characteristic frequencies k_s can be found from Eq. (5-52) for diffraction problems involving circular cylinders, elliptic cylinders, parabolic cylinders, ellipsoids of revolution, and paraboloids of revolution.

Before we conclude this discussion, we should emphasize the fact that these results are useful also in the extension of Olver's studies (Ref. 7) of the zeros of Bessel functions. For example, if we replace Eq. (5-59) by

$$J_\nu'(x_s) + \left(\frac{2}{x_s}\right) Z J_\nu(x_s) = 0 \qquad (5\text{-}73)$$

we can use Eq. (5-60) and Eq. (5-61) to Solve for x_s by letting $\tau_s = \beta_s$, where $Ai'(-\beta_s) = 0$. If we consider the Neumann functions

$Y_\nu(x)$ [sometimes denoted by $N_\nu(x)$] and seek to solve for x_s defined by

$$Y_\nu'(x_s) + \left(\frac{2}{x_s}\right) Z\, Y_\nu(x_s) = 0 \tag{5-74}$$

we again use Eq. (5-60) and Eq. (5-61), but now we must let

$$\tau = \delta_s \tag{5-75}$$

where δ_s denotes the roots of

$$Bi'(-\delta_s) = 0 \tag{5-76}$$

Tables of these roots have been prepared by Miller (Ref. 9) and Olver (Ref. 7).

5.6 The Normalization Integral

Let us now see what representations we can obtain for the normalization integral

$$N_s = N(\lambda_s) = \int_0^\infty \left[F(y, \lambda_s)\right]^2 dy \tag{5-77}$$

which was defined in Eq. (4-20). For complex λ, the function $F(y,\lambda)$ will increase as y increases and the integral of the square of $F(y,\lambda)$ taken over y from 0 to ∞ will diverge. However, for certain assumptions on $f(y)$ in the complex plane, $F(y,\lambda)$ will behave like $w_{1,2}(t - k^{\frac{2}{3}} f_1^{\frac{1}{3}} y)$ for complex y. Under these conditions, the integral will converge on the ray $y = y \exp(\pm\frac{1}{3}\pi)$. Therefore, we can express the normalization integral in the form

$$N_s = \int_0^{\infty \exp(\pm i\pi/3)} [F(y, \lambda)]^2 dy \tag{5-78}$$

where the $\pm$ sign is used with $\exp(\mp ikct)$ time dependance (i.e., with $w_{1,2}(t - u)$. However, if we assume that k has a small imaginary part (for example, Im $k > 0$ for $\exp(-ikct)$ time dependance), then we can continue to use Eq. (5-77).

The techniques for the evaluation of N_s are well known. The classic method consists of considering two solutions $F(y,\lambda_i)$ and $F(y,\lambda_j)$ which are associated with different eigenvalues λ_s. We then assume that $i \neq j$ and write

$$\int_0^\infty \frac{d}{dy}\left[F(y,\lambda_i)\frac{d}{dy}F(y,\lambda_j) - F(y,\lambda_j)\frac{d}{dy}F(y,\lambda_i)\right]dy$$

$$= \left[F(y,\lambda_i)\frac{d}{dy}F(y,\lambda_j) - F(y,\lambda_j)\frac{d}{dy}F(y,\lambda_i)\right]_0^\infty$$

$$= \int_0^\infty \left[F(y,\lambda_i)\frac{d^2}{dy^2}F(y,\lambda_j) - F(y,\lambda_j)\frac{d^2}{dy^2}F(y,\lambda_i)\right]dy$$

$$= (\lambda_j - \lambda_i)\int_0^\infty F(y,\lambda_i)\,F(y,\lambda_j)\,dy \qquad (5\text{-}79)$$

We assume that the behavior of $f(y)$ is such that the evaluation at $y = \infty$ can be replaced by zero. We then have

$$(\lambda_i - \lambda_j)\int_0^\infty F(y,\lambda_i)\,F(y,\lambda_j)\,dy = F(0,\lambda_i)\frac{dF(0,\lambda_j)}{dy} - F(0,\lambda_j)\frac{dF(0,\lambda_i)}{dy} \qquad (5\text{-}80)$$

Since we assume that the boundary condition on $F(y,\lambda_s)$ is of the form given in Eq. (5-1c), namely

$$\frac{dF(0,\lambda)}{dy} + k^{2/3} f_1^{1/3} q\, F(0,\lambda) = 0$$

the right side of Eq. (5-80) is equal to zero and hence we have

$$\int_0^\infty F(y,\lambda_i)\,F(y,\lambda_j)\,dy = 0 \qquad (5\text{-}81)$$

provided $i \neq j$.

For $i = j$ we can use L'Hospital's rule and arrive at

$$N_s = \int_0^\infty \left[F(y, \lambda_s)\right]^2 dy = \left[\frac{\partial F(0, \lambda_i)}{\partial \lambda i} \frac{\partial F(0, \lambda_j)}{\partial y} - F(0, \lambda_j) \frac{\partial^2 F(0, \lambda_i)}{\partial \lambda_i \partial y}\right]_{\lambda_i = \lambda_j = \lambda_s}$$

$$= \frac{\partial F(0, \lambda_s)}{\partial \lambda_s} \frac{\partial F(0, \lambda_s)}{\partial y} - F(0, \lambda_s) \frac{\partial^2 F(0, \lambda_s)}{\partial \lambda_s \partial y} \qquad (5\text{-}82)$$

If we recall that we have defined

$$F(y, \lambda_s) = G(u, t_s)$$

$$y = \beta u = (k^{2/3} f_1^{1/3})^{-1} u$$

$$\lambda_s = k^2 f_o + k^{4/3} f_1^{2/3} t_s$$

we can also express Eq. (5-82) in the form

$$N_s = \left[k^{2/3} f_1^{1/3}\right]^{-1} \int_0^\infty \left[G(u, t_s)\right]^2 du \qquad (5\text{-}83)$$

where

$$\int_0^\infty \left[G(u, t_s)\right]^2 du = \frac{\partial G(0, t_s)}{\partial t_s} \frac{\partial G(0, t_s)}{\partial y} - G(0, t_s) \frac{\partial^2 G(0, t_s)}{\partial t_s \partial y} \qquad (5\text{-}84)$$

Fock (Ref.10) has shown how to arrive at the result given in Eq. (5-82) without the use of a result of the form of Eq. (5-80). Let us observe that from

$$\frac{d^2 F}{dy^2} + \left[-\lambda_s + k^2 f(y)\right] F = 0 \qquad (5\text{-}85)$$

it follows that

$$\frac{\partial^2}{\partial y^2}\left(\frac{\partial F}{\partial \lambda_s}\right) + \left[-\lambda_s + k^2 f(y)\right] \frac{\partial F}{\partial \lambda_s} = F \qquad (5\text{-}86)$$

If we multiply Eq. (5-85) by $\partial F/\partial \lambda_s$ and multiply Eq. (5-86) by F, we can form the difference of the two results and show that

$$F^2(y, \lambda_s) = F(y, \lambda_s) \frac{\partial^2}{\partial y^2}\left(\frac{\partial F(y, \lambda_s)}{\partial \lambda_s}\right) - \frac{\partial F(y, \lambda_s)}{\partial \lambda_s} \frac{\partial^2 F(y, \lambda_s)}{\partial y^2}$$

$$= \frac{\partial}{\partial y}\left[F(y, \lambda_s) \frac{\partial^2 F(y, \lambda_s)}{\partial y \, \partial \lambda_s} - \frac{\partial F(y, \lambda_s)}{\partial \lambda_s} \frac{\partial F(y, \lambda_s)}{\partial y}\right] \qquad (5\text{-}87)$$

The result given in Eq. (5-82) follows immediately from Eq. (5-87).

Fock (Ref. 10) also showed that Eq. (5-82) could be expressed in the form

$$N_s = - F^2(0, \lambda_s) \frac{\partial}{\partial \lambda_s} \left\{\frac{\partial F(0, \lambda_s)}{\partial y} \Big/ F(o, \lambda_s)\right\} \qquad (5\text{-}88)$$

From Eq. (5-1c) we find that the logarithmic derivative in Eq. (5-88) is equal to $-k^{\frac{2}{3}} f_1^{\frac{1}{3}} q$. Therefore, we can write

$$N_s = k^{2/3} f_1^{1/3} F^2(0, \lambda_s) \left[\frac{\partial \lambda_s}{\partial q}\right]^{-1} \qquad (5\text{-}89)$$

From Eq. (5-2) and the property

$$\frac{d\tau}{dq} = \frac{1}{\tau - q^2} \qquad (5\text{-}90)$$

we find that

$$\frac{\partial \lambda_s}{\partial q} = \frac{k^{4/3} f_1^{2/3}}{\tau_s - q^2}\left[1 + \sum_{n=1}^{\infty} \frac{\partial D_n(\tau_s, q)}{\partial \tau_s} \beta^n\right] \qquad (5\text{-}91)$$

This result is most useful when y and y_0 are zero. For this case we can express Eq. (4-13), i.e.,

$$U(x, y; x_0, y_0) = \sum_{s=1}^{\infty} \frac{\Psi^+(x_>, \lambda_s) \Psi^-(x_<, \lambda_s)}{W_s} \frac{F(y, \lambda_s) F(y_0, \lambda_s)}{N_s} \qquad (5\text{-}92)$$

$$U(x,0;x_0,0) = \sum_{s=1}^{\infty} \frac{\Psi^+(x_>, \lambda_s)\,\Psi^-(x_<, \lambda_s)}{k^{\frac{2}{3}} f_1^{\frac{1}{3}}\, W_s(\tau_s - q^2)} \left[1 + \sum_{n=1}^{\infty} \frac{\partial D_n(\tau_s, q)}{\partial \tau_s}\, \beta^n\right] \qquad (5\text{-}93)$$

We can obtain another useful expression for N_s by using Eq. (4-18), i.e.,

$$\left[\frac{\partial F(y, \lambda_s)}{\partial y} + i k Z F(y, \lambda_s)\right]_{y=0} = 0 \qquad (5\text{-}94)$$

to write

$$N_s = - F_1(0, \lambda_s) \frac{\partial}{\partial \lambda_s} \left[\frac{\partial F(0, \lambda_s)}{\partial y} + i k Z F(o, \lambda_s)\right] \qquad (5\text{-}95)$$

If we express $F(y, \lambda_s)$ in the form

$$F(y, \lambda_s) = F_1(y, \lambda_s) = G(u, t_s) = H(u, t_s) = \left(\frac{d\eta}{du}\right)^{-1/2} w_1(-\eta) \qquad (5\text{-}96)$$

we find that by defining

$$F_2(y, \lambda_s) = \left(\frac{d\eta}{du}\right)^{-1/2} w_2(-\eta) \qquad (5\text{-}97)$$

and use of the Wronskian relation

$$w_1(-\eta)\, w_2'(-\eta) - w_1'(-\eta)\, w_2(-\eta) = 2i \qquad (5\text{-}98)$$

that we have

$$F_1(y, \lambda) \frac{dF_2(y, \lambda)}{dy} - F_2(y, \lambda) \frac{dF_1(y, \lambda)}{dy} = - 2i \frac{du}{dy} = - 2i\, k^{2/3} f_1^{1/3} \qquad (5\text{-}99)$$

Therefore, if

$$\left[\frac{dF_1}{dy} + i k Z F_1\right]_{y=0} = 0 \qquad (5\text{-}100)$$

it follows that

$$\frac{dF_2(0, \lambda_s)}{dy} + i k Z F_2(0, \lambda_s) = - \frac{2i k^{2/3} f_1^{1/3}}{F_1(0, \lambda_s)} \qquad (5\text{-}101)$$

If we use Eq. (5-101) in Eq. (5-95), we find that

$$\frac{1}{N_s} = \frac{1}{2i k^{2/3} f_1^{1/3}} \frac{\frac{\partial F_2(0, \lambda_s)}{\partial y} + i k Z F_2(0, \lambda_s)}{\frac{\partial}{\partial \lambda_s}\left[\frac{\partial F_1(0, \lambda_s)}{\partial y} + i k Z F_1(0, \lambda_s)\right]} \qquad (5\text{-}102)$$

If we use Eq. (5-96) and Eq, (5-97) and define

$$q = i \left(\frac{k}{f_1}\right)^{1/3} Z$$

we can express Eq. (5-102) in the form

$$\frac{1}{N_s} = \frac{1}{2i k^{2/3} f_1^{1/3}} \frac{- \left(\frac{d\eta}{du}\right)^{1/2} \left[w_2'(-\eta) + \frac{1}{2} \frac{\dot{\eta}}{\eta^2} w_2(-\eta) - \frac{q}{\eta} w_2(-\eta)\right]}{\frac{\partial}{\partial \lambda_s}\left\{- \left(\frac{d\eta}{du}\right)^{1/2} \left[w_1'(-\eta) + \frac{1}{2} \frac{\dot{\eta}}{\eta^2} w_1(-\eta) - \frac{q}{\eta} w_1(-\eta)\right]\right\}} \qquad (5\text{-}103)$$

where the dots denote differentiation with respect to u. From Eq. (5-100), we have

$$w_1'(-\eta) + \frac{1}{2} \frac{\dot{\eta}}{\eta^2} w_1(-\eta) - \frac{q}{\eta} w_1(-\eta) = 0 \qquad (5\text{-}104)$$

This property permits us to express Eq. (5-103) in the form

$$\frac{1}{N_s} = \frac{1}{2i k^{2/3} f_1^{1/3}} \frac{w_2'(-\eta) + \frac{1}{2} \frac{\dot{\eta}}{\eta^2} w_2(-\eta) - \frac{8}{\eta} w_2(-\eta)}{\frac{\partial}{\partial \lambda_s}\left[w_1'(-\eta) + \frac{1}{2} \frac{\dot{\eta}}{\eta^2} w_1(-\eta) - \frac{8}{\eta} w_1(-\eta)\right]} \qquad (5\text{-}105)$$

From the Wronskian property expressed by Eq. (5-98) we see that for the cases of $q = 0$ and $q = \infty$ that the reciprocal of N_s can be expressed in the elegant forms

$$\left[\frac{1}{N_s}\right]_{q=0} = \frac{1}{k^{2/3} f_1^{1/3}} \left[\frac{1}{\eta\left[w_1(-\eta)\right]^2} \frac{\partial \lambda_s}{\partial \eta}\right]_{\eta = -\tau_s^o}$$

$$= -\frac{1}{k^{2/3} f_1^{1/3}} \frac{1}{\tau_s^o\left[w_1(\tau_s^o)\right]^2} \frac{\partial \lambda_s}{\partial \tau_s^o}$$

$$= -\frac{k^{2/3} f_1^{1/3}}{\tau_s^o\left[w_1(\tau_s^o)\right]^2} \frac{\partial \lambda_s}{\partial \tau_s^o} \qquad (5\text{-}106a)$$

and

$$\left[\frac{1}{N_s}\right]_{q=\infty} = \frac{1}{k^{2/3} f_1^{1/3}} \left[\frac{1}{\left[w'(-\eta)\right]^2} \frac{\partial \lambda_s}{\partial \eta}\right]_{\eta = -\tau_s^\infty}$$

$$= \frac{1}{k^{2/3} f_1^{1/3}} \frac{1}{\left[w_1'(\tau_s^\infty)\right]^2} \frac{\partial \lambda_s}{\partial \tau_s^\infty} \qquad (5\text{-}106b)$$

We can evaluate $w_1(\tau_s^o)$ and $w_1'(\tau_s^\infty)$ by using the values of $Ai(-\beta_s)$ and $Ai'(-\alpha_s)$ given in Tables 2-1 and 2-2. The relations which we need are

$$w_1'(\tau_s^\infty) = -2\sqrt{\pi} \exp(-i\pi/6)\, Ai'(-\alpha_s) \qquad (5\text{-}107a)$$

$$w_1(\tau_s^o) = 2\sqrt{\pi} \exp(i\pi/6)\, Ai(-\beta_s) \qquad (5\text{-}107b)$$

$$\tau_s^\infty = \alpha_s \exp(i\pi/3) \qquad (5\text{-}107c)$$

$$\tau_s^o = \beta_s \exp(i\pi/3) \qquad (5\text{-}107d)$$

We can then write

$$\left[\frac{1}{N_s}\right]_{q=0} = \frac{1}{k^{2/3} f_1^{1/3}} \frac{1}{4\pi} \frac{1}{\beta_s\left[Ai(-\beta_s)\right]^2} \frac{\partial \lambda_s}{\partial \beta_s} \qquad (5\text{-}108a)$$

$$\left[\frac{1}{N_s}\right]_{q=\infty} = \frac{1}{k^{2/3} f_1^{1/3}} \frac{1}{4\pi} \frac{1}{\left[Ai'(-\alpha_s)\right]^2} \frac{\partial \lambda_s}{\partial \alpha_s} \qquad (5\text{-}108b)$$

Olver (Ref. 11) has shown that the roots α_s and β_s of the Airy functions possess an interesting and a useful property. Since s takes on only integral values, we will have to give a meaning to the Olver relations

$$\frac{d\beta_s}{ds} = \frac{1}{\beta_s\left[Ai(-\beta_s)\right]^2} \tag{5-109a}$$

$$\frac{d\alpha_s}{ds} = \frac{1}{\left[Ai'(-\alpha_s)\right]^2} \tag{5-109b}$$

Let $\alpha(s)$ be a continuous function of s, and let the Airy function have $\alpha(s)$ as its argument in the following sense

$$y(s) = Ai[-\alpha(s)]$$

Then the property displayed as Eq. (5-109b) is to be interpreted in the sense

$$\left[\frac{d\alpha(s)}{ds}\right]_{s=n} = \frac{1}{\{[-dy/d\alpha]^2\}_{s=n}} = \frac{1}{\{Ai'[-\alpha(n)]\}^2}$$

where n is an integer.

The Olver relations permit us to express both Eqs. (5-106a) and (5-106b) in the single relation which includes both the case $q = 0$ and the case $q = \infty$.

$$\frac{1}{N_s} = \frac{1}{4\pi\, k^{2/3} f_1^{1/3}} \frac{\partial \lambda_s}{\partial s} \tag{5-110}$$

The Olver relation can be generalized to include the case of the roots τ_s defined by

$$w_1'(\tau_s) - q\, w_1(\tau_s) = 0$$

It can be shown that the generalized Olver relation for these roots is of the form

$$\frac{\partial \tau_s}{\partial s} = \frac{4\pi}{\left[w_1'(\tau_s)\right]^2 - \tau_s\left[w_1(\tau_s)\right]^2} \tag{5-111}$$

If we use Eq. (5-111) in the representation

$$\frac{1}{N_s} = \frac{1}{k^{2/3} f_1^{1/3}} \frac{1}{\left[w_1'(\tau_s)\right]^2 - \tau_s\left[w_1(\tau_s)\right]^2} \frac{\partial \lambda_s}{\partial \tau_s} \tag{5-112}$$

we see that Eq. (5-110) holds for arbitrary values of q.

If we use Eq. (5-2) along with Eq, (5-112), we can express Eq. (5-92) in the form

$$U(x, y; x_o, y_o) = k^{2/3} f_1^{1/3} \sum_{s=1}^{\infty} \left\{ \frac{\Psi^+(x_>, \lambda_s)\, \Psi^-(x_<, \lambda_s)}{W_s} \right.$$

$$\left. \frac{F(y, \lambda_s)\, F(y_o, \lambda_s)}{\left[w'(\tau_s)\right]^2 - \tau_s\left[w(\tau_s)\right]^2} \left[1 + \sum_{n=1}^{\infty} \frac{\partial D_n(\tau_s, q)}{\partial \tau_s} \beta^n \right] \right\} \tag{5-113}$$

where $F(y,\lambda_s)$ is assumed to be defined as in Eq. (5-96)

Eq. (5-110) can be used to express $U(x,y,;x_0,y_0)$ in the form

$$U(x, y; x_o, y_o) = \frac{1}{4\pi k^{2/3} f_1^{1/3}} \sum_{s=1}^{\infty} \left\{ \frac{\Psi^+(x_>, \lambda_s)\, \Psi^-(x_<, \lambda_s)}{W_s} \right.$$

$$\left. F(y, \lambda_s)\, F(y_o, \lambda_s) \frac{\partial \lambda_s}{\partial s} \right\} \tag{5-114}$$

This form is very convenient for use with the Euler-Maclaurin summation formula

$$\left(\sum_{s=N}^{\infty} - \int_{s=N}^{\infty} ds \right) f(s) = \frac{1}{2} f(N) - \frac{1}{12} \frac{d}{dN} f(N) + \frac{1}{720} \frac{d^3}{dN^3} f(N) - \frac{1}{30240} \frac{d^5}{dN^5} f(N)$$

$$+ \cdots (-1)^n \frac{B_n}{(2n)!} \frac{d^{2n-1}}{dN^{2n-1}} f(N) + \cdots \tag{5-115}$$

where B_n denotes the Bernouli numbers

$$B_1 = \frac{1}{6}\ ,\ B_2 = \frac{1}{30}\ ,\ B_3 = \frac{1}{42}\ ,\ B_4 = \frac{1}{30}\ ,\ B_5 = \frac{5}{66}\ ,\ B_6 = \frac{691}{2730}\ ,\ B_7 = \frac{7}{6},\ \ldots$$

or the Gregory formula of numerical integration

$$\left(\sum_{s=N}^{\infty} - \int_{s=N}^{\infty} ds\right) f(s) = \frac{1}{2} f(N) - \frac{1}{12}\,\Delta f(N) + \frac{1}{24}\,\Delta^2 f(N) - \frac{19}{720}\,\Delta^3 f(N)$$
$$+ \frac{3}{160}\,\Delta^4 f(N) - \frac{863}{60480}\,\Delta^5 f(N) + \frac{275}{24192}\,\Delta^6 f(N)$$
$$- \frac{33953}{3628800}\,\Delta^7 f(N) + \frac{8183}{1036800}\,\Delta^8 f(N)$$
$$- \frac{3250433}{479001600}\,\Delta^9 f(N) + \frac{4671}{788480}\,\Delta^{10} f(N)$$
$$- \frac{13695779093}{2615348736000}\,\Delta^{11} f(N) + \cdots \qquad (5\text{-}116)$$

where the foward differences are defined by

$$\Delta f(N) = f(N+1) - f(N)$$
$$\Delta^2 f(N) = f(N+2) - 2f(N+1) + f(N)$$
$$\Delta^{n+1} f(N) = \Delta^n f(N+1) - \Delta^n f(N)$$

Either of Eqs. (5-115) or (5-116) can be used to express the wave function $U(x,y;x_0,y_0)$ in the form

$$U(x,y;x_0,y_0) = \frac{1}{4\pi k^{2/3} f_1^{1/3}} \int_{k^2 f_0}^{\infty} \frac{\Psi^+(x_>,\lambda)\,\Psi^-(x_<,\lambda)}{W(\lambda)}\, F(y,\lambda)\, F(y_0,\lambda)\, d\lambda$$
$$- \frac{1}{4\pi k^{2/3} f_1^{1/3}} \int_{k^2 f_0}^{\lambda_N} \frac{\Psi^+(x_>,\lambda)\,\Psi^-(x_<,\lambda)}{W(\lambda)}\, F(y,\lambda)\, F(y_0,\lambda)\, d\lambda$$
$$+ \frac{1}{4\pi k^{2/3} f_1^{1/3}} \sum_{s=1}^{N-1} \frac{\Psi^+(x_>,\lambda_s)\,\Psi^-(x_<,\lambda_s)}{W_s}\, F(y,\lambda_s)\, F(y_0,\lambda_s) \frac{d\lambda_s}{ds}$$
$$+ \frac{1}{4\pi k^{2/3} f_1^{1/3}} \left(\sum_{s=N}^{\infty} - \int_{s=N}^{\infty} ds\right) \frac{\Psi^+(x_>,\lambda_s)\,\Psi^-(x_<,\lambda_s)}{W_s}\, F(y,\lambda_s)\, F(y_0,\lambda_s)\, \frac{d\lambda_s}{ds} \qquad (5\text{-}117)$$

The limits on the integral of the form

$$\int_{k^2 f_o}^{\infty} \left[\cdots \right] d\lambda$$

are used as a shorthand way of specifying the path of integration. The limit ∞ which appears as the upper limit on this integral is to be interpreted in the sense

$$\int_{k^2 f_o}^{\infty} \left[\cdots \right] d\lambda = \lim_{s \to \infty} \int_{k^2 f_o}^{\lambda_s} \left[\cdots \right] d\lambda = \int_{c} \left[\cdots \right] d\lambda$$

where $\underline{c}$ denotes a contour which starts at $\lambda = k^2 f_0$ and goes to infinity along the root locus along which one finds the eigenvalues λ_s. With an $\exp(-i\omega t)$ time dependance this will usually be a contour which starts at a point $\lambda = k^2 f_0$ on the positive real axis and then passes upward into the first quadrant and bends so as to pass into the second quadrant and go off to infinity (with the imaginary part of $\lambda > 0$) along a locus such that the angle between a point on the locus and the negative real λ-axis tends to zero.

The only case of such a locus which has been extensively studied is that of the Hankel functions. Magnus and Kotin (Ref. 12) have shown that for the roots ν_s defined by

$$H_{\nu_s}^{(1)}(z) = H_{\nu_s}^{(1)}(re^{i\phi}) = 0$$

that

$$\nu_s = \alpha_s + i\beta_s$$

where

$$\alpha_s = \pi \left(\frac{\pi}{2} - \phi\right) \left(s + \frac{1}{4}\right) \left[\log \frac{(s + 1/4)\,\pi}{er}\right]^{-2}$$

$$\beta_s = \pi \left(s + 1/4\right) \left[\log \frac{(s + 1/4)\,\pi}{er}\right]^{-1}$$

Let us now make some remarks concerning the integral

$$U_f(x, y; x_o, y_o) = \frac{1}{4\pi k^{2/3} f_1^{1/3}} \int_{k^2 f_o}^{\infty} \frac{\Psi^+(x_>, \lambda)\, \Psi^-(x_<, \lambda)}{W(\lambda)} F(y, \lambda)\, F(y_o, \lambda)\, d\lambda \quad (5\text{-}118)$$

This integral can generally be evaluated by approximating it by a Fresnel integral when

$$\delta = \left\{ \int_{x_<}^{x_>} \frac{du}{\sqrt{g(u) + f_o}} - \left(\int_0^{y} + \int_0^{y_o} \right) \frac{dv}{\sqrt{f(v) - f_o}} \right\} > 0 \quad (5\text{-}119)$$

For $\delta < 0$, we can generally write

$$U_f(x, y; x_o, y_o) = \frac{1}{4\pi k^{2/3} f_1^{1/3}} \int_{-\infty}^{\infty} [\cdots]\, d\lambda - \frac{1}{4\pi k^{2/3} f_1^{1/3}} \int_{-\infty}^{k^2 f_o} [\cdots] d\lambda \quad (5\text{-}120)$$

where the integral

$$\int_{-\infty}^{\infty} [\cdots] d\lambda$$

can be evaluated by the method of stationary phase and the integral

$$\int_{-\infty}^{k^2 f_o} [\cdots]\, d\lambda$$

can be approximated by a Fresnel integral.

The stationary phase evaluation of the integral taken between the limits $-\infty$ and $+\infty$ generally yields the "free space field" which is present when there is no reflecting surface at $y = 0$.

Before leaving the subject of these integral representations for the wave function $U(x,y;x_0,y_0)$, we should remark that it is also possible to express this function in the form of a Fourier integral

$$U(x, y; x_o, y_o) = \frac{1}{4\pi k^{2/3} f_1^{1/3}} \int_{-\infty}^{\infty} \frac{\Psi^+(x_>, \lambda)\, \Psi^-(x_<, \lambda)}{W(\lambda)} \left[F_1(y_>, \lambda)\, F_2(y_<, \lambda) - \frac{\frac{\partial F_2(0,\lambda)}{\partial y} + ikZ\, F_2(0,\lambda)}{\frac{\partial F_1(0,\lambda)}{\partial y} + ikZ\, F_1(0,\lambda)} F_1(y, \lambda)\, F_1(y_o, \lambda) \right] d\lambda \qquad (5\text{-}121)$$

If we evaluate this expression in the form of the residue series given in Eq. (5-92) we can readily verify the result given in Eq. (5-102).

5.7 Direct Integration of N_s

In our previous discussions, we arrived at some elegant representations for the normalization integral. Let us now consider an alternative procedure which is more direct and which contains some interesting analysis.

We want to evaluate the integral

$$\int_0^{\infty} \left[G(u, t_s) \right]^2 du = k^{2/3} f_1^{1/3} \int_0^{\infty} \left[F(y, \lambda_s) \right]^2 dy \qquad (5\text{-}122)$$

where $G(u, t_s)$ can be obtained from Eq. 4-148). Thus

$$G(u, t) = w(t - u) - \frac{\beta h_2}{15} \left[(3u + 2t)\, w(t - u) + (3u^2 + 4ut + 8t^2)\, w'(t - u) \right] \qquad (5\text{-}123)$$

Let us define

$$\zeta = t - u \quad , \quad d\zeta = - du$$

and observe that

$$\int_0^{\infty} \left[G(u, t) \right]^2 du = \int_{-\infty}^{t} w^2(\zeta)\, d\zeta - \frac{2\beta h_2}{15} \int_{-\infty}^{t} \left\{ (5t - 3\zeta)\, w^2(\xi) + (15t^2 - 10\, \zeta t + 3\zeta^2)\, w(\zeta)\, w'(\zeta) \right\} d\zeta \qquad (5\text{-}124)$$

Since

$$\frac{d}{d\zeta}\left\{\zeta w^2(\zeta) - w'^2(\zeta)\right\} = w^2(\zeta) + 2\zeta w' w - 2\zeta\, w\, w' = w^2(\zeta) \tag{5-125a}$$

$$\frac{d}{d\zeta}\left\{\frac{1}{3}\zeta^2 w^2 - \frac{1}{3}\zeta\, w'^2(\zeta) + \frac{1}{3} w\, w'\right\} = \frac{2\zeta}{3} w^2 + \frac{2\zeta^2}{3} w\, w' - \frac{1}{3} w'^2 - \frac{2}{3}\zeta^2 w\, w' + \frac{1}{3} w'^2 + \frac{1}{3}\zeta\, w^2 = \zeta\, w^2(\zeta) \tag{5-125b}$$

and

$$\frac{d}{d\zeta}\left\{\frac{1}{6}\zeta^2 w^2(\zeta) + \frac{1}{3}\zeta\, w'^2(\zeta) - \frac{1}{3} w(\zeta)\, w'(\zeta)\right\} = \frac{1}{3}\zeta w^2(\zeta) + \frac{1}{3}\zeta^2 w\, w' + \frac{1}{3} w'^2 + \frac{2}{3}\zeta^2 w\, w' - \frac{1}{3} w'^2 - \frac{1}{3}\zeta\, w^2 = \zeta^2 w(\zeta)\, w'(\zeta) \tag{5-125c}$$

we have the following integrals

$$\int w^2(\zeta)\, d\zeta = \zeta\, w^2(\zeta) - w'^2(\zeta) \tag{5-126a}$$

$$\int \zeta\, w^2(\zeta)\, d\zeta = \frac{1}{3}\zeta^2 w^2(\zeta) - \frac{1}{3}\zeta\, w'^2(\zeta) + \frac{1}{3} w(\zeta)\, w'(\zeta) \tag{5-126b}$$

$$\int w'(\zeta)\, w(\zeta)\, d\zeta = \frac{1}{2} w^2(\zeta) \tag{5-126c}$$

$$\int \zeta^2 w(\zeta)\, w'(\zeta)\, d\zeta = \frac{1}{6}\zeta^2 w^2(\zeta) + \frac{1}{3}\zeta\, w'^2(\zeta) - \frac{1}{3} w(\zeta)\, w'(\zeta) \tag{5-126d}$$

Therefore, we find that

$$\int_0^\infty \left[G(u, t)\right]^2 du = t\, w^2(t) - w'^2(t) - \frac{2\beta h_2}{15}\left\{5t\,(t\, w^2(t) - w'^2(t)) - 3\left[\frac{1}{3} t^2 w^2(t) - \frac{1}{3} t\, w'^2(t) + \frac{1}{3} w(t)\, w'(t)\right]\right. \left.\frac{15}{2} t^2 w^2(t) - 10t\frac{w'^2(t)}{2} + 3\left[\frac{1}{6} t^2 w^2(t) + \frac{1}{3} t\, w'^2(t) - \frac{1}{3} w(t) w'(t)\right]\right\}$$

$$= w^2(t)\,(t - q^2) - \frac{2\beta h_2}{15} w^2(t)\left\{12t^2 - 8tq^2 - 2q\right\} + \cdots \tag{5-127}$$

since

$$w_1'(t) - q\, w_1(t) = 0$$

We can readily verify this result by using Eq. (5-82) to write

$$\int_0^\infty \left[G(u,\ t)\right]^2 du = \left[\frac{\partial G}{\partial t}\frac{\partial G}{\partial u} - G\frac{\partial^2 G}{\partial t \partial u}\right]_{u=0} \tag{5-128}$$

where

$$\left[\frac{\partial G}{\partial t}\ \frac{\partial G}{\partial u}\right]_{u=0} = -w'^2(t) + \frac{\beta h_2}{15}\left[-w\,w' + 16t\,w'^2 + 16\,t^3\,w\,w'\right] \tag{5-129a}$$

$$\left[-G\frac{\partial^2 G}{\partial t \partial u}\right]_{u=0} = t\,w^2(t) + \frac{\beta h_2}{15}\left\{5w\,w' - 24\,t^2 w^2 - 16\,t^3\,w\,w'\right\} \tag{5-129b}$$

It is readily verified that these results confirm Eq. (5-127).

In his paper, Fock (Ref. 10) defined D(t) by means of

$$\int_0^\infty \left[G(u,\ t)\right]^2 du = \left[G(o,\ t)\right]^2 D(t) \tag{5-130}$$

Since

$$\left[G(0,\ t)\right]^2 = \left\{w(t) - \frac{\beta h_2}{15}\left[2t\,w(t) + 8t^2\,w'(t)\right]\right\}^2 = w^2(t)\left\{1 - \frac{\beta h_2}{15}(2t + 8t^2 q)\right\} + \cdots \tag{5-131}$$

we find that

$$D(t) = \frac{(t - q^2) - \frac{2\beta h_2}{15}(12t^2 - 8t\,q^2 - 2q) + \cdots}{1 - \frac{2\beta h_2}{15}(2t + 8t^2\,q) + \cdots}$$

$$= (t - q^2) + \beta h_2\left\{\frac{16}{15}\,t^2 q(t - q^2) - \frac{4}{3}\,t\,(t - q^2) - \frac{8}{15}\,tq^2 + \frac{4}{15}q\right\} \tag{5-132}$$

Since Fock gave D(t) to be [see his Eq. 5.09]

$$D(t) = (t - q^2)\,(1 - 4/3\,\beta h_2 t) + 2/3\,\beta h_2\,q \tag{5-133}$$

we must conclude that his result is in error.

5.8 Eigenvalues Derived from the Langer Approximation

In this section we want to discuss a technique which may be of some value in obtaining more useful numerical techniques for the determination of the eigenvalues. Let us assume that the function G(u,t) which satisfies Eq. (4-77), namely

$$\frac{d^2G}{du^2} + [(-t + u) + \beta h_2 u^2 + \beta^2 h_3 u^3 + \cdots] G = 0 \qquad (5\text{-}134)$$

is to be approximated by an expression of the form

$$G(u, t) \sim \left(\frac{d\eta_o}{du}\right)^{-\frac{1}{2}} w_{1,2}(-\eta_o) \qquad (5\text{-}135)$$

where

$$\tfrac{2}{3}\eta_o^{3/2} = \int_{u_1}^{u} \sqrt{(v - t) + \beta h_2 v^2 + \beta^2 h_3 v^3 + \cdots}\, dv \qquad (5\text{-}136a)$$

for $u > u_1$, and

$$\tfrac{2}{3}(-\eta_o)^{3/2} = \int_{u}^{u_1} \sqrt{(t - v) - \beta h_2 v^2 - \beta h_3 v^3 - \cdots}\, dv \qquad (5\text{-}136b)$$

for $u < u_1$, where u_1 is the turning point defined by

$$t - u_1 - \beta h_2 u_1^2 - \beta^2 h_3 u_1^3 - \cdots = 0 \qquad (5\text{-}137)$$

In Section 6 we shall refer to this type of approximation as the "Langer approximation."

Let τ denote the root of

$$w_{1,2}(\tau) = 0 \qquad (5\text{-}138)$$

Then, in order to solve for the eigenvalue t defined by

$$G(0, t) \sim \left[\left(\frac{d\eta_o}{du}\right)^{-\frac{1}{2}} w_{1,2}(-\eta_o)\right]_{u=0} \qquad (5\text{-}139)$$

we must invert the equation

$$\tfrac{2}{5}\tau^{3/2} = \int_0^{u_1} \sqrt{t - u - \beta h_2 u^2 - \beta^2 h_3 u^3 - \beta^3 h_4 u^4 - \beta^4 h_5 u^5 - \cdots}\; du \qquad (5\text{-}140)$$

and solve for t in terms of τ. In order to carry out these operations, let us define

$$x = t - u - \beta h_2 u^2 - \beta^2 h_3 u^3 + \cdots \qquad (5\text{-}141)$$

and express the integral in the form

$$\tfrac{2}{5}\tau^{3/2} = \int_0^t \sqrt{x}\,(-\frac{du}{dx})\,dx \qquad (5\text{-}142)$$

Since

$$\begin{aligned} u = {} & (t-x) - \beta h_2 (t-x)^2 - \beta^2 (h_3 - 2h_2^2)(t-x)^3 \\ & - \beta^3 (h_4 - 5h_2 h_3 + 5h_2^3)(t-x)^4 \\ & - \beta^4 (h_5 - 6h_2 h_4 - 3h_3^2 + 21h_2^2 h_3 - 14h_2^4)(t-x)^5 + \cdots \end{aligned} \qquad (5\text{-}143)$$

we find that

$$\begin{aligned} -\frac{du}{dx} = {} & 1 - 2\beta h_2 (t-x) - 3\beta^2 (h_3 - 2h_2^2)(t-x)^2 \\ & - 4\beta^3 (h_4 - 5h_2 h_3 + 5h_2^3)(t-x)^3 \\ & - 5\beta^4 (h_5 - 6h_2 h_4 - 3h_3^2 + 21h_2^2 h_3 - 14h_2^4)(t-x)^4 + \cdots \end{aligned} \qquad (5\text{-}144)$$

Therefore, if we use the results

$$\int_0^t \sqrt{x}\,dx = \tfrac{2}{3} t^{3/2}$$

$$\int_0^t \sqrt{x}\,(t-x)\,dx = \frac{4}{15} t^{5/2}$$

$$\int_0^t \sqrt{x}\,(t-x)^2\,dx = \frac{16}{105} t^{7/2}$$

$$\int_0^t \sqrt{x}\,(t-x)^3\,dx = \frac{172}{315} t^{9/2}$$

$$\int_0^t \sqrt{x}\,(t-x)^4\,dx = \frac{256}{3465} t^{11/2}$$

we arrive at

$$\tfrac{2}{3}\tau^{3/2} = \tfrac{2}{3}t^{3/2} - \frac{8}{15}\beta h_2 t^{5/2} - \frac{16}{35}\beta^2 (h_3 - 2h_2^2) t^{7/2} - \frac{128}{315}\beta^3 (h_4 - 5h_2 h_3 + 5h_2^3) t^{9/2} - \frac{256}{693}\beta^4 (h_5 - 6h_2 h_4 - 3h_3^2 + 21h_2^2 h_3 - 14h_2^4) t^{11/2} \quad (5\text{-}145)$$

In order to invert Eq. (5-145), we assume that t can be represented by a series of the form

$$t = \tau + \beta L_1 + \beta^2 L_2 + \beta^3 L_3 + \beta^4 L_4 + \cdots \quad (5\text{-}146)$$

we can then use the binomial theorem to form the following expressions:

$$t^{3/2} = \tau^{3/2}\left\{1 + \beta\left(\frac{3}{2}\frac{L_1}{\tau}\right) + \beta^2\left(\frac{3}{2}\frac{L_2}{\tau} + \frac{3}{8}\frac{L_1^2}{\tau^2}\right) + \beta^3\left(\frac{3}{2}\frac{L_3}{\tau} + \frac{3}{4}\frac{L_1 L_2}{\tau^2} - \frac{1}{16}\frac{L_1^3}{\tau^3}\right) + \beta^4\left(\frac{3}{2}\frac{L_4}{\tau} + \frac{3}{4}\frac{L_1 L_3}{\tau^2} + \frac{3}{8}\frac{L_2^2}{\tau^2} - \frac{3}{16}\frac{L_1^2 L_2}{\tau^3} + \frac{3}{128}\frac{L_1^4}{\tau^4} + \cdots\right\} \quad (5\text{-}147a)$$

$$t^{5/2} = \tau^{5/2}\left\{1 + \beta\left(\frac{5}{2}\frac{L_1}{\tau}\right) + \beta^2\left(\frac{5}{2}\frac{L_2}{\tau} + \frac{15}{8}\frac{L_1^2}{\tau^2}\right) + \beta^3\left(\frac{5}{2}\frac{L_3}{\tau} + \frac{15}{4}\frac{L_1 L_2}{\tau^2} + \frac{5}{16}\frac{L_1^3}{\tau^3}\right) + \cdots\right\} \quad (5\text{-}147b)$$

$$t^{7/2} = \tau^{7/2}\left\{1 + \beta\left(\frac{7}{2}\frac{L_1}{\tau}\right) + \beta^2\left(\frac{7}{2}\frac{L_2}{\tau} + \frac{35}{8}\frac{L_1^2}{\tau^2}\right) + \cdots\right\} \quad (5\text{-}147c)$$

$$t^{9/2} = \tau^{9/2}\left\{1 + \beta\left(\frac{9}{2}\frac{L_1}{\tau}\right) + \cdots\right\} \quad (5\text{-}147d)$$

If we now use Eq. (5-147) in Eq. (5-145), we find that

$$\frac{2}{3}t^{3/2} = \frac{2}{3}\tau^{3/2} + \beta\left(L_1\tau^{1/2} - \frac{8}{15}h_2\tau^{5/2}\right) + \beta^2\left(L_2\tau^{1/2} + \frac{1}{4}L_1{}^2\tau^{-1/2}\right.$$
$$- \frac{4}{3}h_2L_1\tau^{3/2} - \frac{16}{35}h_3\tau^{7/2} + \frac{32}{35}h_2{}^2\tau^{7/2}\Big) + \beta^3\left(L_3\tau^{1/2}\right.$$
$$+ \frac{1}{2}L_1L_2\tau^{-1/2} - \frac{1}{24}L_1{}^3\tau^{-3/2} - \frac{4}{3}h_2L_2\tau^{3/2} - h_2L_1{}^2\tau^{1/2}$$
$$- \frac{8}{5}h_3L_1\tau^{5/2} + \frac{16}{5}h_2{}^2L_1\tau^{5/2} + \frac{688}{63}h_2h_3\tau^{9/2} - \frac{688}{315}h_4\tau^{9/2}$$
$$\left. - \frac{688}{63}h_2{}^3\tau^{9/2}\right) + \cdots \qquad (5\text{-}148)$$

Since the coefficients of β^n, $n \geq 1$, must vanish, we find that

$$L_1 = \frac{8}{15}h_2\tau^2 \qquad (5\text{-}149a)$$

$$L_2 = \left(\frac{16}{35}h_3 - \frac{48}{175}h_2{}^2\right)\tau^3 \qquad (5\text{-}149b)$$

$$L_3 = \left(\frac{128}{315}h_4 - \frac{11968}{17325}h_2h_3 + \frac{22912}{70875}h_2{}^3\right)\tau^4 \qquad (5\text{-}149c)$$

If we compare these value of $L_n(\tau)$ with the values $D_n(\tau)$ for $q = \infty$ in Eq. (5-4) we find that these are <u>precisely</u> the terms which contain the highest power of τ. The same thing is true concerning Eq. (5-7) in which $D_n(\tau)$ are given for $q = 0$. This result suggests a valuable means of extending the computation of λ_s when s (or mod τ) is relatively large. Let T_s denote the solution of Eq. (5-145). Use the expansion

$$T_s = \tau_s + \beta L_1(\tau_s) + \beta^2 L_2(\tau_s) + \beta^3 L_3(\tau_s) + \cdots \qquad (5\text{-}150)$$

to compute an approximate value of T_s. Then use this as an approximation in Eq. (5-145) and use numerical methods to fins a more accurate value of T_s. Then compute λ_s by replacing Eq. (5-2) by the equation

$$\lambda_s = k^2 f_o + k^{4/3} f_1{}^{2/3}\left\{T_s + \sum_{n=1}^{\infty}\left[D_n(\tau_s, q) - L_n(\tau_s)\right]\beta^n\right\} \qquad (5\text{-}151)$$

REFERENCES FOR SECTION 5

1. Air Force Cambridge Research Center, The Zeros and Maxima of the Airy Function and Its First Derivative to 25 Significant Figures, by. M. E. Sherry, Report AFCRC-TR-59-135, ASTIA Document No. AD 214568, Bedford, Mass., April, 1959

2. C. L. Pekeris, "Asymptotic Solutions for the Normal Modes in the Theory of Microwave Propagation," J. Appl. Phys., Vol. 17, 1946, pp. 1108-1124

3. I, Imai, "Asymptotic Solutions of Ordinary Linear Differential Equations of the Second Order," Phys. Rev., Vol. 80, 1950, p. 1112

4. I. Imai, "A Refinement of the WKB Method and Its Application to the Electromagnetic Wave Theory," IRE Trans. on Antennas and Propagation, Vol. AP-4, July, 1956, pp. 233-239

5. H. Bremmer, "On the Theory of Wave Propagation Through a Concentrically Stratified Troposphere with a Smooth Profile. Part I. Discussion of the Extended W.K.B. Approximation," J. Research NBS, Vol. 64D, Sept. -Oct., 1960, pp. 467-482

6. F. W. J. Olver, "A Further Method for the Evaluation of Zeros of Bessel Functions, and Some New Asymptotic Expansions for Zeros of Functions of Large Order," Proc. Cambridge Phil. Soc., Vol. 47, 1951, pp. 699-712

7. F. W. J. Olver, "Definitions, Formulae, and Methods," Bessel Functions, Part III, Zeros and Associated Values, Royal Society Mathematical Tables, Vol. 7, Cambridge, Cambridge University Press, 1960

8. Radiation Laboratory, University of Michigan, Exact Near Field and Far Field Solution for the Back-Scattering of a Pulse from a Perfectly Conducting Sphere, by V. H. Weston, Report RADC TN-59-163, ASTIA Document No. 215486, Contract No. AF 30(602)-1853, Ann Arbor, Mich., 1959

9. J. C. P. Miller, The Airy Integral, British Association Mathematical Tables, Part-Volume B, Cambridge, Cambridge University Press, 1946

10. V. A. Fock, "Propagation of the Direct Wave Around the Earth with Due Account for Diffraction and Propagation," Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965. (This classic paper was first published in Bulletin de l'Academie des Sciences de l'URSS, ser. phys., Vol. 12, No. 2, 1948, pp. 81-97.)

11. F. W. J. Olver, "The Asymptotic Expansion of Bessel Functions of Large Order," Trans. Roy. Soc. (London), Vol. 247A, 1954, pp. 328-368

12. W. Magnus and L. Kotin, "The Zeros of the Hankel Function as a Function of Its Order," Numerische Mathematik, Vol. 2, 1960, pp. 228-244

Section 6

HEIGHT GAIN FUNCTIONS FOR LARGE HEIGHTS

6.1 The Langer Asymptotic Estimate

The representation

$$F(y, \lambda_s) = \left(\frac{d\eta}{du}\right)^{-\frac{1}{2}} w_1(-\eta) \tag{6-1}$$

where η is defined by Eq. (4-118), i.e,

$$\eta = u - t_s + \frac{\beta h_2}{15}\left[3u^2 + 4ut - 8t^2\right] + O(\beta^2) \tag{6-2}$$

is only useful for small values of u. Since

$$y = \frac{1}{k^{2/3} f_1^{1/3}}\, u$$

this requires that we restrict the use of this form of representation for small heights. However, for large heights we can still use Eq. (6-1) if we define η by means of Eq. (4-51). Thus, if we let

$$\frac{2}{3}\zeta^{3/2} = \int_{y_1}^{y} \sqrt{k^2 f(v) - \lambda}\; dv \tag{6-3}$$

we find that

$$\eta = \zeta + \left(\frac{3}{2}\right)^{2/3} \frac{\alpha}{k^{4/3}} + \frac{1}{3}\,\frac{\alpha_0}{k^2}\,\zeta + \frac{1}{5}\left(\frac{2}{3}\right)^{2/3} \frac{\alpha_1}{k^{8/3}}\,\zeta^2 + O\left(k^{-10/3}\right) \tag{6-4}$$

where α_0, α_1, and α_2 are defined in Eqs. (4-37) through (4-39).

The approximation

$$\eta \approx \zeta \tag{6-5}$$

is often called the Langer asymptotic estimate in recognition of Langer's (Ref. 1) research on problems of this type.

The Harvard University Computation Laboratory tables (Ref. 2) can be used to evaluate $w_{1,2}(-\eta)$ for mod $\eta < 6$. These tables give eight decimal values of

$$h_{1,2}(x + iy) = \frac{\sqrt[6]{12}}{\sqrt{\pi}} \exp\left(\mp i \frac{2}{3}\pi\right) w_{1,2}(-x - iy) \qquad (6\text{-}6)$$

and its derivatives for $\Delta x = \Delta y = 0.1$ and $x^2 + y^2 < 36$. The relation

$$w_{1,2}(x + iy) = \frac{\sqrt{\pi}}{\sqrt[6]{12}} \exp\left(\pm i \frac{2}{3}\pi\right) \left[\text{complex conjugate of } h_{2,1}(-x + iy)\right] \qquad (6\text{-}7)$$

is valuable in connection with the use of these tables since the tabulation of $h_{1,2}(x + iy)$ and $h_{1,2}'(x + iy)$ only for $y > 0$.

For mod $\eta >> 1$, we can compute $w_1(-\eta)$ by using the phase and amplitude functions defined and tabluated by Miller (Ref. 3). Thus, we let

$$w_{1,2}(-\eta) = \sqrt{\pi}\, F(-\eta) \exp\left[\pm i \chi(-\eta)\right] \qquad (6\text{-}8a)$$

$$w'_{1,2}(-\eta) = \sqrt{\pi}\, G(-\eta) \exp\left[\pm i \psi(-\eta)\right] \qquad (6\text{-}8b)$$

and use for F, G, χ, and ψ the asymptotic expansions given by Miller.

$$\left|F(-\eta)\right|^2 \sim \frac{1}{\pi\eta^{1/2}}\left(1 - \frac{1\cdot3\cdot5}{1!\ 96}\frac{1}{\eta^3} + \frac{1\cdot3\cdot5\cdot7\cdot9\cdot11}{2!\ 96^2}\frac{1}{\eta^6} - \frac{1\cdot3\cdot5\cdot7\cdot9\cdot11\cdot13\cdot15\cdot17}{3!\ 96^3}\frac{1}{\eta^9} + \cdots\right) \qquad (6\text{-}9a)$$

$$\left|G(-\eta)\right|^2 \sim \frac{1}{\pi}\eta^{1/2}\left(1 + \frac{1\cdot3}{1!\ 96}\frac{7}{\eta^3} - \frac{1\cdot3\cdot5\cdot7\cdot9}{2!\ 96^2}\frac{13}{\eta^6} + \frac{1\cdot3\cdot5\cdot7\cdot9\cdot11\cdot13\cdot15}{3!\ 96^3}\frac{19}{\eta^9} - \cdots\right) \qquad (6\text{-}9b)$$

$$\chi(-\eta) - \frac{1}{4}\pi \sim \frac{2}{3}\eta^{3/2}\left(1 - \frac{5}{32}\frac{1}{\eta^3} + \frac{1105}{6144}\frac{1}{\eta^6} - \frac{82825}{65536}\frac{1}{\eta^9} + \frac{1282031525}{58720256}\frac{1}{\eta^{12}} - \cdots\right) \qquad (6\text{-}9c)$$

$$\psi(-\eta) + \frac{1}{4}\pi \sim \frac{2}{3}\eta^{3/2}\left(1 + \frac{7}{32}\frac{1}{\eta^3} - \frac{1463}{6144}\frac{1}{\eta^6} + \frac{4\,95271}{3\,27680}\frac{1}{\eta^9} - \frac{206530429}{8388608}\frac{1}{\eta^{12}} + \cdots\right) \qquad (6\text{-}9d)$$

The asymptotic expansions given in Eq. (6-9) can be used for complex values of η provided

$$|\arg \eta| < \frac{2\pi}{3}$$

Miller showed that

$$\frac{d\chi(-\eta)}{d\eta} = \frac{1}{\pi F^2(-\eta)} \quad , \quad \frac{d\psi(-\eta)}{d\eta} = \frac{\eta}{\pi G^2(-\eta)} \tag{6-10}$$

We can use these properties to write

$$w_{1,2}(-\eta) = \left(\frac{d\chi}{d\eta}\right)^{-1/2} \exp\left[\pm i\,\chi\right] \tag{6-11}$$

$$w'_{1,2}(-\eta) = \sqrt{\eta}\ \left(\frac{d\psi}{d\eta}\right)^{-1/2} \exp\left[\pm i\psi\right] \tag{6-12}$$

From Eq. (6-1) we deduce the result

$$F_{1,2}(y, \lambda) = \left(\frac{d\chi}{du}\right)^{-1/2} \exp\left[\pm i\chi\right] = k^{1/3} f_1^{1/6} \left(\frac{d\chi}{dy}\right)^{-1/2} \exp\left[\pm i\,\chi\right] \tag{6-13}$$

where $\chi = \chi(-\eta) = \chi\{-\eta[\zeta(y)]\}$ is defined in terms of η by means of Eq. (6-9c) and η is defined in terms of ζ by Eq. (6-4) and ζ is defined in terms of y by Eq. (6-3). Thus, we find that

$$\begin{aligned}\chi\left[-\eta\ (\zeta)\right] - \frac{\pi}{4} \sim &\left[\frac{2}{3}\zeta^{3/2} - \frac{5}{48}\frac{1}{\zeta^{3/2}} + \frac{1105}{9216}\frac{1}{\zeta^{9/2}}\right] + \left(\frac{3}{2}\right)^{2/3}\left[\zeta^{1/2} + \frac{5}{32}\zeta^{-5/2}\right. \\ &\left. - \frac{1105}{2048}\zeta^{-11/2}\right]\frac{\alpha}{k^{4/3}} + \left[\frac{1}{3}\zeta^{3/2} + \frac{5}{96}\zeta^{-3/2} - \frac{1105}{6144}\zeta^{-9/2}\right]\frac{\alpha_o}{k^{6/3}} \\ &+ \left[\left(\frac{1}{5}\zeta^{5/2} + \frac{1}{32}\zeta^{-1/2} - \frac{221}{2048}\zeta^{-7/2}\right)\alpha_1 + \left(\frac{9}{16}\zeta^{-1/2}\right.\right. \\ &\left.\left. - \frac{225}{512}\zeta^{-7/2}\right)\alpha^2\ \left(\frac{2}{3}\right)^{2/3}\frac{1}{k^{8/3}} + \cdots\end{aligned} \tag{6-14}$$

If we use the first term

$$\chi\left[-\eta\ (\zeta)\right] - \frac{\pi}{4} \sim \frac{2}{3}\zeta^{3/2} = \int_{y_1}^{y} \sqrt{k^2 f(v) - \lambda}\ dv \tag{6-15}$$

we find that

$$F_{1,2}(y,\lambda) \sim \frac{k^{1/3} f_1^{1/6} \exp(\pm \frac{\pi}{4})}{\sqrt[4]{k^2 f(y) - \lambda}} \exp\left[\pm i \int_{y_1}^{y} \sqrt{k^2 f(v) - \lambda}\, dv \right] \qquad (6\text{-}16)$$

This is the well-known W.K.B. approximation of the physicists or the Jeffrey's approximation of some of the English applied mathematicians.* Therefore, we see that Eq. (6-14) provides an extension of these classical approximations.

6.2 The W.K.B. Approximation

We do not need to solve first for η before arriving at Eq. (6-14) Since $\exp(\pm i\chi)$ is a solution of

$$\frac{d^2 Z}{d\chi^2} + Z = 0 \qquad (6\text{-}17)$$

and $F(y,\lambda)$ is a solution of

$$\frac{d^2 F}{dy^2} + \left[-\lambda + k^2 f(y) \right] F = 0 \qquad (6\text{-}18)$$

we can use Eq. (4-51) to show that $\chi(y)$ is a solution of

$$(\dot{\chi})^2 - \frac{3}{4}\left(\frac{\ddot{\chi}}{\dot{\chi}}\right)^2 + \frac{1}{2}\frac{\dddot{\chi}}{\dot{\chi}} = k^2 f(y) - \lambda \qquad (6\text{-}19)$$

where the dots denote differentiation with respect to y. From Eq. (6-19) we find that

$$(\dot{\chi})^2 = \left[k^2 f(y) - \lambda \right] - \frac{1}{4}\frac{k^2 \ddot{f}(y)}{\left[k^2 f(y) - \lambda \right]} + \frac{5}{16}\left[\frac{k^2 \dot{f}(y)}{k^2 f(y) - \lambda} \right]^2 + O(k^{-2}) \qquad (6\text{-}20a)$$

or

$$\dot{\chi} = \frac{d\chi}{dy} = \sqrt{k^2 f(y) - \lambda} - \frac{1}{8}\frac{k^2 \ddot{f}(y)}{\left[k^2 f(y) - \lambda \right]^{3/2}} + \frac{5}{32}\frac{k^4 \left[\dot{f}(y) \right]^2}{\left[k^2 f(y) - \lambda \right]^{5/2}} + O(k^{-3}) \qquad (6\text{-}20b)$$

---

*For one of the most recent publications which deals with the W.K.B. approximation, see N. Fröman and P. O. Fröman, JWKB Approximation: Contributions to the Theory, Amsterdam, North-Holland Publishing Co., 1965.

This equation can be integrated by writing

$$\chi(y)=\int_{y_0}^{y} \dot{\chi}(z)dz + \chi(y_0) \tag{6-21}$$

where y_0 is arbitrary. However, for comparison with Eq. (6-14), we should take $y_0 = y_1$, where y_1 is the turning point.

If we reconsider the result

$$F_{1,2}(y,\lambda) = (d\eta/du)^{-1/2} w_{1,2}(-\eta) = (d\chi/du)^{-1/2}\exp(\pm i\chi) \tag{6-22}$$

we see that Eq. (6-9c) allows us to calculate χ if we know η. Therefore, let us now show how to determine η when we know χ. We do this by inverting Eq. (6-9c). Let us first define γ by letting

$$\chi = \frac{\pi}{4} + \gamma \tag{6-23}$$

and then write

$$F_{1,2}(y,\lambda) = \underbrace{\left(\frac{d\gamma}{du}\right)^{-1/2} \exp\left(\pm i\frac{\pi}{4}\right)\exp(\pm i\gamma)}_{\substack{\text{extended}\\ \text{Jeffreys form}}} = \underbrace{\left(\frac{d\eta}{du}\right)^{-1/2} w_{1,2}(-\eta)}_{\substack{\text{extended}\\ \text{Langer form}}} \tag{6-24}$$

From Eq. (6-9c), we find that

$$\gamma = \frac{2}{3}\eta^{3/2}\left\{1 - \frac{5}{32}\frac{1}{\eta^3} + \frac{1105}{6144}\frac{1}{\eta^6} - \frac{82825}{65536}\frac{1}{\eta^9} + \frac{12820315 25}{58720256}\frac{1}{\eta^{12}} + \cdots\right\} \tag{6-25}$$

If we let

$$\frac{2}{3}\eta^{3/2} = \gamma + \frac{A_1}{\gamma} + \frac{A_2}{\gamma^3} + \cdots$$

we can show that

$$\frac{2}{3}\eta^{3/2} \sim \gamma + \frac{5}{72}\frac{1}{\gamma} - \frac{1255}{31104}\frac{1}{\gamma^3} + \frac{272075}{2239488}\frac{1}{\gamma^5} + \cdots \tag{6-26a}$$

or

$$\eta = \left(\frac{3}{2}\gamma\right)^{2/3} + \frac{5}{48}\left(\frac{3}{2}\gamma\right)^{-4/3} - \frac{295}{2304}\left(\frac{3}{2}\gamma\right)^{-10/3} + \cdots \qquad (6\text{-}26b)$$

This important result permits us to change from the extended Jeffreys form to the extended Langer form. On the other hand, Eq. (6-25) permits us to change from the extended Langer form to the extended Jeffreys form. These results are elegant statements of a procedure which is suggested in a remark made by Cherry (Ref. 4, p. 250, footnote) and which was used (in a different form) by Olver (Ref. 5, p. 341).

In Eq. (4-29) we showed that η is a solution of

$$\eta\,(\dot{\eta})^2 - \frac{3}{4}\left(\frac{\ddot{\eta}}{\dot{\eta}}\right)^2 + \frac{1}{2}\frac{\dddot{\eta}}{\dot{\eta}} = k^2 f(y) - \lambda \qquad (6\text{-}27)$$

It may be convenient in solving this equation to let

$$\xi = \frac{2}{3}\eta^{3/2}$$

and observe that $\xi(y)$ is a solution of

$$(\dot{\xi})^2 - \frac{3}{4}\left(\frac{\ddot{\xi}}{\dot{\xi}}\right)^2 + \frac{1}{2}\frac{\dddot{\xi}}{\dot{\xi}} + \frac{5}{36}\left(\frac{\dot{\xi}}{\xi}\right)^2 = k^2 f(y) - \lambda \qquad (6\text{-}28)$$

where the dots denote differentiation with respect to y. Eq. (6-28) is remarkably similar to Eq. (6-19), the only difference being the term containing the square of the logarithmic derivative of ξ.

We should also observe that Imai (Ref. 6, Eq. (19)) has given a result which can be interpreted as an extended Jeffreys' approximation. The result is of the form

$$F_{1,2}(y,\lambda) \sim \frac{k^{1/3} f_1^{1/6}}{\sqrt{k^2 f(y) - \lambda}} \exp\left(\pm i\frac{\pi}{4}\right) \exp\left[\pm ikz \pm \frac{1}{2ik} Q_1(z) + \frac{1}{4k^2} Q(z)\right] \qquad (6\text{-}29)$$

where

$$z = \int_{y_1}^{y} \sqrt{f^2(v) - \lambda k^{-2}}\; dv \qquad (6\text{-}30a)$$

$$Q_1(z) = \int_0^z \left[Q(v) + \frac{5}{36}\frac{1}{v^2}\right] dv + \frac{5}{36}\frac{1}{z} \qquad (6\text{-}30b)$$

$$Q(z) = -\left(\frac{5}{36}\right)\frac{1}{z^2} + \alpha\, z^{-2/3} + \alpha_0 + \alpha_1\, z^{2/3} + \cdots) \qquad (6\text{-}30c)$$

and α, α_0, and α_1 are defined by Eq. (4-37), Eq. (4-38), and Eq. (4-39), respectively. If we let

$$\gamma \sim kz - \frac{1}{2k}\, Q_1(z) \qquad (6\text{-}31)$$

and observe that

$$\frac{d\gamma}{d(kz)} \sim 1 - \frac{1}{2k^2}\, Q(z) \quad , \qquad \frac{d(kz)}{dy} = \sqrt{k^2 f(y) - \lambda}$$

we can show that

$$\left(\frac{d\gamma}{dy}\right)^{-1/2} = \frac{1}{\sqrt[4]{k^2 f(y) - \lambda}}\left[1 - \frac{1}{2k^2}\, Q(z)\right]^{-1/2} \sim \frac{1}{\sqrt[4]{k^2 f(y) - \lambda}}\left[1 + \frac{1}{4k^2}\, Q(z)\right]$$

$$\sim \frac{1}{\sqrt[4]{k^2 f(y) - \lambda}}\, \exp\left[\frac{1}{4k^2}\, Q(z)\right] \qquad (6\text{-}32)$$

Therefore, we can rewrite Imai's result in the concise form

$$F_{1,2}(y,\lambda) \sim k^{1/3} f_1^{1/6} \left(\frac{d\gamma}{dy}\right)^{-1/2} \exp\left(\pm i\,\frac{\pi}{4}\right) \exp\left(\pm i\,\gamma\right) \qquad (6\text{-}33)$$

if we let γ be defined by Eq. (6-31). In order to use these results, we must require that

$$\left|\arg \gamma\right| < \pi$$

6.3 Some Further Forms for $F_{1,2}(y,\lambda)$

A study of a classic memoir by Lorenz (Ref. 7) has led the present author to see that the methods which this 19th century author used to obtain asymptotic estimates for the Bessel functions provides a suggestion for an alternative approach to the computation of $F_{1,2}(y,\lambda)$. Let

$$F_{1,2}(y,\lambda) = \sqrt{\pm 1\, A(y)}\ \exp(\pm i\gamma) \tag{6-34}$$

If we let primes denote differentiation with respect to y, we can show that the function A(y) has the properties

$$A(y) = F_1(y,\lambda)\, F_2(y,\lambda) = F_1 F_2$$

$$A'(y) = F_1'\, F_2 + F_1\, F_2'$$

$$A''(y) = F_1''\, F_2 + F_1\, F_2'' + 2F_1'\, F_2' = 2\left[k^2 f(y) - \lambda\right] A(y) + 2F_1' F_2'$$

$$A'''(y) = 2\left[k^2 f(y) - \lambda\right] A'(y) + 2k^2 f'(y)\, A(y) + 2F_1' F_2'' + 2F_1''\, F_2'$$

$$= 2\left[k^2 f(y) - \lambda\right] A'(y) + 2k^2 f'(y)\, A(y) + 2\left[k^2 f(y) - \lambda\right] A'(y)$$

Therefore, we see that A(y) is a solution of the third-order differential equation

$$\frac{d^3 A}{dy^3} + 4\left[\lambda - k^2 f(y)\right] \frac{dA}{dy} - 2k^2 f'(y)\, A(y) = 0 \tag{6-35}$$

Differentiation of the relation

$$\exp(i\, 2\,\gamma) = \frac{F_1(y,\lambda)}{F_2(y,\lambda)}$$

and use of the Wronskian relation $W(F_1, F_2)$ defined by Eq. (5-99), namely

$$W(F_1, F_2) = F_1(y,\lambda) \frac{dF_2(y,\lambda)}{dy} - F_2(y,\lambda) \frac{dF_1(y,\lambda)}{dy} = -2i\, k^{2/3} f_1^{1/3}$$

leads to a first-order differential equation for $\gamma(y)$ which is of the form

$$\frac{d\gamma}{dy} = \frac{\frac{dF_1}{dy} F_2 - \frac{dF_2}{dy} F_1}{2i\, F_1 F_2} = \frac{k^{2/3} f_1^{1/3}}{A(y)} \tag{6-36}$$

We can find an interesting special case of these relations in the introduction of Miller's (Ref. 3) table of the Airy functions.

Miller showed that if a,b, and c were arbitrary constants, that the function

$$Z(x) = a\,Ai^2(x) + b\,Ai(x)\,Bi(x) + c\,Bi^2(x) \tag{6-37}$$

was a solution of the third-order differential equation

$$\frac{d^3Z}{dx^2} - 4x\frac{dZ}{dx} - 2Z = 0 \tag{6-38}$$

We have already cited some results taken from Miller's work which is related to these ideas employed by Lorenz. For example, since the Airy functions are solutions of

$$\frac{d^2w_{1,2}(t)}{dt^2} = t\,w_{1,2}(t)$$

we can use the definitions

$$w_{1,2}(t) = \sqrt{\pi}\,[Bi(t) \pm iAi(t)] \tag{6-39}$$

to show that

$$Z(t) = w_1(t)w_2(t) = \pi\{[Bi(t)]^2 + [Ai(t)]^2\} = \pi F^2(t)$$

is a solution of Eq. (6-37) provided we merely change the indepedent variable from x to t. We readily see that Eq. (6-10) provides an example of Eq. (6-36) if we compare Eq. (6-39) with Eq. (6-11). If we compare the Lorenz form in Eq. (6-34) with the extended Jeffreys form in Eq. (6-34) we see that they differ only in the manner in which the factor multiplying $\exp(\pm i\gamma)$ are defined; however, Eq. (6-36) even provides a relationship between these factors. The basic difference in the Lorenz formulation is that the third-order differential equation in Eq. (6-35) is to be solved, and then the "phase factor" determined by integrating the first-order equation given in Eq. (6-36). From Eq. (6-23) we can show that $\gamma(y)$ is a solution of the non-linear, third-order differential equation

$$(\dot{\gamma})^2 - \frac{3}{4}\left(\frac{\ddot{\gamma}}{\dot{\gamma}}\right)^2 + \frac{1}{2}\left(\frac{\dddot{\gamma}}{\dot{\gamma}}\right) = k^2 f(y) - \lambda \tag{6-40}$$

From the experience of the present author, it appears that if one is interested in deriving asymptotic expansions that the results are arrived at with the greatest ease by solving Eq. (6-40) by assuming a form for the solution (such as we have done in Eq. (4-115) in order to solve Eq. (4-114) and using the nonlinear, third-order differential equation to determine the unknown coefficients. The amplitude (which we should put in quotation marks as "amplitude" since the function will generally be a complex-valued function) can then be determined by computing the derivative $d\Upsilon/du$ as shown in the extended Jeffreys form of Eq. (6-24). However, from the standpoint of obtaining the solutions for $F_{1,2}(y,\lambda)$ by means of numerical methods there are advantages associated with using the form given in Eq. (6-34) and in starting the calculations by solving for $A(y)$ from the linear third-order differential equation of Eq. (6-35). The "phase" function $\Upsilon(y)$ is then to be determined by integrating the first-order differential equation given as Eq. (6-36).

Miller (Ref. 8) has made extensive use of Eq. (6-35) in his studies of the parabolic cylinder functions which satisfy the differential equation

$$\frac{d^2f}{dx^2} + (-a + \frac{1}{4}x^2)\, f = 0 \qquad (6\text{-}41)$$

The use of Eq. (6-35) and Eq. (6-36) is most attractive when λ is real and positive, and $k^2f(y)$ is real, and $\lambda > k^2f(y)$. Under these conditions, both $A(y)$ and $\Upsilon(y)$ are real.

Lorenz (Ref. 7) also gave a method (for the study of the Bessel functions) which suggests a method which can be useful when λ is real and $[-\lambda + k^2f(y)] < 0$ for the values of y of interest. In this region we can show that for y not near the turning point y_1 that $F_{1,2}(y,\lambda)$ has the asymptotic behavior

$$F_{1,2}(y,\lambda) \sim \frac{k^{1/3} f_1^{1/6}}{\sqrt[4]{\lambda - k^2f(y)}} \exp\left[\int_{y_1}^{y} \sqrt{\lambda - k^2f(v)}\; dv\right] \qquad (6\text{-}42)$$

Since this result does not permit us to distinguish between F_1 and F_2, we shall illustrate the idea set forth by Lorenz by considering the analogous situation with the Airy functions as a model of the method. Let us define two functions $U(y,\lambda)$ and $V(y,\lambda)$ by means of the relation

$$2F_{1,2}(y,\lambda) = U(y,\lambda) \pm i\, V(y,\lambda) \tag{6-43}$$

so that

$$U(y,\lambda) = \frac{F_1(y,\lambda) - F_2(y,\lambda)}{2\,i} \tag{6-44a}$$

$$V(y,\lambda) = \frac{F_1(y,\lambda) + F_2(y,\lambda)}{2} \tag{6-44b}$$

The corresponding Airy function formulae are

$$2\,w_{1,2}(z) = u(z) \pm i\, v(z) \tag{6-45}$$

$$v(z) = \frac{w_1(z) - w_2(z)}{2\,i} = \sqrt{\pi}\ \mathrm{Ai}(z) \tag{6-46a}$$

$$u(z) = \frac{w_1(z) + w_2(z)}{2} = \sqrt{\pi}\ \mathrm{Bi}(z) \tag{6-46b}$$

For y near the turning point y_1, we use the function η defined by Eq. (4-118) and write

$$U(y,\lambda) = k^{1/3} f_1^{1/6} \left(\frac{d\eta}{dy}\right)^{-1/2} v(-\eta) \tag{6-47a}$$

$$V(y,\lambda) = k^{1/3} f_1^{1/6} \left(\frac{d\eta}{dy}\right)^{-1/2} u(-\eta) \tag{6-47b}$$

where η is negative and real when $[-\lambda + k^2 f(y)]$ is real and negative.

Miller (Ref. 3) showed that for $\arg x < \frac{2}{3}\pi$,

$$v(x) = \sqrt{\pi}\ \mathrm{Ai}(z) \sim \frac{1}{2}\, x^{-1/4}\, e^{-\xi} \left(1 - \frac{3\cdot 5}{1!\ 216}\,\frac{1}{\xi} + \frac{5\cdot 7\cdot 9\cdot 11}{2!\,(216)^2}\,\frac{1}{\xi^2} - \frac{7\cdot 9\cdot 11\cdot 13\cdot 15\cdot 17}{3!\ (216)^3}\,\frac{1}{\xi^3} + \cdots\right) \tag{6-48a}$$

$$u(x) = \sqrt{\pi}\ \mathrm{Bi}(z) \sim x^{-1/4}\, e^{\xi} \left(1 + \frac{3\cdot 5}{1!\ 216}\ \frac{1}{\xi} + \frac{5\cdot 7\cdot 9\cdot 11}{2!\ (216)^2}\ \frac{1}{\xi^2} + \frac{7\cdot 9\cdot 11\cdot 13\cdot 15\cdot 17}{3!\ (216)^3}\ \frac{1}{\xi^3} + \cdots\right) \qquad (6\text{-}48b)$$

where

$$\xi = \frac{2}{3} x^{3/2}$$

However, the form of asymptotic expansion which is suggested by a study of Lorenz's memoir (Ref. 7) is dependent upon the introduction of the auxiliary functions $r(x)$ and $\mu(x)$ by means of

$$2v(x) = 2\sqrt{\pi}\ \mathrm{Ai}(x) = \sqrt{r(x)}\ \exp\left[-\mu(x)\right] = \left(\frac{d\mu}{dx}\right)^{-1/2} \exp\left[-\mu(x)\right]$$

$$u(x) = \sqrt{\pi}\ \mathrm{Bi}(x) = \sqrt{r(x)}\ \exp\left[\mu(x)\right] = \left(\frac{d\mu}{dx}\right)^{-1/2} \exp\left[\mu(x)\right]$$

Since

$$r(x) = 2v(x)u(x) = 2\pi \mathrm{Ai}(x)\mathrm{Bi}(x)$$

we see that $r(x)$ is a solution of Eq. (6-38) and that

$$\frac{d\mu(x)}{dx} = \frac{1}{r(x)} \qquad (6\text{-}50)$$

or

$$(\dot{\mu})^2 + \frac{3}{4}\left(\frac{\ddot{\mu}}{\dot{\mu}}\right)^2 - \frac{1}{2}\ \frac{\dddot{\mu}}{\dot{\mu}} = x \qquad (6\text{-}51)$$

where the dots denote differentiation with respect to x. We can show that $r(x)$ and $\mu(x)$ possess asymptotic expansions of the

$$r(x) \sim \frac{1}{\sqrt{x}} \left\{1 + \frac{1\cdot 3\cdot 5}{1!\,96}\ \frac{1}{x^3} + \frac{1\cdot 3\cdot 5\cdot 7\cdot 9\cdot 11}{2!\ (96)^2}\ \frac{1}{x^6} + \frac{1\cdot 3\cdot 5\cdot 7\cdot 9\cdot 11\cdot 13\cdot 15\cdot 17}{3!\ (96)^3}\ \frac{1}{x^9} + \cdots\right\} \qquad (6\text{-}52a)$$

$$\mu(x) \sim \frac{2}{3} x^{3/2} \left[1 + \frac{5}{32}\frac{1}{x^3} + \frac{1105}{6144}\ \frac{1}{x^6} + \frac{82825}{65536}\ \frac{1}{x^9} + \cdots\right] \qquad (6\text{-}52b)$$

In a similar way, we can show that

$$2v'(x) = 2\sqrt{\pi}\ \mathrm{Ai}'(x) = -\sqrt{p(x)}\ \exp\left[-\lambda(x)\right]$$

$$u'(x) = \sqrt{\pi}\ \mathrm{Bi}'(x) = \sqrt{p(x)}\ \exp\left[\lambda(x)\right]$$

where

$$\frac{d\lambda}{dx} = \frac{x}{p(x)} \tag{6-53}$$

and $p(x)$ is a solution of the third-order differential equation

$$x^2 \frac{d^3p}{dx^3} - 3x \frac{d^2p}{dx^2} + 3 \frac{dp}{dx} - 4x^3 \frac{dp}{dx} + 2x^2 p = 0 \tag{6-54}$$

We can show that $p(x)$ and $\lambda(x)$ possess asymptotic expansions of the form

$$p(x) \sim x^{1/2}\left\{1 - \frac{1\cdot 3}{1!\,96}\frac{7}{x^3} \frac{1\cdot 3\cdot 5\cdot 7\cdot 9}{2!\,96^2}\frac{13}{x^6} - \frac{1\cdot 3\cdot 5\cdot 9\cdot 11\cdot 13\cdot 15}{3!\,96^3}\frac{1}{x^9} \cdots\right\} \tag{6-55a}$$

$$\lambda(x) \sim \frac{2}{3}x^{3/2}\left\{1 - \frac{7}{32}\frac{1}{x^3} - \frac{1463}{6144}\frac{1}{x^6} - \frac{495291}{327680}\frac{1}{x^9} - \frac{206530429}{8388608}\frac{1}{x^{12}} + \cdots\right\} \tag{6-55b}$$

Let us now turn to Eq. (6-44) and obtain generalizations of Eq. (6-48) by making the definitions

$$2V(y,\lambda) = \sqrt{R(y)}\ \exp[-\Gamma(y)] \tag{6-56a}$$

$$U(y,\lambda) = \sqrt{R(y)}\ \exp[+\Gamma(y)] \tag{6-56b}$$

The function $R(y)$ will be a solution of Eq. (6-35) and $\Gamma(y)$ will be a solution of

$$\dot{\Gamma} = \frac{d\Gamma}{dy} = \frac{k^{2/3} f_1^{1/3}}{R(y)} \tag{6-57}$$

or

$$(\dot{\Gamma})^2 + \frac{3}{4}\left(\frac{\ddot{\Gamma}}{\dot{\Gamma}}\right)^2 - \frac{1}{2}\frac{\dddot{\Gamma}}{\dot{\Gamma}} = k^2 f(y) - \lambda \tag{6-58}$$

The procedures which we have derived from the methods employed by Lorenz were used extensively by Nicholson (Ref. 9). The casual

manner in which Lorenz used these procedures suggests that they may have been well-known methods for analysts of the late nineteenth century. One of the most extensive uses of these methods in recent years has been in the study of the parabolic cylinder functions by Miller (Ref. 8).

Watson (Ref. 10, p. 224) says of the method used in Lorenz's memoir: "It is not easy to estimate exactly the magnitude or the sign of the remainder after any number of terms in these asymptotic expansions when this method is used." This is still a very valid criticism. Nevertheless, these methods permit one to readily "grind out" successive terms in the asymptotic expansions. These series are often found to be useful for computational purposes. The value of these expansions can be greatly enhanced if one follows the suggestions of Miller (Ref. 11) and seeks to determine converging factors. Slater (Ref. 12) has had some outstanding success with Miller's converging factors in the study of the confluent hypergeometric functions which are solutions of Kummer's equation

$$x\frac{d^2y}{dx^2} + (b - x)\frac{dy}{dx} - a\,y = 0$$

If we define $z(x)$ by means of

$$y(x) = x^{-b/2} \exp(x/2)\; z(x)$$

we see that these studies involve Whittaker's equation

$$\frac{d^2w}{dx^2} + \left[-\frac{1}{4} + \frac{b - 2a}{2x} - \frac{b(b-2)}{4x^2}\right] z = 0$$

Since Whittaker's equation is of the same form as the height gain function, some of the methods and results which have been discussed above can be used in the study of this function. In quantum physics, this equation leads to the Coulomb wave functions for which there is an extensive literature.

REFERENCES FOR SECTION 6

1. R. E. Langer, "On Asymptotic Solutions of Ordinary Differential Equations with Applications to Bessel Functions of Large Order," Trans. Amer. Math. Soc., Vol. 33, 1931, pp. 23-64

2. Harvard University Computation Laboratory, Tables of the Modified Hankel Functions of Order One-Third and of Their Derivatives, Annals of the Computation Laboratory of Harvard University, Vol. II, Cambridge, Mass., Harvard University Press, 1945. (The "Introduction" of this book was written by W. E. Furry and H. A. Arnold.)

3. J. C. P. Miller, The Airy Integral, British Association Mathematical Tables, Part-Volume B, Cambridge, Cambridge University Press, 1946

4. T. M. Cherry, "Uniform Asymptotic Formulas for Functions with Transition Points," Trans. Amer. Math. Soc., Vol. 68, 1950, pp. 224-257

5. F. W. J. Olver, "The Asymptotic Expansion of Bessel Functions of Large Order," Phil. Trans. Roy. Soc. (London), Vol. 247A, 1954, pp. 328-368

6. I. Imai, "A Refinement of the WKB Method and Its Application to the Electromagnetic Wave Theory," IRE Trans. on Antennas and Propagation, Vol. AP-4, July, 1956, pp. 233-239

7. L. Lorenz, "Sur la Lumiere Reflechie et Refractee par une Sphere Transparente," Oeuvres Scientifiques de L. Lorenz, Vol. 1, New York, Johnson Reprint Corp., 1964, pp. 405-502

8. J. C. P. Miller, Tables of Weber Parabolic Cylinder Functions, London, Her Majesty's Stationery Office, 1955

9. J. W. Nicholson, "The Asymptotic Expansions of Bessel Functions," Phil. Mag. (Series 6), Vol. 19, 1910, pp. 228-249

10. G. N. Watson, A Treatise on the Theory of Bessel Functions, New York, Cambridge University Press, 1952

11. J. C. P. Miller, "A Method for the Determination of Converging Factors," Proc. Cambridge Phil. Soc., Vol. 48, 1952, pp. 243-254

12. L. J. Slater, Confluent Hypergeometric Functions, New York, Cambridge University Press, 1960

Section 7

ASYMPTOTIC ESTIMATES FOR $U(x,y;x_0,y_0)$

7.1 The Diffraction Functions

In Section 1 we have discussed a mathematical model for the diffraction of waves by a convex surface which involved a function $D(\xi,\zeta,\zeta_0,q)$ which could be represented by a series of the form

$$D(\xi,\zeta,\zeta_0,q) = i\sum_{s=1}^{\infty} \frac{\exp(i\xi\tau_s)\, w_1(\tau_s-\zeta)\, w_1(\tau_s-\zeta_0)}{\tau_s\,[w_1(\tau_s)]^2 - [w_1'(\tau_s)]^2} \tag{7-1}$$

in which the functions $w_1(z)$ and $w_1'(z)$ are the Airy function

$$w_1(z) = \frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty e^{i\frac{1}{3}\pi}} \exp(\tfrac{1}{3}\omega^3 - \omega z)\,d\omega \tag{7-2}$$

and its derivative

$$w_1'(z) = \frac{dw_1(z)}{dz}$$

and the quantitites τ_s are the roots of the equation

$$w_1'(\tau_s) - q\, w_1(\tau_s) = 0 \tag{7-3}$$

We can refer to Eq. (7-1) as the van der Pol-Bremmer formula since this function was first extensively used in their classic papers (Refs. 1 and 2) which appeared during the late 1930's. Pryce (Ref. 3) and Fock (Ref. 4) also expressed $D(\xi,\zeta,\zeta_0,q)$ in the form of a Fourier integral

$$D(\xi,\zeta,\zeta_0,q) = \frac{1}{2\pi}\int_{-\infty}^{\infty} \exp(i\xi t)F(t,\zeta,\zeta_0,q)\,dt \tag{7-4a}$$

where

$$F(t,\zeta,\zeta_0,q) = w_1(t-\zeta_>)v(t-\zeta_<) - \frac{v'(t)-qv(t)}{w_1'(t)-qw_1(t)} w_1(t-\zeta)w_1(t-\zeta_0)$$

$$= \frac{i}{2}w_1(t-\zeta_0)\left[w_2(t-\zeta) - \frac{w_2'(t)-qw_2(t)}{w_1'(t)-qw_1(t)} w_1(t-\zeta)\right] \qquad (7\text{-}4b)$$

where v(t) is the Airy integral

$$v(t) = \frac{1}{\sqrt{\pi}}\int_0^\infty \cos(\tfrac{1}{3}x^3 - xt)dx = \frac{w_1(t) + w_2(t)}{2i} \qquad (7\text{-}5)$$

We have discussed three limiting forms of Eq.(7-1) and three families of two-parameter functions have been introduced and have been referred to as "diffraction functions."

The attenuation function was defined by

$$V_0(\xi,q) = \frac{1}{2}\sqrt{\frac{-i\xi}{\pi}}\int_{-\infty}^{\infty} \exp(i\xi t)\frac{w_1(t)}{w_1'(t)-qw_1(t)}\,dt \qquad (7\text{-}6a)$$

$$= \sqrt{i\pi\xi}\sum_{s=1}^{\infty}\frac{\exp(i\xi\tau_s)}{\tau_s - q^2} \qquad (7\text{-}6b)$$

The current distribution function was defined by

$$V_1(\xi,q) = \frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty}\frac{\exp(i\xi t)}{w_1'(t)-qw_1(t)}\,dt \qquad (7\text{-}7a)$$

$$= i2\sqrt{\pi}\sum_{s=1}^{\infty}\frac{\exp(i\xi\tau_s)}{(\tau_s - q^2)w_1(\tau_s)} \qquad (7\text{-}7b)$$

The reflection coefficient function was defined by

$$V_{11}(\xi,q) = -\sqrt{i/\pi}\int_{-\infty}^{\infty} \exp(i\xi t)\frac{v'(t)-qv(t)}{w_1'(t)-qw_1(t)}\,dt \qquad (7\text{-}8a)$$

$$= -2\sqrt{-i\pi}\sum_{s=1}^{\infty}\frac{\exp(i\xi\tau_s)}{(\tau_s - q^2)[w_1(\tau_s)]^2} \qquad (7\text{-}8b)$$

$$= V_2(\xi,q) + \frac{\sqrt{i}}{2\sqrt{\pi}\,\xi} \qquad (7\text{-}8c)$$

These functions can serve as "models" for the defining of special functions which can be derived from the Green's function which was defined in Eq. (4-13), namely

$$U(x, y; x_0, y_0) = \sum_{s=1}^{\infty}\frac{\Psi^+(x_>,\lambda_s)\,\Psi^-(x_<,\lambda_s)}{W_s}\,\frac{F(y,\lambda_s)\,F(y_0,\lambda_s)}{N_s} \qquad (7\text{-}9)$$

However, we shall more often be interested in actually employing the functions $D(\xi,\zeta,\zeta_0,q)$, $V_0(\xi,q)$, $V_1(\xi,q)$, $V_{11}(\xi,q)$, and $V_2(\xi,q)$ as asymptotic approximations to $U(x,y,;x_0,y_0)$.

7.2 The Reflection Formula

An exception to our aim of actually employing the functions defined in Eqs. (7-1) through (7-8) to approximate Eq. (7-9) is the case of the obtaining of a representation for the reflected wave which corresponds to the result obtained from geometrical optics. In this case, we shall use the theory associated with the function $D(\xi,\zeta,\zeta_0,q)$ as a "model" for how to proceed with the more general function $U(x,y;x_0,y_0)$.

According to Eq. (1-42), we have

$$D_0(\xi,\zeta,\zeta_0) = \frac{1}{2\pi}\int_{-\infty}^{\infty}\exp(i\xi t)\,w_1(t-\zeta_>)v(t-\zeta_<) = \frac{\sqrt{i}}{2\sqrt{\pi\xi}}\exp(-i\Omega) \qquad (7\text{-}10a)$$

where

$$\Omega = \Omega(\xi,\zeta,\zeta_0) = \frac{\xi^3}{12} - \frac{\xi}{2}(\zeta + \zeta_0) - \frac{(\zeta - \zeta_0)^2}{4\xi} \qquad (7\text{-}10b)$$

If we decompose $D(\xi,\zeta,\zeta_0,q)$ into the components

$$D(\xi,\zeta,\zeta_0,q) = D_0(\xi,\zeta,\zeta_0) + R(\xi,\zeta,\zeta_0,q) \qquad (7\text{-}11a)$$

we can use the fact that $D_0(\xi,\zeta,\zeta_0)$ is the field which exists when the "reflecting surface" does not exist to refer to the portion $R(\xi,\zeta,\zeta_0,q)$ as the "reflected wave." We can use the first of the equations in Eq. (7-4b) to arrive at the explicit form for $R(\xi,\zeta,\zeta_0,q)$ which is given in Eq. (1-43). However, we are interested at this point in our discussion in making a different decomposition of $D(\xi,\zeta,\zeta_0,q)$ by deforming the path of integration and by using the second of the equations in Eq. (7-4b). Thus, we define

$$D(\xi,\zeta,\zeta_0,q) = D_i(\xi,\zeta,\zeta_0) + D_r(\xi,\zeta,\zeta_0,q) \qquad (7\text{-}11b)$$

where

$$D_i(\xi,\zeta,\zeta_0) = \frac{i}{4\pi}\int_C \exp(i\xi t) w_1(t-\zeta_>) w_2(t-\zeta_<)\,dt \qquad (7\text{-}12a)$$

$$D_r(\xi,\zeta,\zeta_0,q) = \frac{-i}{4\pi}\int_C \exp(i\xi t) R(t,q) w_1(t-\zeta) w_1(t-\zeta_0)\,dt \qquad (7\text{-}12b)$$

$$R(t,q) = \frac{w_2'(t) - q w_2(t)}{w_1'(t) - q w_1(t)} \qquad (7\text{-}12c)$$

The evaluation of $D_i(\xi,\zeta,\zeta_0)$ and $D_r(\xi,\zeta,\zeta_0,q)$ by means of the method of stationary phase has been discussed in considerable detail by Fock (Ref. 4, pp. 241-246, under the heading "The Reflection Formula."). The contour C associated with the integrals is to be chosen so as to permit the application of the method of stationary phase to the evaluation of the integrals.

Fock's analysis leads to the results

$$D_i(\xi,\zeta,\zeta_0) \approx D_0(\xi,\zeta,\zeta_0) = \frac{\sqrt{i}}{2\sqrt{\pi\xi}} \exp(-i\Omega) \tag{7-13a}$$

where Ω is defined in Eq. (7-10b), and

$$D_r(\alpha+\alpha_0, \alpha^2+2\alpha\beta, \alpha_0{}^2+2\alpha_0\beta_0, q) \xrightarrow[\beta>>1]{}$$

$$- \frac{\sqrt{i}}{2\sqrt{\pi}} \frac{q - i\beta}{q + i\beta} \frac{\sqrt{\beta}}{\sqrt{2\alpha\alpha_0+\beta(\alpha+\alpha_0)}} \exp(i\omega^*) \tag{7-13b}$$

where

$$\omega^* = \frac{2}{3}(\alpha^3+\alpha_0{}^3) + 2\beta(\alpha^2+\alpha_0{}^2) + \beta^2(\alpha + \alpha_0) \tag{7-13c}$$

Eq. (7-13c) is merely a restatement of the result which we have already given as Eq. (1-44).

With the background which we have sketched above, we can now turn to the case of the Green's function $U(x,y;x_0,y_0)$. If we consider the Fourier integral representation given in Eq. (5-121) we see that the counterpart of $D_r(\xi,\zeta,\zeta_0,q)$ which is defined in Eq. (7-12b) is the integral

$$U_r(x, y; x_0, y_0) = - \frac{1}{4\pi k^{2/3} f_1^{1/3}} \int_P \frac{\Psi^+(x_>, \lambda)\, \Psi^-(x_<, \lambda)}{W(\lambda)}$$

$$\frac{F_2'(0, \lambda) + i k Z F_2(0, \lambda)}{F_1'(0, \lambda) + i k Z F_1(0, \lambda)} F_1(y, \lambda) F_1(y_0, \lambda)\, d\lambda \tag{7-14}$$

We define

$$\lambda = k^2\mu$$

and let y_1 denote the "turning point" at which

$$f(y_1) = \mu$$

If we recall the Wronskian property defined by Eq. (4-19)

$$W_s = W_s(\lambda) = \psi^+ \frac{d\psi^-}{dx} - \psi^- \frac{d\psi^+}{dx}$$

and use $\lambda = k^2\mu$, we see that the asymptotic estimate (WKB approximation) of Eq. (4-16), namely

$$\psi^{\pm}(x, \lambda) \xrightarrow[k\to\infty]{} \frac{\text{constant}}{\sqrt[4]{k^2 g(x) + \lambda}} \exp\left[\pm i \int_0^x \sqrt{k^2 g(u) + \lambda}\ du\right]$$

leads to the asymptotic result

$$\frac{\psi^+(x_>, \lambda)\, \psi^-(x_<, \lambda)}{W(\lambda)} \xrightarrow[k\to\infty]{} \frac{\exp\left[i k \int_{x_<}^{x_>} \sqrt{g(u) + \mu}\ du\right]}{-2i k \sqrt[4]{g(x) + \mu}\ \sqrt[4]{g(x_o) + \mu}} \tag{7-15}$$

If we use the leading term of Eq. (6-29) for $F_{1,2}(y,\lambda)$, namely

$$F_{1,2}(y,\lambda) \approx \frac{k^{\frac{1}{3}} \sqrt{\pm i f_1}^{\frac{1}{3}}}{\sqrt{\sqrt{k^2 f(y) - \lambda}}} \exp\left(\pm i \int_{y_1}^{y} \sqrt{k^2 f(v) - \lambda}\ dv\right) \tag{7-16}$$

and use $\lambda = k^2\mu$, we find that

$$\frac{F_2'(0, \lambda) + i k Z F_2(0, \lambda)}{F_1'(0, \lambda) + i k Z F_1(0, \lambda)} \xrightarrow[k\to\infty]{} -i \frac{Z - \sqrt{f_o - \mu}}{Z + \sqrt{f_o - \mu}} \exp\left[-i 2k \int_{y_1}^{0} \sqrt{f(v) - \mu}\ dv\right] \tag{7-17}$$

and

$$F_1(y, \lambda)\, F_1(y_o, \lambda) \xrightarrow[k\to\infty]{} i \frac{k^{-1/3} f_1^{1/3}}{\sqrt{\sqrt{f(y) - \mu}\ \sqrt{f(y_o) - \mu}}} \exp\left[i k \left(\int_{y_1}^{y_o} + \int_{y_1}^{y}\right) \sqrt{f(v) - \mu}\ dv\right] \tag{7-18}$$

where we recall that $f_0 = f(0)$ and $f_1 = f'(0)$. We can now use Eqs. (7-15) through (7-18) to show that these asymptotic results permit us to express Eq. (7-14) in the form

$$U_r(x, y; x_o, y_o) \xrightarrow[k\to\infty]{} -\frac{i}{8\pi} \int_P A(\mu) \exp\left[i k \phi(\mu)\right] d\mu \tag{7-19}$$

where

$$A(\mu) = \frac{Z - \sqrt{f_o - \mu}}{Z + \sqrt{f_o + \mu}} \; \frac{1}{\sqrt[4]{g(x) + \mu}\,\sqrt[4]{g(x_o) + \mu}\,\sqrt[4]{f(y) - \mu}\,\sqrt[4]{f(y_o) - \mu}} \tag{7-20}$$

$$\phi(\mu) = \int_{x_<}^{x_>} \sqrt{g(u) + \mu}\, du + \int_0^{y} \sqrt{f(v) - \mu} + \int_0^{y_o} \sqrt{f(v) - \mu}\, dv \tag{7-21}$$

and the symbol P on the integral sign denotes the fact that the path of the integral is to pass through the point of stationary phase. According to the theory of the method of stationary phase,

$$\int g(z) \exp [i\, k\, f(z)]\, dz \sim \sqrt{\frac{2\pi}{-k\, f''(z_o)}}\; g(z_o) \exp [i\, k\, f(z_o) - i\tfrac{\pi}{4}] \tag{7-22}$$

where z_0 is defined by the condition

$$f_1'(z_0) = 0$$

and in the form given in Eq. (7-22) we assume that $f''(z_0) < 0$. Therefore, we find that

$$U_r(x,y;x_o,y_o) \xrightarrow[k \to \infty]{} - \frac{1}{\sqrt{-32\, \pi\, k\, \phi''(\mu_o)}}\; A(\mu_o) \exp [i\, k\, \phi(\mu_o) + i\tfrac{\pi}{4}] \tag{7-23}$$

where the point of stationary phase μ_0 is defined by

$$\phi'(\mu_o) = \frac{1}{2} \int_{x_<}^{x_>} \frac{1}{\sqrt{g(u) + \mu_o}}\, du - \int_0^{y} \frac{1}{\sqrt{f(v) - \mu_o}}\, dv - \int_0^{y_o} \frac{1}{\sqrt{f(v) - \mu_o}}\, dv = 0 \tag{7-24}$$

The second derivative of $\phi(\mu)$ is of the form

$$\phi''(\mu) = -\frac{1}{4}\left[\int_{x_<}^{x_>} [g(u) + \mu]^{-3/2}\, du + \int_0^{y} [f(v) - \mu]^{-3/2} dv + \int_0^{y_o} [f(v) - \mu]^{-3/2} dv\right]$$

For the case $\mu = \mu_0$ we can express $\phi''(\mu)$ in the form

$$\phi''(\mu_o) =$$

$$-\frac{1}{4\mu_o}\left[\int_0^{y} \frac{f(v)}{[f(v) - \mu_o]^{3/2}}\, dv + \int_0^{y_o} \frac{f(v)}{[f(v) - \mu_o]^{3/2}}\, dv - \int_{x_<}^{x_>} \frac{g(u)}{[g(u) + \mu_o]^{3/2}}\, du\right] \tag{7-25}$$

If we collect these results, we see that Eq.(7-23) informs us that the evaluation by means of the method of stationary phase has led us to the approximation

$$U_r(x,y;x_0,y_0) \xrightarrow[k \to \infty]{} \frac{\sqrt{f_0-\mu_0} - Z}{\sqrt{f_0-\mu_0} + Z} F(x,y,;x_0,y_0) \exp[iP(x,y,x_0,y_0)] \quad (7\text{-}26)$$

where

$$[F(x,y,;x_0,y_0)]^2 = \frac{\left\{\frac{8\pi k}{\mu_0}\left[\left(\int_0^y + \int_0^{y_0}\right)\frac{f(v)}{[f(v)-\mu_0]^{3/2}}dv - \int_{x_<}^{x_>}\frac{g(u)}{[g(u)+\mu_0]^{3/2}}du\right]\right\}^{-1}}{\sqrt{[g(x)+\mu_0][g(x_0)+\mu_0][f(y)-\mu_0][f(y_0)-\mu_0]}}$$

and

$$P(x,y,;x_0,y_0) = \left(\int_{x_<}^{x_>}\sqrt{g(u)+\mu_0}\,du + \int_0^y \sqrt{f(v)-\mu_0} + \int_0^{y_0}\sqrt{f(v)-\mu_0}\,dv\right) + \frac{\pi}{4}$$

Further discussions of the result described by Eq. (7-26) can be found in Refs. 5 through 11. In the above discussion we have merely attempted to sketch the manner in which the WKB approximations are employed to obtain this reflection formula for the field in the lighted region and well away from the "horizon." In this report we are primarily concerned with the applications of the diffraction functions which make it possible to describe the transition across the "horizon".

However, before returning to the applications of the diffraction functions, we will pause to make some remarks concerning the use of the WKB approximation in the case of inhomogeneous media. The author is of the opinion that for many of the practical problems which confront the microwave engineer that the changes in the index

of refraction are often too rapid to permit one to obtain a result of the desired accuracy by means of the WKB approximation. Since the extensions of the WKB approximation to obtain higher-order corrections generally involves the use of asymptotic series (such as the series we gave as Eq. (4-42)), the theory is difficult to "translate" into practical results and the use of these classical results has been rather limited.

The future work on the height gain function will probably be very much influenced by the techniques which are developed for the treatment of the problem of propagation of waves through inhomogeneous layers. Rydbeck (Ref. 12) has laid the foundation for a method which is attractive from the point of view of numerical analysis. Tischer (Ref. 13) has presented a brief discussion of this method in his monograph on space communications. There is already a sizeable literature upon the theory behind these methods. Schelkunoff (Ref. 14) and Bremmer (Ref. 15) have written papers that indicate how one can begin to tackle the problem of a medium in which the rate of change of the index of refraction plays an important role in the obtaining of numerical results. Bremmer (Ref. 15) showed that the WKB approximation could be interpreted as the first term in a series in which the successive terms are classified according to the number of "reflections" which they have undergone. This series has become known as the "Bremmer series" and the reader is cautioned to not confuse the "Bremmer series" discussed by Atkinson (Ref. 16) and Kay (Ref 17) with the series described by Eq. (7-1) which the present author sometimes refers to as the "van der Pol-Bremmer formula."

The basic idea which is contained in Refs. 12 through 17 is that we should seek to determine the "total" solution of height gain differential equation instead of trying to find the two solutions $F_{1,2}(y,\lambda)$ and seek representations for each of these solutions.

To illustrate the difference in approach which may be expected to play an important role in future work, let us observe that for $y_0 > y$ that Eq. (5-121) contains a function

$$F(y,\lambda) = F_2(y,\lambda) - \frac{\dfrac{\partial F_2(0,\lambda)}{\partial y} + ikZ\, F_2(0,\lambda)}{\dfrac{\partial F_1(0,\lambda)}{\partial y} + ikZ\, F_1(0,\lambda)} F_1(y,\lambda) \tag{7-27}$$

which is a solution of the height gain differential equation

$$\frac{d^2F(y,\lambda)}{dy^2} + [-\lambda + k^2 f(y)]\, F(y,\lambda) = 0 \tag{7-28a}$$

which satisfies the boundary condition

$$\frac{\partial F(0,\lambda)}{\partial y} + ikZ\, F(0,\lambda) = 0 \tag{7-28b}$$

If we use the approximations of Eq. (7-17) and Eq. (7-18) to obtain an asymptotic result for the quantity

$$H(y,y_0,\lambda,k) = F(y,\lambda)F_1(y_0,\lambda) \tag{7-29}$$

we arrive at the result (with $\lambda = k^2\mu$)

$$H(y,y_0,\mu,k) \approx H_0(y,y_0,\mu,k) + H_1(y,y_0,\mu,k) \tag{7-30a}$$

where

$$H_0(y,y_0,\mu,k) = \frac{k^{-\frac{1}{3}} f_1^{\frac{1}{3}}}{\sqrt{\sqrt{f(y)-\mu}\,\sqrt{f(y_0)-\mu}}} \exp\left(ik\int_y^{y_0} \sqrt{f(v)-\mu}\, dv\right) \tag{7-30b}$$

$$H_1(y,y_0,\mu,k) = \frac{Z - \sqrt{f_0-\mu}}{Z + \sqrt{f_0-\mu}} \frac{k^{-\frac{1}{3}} f_1^{\frac{1}{3}}}{\sqrt{\sqrt{f(y)-\mu}\,\sqrt{f(y_0)-\mu}}} \exp(ik\Lambda) \tag{7-30c}$$

and

$$\Lambda = \Lambda(y,y_0,\mu) = \left(\int_0^y + \int_0^{y_0}\right)\sqrt{f(v) - \mu}\, dv \tag{7-30d}$$

We can interpret Eq. (7-30c) as a wave which has been reflected from the surface y = 0 with a reflection coefficient

$$R(Z,f_0,\mu) = \frac{Z - \sqrt{f_0 - \mu}}{Z + \sqrt{f_0 - \mu}},$$

a "diverging factor"

$$S(y,y_0,\mu,k) = \frac{k^{-\frac{1}{3}} f_1^{\frac{1}{3}}}{\sqrt{f(y)-\mu}\,\sqrt{f(y_0)-\mu}}$$

and a phase shift which is represented by the factor $\exp(ik\Lambda)$. These considerations can be shown to permit one to interpret these results on the basis of a "geometrical optics" of inhomogeneous media. However, if the function f(y) varies "rapidly" we should expect that the effect of traversing the inhomogeneous medium should involve some "internal reflections" in the medium as well as the "phase shift." If these effects are to be included in Eq. (7-27), one may be forced to obtain some extremely accurate solutions of $F_1(y,\lambda)$ and $F_2(y,\lambda)$. The author's experience with this problem has led him to favor seeking to solve for an approximation to $F(y,\lambda)$, as defined in Eq. (7-28), in the case of media in which these "internal reflections" are important.

In order to illustrate the techniques which can be learned from a study of Refs. 12 through 17, let us consider the problem of finding a solution of the differential equation

$$\frac{d^2U}{dx^2} + p^2(x)U = 0 \qquad (7\text{-}31)$$

under the condition that p(x) is continuous function of the variable x. We assume that $p(x) = p_0$ = constant when $x < 0$. Then, if the time dependence $\exp(-i\omega t)$ is suppressed, an incident plane wave is given by $\exp(ip_0x)$ in the region $x < 0$. For the sake of

simplicity, we shall assume that the medium to the right of x = 0 extends to infinity. The theory can be readily extended to the case in which

$$\frac{dU}{dx} + iZU = 0$$

at $x = x_o > 0$. However, this complication "clutters up" the theory and our aim at this point is merely to point the way for further work in this type of treatment of Eq. (7-31).

Let us assume that in the region x > 0 the total field U(x) can be represented as the sum of two terms

$$U(x) = V(x) + W(x) \tag{7-32}$$

where V(x) is the contribution of all waves moving foward which have arisen from points to the left of x, and W(x) represents the total of all backward waves starting from points to the right of x. From the theory of the WKB approximation, we know that the function

$$Y(x) = \frac{1}{\sqrt{p(x)}} \exp\left(i\int_0^x p(s)ds\right) \tag{7-33a}$$

represents a transmitted wave propagating in the foward direction, and

$$Z(x) = \frac{1}{\sqrt{p(x)}} \exp\left(-i\int_0^x p(s)ds\right) \tag{7-33b}$$

represents a wave propagating in the backward direction. We make the observation that Y(x) and Z(x) satisfy the first-order differential equations

$$Y' = \left(+ip - \frac{p'}{2p}\right)Y \tag{7-34a}$$

$$Z' = \left(-ip - \frac{p'}{2p}\right)Z \tag{7-34b}$$

The "trick" which is found in Refs. 12 through 17 is that of

observing that the functions V(x) and W(x) are solutions of the coupled first-order differential equations

$$V' = (+ip - \frac{p'}{2p})V + \frac{p'}{2p} W \tag{7-35a}$$

$$W' = (-ip - \frac{p'}{2p})W + \frac{p'}{2p} V \tag{7-35b}$$

We can readily verify that this pair of equations is equivalent to Eq. (7-31) by observing that

$$U' = V' + W_1' = ip(V - W)$$

and

$$U'' = ip'(V - W) + ip(V' - W') = - p^2 U$$

If we define $\Psi(x)$ to be the phase factor

$$\Psi(x) = \int_0^x p(s)ds \tag{7-36}$$

we can replace the pair of coupled differential equations given above as Eq. (7-35) by the coupled integral equations

$$W(x) = - \frac{\exp[-i\Psi(x)]}{\sqrt{p(x)}} \int_x^\infty \exp[+i\Psi(t)] \frac{p'(t)}{2\sqrt{p(t)}} V(t)dt \tag{7-37a}$$

$$V(x) = \frac{\exp[+i\Psi(x)]}{\sqrt{p(x)}} \left[A + \int_0^x \exp[-i\Psi(t)] \frac{p'(t)}{2\sqrt{p(t)}} W(t)dt \right] \tag{7-37b}$$

The constant A which appears in Eq. (7-37b) is to be determined by the boundary conditions at the interface x = 0. If we assume that the field for x < 0 is of the form

$$U(x) = \exp(+ip_0 x) + R \exp(-ip_0 x) \tag{7-38}$$

If we require that both U(x) and its derivative be continuous at x = 0, we readily see that

$$A = \sqrt{p(0)} = \sqrt{p_0} \tag{7-39a}$$

$$R = -\frac{1}{\sqrt{p_0}} \int_0^\infty \exp[+i\Psi(t)] \frac{p'(t)}{2\sqrt{p(t)}} V(t)dt \tag{7-39b}$$

The author has not been able to find any literature which deals with the numerical integration of Eq. (7-37) on an electronic computer. Atkinson (Ref. 16) and Kay (Ref. 17) have discussed certain iteration processes which lead to solutions. One of the chief applications of this theory appears to be that of introducing the concept of a "reflection coefficient" which is defined as the ratio of W(x) to V(x), namely

$$R(x) = \frac{W(x)}{V(x)}$$

It is customary to approximate V(x) by neglecting the integral in Eq. (7-37b). This leads to the approximation

$$R(x) \approx -\exp[-i2\Psi(x)] \int_x^\infty \exp[+i2\Psi(t)] \frac{p'(t)}{2p(t)} dt \tag{7-40}$$

Our discussion of the replacement of the second-order equation of the form of Eq. (7-31) by the coupled first-order differential equations has taken us quite far astray from our main purpose which was that of finding approximations for the reflected wave which was defined by Eq. (7-14). However, the practical importance of problems involving inhomogeneous media surrounding convex surfaces (such as the ionized layers surrounding a re-entry body) cannot be overemphasized. The WKB approximation has only limited applicability in some of these problems and in order to enlarge the scope of the applicability of our diffraction functions we need to

be aware of new methods which may offer the prospect of extending the theory to a wider range of problems.

The author would like to direct attention to the fact that the presence of the reciprocal of $p(x)$ in Eq. (7-35) indicates that the type of decomposition of $U(x)$ which is given in Eq. (7-32) can only be used "far away" from the turning points at which $p(x)$ vanishes.

7.3 The Diffraction Formula

Let us now turn to the problem of expressing the Green's function $U(x,y;x_0,y_0)$ as defined in Eq. (7-9) in terms of the diffraction function $D(\xi,\zeta,\zeta_0,q)$ as defined in Eq. (7-1). The first step will be that of using the WKB estimates in Eq. (4-16) and the Wronskian definition in Eq. (4-19) to show that

$$\frac{\Psi^+(x_>,\lambda_s)\Psi^-(x_<,\lambda_s)}{W_s} \xrightarrow[k\to\infty]{} \frac{\exp\left[i\int_{x_<}^{x_>}\sqrt{[k^2g(u)+\lambda_s]}\,du\right]}{-i2\sqrt{\sqrt{[k^2g(x)+\lambda_s]}\,\sqrt{[k^2g(x_0)+\lambda_s]}}} \tag{7-41}$$

We then recall Eq. (5-2)

$$\lambda_s = k^2 f_0 + (k^{\frac{2}{3}} f_1^{\frac{1}{3}})^2 t_s$$

and observe that

$$\begin{aligned}\sqrt{[k^2g(x)+\lambda_s]} &= k\sqrt{[g(x) + f_0 + (f_1/k)^{\frac{2}{3}}t_s]} \\ &\approx k\sqrt{[g(x)+f_0]} + \frac{1}{2}\frac{k^{\frac{1}{3}}f_1^{\frac{2}{3}}t_s}{\sqrt{[g(x)+f_0]}} + \cdots\end{aligned} \tag{7-42}$$

Let us now use the leading term of Eq. (7-42) in the denominator of Eq. (7-41) and retain the first two terms in the phase factor. We then have the approximation

$$\frac{\Psi^+(x_>, \lambda_s)\,\Psi^-(x_<, \lambda_s)}{W_s} \xrightarrow[k\to\infty]{}$$

$$\frac{\exp\left\{ik\int_{x_<}^{x_>}\sqrt{g(u)+f_o}\,du + i\frac{1}{2}t_s k^{1/3} f_1^{2/3}\int_{x_<}^{x_>}\frac{1}{\sqrt{g(u)+f_o}}\,du\right\}}{-i2k\sqrt[4]{g(x)+f_o}\;\sqrt[4]{g(x)+f_o}} \tag{7-43}$$

We now recall from Eq. (5-2) that

$$t_s = \tau_s + \sum_{n=1}^{\infty} D_n(\tau_s, q)\beta^n \quad , \quad \beta = k^{-\frac{2}{3}} f_1^{-\frac{1}{3}} \tag{7-44}$$

and from Eq. (5-112) that

$$\frac{1}{N_s} = \frac{k^{\frac{2}{3}} f_1^{\frac{1}{3}}}{[w_1'(\tau_s)]^2 - \tau_s[w_1(\tau_s)]^2}\,\frac{dt_s}{d\tau_s} \tag{7-45}$$

Let us now introduce several definitions which will simplify the writing out of the formulae which are to follow. Let

$$\gamma = \gamma(x, x_0) = \int_{x_<}^{x_>}\sqrt{g(u)+f_o} \tag{7-46}$$

$$\xi = \xi(x, x_0) = \frac{1}{2} k^{\frac{1}{3}} f_1^{\frac{2}{3}}\int_{x_<}^{x_>}\frac{1}{\sqrt{g(u)+f_o}}\,du \tag{7-47}$$

$$U(x, y; x_0, y_0) \xrightarrow[k\to\infty]{} \frac{i}{2}\frac{(f_1/k)^{\frac{1}{3}}\exp(ik\gamma)}{\sqrt[4]{g(x)+f_o}\;\sqrt[4]{g(x_o)+f_o}}\sum_{s=1}^{\infty}\left\{\exp(i\xi t_s)\right.$$

$$\left.\frac{F(y, \lambda_s)\,F(y_o, \lambda_s)}{\tau_s[w_1(\tau_s)]^2 - [w_1'(\tau_s)]^2}\left[1 + \sum_{n=1}^{\infty}\frac{\partial D_n(\tau_s)}{\partial \tau_s}\beta^n\right]\right\} \tag{7-48}$$

From the first several terms of Eq. (4-148) we find that we can

express the height gain function $F(y,\lambda)$ for $y \to 0$ and $k \to \infty$ in the form

$$F(y,\lambda) \sim w(t-u) - \frac{\beta h_2}{15}\left[(3u+2t)\,w(t-u) + (3u^2+4ut+8t^2)\,w'(t-u)\right] + \cdots \quad (7\text{-}49)$$

where

$$u = k^{2/3} f_1^{1/3}\, y \quad , \quad \lambda = k^2 f_o + k^{4/3} f_1^{2/3}\, t \quad , \quad \beta = k^{-2/3} f_1^{-1/3}$$

$$t = \tau + D_1(\tau)\,\beta + D_2(\tau)\,\beta^2 + \cdots$$

and the Airy function $w(z)$ is understood to be $w_1(z)$. For sufficiently large value of k, we have the parameter $\beta \to 0$ and we can introduce the further approximations

$$t_s \approx \tau_s \quad (7\text{-}50)$$

$$F(y_o,\lambda)F(y,\lambda) \approx w_1(\tau_s - \mu^2)\,w_1(\tau_s - \mu_o^2) \quad (7\text{-}51)$$

where

$$\mu^2 = k^{\frac{2}{3}} f_1^{\frac{1}{3}} y \quad (7\text{-}52a)$$

$$\mu_o^2 = k^{\frac{2}{3}} f_1^{\frac{1}{3}} y_o \quad (7\text{-}52b)$$

If we now use Eq. (7-50) and Eq. (7-51) in Eq. (7-48), and recall that we have $\beta \to 0$, we find that

$$U(x,y;x_o,y_o) \xrightarrow[\substack{y \to 0 \\ y_o \to 0}]{} \frac{(f_1/k)^{\frac{1}{3}} \exp(ik\gamma)}{2\sqrt[4]{g(x)+f_o}\;\sqrt[4]{g(x_o)+f_o}}\, D(\xi,\mu^2,\mu_o^2,q) \quad (7\text{-}53)$$

This result should only be used in the thin boundary layer for which both μ^2 and μ_o^2 are small (where "small" probably means that we should require that these quantities be less than unity).

For larger heights, we should use the WKB approximation given in Eq. (6-16), namely

$$F_1(y,\lambda) \sim \frac{k^{1/3} f_1^{1/6}}{\sqrt[4]{k^2 f(y) - \lambda}} \exp\left(i\frac{\pi}{4}\right) \exp\left[i \int_{y_1}^{y} \sqrt{k^2 f(v) - \lambda}\ dv \right] \qquad (7\text{-}54)$$

where

$$\lambda = k^2 f_o + k^{4/3} f_1^{2/3} t$$

From Eqs. (5-42) and (5-45) we see that

$$y_1 \xrightarrow[\beta \to 0]{} \beta^2 \tau_s$$

and therefore we will replace the lower limit in Eq. (7-54) by 0 when we consider the case $\beta \to 0$. We also take

$$\lambda_s \approx k^2 f_o + (k^2 f_1)^{\frac{2}{3}} \tau_s$$

and observe that the radicals can be approximated by

$$\sqrt{k^2 f(y) - \lambda_s} \approx k\sqrt{f(y) - f_o} + \frac{1}{2} \frac{k^{\frac{1}{3}} f_1^{\frac{2}{3}} \tau_s}{\sqrt{f(y) - f_o}} + \cdots$$

We can then approximate Eq. (7-54) by

$$F_1(y,\lambda_s) \approx \left(\frac{f_1}{k}\right)^{1/6} \frac{\exp(i\pi/4)}{\sqrt[4]{f(y) - f_o}} \exp\left\{ ik \int_0^y \sqrt{f(v) - f_o}\ dv - i\frac{1}{2} \tau_s k^{\frac{1}{3}} f_1^{\frac{2}{3}} \int_0^y \frac{dv}{\sqrt{f(v) - f_o}} \right\} \qquad (7\text{-}55)$$

We can then express $U(x,y;x_o,y_o)$ in the form

$$U(x,y,x_o,y_o) \xrightarrow[\substack{\beta \to 0 \\ y_o \to \infty}]{} \left\{ \frac{i(f_1/k)^{\frac{1}{3}} \exp(ik\gamma + ik\Lambda_o + i\frac{\pi}{4})}{\sqrt[4]{g(x) + f_o}\ \sqrt[4]{g(x_o) + f_o}\ \sqrt[4]{f(y_o) - f_o}} \sum_{s=1}^{\infty} \frac{\exp(i\zeta\tau_s)\ w_1(\tau_s - y)}{\tau_s [w_1(\tau_s)]^2 - [w_1'(\tau_s)]} \right\} \qquad (7\text{-}56)$$

where Λ_o and ζ are defined by

$$\Lambda_0 = \Lambda(y_0) = \int_0^{y_0} \sqrt{f(v) - f_0}\, dv \tag{7-57a}$$

$$\zeta = \tfrac{1}{2} k^{\frac{1}{3}} f_1^{\frac{2}{3}} \left[\int_{x_<}^{x_>} \frac{1}{\sqrt{g(u) + f_0}}\, du - \int_0^{y_0} \frac{1}{\sqrt{f(v) - f_0}}\, dv \right] \tag{7-57b}$$

Since $w_1'(\tau_s) = q w_1(\tau_s)$, we observe that

$$\tau_s [w_1(\tau_s)]^2 - [w_1'(\tau_s)]^2 = (\tau_s - q^2)[w_1(\tau_s)]^2 \tag{7-58}$$

and, therefore, $V_1(\zeta, q)$ as defined by Eq. (7-7b) can also be expressed in the form

$$V_1(\zeta, q) = i\, 2\sqrt{\pi} \sum_{s=1}^{\infty} \frac{\exp(i \zeta \tau_s)}{(\tau_s - q^2)\, w_1(\tau_s)} \tag{7-59}$$

Therefore, if we let $y = 0$ in Eq. (7-56), we obtain

$$U(x, 0; x_0, y_0) \xrightarrow[y_0 \to \infty]{\beta \to 0} \left[\frac{(f_1/k)^{\frac{1}{3}} \exp(ik\gamma + ik\Lambda_0 + i\frac{\pi}{4})}{2\sqrt{\pi}\, \sqrt[4]{g(x) + f_0}\, \sqrt[4]{g(x_0) + f_0}\, \sqrt[4]{f(y_0) - f_0}} \right] V_1(\zeta, q) \tag{7-60}$$

When both y and y_0 are large, we use Eq. (7-55) for both $F(y_0, \lambda_s)$ and $F(y, \lambda_s)$ and arrive at

$$U(x, y; x_0, y_0) \xrightarrow[\substack{y_0 \to \infty \\ y \to \infty}]{\beta \to 0} -\left[\frac{(f_1/k)^{\frac{1}{3}} \exp(ik\sigma)}{\sqrt{[\sqrt{g(x) + f_0}\, \sqrt{g(x_0) + f_0}\, \sqrt{f(y) - f_0}\, \sqrt{f(y_0) - f_0}\,]}} \right.$$

$$\left. \sum_{s=1}^{\infty} \frac{\exp(i \eta \tau_s)}{\tau_s [w_1(\tau_s)]^2 - [w_1'(\tau_s)]^2} \right]$$

$$= \frac{(f_1/k)^{\frac{1}{3}} \exp(ik\sigma + \frac{\pi}{4})}{2\sqrt{\pi [\sqrt{g(x) + f_0}\, \sqrt{g(x_0) + f_0}\, \sqrt{f(y) - f_0}\, \sqrt{f(y_0) - f_0}\,]}}\, V_{11}(\eta, q) \tag{7-61}$$

where Eq. (7-58) has been used in Eq. (7-8b) in order to show that the reflection function can be written in the form

$$V_{11}(\eta,q) = i2\sqrt{i\pi} \sum_{s=1}^{\infty} \frac{\exp(i\eta\tau_s)}{\tau_s[w_1(\tau_s)]^2 - [w_1'(\tau_s)]^2} \qquad (7\text{-}62)$$

and where σ is defined by

$$\sigma = \int_{x_<}^{x_>} \sqrt{g(u) + f_o}\, du + \int_0^y \sqrt{f(v) - f_o}\, dv + \int_0^{y_o} \sqrt{f(v) - f_o}\, dv \qquad (7\text{-}63)$$

and the argument η is defined in terms of a quantity δ by means of

$$\eta = \tfrac{1}{2} k^{1/3} f_1^{2/3} \int_{x_<}^{x_>} \frac{du}{\sqrt{g(u) + f_o}} - \tfrac{1}{2} k^{1/3} f_1^{2/3} \left(\int_0^y + \int_0^{y_o} \right) \frac{dv}{\sqrt{f(v) - f_o}}$$

$$= \tfrac{1}{2} k^{1/3} f_1^{2/3} \delta \qquad (7\text{-}64)$$

and δ is defined by

$$\delta = \int_{x_<}^{x_>} \frac{du}{\sqrt{g(u) + f_o}} - \int_0^y \frac{dv}{\sqrt{f(v) - f_o}} - \int_0^{y_o} \frac{dv}{\sqrt{f(v) - f_o}} \qquad (7\text{-}65)$$

The physical significance of the parameters σ and δ has been discussed by Friedlander (Ref. 6). The path of the diffracted rays is depicted in Fig. 7-1. Friedlander showed that η is the geodesic distance of (x,y) from (x_0,y_0) measured along the descending branch (PA) of the glancing ray, the segment (AB) along the boundary, and the diffracted ray (BQ). The variable δ is the geodesic distance from A to B, i.e., the distance the wave travels along the surface. Friedlander showed that it was possible to also express δ in the form

$$\delta = \int_{x_1}^{x_2} \frac{du}{\sqrt{g(u) + f_o}} \qquad (7\text{-}66)$$

where x_1 and x_2 are defined in Fig. 7-1. Similar discussions have been given by Seckler and Keller (Ref. 7) and by Jones (Refs. 8 and 9).




Fig. 7-1 Ray Paths in Inhomogeneous Medium

These geometrical interpretations for the parameters appearing in the asymptotic estimate for $U(x,y;x_0,y_0)$ are made possible by the use of the asymptotic estimate for the height gain function which is given as Eq. (7-55). This estimate was used in the above form by Friedlander (Ref. 6). However, an analogous result had already been used by Fock (Ref. 4, Eqs. (6.10) and (6.11, p. 274) in his classic paper of 1948.

Since δ as defined by Eq. (7-66) is the same as γ defined by Eq. (7-46), we observe that for $y = y_0 = 0$ that we can use the asymptotic estimate

$$U(x,0;x_0,0) \xrightarrow[\beta \to \infty]{} \frac{i(f_1/k)^{\frac{1}{3}}\exp(ik\sigma)}{\sqrt[4]{g(x)+f_0}\,\sqrt[4]{g(x_0)+f_0}} \sum_{s=1}^{\infty} \frac{\exp(i\eta\tau_s)\,[w_1(\tau_s)]^2}{\tau_s[w_1(\tau_s)]^2 - [w_1'(\tau_s)]^2} \quad (7\text{-}67)$$

along with the property in Eq. (7-58) to show that

$$U(x,0;x_0,0) \xrightarrow[\beta \to \infty]{} \frac{(f_1/k)^{\frac{1}{3}} \exp(ik\sigma + \frac{\pi}{4})}{\sqrt{\pi\eta[\sqrt{g(x)+f_0}\sqrt{g(x_0)+f_0}]}} V_0(\eta,q) \qquad (7\text{-}68)$$

where $V_0(\eta,q)$ is the attenuation function defined in Eq. (7-6), and Eq. (7-63) reduces to

$$\sigma = \int_{x_1}^{x_2} \sqrt{g(u) + f_0}\, du$$

For the case $y = 0$, we observe that

$$\sigma = \Upsilon + \Lambda_0$$

where Υ and Λ_0 are defined by Eqs. (7-46) and (7-57a), respectively. For this case, we also observe that ζ as defined by Eq. (7-57b) is equivalent to η as defined by Eq. (7-64). Therefore, we can express Eq. (7-60) in the alternative form

$$U(x,0;x_0,y_0) \xrightarrow[\substack{y_0 \to \infty}]{\beta \to 0} \frac{(f_1/k)^{\frac{1}{3}} \exp(ik\sigma + \frac{\pi}{4})}{2\sqrt{\pi\sqrt{g(x)+f_0}\sqrt{g(x_0)+f_0}\sqrt{f(y_0)-f_0}}} V_1(\eta,q) \qquad (7\text{-}69)$$

The forms in which we have put Eqs. (7-61), (7-68) and (7-69) reveal that in each case where we have used the two-parameter functions $V_0(\eta,q)$, $V_1(\eta,q)$, and $V_{11}(\eta,q)$ that the parameter η is related to the distance that the wave has traveled along the interface at $y = 0$, whereas the parameter σ which has appeared in the exponential factor that multiplies these special functions is related to the geodesic distance that the wave has traveled along the ray path between the source at (x_0,y_0) and the receiver at (x,y).

Although we have obtained some useful special results in the course of the above discussion, we do not have a formula which is valid for arbitrary values of y and y_0 as $\beta \to 0$. In Eq. (7-53) we have obtained a result for the boundary layer in which <u>both</u> y and y_0

are assumed to be small. In Eq. (7-69) we required that $y = 0$ and $y_0 \to \infty$, whereas in Eq. (7-61) we required that <u>both</u> $y \to \infty$ and $y_0 \to \infty$. Let us now show how to obtain a result which contains all of the results given above as special cases.

7.4 A Digression on Hankel Functions

Before introducing the type of result which we want to employ in place of Eq. (7-55) or Eq. (7-51), let us consider the special case of the Hankel function for which Eq. (7-55) yields the asymptotic estimate

$$H_\nu^{(1)}(kr) \sim \sqrt{\frac{-2i}{\pi k\sqrt{r^2 - a^2}}}\ \exp\left[ik\left(\sqrt{r^2 - a^2} - a\tan^{-1}\frac{\sqrt{r^2 - a^2}}{a}\right) - it\left(\frac{ka}{2}\right)^{1/3}\tan^{-1}\frac{\sqrt{r^2 - a^2}}{a}\right] \tag{7-70}$$

where

$$\nu^2 = (ka)^2 + 4\left(\frac{ka}{2}\right)^{4/3} t \tag{7-71}$$

The corresponding special form taken by Eq. (7-49) is

$$H_\nu^{(1)}(kr) = -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}\left\{w_1(t-u) - \frac{1}{30}\left(\frac{2}{ka}\right)^{2/3}\left[(3u + 2t)\,w_1(t-u) + (3u^2 + 4ut + 8t^2)\,w_1'(t-u)\right] + O\left[\left(\frac{2}{ka}\right)^{4/3}\right]\right\} \tag{7-72}$$

where

$$u = \left(\frac{2}{ka}\right)^{1/3} ka\log\frac{r}{a} = \left(\frac{ka}{2}\right)^{2/3}\log\left(\frac{r}{a}\right)^2 \tag{7-73}$$

The results in Eq. (7-72) are new since in the work of van der Pol, Bremmer, Fock, and other writers it has been customary to replace the Hankel functions $H_\nu^{(1)}(kr)$, $H_\nu^{(1)\prime}(kr)$, where

$$\nu = (ka) + (ka/2)^{1/3}\,\tau \tag{7-74}$$

by the asymptotic approximations

$$H_\nu^{(1)}(kr) \sim -\frac{i}{\sqrt{\pi}} \left(\frac{2}{ka}\right)^{1/3} w_1(\tau - y) \tag{7-75}$$

$$H_\nu^{(1)\prime}(kr) \sim \frac{i}{\sqrt{\pi}} \left(\frac{2}{ka}\right)^{2/3} w_1'(\tau - y) \tag{7-76}$$

where

$$y = \left(\frac{2}{ka}\right)^{1/3} k(r - a) \tag{7-77}$$

Since

$$u = \left(\frac{2}{ka}\right)^{1/3} ka \log \frac{r}{a} = \left(\frac{2}{ka}\right)^{1/3} ka \left[\frac{r - a}{a} - \frac{1}{2} \frac{(r - a)^2}{a^2} + \cdots\right] \tag{7-78}$$

we can express Eq. (7-72) in the alternative form

$$H_\nu^{(1)}(kr) = -\frac{i}{\sqrt{\pi}} \left(\frac{2}{ka}\right)^{1/3} \left\{ w_1(t - y) - \frac{1}{60} \left(\frac{2}{ka}\right)^{2/3} [(6y + 4t)\, w_1(t - y) + (9y^2 + 8y\,t + 16\,t^2)\, w_1'(t - y)] + O\left[(2/ka)^{4/3}\right] \right\} \tag{7-79}$$

A comparison of Eq. (7-71) and Eq. (7-74) leads to

$$t = \tau + \frac{1}{4} \left(\frac{2}{ka}\right)^{2/3} \tau^2 \tag{7-80}$$

Therefore, Eq. (7-79) can be replaced by

$$H_\nu^{(1)}(kr) = -\frac{i}{\sqrt{\pi}} \left(\frac{2}{ka}\right)^{1/3} \left\{ w_1(\tau - y) - \frac{1}{60} \left(\frac{2}{ka}\right)^{2/3} [(6y + 4\tau)\, w_1(\tau - y) + (9y^2 + 8y\tau + \tau^2)\, w_1'(\tau - y)] + O\left[(2/ka)^{4/3}\right] \right\} \tag{7-81}$$

Although Eq. (7-81) is an extension of the classical approximation described by Eq. (7-75), the result is of limited usefulness unless ka is very large and (r - a) is small, or, more specifically,

$$r = a \left\{1 + O[(2/ka)^{2/3}]\right\} \tag{7-82}$$

The results in Eqs. (7-49) and (7-72) are also useful only in the "boundary layer" defined by Eq. (7-82).

If we use the property

$$w_1(t-y) \xrightarrow[y \gg t]{} \frac{1}{4\sqrt{y-t}} \exp\left[i\frac{2}{3}(y-t)^{3/2} + i\frac{\pi}{4}\right] \tag{7-83a}$$

we obtain

$$w_1(t-y) \xrightarrow[y \to \infty]{} \frac{1}{\sqrt[4]{y}} \exp\left[i\left(\frac{2}{3}y^{3/2} - y^{1/2}\,t + \frac{\pi}{4}\right)\right] \tag{7-83b}$$

We then find that according to the "boundary layer" approximation of Eq. (7-75) that

$$H_\nu^{(1)}(kr) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3} w_1(\tau - y) \xrightarrow[y \to \infty]{} \frac{1}{\sqrt{\pi}\sqrt{y}}\left(\frac{2}{ka}\right)^{1/3} \exp\left[i\frac{2}{3}y^{3/2} - i\,y^{1/2}\,\tau - i\frac{\pi}{4}\right] \tag{7-84}$$

or

$$H_\nu^{(1)}(kr) \sim \sqrt{\frac{-2i}{\pi k\sqrt{2a(r-a)}}} \exp\left\{i\frac{1}{3}ka\left[\frac{2(r-a)}{a}\right]^{3/2} - i\tau\left(\frac{ka}{2}\right)^{1/3}\sqrt{\frac{2(r-a)}{a}}\right\} \tag{7-85}$$

This result does not agree with Eq. (7-70). However, if in Eq. (7-70) we use the approximations

$$\sqrt{r^2-a^2} = \sqrt{(r+a)(r-a)} \sim \sqrt{2a(r-a)}$$

$$k\left(\sqrt{r^2-a^2} - a\tan^{-1}\frac{\sqrt{r^2-a^2}}{a}\right) \sim \frac{k}{3}\frac{(r^2-a^2)^{3/2}}{a^2} \sim \frac{ka}{3}\left[\frac{2(r-a)}{a}\right]^{3/2}$$

$$\tan^{-1}\frac{\sqrt{r^2-a^2}}{a} \sim \frac{\sqrt{r^2-a^2}}{a} \sim \sqrt{2\frac{r-a}{a}}$$

we see that Eq. (7-70) "joins" onto the result derived from Eq. (7-75). However, these results serve to emphasize the fact that Eq. (7-75) is only useful in the "boundary layer" defined by Eq. (7-82).

A study of these results reveals that if we merely replace the variable y by a new parameter

$$\zeta^2 = \left(\frac{ka}{2}\right)^{2/3}\frac{r^2-a^2}{a^2} \tag{7-86}$$

we obtain a modified approximation which has the property

$$H_\nu^{(1)}(kr) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3} w_1(\tau - \zeta^2)$$
$$\xrightarrow[\zeta \to \infty]{} \sqrt{\frac{-2i}{\pi k\sqrt{r^2 - a^2}}}\ \exp\left[i\frac{ka}{3}\left(\frac{r^2 - a^2}{a^2}\right)^{3/2} - i\left(\frac{ka}{2}\right)^{1/3}\tau\left(\frac{r^2 - a^2}{a^2}\right)^{1/2}\right] \qquad (7\text{-}87)$$

This result has the advantage of leading to the same amplitude as given by Eq. (7-70). However, the phase is only an approximation to the phase in the exponent of Eq. (7-70). The phase can be readily corrected by observing that for given (finite) values of ζ and τ,

$$\exp\left[ik\left(\sqrt{r^2-a^2} - a\tan^{-1}\frac{\sqrt{r^2-a^2}}{a}\right) - i\left(\frac{ka}{2}\right)^{1/3}\tau\tan^{-1}\left(\frac{\sqrt{r^2-a^2}}{a}\right) - i\frac{2}{3}\zeta^3 + i\zeta\tau\right]$$
$$= \exp\left\{ i\left[\left(\frac{2}{ka}\right)^{2/3}\left(-\frac{2}{5}\zeta^5 + \frac{1}{3}\tau\zeta^3\right) + \left(\frac{2}{ka}\right)^{4/3}\left(\frac{2}{7}\zeta^7 - \frac{1}{5}\tau\zeta^5\right) + \cdots\right]\right\}$$
$$= 1 + O\left[\left(\frac{2}{ka}\right)^{2/3}\right] \qquad (7\text{-}88)$$

Therefore, the modified asymptotic estimate

$$H_\nu^{(1)}(kr) \sim \frac{1}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3} \exp(i\Phi)\, w_1(\tau - \zeta^2) \qquad (7\text{-}89)$$

where

$$\Phi = k\sqrt{r^2-a^2} - ka\tan^{-1}\frac{\sqrt{r^2-a^2}}{a} - \left(\frac{ka}{2}\right)^{1/3}\tau\tan^{-1}\left(\frac{\sqrt{r^2-a^2}}{a}\right) - \frac{2}{3}\zeta^3 + \zeta\tau \qquad (7\text{-}90)$$

leads to Eq. (7-70) for $\zeta \to \infty$ and to Eq. (7-75) for $\zeta \to 0$.

If we use the property

$$\zeta^2 = y + \frac{1}{4}\left(\frac{2}{ka}\right)^{2/3} y^2 \qquad (7\text{-}91)$$

we can use Eq. (7-81) to obtain the result

$$H_\nu^{(1)} \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}\left\{ w_1(\tau - \zeta^2) + \left(\frac{2}{ka}\right)^{2/3}\left[\left(-\frac{1}{15}\tau - \frac{1}{10}\zeta^2\right) w_1(\tau - \zeta^2) + \left(-\frac{\tau^2}{60} - \frac{2}{15}\tau\zeta^2 + \frac{2}{5}\zeta^4\right) w_1'(\tau - \zeta^2)\right] + O\left[\left(\frac{2}{ka}\right)^{4/3}\right]\right\} \qquad (7\text{-}92)$$

$$w'_1(\tau - \zeta^2) \xrightarrow[\zeta \to \infty]{} -i\sqrt{\zeta^2-\tau}\left[1 + i\frac{1}{4(\zeta^2-\tau)^{3/2}} - \frac{5}{32(\zeta^2-\tau)^3} + \cdots\right] w_1(\tau-\zeta^2)$$

$$= -i\left[\zeta - \frac{\tau}{2\zeta} + i\frac{1}{4\zeta^2} - \frac{1}{8}\frac{\tau^2}{\zeta^3} + i\frac{\tau}{4\zeta^4} - \frac{1}{16}\frac{\tau^3}{\zeta^5} - \frac{5}{32}\frac{1}{\zeta^5} + \cdots\right] w_1(\tau - \zeta^2) \qquad (7\text{-}93)$$

we can show that

$$H_\nu^{(1)}(kr) \xrightarrow[\zeta \to \infty]{} -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}\left\{1 + \left(\frac{2}{ka}\right)^{2/3}\left[-i\frac{2}{5}\zeta^5 + i\frac{1}{3}\tau\zeta^3 + i\frac{1}{16}\frac{1}{\zeta} + \cdots\right] + O\left[\left(\frac{2}{ka}\right)^{4/3}\right]\right\} w_1(\tau - \zeta^2) \qquad (7\text{-}94)$$

The presence of the positive powers of ζ in the coefficient of $(ka/2)^{-\frac{2}{3}}$ reveals that this form of representation for $H_\nu^{(1)}(kr)$ is only useful for moderate values of ζ. However, if we write Eq. (7-90) in the form

$$\Phi = 2\left(\frac{ka}{2}\right)^{2/3}\zeta - 2\left(\frac{ka}{2}\right)\tan^{-1}\left[\left(\frac{2}{ka}\right)^{1/3}\zeta\right] - \tau\left(\frac{ka}{2}\right)^{1/3}\tan^{-1}\left[\left(\frac{2}{ka}\right)^{1/3}\zeta\right] - \frac{2}{3}\zeta^3 + \zeta\tau$$

$$= \left(\frac{2}{ka}\right)^{2/3}\left(-\frac{2}{5}\zeta^5 + \frac{1}{3}\tau\zeta^3\right) + \left(\frac{2}{ka}\right)^{4/3}\left(\frac{2}{7}\zeta^7 - \frac{1}{5}\tau\zeta^5\right) + \cdots \qquad (7\text{-}95)$$

we find that

$$\exp(i\Phi) = 1 + \left(\frac{2}{ka}\right)^{2/3}\left[-i\frac{2}{5}\zeta^5 + i\frac{1}{3}\tau\zeta^3\right] + O\left[\left(\frac{2}{ka}\right)^{4/3}\right] \qquad (7\text{-}96)$$

Therefore, if we replace Eq. (7-94) by

$$H_\nu^{(1)}(ka) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}\exp(i\Phi)\left\{1 + \left(\frac{2}{ka}\right)^{2/3}\left(i\frac{1}{16}\frac{1}{\zeta} + \cdots\right) + O\left[\left(\frac{2}{ka}\right)^{4/3}\right]\right\} w_1(\tau - \zeta^2) \qquad (7\text{-}97)$$

we find that

$$H_\nu^{(1)}(ka) \xrightarrow[\zeta \to \infty]{} -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}\exp(i\Phi)\left\{1 + O\left[\left(\frac{2}{ka}\right)^{4/3}\right]\right\} w_1(\tau - \zeta^2) \qquad (7\text{-}98)$$

This result shows that not only do we get the correct phase for the leading term by using Eq. (7-89), but we also obtain a result

for $\zeta \to \infty$ in which the corrections of order $(ka/2)^{-\frac{2}{3}}$ vanish.

Eq. (7-97) cannot be used for small values of ζ because of the presence of the inverse powers of ζ. However, the alternative form

$$H_\nu^{(1)}(kr) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3} \exp[i\Phi]\left\{w_1(\tau-\zeta^2) + \left(\frac{2}{ka}\right)^{2/3}\left[\left(i\frac{2}{5}\zeta^5 - i\frac{1}{3}\tau\zeta^3 - \frac{1}{15}\tau - \frac{1}{10}\zeta^2\right) w_1(\tau-\zeta^2) + \left(-\frac{\tau^2}{60} - \frac{2}{15}\tau\zeta^2 + \frac{2}{5}\zeta^4\right) w_1'(\tau-\zeta^2)\right] + \cdots\right\} \tag{7-99}$$

can be used for both large and small values of ζ. For large values of ζ, the form in Eq. (7-97) may be unsatisfactory because it may be more convenient to replace $w_1(\tau - \zeta^2)$ by an asymptotic expansion. We can obtain such a result from Eq. (7-99) by using the expansions

$$w_1(\tau-\zeta^2) \xrightarrow[\zeta \gg \tau]{} \frac{1}{\sqrt{\zeta^2-\tau}} \exp\left\{i\frac{2}{3}(\zeta^2-\tau)^{3/2} + i\frac{\pi}{4}\right\}\left\{1 - \frac{i\,3\cdot5}{1!216}\left(\frac{3}{2}\right)\frac{1}{(\zeta^2-\tau)^{3/2}} - \frac{5\cdot7\cdot9\cdot11}{2!\,(216)^2}\left(\frac{3}{2}\right)^2\frac{1}{(\zeta^2-\tau)^{3/2}} + \cdots\right\}$$

$$\xrightarrow[\zeta \to \infty]{} \frac{\exp\left(i\frac{2}{3}\zeta^3 - i\tau\zeta + i\frac{\pi}{4}\right)}{\sqrt{\zeta}}\left\{1 + \frac{i\tau^2}{4\zeta} + \frac{1}{\zeta^2}\left(\frac{\tau}{4} - \frac{\tau^4}{32}\right) + \frac{1}{\zeta^3}\left(-\frac{\tau^6}{384} + \frac{5}{48}\tau^3 - \frac{5}{48}\right) + \frac{1}{\zeta^4}\left(\frac{\tau^8}{6144} - \frac{7\tau^5}{384} + \frac{35}{192}\tau^2\right) + \frac{i}{\zeta^5}\left(\frac{\tau^{10}}{122880} - \frac{\tau^7}{512} + \frac{35}{512}\tau^4 - \frac{35}{192}\tau\right) + O\left(\frac{1}{\zeta^6}\right)\right\} \tag{7-100}$$

and

$$w_1'(\tau-\zeta^2) \xrightarrow[\zeta \to \infty]{\zeta \gg \tau} \sqrt{\zeta}\exp\left(i\frac{2}{3}\zeta^3 - i\tau\zeta - i\frac{\pi}{4}\right)\left\{1 + \frac{i\tau^2}{4\zeta} - \frac{1}{\zeta^2}\left(\frac{\tau^4}{32} + \frac{\tau}{4}\right) + \frac{i}{\zeta^3}\left(-\frac{\tau^6}{384} - \frac{\tau^3}{48} + \frac{7}{48}\right) + \frac{1}{\zeta^4}\left(\frac{\tau^8}{6144} - \frac{\tau^5}{384} - \frac{25}{192}\tau^2\right) + \frac{i}{\zeta^5}\left(\frac{\tau^{10}}{122880} - \frac{\tau^7}{1536} - \frac{35}{1536}\tau^4 + \frac{35}{192}\tau\right)\right.$$

(Eq. concluded on next page.)

$$+\frac{1}{\zeta^6}\left(-\frac{\tau^{12}}{2949120}+\frac{5}{73728}\tau^9+\frac{7}{6144}\tau^6-\frac{245}{2304}\tau^3+\frac{455}{4608}\right)$$
$$+O\left(\frac{1}{\zeta^7}\right)\Bigg\} \qquad (7\text{-}101)$$

In the discussion between Eq. (7-75) and Eq. (7-99), we have used τ as defined by Eq. (7-74) in order to conform with the usage of Fock, Wait, and other authors. However, for the sake of completeness, we will also set down the Hankel function results which involve t as defined in Eq. (7-71). In order to do this, we use Eq. (7-80) to write

$$\tau = 2\left(\frac{ka}{2}\right)^{2/3}\left[\sqrt{1+\left(\frac{2}{ka}\right)^{2/3}t}-1\right] = t-\frac{1}{4}\left(\frac{2}{ka}\right)^{2/3}t^2+O\left[\left(\frac{2}{ka}\right)^{4/3}\right] \qquad (7\text{-}102)$$

We then obtain from Eq. (7-72) the result

$$H_\nu^{(1)}(kr) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3}\Bigg\{w_1(t-\zeta^2)-\frac{1}{30}\left(\frac{2}{ka}\right)^{2/3}\left[(3\zeta^2+2t)\,w_1(t-\zeta^2)\right.$$
$$\left.+(-12\zeta^4+4\zeta^2t+8t^2)\,w_1'(t-\zeta^2)+O\left[\left(\frac{2}{ka}\right)^{4/3}\right]\right\} \qquad (7\text{-}103)$$

We can also obtain this result from Eq. (7-72) by using the property

$$u=\left(\frac{ka}{2}\right)^{2/3}\log\left(\frac{r}{a}\right)^2=\zeta^2-\frac{1}{2}\left(\frac{2}{ka}\right)^{2/3}\zeta^4+\cdots \qquad (7\text{-}104)$$

In order to improve Eq. (7-103) for the case when ζ is very large, we define

$$\Psi = k\sqrt{r^2-a^2}-ka\tan^{-1}\frac{\sqrt{r^2-a^2}}{a}-t\left(\frac{ka}{2}\right)^{1/3}\tan^{-1}\frac{\sqrt{r^2-a^2}}{a}-\frac{2}{3}\zeta^3+\zeta t$$
$$=\left(\frac{2}{ka}\right)^{2/3}\left(-\frac{2}{5}\zeta^5+\frac{1}{3}t\zeta^3\right)+O\left[\left(\frac{2}{ka}\right)^{4/3}\right] \qquad (7\text{-}105)$$

and obtain the modified result

$$H_\nu^{(1)}(kr) \sim -\frac{i}{\sqrt{\pi}}\left(\frac{2}{ka}\right)^{1/3} \exp(i\Psi)\Big\{w_1(t-\zeta^2) + \left(\frac{2}{ka}\right)^{2/3}\left[\left(i\frac{2}{5}\zeta^5 - i\frac{1}{3}t\zeta^3 - \frac{1}{15}t - \frac{1}{10}\zeta^2\right) w_1(t-\zeta^2) + \left(-\frac{4}{15}t^2 - \frac{2}{15}t\zeta^2 + \frac{2}{5}\zeta^4\right) w_1'(t-\zeta^2)\right] + O\left[\left(\frac{2}{ka}\right)^{4/3}\right]\Big\} \tag{7-106}$$

These results which we have given for the Hankel function have proven to be very useful in the case of the diffraction of waves by spheres and circular cylinders. We want to now use these concepts to obtain a more useful form for the more general height gain function $F(y,\lambda)$.

7.5 The Extended Height Gain Approximation

The form taken by Eq. (7-99) can serve as a model for an extended result for $F_1(y,\lambda)$ which behaves like Eq. (7-49) for small values of y and which behaves like Eq. (7-55) for large values of y. The first step is to start from the asymptotic estimate of Eq. (7-49) and observe that

$$F_1(y,\lambda) \sim w_1(t-u) = w_1(t - k^{2/3}f_1^{1/3}) \xrightarrow[y \to \infty]{} \left(\frac{f_1}{k}\right)^{1/6} \frac{\exp(i\frac{\pi}{4})}{\sqrt[4]{f_1 y}} \exp\left\{i\frac{2}{3}k f_1^{1/2} y^{3/2} - i t k^{1/3} f_1^{1/6} y^{1/2}\right\} \tag{7-107}$$

Since the factor multiplying the exponent in Eq. (7-55) differs from that in Eq. (7-107), we take a hint from Eq. (7-86) and define

$$\zeta^2 = \left(\frac{k}{f_1}\right)^{2/3}[f(y) - f_o] = u + \beta h_2 u^2 + \beta^2 h_3 u^3 + O(\beta^3) \tag{7-108a}$$

or

$$u = \zeta^2 - \beta h_2 \zeta^4 + O(\beta^2) \tag{7-108b}$$

Next we take a hint from Eq. (7-105) and observe that the exponent in Eq. (7-55) is of the form

$$k \int_0^y \sqrt{f(v) - f_o}\, dv - \frac{1}{2} k^{1/3} f_1^{2/3}\, t \int_0^y \frac{dv}{\sqrt{f(v) - f_o}} = \frac{2}{3}\zeta + \beta\left(-\frac{4}{5} h_2 \zeta^5 + \frac{2}{3} h_2 \zeta^3 t\right) + O(\beta^2) \tag{7-109}$$

Therefore, by defining

$$\Psi = k \int_0^y \sqrt{f(v) - f_o}\, dv - \frac{1}{2} k^{1/3} f_1^{2/3}\, t \int_0^y \frac{dv}{\sqrt{f(v) - f_o}} - \frac{2}{3}\zeta^3 + t\zeta \tag{7-110a}$$

$$= 2\beta h_2 \left(-\frac{2}{5}\zeta^5 + \frac{1}{3} t \zeta^3\right) + O(\beta^2) \tag{7-110b}$$

and observing that

$$w_1(t - u) = w_1(t - \zeta^2) + \beta h_2 \zeta^4 w_1'(t - \zeta^2) + O(\beta^2) \tag{7-111}$$

we can modify Eq. (7-49) to obtain the representation

$$F_1(y, \lambda) = \exp(i\Psi)\left[w_1(t - \zeta^2) + 2\beta h_2 \left[\left(i\frac{2}{5}\zeta^5 - i\frac{1}{3} t\zeta^3 - \frac{1}{15} t - \frac{1}{10}\zeta^2\right) w_1(t-\zeta^2) + \left(-\frac{4}{15} t^2 - \frac{2}{15} t\zeta^2 + \frac{2}{5}\zeta^4\right) w_1'(t - \zeta^2)\right] + O(\beta^2)\right] \tag{7-112}$$

This extended form of the asymptotic expansion for $F_1(y,\lambda)$ is very useful in the problem of finding representation for the Green's function $U(x,y;x_0,y_0)$ which involve the van der Pol-Bremmer formula $D(\xi,\zeta,\zeta_0,q)$. Thus, we can now replace Eq. (7-53) by

$$U(x, y; x_o, y_o) \sim \left(\frac{f_1}{k}\right)^{1/3} \frac{\exp(i k \sigma)}{2\sqrt[4]{g(x) + f_o}\ \sqrt[4]{g(x_o) + f_o}}\, E(\eta, \gamma^2, \gamma_o{}^2, q) \tag{7-113}$$

where

$$E(\eta, \gamma^2, \gamma_o{}^2, q) = \exp\left[-i\frac{2}{3}(\gamma^3 + \gamma_o{}^3)\right] D(\xi, \gamma^2, \gamma_o{}^2, q) \tag{7-114}$$

and

$$\xi = \eta + \gamma + \gamma_o \tag{7-115a}$$

$$\gamma^2 = \left(\frac{k}{f_1}\right)^{2/3} [f(y) - f_o] \qquad (7\text{-}115b)$$

$$\gamma_o^2 = \left(\frac{k}{f_1}\right)^{2/3} [f(y) - f_o] \qquad (7\text{-}115c)$$

and σ and η are defined in Eqs. (7-63) and (7-64), respectively.

The result contained in Eq. (7-113) is the desired extension of the previous results. For example, the limiting form

$$E(\eta, \gamma^2, \gamma_o^2, q) \xrightarrow[\gamma_o \to \infty]{\gamma \to \infty} \frac{1}{2} \sqrt{\frac{1}{\pi \gamma \gamma_o}}\ V_{11}(\eta, q) \qquad (7\text{-}116)$$

leads to Eq. (7-61). The limiting form

$$E(\eta, 0, \gamma_o^2, q) \xrightarrow[\gamma_o \to \infty]{} \frac{1}{2} \sqrt{\frac{1}{\pi \gamma_o}}\ V_1(\eta, q) \qquad (7\text{-}117)$$

leads to Eq. (7-69). If we use the approximation

$$f(y) \approx f_o + f_1 y$$

in Eqs. (7-115b) and (7-115c) we see that

$$\gamma^2 \approx \mu^2$$

$$\gamma_o^2 \approx \mu_o^2$$

the boundary layer approximation of Eq. (7-53) follows from the more general result in Eq. (7-113).

7.6 Extension of Bremmer's Results

The approximation which we have given above as Eq. (7-113) is one of the key results of our investigation. This result permits us have an asymptotic estimate which is valid for arbitrary values of the height parameters provided the parameter k is large enough, i.e., $\beta = k^{-\frac{2}{3}} f_1^{-\frac{1}{3}} \to 0$. Eq. (7-113) is only the leading term in

an asymptotic expansion of $U(x,y;x_0,y_0)$ in powers of β. We have not attempted to include above the further terms in the expansion but it is readily seen that we have laid out the necessary formulae from which such an expansion can be obtained. The further terms in the expansion will involve various derivatives (and integrals) of the basic function $D(\xi,\zeta,\zeta_0,q)$ which was defined in Eq. (7-1). In a later phase of the present research work we will give the explicit forms for these corresctions along with numerical work related to finding the deriva tives and the integrals of $D(\xi,\zeta,\zeta_0,q)$ which are required.

We would like to conclude this discussion by indicating how one can modify the theory as given by Bremmer (Ref. 18) for the propagation of waves through a concentrically stratified troposphere. The results that we wish to give could be obtained directly from Eq. (7-113). However, we have chosen to employ the notation used by Bremmer and to assume that the reader who wishes to follow the discussion which we now undertake will have a copy of Bremmer's paper before him.

We start by making a few changes in the notation employed by Bremmer. We define

$$n_o = n_{eff}(a) = n(a) \tag{7-118a}$$

$$a_o = a_{eff} = a\,\frac{n(a)}{n(a) + a\,n'(a)} \tag{7-118b}$$

$$n(r) = n_{eff}(r) \tag{7-118c}$$

for convenience in writing out certain equations for which Bremmer employs "eff" as subscripts. We then replace Eq. (58) of Bremmer's paper with the asymptotic estimate

$$u_\ell(r) \sim \frac{2}{3}\eta_\infty^{3/2} \tag{7-119a}$$

where

$$\frac{2}{3}\eta_\infty^{3/2} = k\int_a^r \frac{1}{\rho}\sqrt{\rho^2 n^2(\rho) - a^2 n^2(a)}\,d\rho - \left(\frac{k n_o a^3}{2a_o^2}\right)^{\frac{1}{3}} \theta(r)\, t_s \qquad (7\text{-}119b)$$

$$\theta(r) = n_o a \int_a^r \frac{1}{\rho}\frac{1}{\sqrt{\rho^2 n^2(\rho) - a^2 n^2(a)}}\,d\rho \qquad (7\text{-}119c)$$

The results which we have given in Eq. (7-119) are directly related to the form of the exponent which is used in the modified WKB approximation which we gave above as Eq. (7-55). In fact, we now replace Bremmer's WKB approximation for his height gain function by the result

$$f_\ell(r) \sim \frac{1}{r\sqrt[4]{n^2(r) - \frac{a^2}{r^2}\,n^2(a)}}\exp\left(i\frac{2}{3}\eta_\infty^{3/2}\right) \qquad (7\text{-}120)$$

which is essentially our Eq. (7-55). This result is to be used when

$$k\int_a^r \frac{1}{\rho}\sqrt{\rho^2 n^2(\rho) - a^2 n^2(a)}\,d\rho \gg 1 \qquad (7\text{-}121)$$

which requires $r > a$. For $r \to a$, we use the asymptotic approximation which involves the Airy function, namely

$$f_\ell(r) \sim \sqrt{\frac{i}{ar}}\left(\frac{k a_o}{2n_o^2}\right)^{1/6} w_1\left(t_s - \zeta^2\right) \qquad (7\text{-}122)$$

where

$$\zeta^2 = \left(\frac{k n_o a_o}{2}\right)^{2/3}\frac{r^2 n^2(r) - a^2 n^2(a)}{a^2 n^2(a)} \qquad (7\text{-}123)$$

Since

$$w_1\left(t_s - \zeta^2\right) \xrightarrow[\zeta \to \infty]{} \frac{1}{\sqrt{\zeta}}\exp\left(i\frac{2}{3}\zeta^3 - i\zeta t_s - i\frac{\pi}{4}\right) \qquad (7\text{-}124)$$

we obtain the result

$$f_\ell(r) \xrightarrow[\zeta \to \infty]{} \frac{1}{r\sqrt[4]{n^2(r) - \frac{a^2}{r^2}\, n^2(a)}} \exp\left(i\frac{2}{3}\zeta^3 - i\zeta t_s\right) \qquad (7\text{-}125)$$

Since this result does not agree with that given as Eq. (7-120), it is proposed that Eq. (7-122) be replaced by the modified form

$$f_\ell(r) \sim \sqrt{\frac{1}{ar}}\left(\frac{ka_o}{2n_o^2}\right)^{1/6} \exp\left[i\frac{2}{3}\eta_\infty^{3/2} - i\frac{2}{3}\zeta^3 + i\zeta t_s\right] w_1\left(t_s - \zeta^2\right) \qquad (7\text{-}126)$$

which contains both the properties of Eq. (7-120) and Eq. (7-122). We observe that

$$f_\ell(a) \sim \frac{\sqrt{1}}{a}\left(\frac{ka_o}{2n_o^2}\right)^{1/6} w_1\left(t_s\right) \qquad (7\text{-}127)$$

and therefore the height gain ratio given as Eq. (59) by Bremmer is to be replaced by

$$\frac{f_\ell(r)}{f_\ell(a)} \sim \frac{a}{r} \exp\left[i\frac{2}{3}\eta_\infty^{3/2} - i\frac{2}{3}\zeta^3 + i\zeta t_s\right] \frac{w_1\left(t_s - \zeta^2\right)}{w_1\left(t_s\right)} \qquad (7\text{-}128)$$

Let us now turn to the problem of evaluating the expression referred to in Bremmer's paper as Eq. (38). The quantity to be evaluated is

$$\frac{\partial}{\partial \ell}\left\{\frac{\frac{\partial}{\partial r}\left(r f_\ell\right)}{r f_\ell}\right\} = -\frac{2ka n_o}{a^2 f_\ell^2(a)} \int_a^\infty f_\ell^2(r)\, dr \qquad (7\text{-}129)$$

and we will use the approximations

$$f_\ell(r) \sim \frac{\sqrt{1}}{a}\left(\frac{ka_o}{2n_o^2}\right)^{1/6} w_1\left(t_s - \zeta_o^2\right) \qquad (7\text{-}130a)$$

$$\zeta_o^2 = \left(\frac{2}{kn_o a_o}\right)^{1/3} kn_o y = \left(\frac{2k^2 n_o^2}{a_o}\right)^{1/3} y \quad , \; y = r - a \qquad (7\text{-}130b)$$

which are valid for the situation $r \to a$. From the integral which we designated as Eq. (5-126a) we find that

$$\int_0^\infty w_1^2\left(t_s - \alpha y\right)\, dy = \frac{1}{\alpha}\left[t_s w_1^2\left(t_s\right) - w_1'^2\left(t_s\right)\right] \qquad (7\text{-}131)$$

Therefore, we can use the approximate result

$$\int_a^\infty f_\ell^2(r)\, dr \sim \left[\frac{\sqrt{I}}{a}\left(\frac{ka_o}{2n_o^2}\right)^{1/6}\right]^2 \int_0^\infty w_1^2\left[t_s - \left(\frac{2k^2n_o^2}{a_o}\right)^{1/3} y\right] dy$$

$$= \left(\frac{a_o}{2k^2n_o^2}\right)^{1/3}\left[\frac{\sqrt{I}}{a}\left(\frac{ka_o}{2n_o^2}\right)^{1/6}\right]^2 \left\{t_s\left[w_1\left(t_s\right)\right]^2 - \left[w_1'\left(t_s\right)\right]^2\right\} \qquad (7\text{-}132)$$

to show that

$$\frac{\partial}{\partial \ell}\left\{\frac{\frac{\partial}{\partial r}\left(rf_\ell\right)}{rf_\ell}\right\} \sim -\frac{2kn_o}{a}\left(\frac{a_o}{2k^2n_o^2}\right)^{1/3} \frac{t_s\left[w_1\left(t_s\right)\right]^2 - \left[w_1'\left(t_s\right)\right]^2}{\left[w_1\left(t_s\right)\right]^2}$$

$$= \frac{2^{2/3}k^{1/3}n_o^{1/3}a_o^{1/3}}{a}\left[-t_s + q^2\right] \qquad (7\text{-}133)$$

where t_s is defined by the familiar equation.*

$$w_1'\left(t_s\right) - q\, w_1\left(t_s\right) = 0$$

where we must let q be related to Bremmer's notation by means of the relation

---

*The present author suspects that Bremmer's Eq. (38) and all results based upon it are in error. We need to observe that in Bremmer's notation

$$-t_s = \left[\frac{3}{2}\, u_\ell(a)\right]^{2/3}$$

and therefore a comparison of our Eq. (7-133) with Bremmer's Eq. (38) suggests that $n_{eff}(a)$ should be replaced by $[n_{eff}(a)]^{\frac{1}{3}} = n_0^{\frac{1}{3}}$.

$$q = \frac{\Gamma}{a}\left\{\frac{a_o}{2k^2 n_o^2}\right\}^{1/3} = \frac{\Gamma}{kn_o a}\left\{\frac{kn_o a_o}{2}\right\}^{1/3} \tag{7-134}$$

Let us observe that Bremmer's Eq. (41) appears as

$$\ell \sim kn_o a + \left(\frac{kn_o a^3}{2a_o^2}\right)^{1/3} t_s \tag{7-135}$$

in our notation. We can then use the results which we have given above to rewrite Bremmer's Eq. (42) in the <u>extended</u> form

$$\Pi(r,\theta) \sim \left(\frac{B}{Cb}\right)\sqrt{\frac{i8\pi kn_o a}{rb\sin\theta}}\ \frac{1}{2}\left(\frac{2}{kn_o a_o}\right)^{1/3} \exp\left[ikL - i\omega t\right] E\left(\xi, \zeta^2(r), \zeta^2(b), q\right) \tag{7-136}$$

where $E(\xi,\zeta^2(r),\zeta^2(b),q)$ is the function defined above as Eq. (7-114) and L is the path length based upon the geometrical optics of inhomogeneous media

$$L = n_o a\theta + \left[\left(\int_a^r + \int_a^b\right)\frac{1}{\rho}\sqrt{\rho^2 n^2(\rho) - a^2 n^2(a)}\ d\rho\right] \tag{7-137}$$

and the parameters ξ and ζ are defined by

$$\xi = \left(\frac{kn_o a^3}{2a_o^2}\right)^{1/3}\left[\theta - an_o\left[\int_a^r + \int_a^b\right]\frac{d\rho}{\rho\sqrt{\rho^2 n^2(\rho) - a^2 n^2(a)}}\right] \tag{7-138}$$

$$\zeta^2(r) = \left(\frac{kn_o a_o}{2}\right)^{2/3}\ \frac{r^2 n^2(r) - a^2 n^2(a)}{a^2 n^2(a)} \tag{7-139}$$

In Bremmer's paper the concept of a "critical altitude" is introduced and certain expressions are derived which are to be used in place of his Eq. (42) when the source or the receiver lie above

this critical height. A comparison of Bremmer's discussion with the discussion which we have given above will reveal that Bremmer's Eq. (60) is the analog of our Eq. (7-69) and his Eq. (61) is the analog of our Eq. (7-61). Just as we have caused Eqs. (7-61) and (7-69) to follow as special cases of the more general result given in Eq. (7-113), so also the results of Bremmer's Eqs. (60) and (61) follow as special cases of the more general result which we have given above as Eq. (7-136). All of these results which we have based upon the discussion in Bremmer's paper correspond to results which are valid in the vicinity of, and below, the horizon. This is also true of our Eq. (7-136)

Further research is needed to obtain an extension of the results which involve the function $D(\xi,\zeta,\zeta_0,q)$ which will be valid within the lighted region and which will pass over smoothly to agree with the results of the method of stationary phase as outlined in Section 7.2. The results which we have discussed in Section 3 will provide some "checks" upon the extended theory when the research is completed.

REFERENCES FOR SECTION 7

1. B. van der Pol and H. Bremmer, "Diffraction of Electromagnetic Waves from an Electrical Point Source Round a Finitely Conducting Sphere with Applications of Radio-Telegraphy and the Theory of the Rainbow," Part I., Phil. Mag., Vol. 24, 1937, pp. 141-176; Part II, Phil. Mag., Vol. 24, 1937, pp.825-864; Part III, Phil. Mag., Vol. 25, 1938, pp. 817-837

2. B. van der Pol and H. Bremmer, "Further Note on the Propagation of Radio Waves Over a Finitely Conducting Spherical Earth," Phil. Mag., Vol. 27, 1939, pp. 261-275

3. M. H. L. Pryce, "The Diffraction of Radio Waves by the Curvature of the Earth," Advances in Phys., Vol. 2, 1953, pp. 67-95. (This paper is based upon a manuscript written by Pryce in the Fall of 1941. The present author has a reproduction of the original manuscript and has, with the permission of Prof. Pryce, prepared a typewritten copy of the hand-written notes.)

4. V. A. Fock, Electromagnetic Diffraction and Propagation Problems, New York, Pergamon Press, 1965.

5. Institute of Mathematical Sciences, New York University, Propagation of a Pulse in an Inhomogeneous Medium, by F. G. Friedlander, Report EM-76, Contract AF 19(122)-42, New York, 1955

6. F. G. Friedlander, Sound Pulses, Cambridge, Cambridge University Press, 1958. (Parts of Chapter 6, "Some Other Diffraction Problems," have been obtained by making a condensation of Ref. 5.)

7. B. D. Seckler and J. B. Keller, "Geometrical Theory of Diffraction in Inhomogeneous Media," J. Acoust. Soc. Am., Vol. 31, 1959, pp. 192-205

8. B. D. Seckler and J. B. Keller, "Asymptotic Theory of Diffraction in Inhomogeneous Media," J. Acoust, Soc. Am., Vol. 31, 1959, pp. 206-216

9. D. S. Jones, "High-Frequency Refraction and Diffraction in General Media," Phil. Trans. Roy. Soc. (London), Vol. 255A, 1963, pp. 343-387

10. D. S. Jones, The Theory of Electromagnetism, New York, Macmillan, 1964

11. L. B. Felsen, "Quasi-Optic Diffraction," Proc. Symposium on Quasi-Optics, New York, Polytechnic Press, 1964. (See, in particular, the section entitled "Inhomogeneous Dielectric and Related Problems" and the excellent bibliography.)

12. O. E. H. Rydbeck, "On the Propagation of Waves in an Inhomogeneous Medium," Trans. Chalmers University of Technology, No. 74 (Reports from the Research Laboratory of Electronics No. 7), 1948

13. F. J. Tischer, "Propagation through Nonhomogeneous Layers," Basic Theory of Space Communications, Princeton, Van Nostrand, 1965, pp. 262-271

14. S. A. Schelkunoff, "Remarks Concerning Wave Propagation in Stratified Media," Symposium on the Theory of Electromagnetic Waves, New York, Interscience, 1951, pp. 181-191

15. H. Bremmer, "The W.K.B. Approximation as the First Term of a Geometrical-Optical Series," Symposium on the Theory of Electromagnetic Waves, New York, Interscience, 1951, pp. 169-179

16. F. V. Atkinson, "Wave Propagation and the Bremmer Series," J. Math. Anal. and Applic., Vol. 1, 1960, pp. 255-276

17. Courant Institute of Mathematical Sciences, New York University, Some Remarks Concerning the Bremmer Series, Report EM-173, Contract No. AF 19(604)3495, New York, March, 1962

18. H. Bremmer, "On the Theory of Wave Propagation Through a Concentrically Stratified Troposphere with a Smooth Profile. Part I. Discussion of the Extended W.K.B. Approximation," J. Research NBS, Vol. 64D, Sept.-Oct., 1960, pp. 467-482

Section 8
CHEBYSHEV POLYNOMIAL REPRESENTATIONS

8.1 Introduction

The time-span of the present contract was not adequate for the author to be able to develop the representations in terms of Chebyshev polynomials which he envisioned at the start of this program. The author's increasing awareness of the necessity to deal with functions defined in the complex-plane has dampened a little his orginal enthusiasm for the utility of the Chebyshev polynomials. However, the conviction that the Chebyshev polynomials will play an important role in future work has not been undermined in spite of the growing awareness of the fact that the use of these polynomials does not provide a panecea for all of the representation problems that confront the numerical analyst in the problems that arise in diffraction theory.

The past decade has witnessed a rapid growth of interest in a branch of applied mathematics which will probably come to be known as the constructive theory of functions.* Until recently, there appeared to be much more interest in this subject displayed by the papers appearing in Russian mathematical journals than in the papers being presented for publication in journals whose authors were predominately from the United States. This fact can probably be traced to the fact that the Russian students of Chebyshev and Bernstein found their motives for the study of this class of problems through inspiration from their teachers who had

*An excellent introduction to the subject is to be found in the monograph: J. Todd, Introduction to the Constructive Theory of Functions, Basel, Birkhauser, 1963

found these problems to be fascinating. The recent upsurge of interest in the United States would seem to be directly related to the fact that the digital computer has given a "new life" to this "more-or-less classical" branch of mathematics.

The basic problem is that of constructing approximate representations of functions in terms of simpler functions. To the rapidly expanding literature in English, there have recently been added a number of English translations of standard works which originally appeared in Russian.*

In this section, we are interested in only a very narrow spectrum of the topics in the theory of the construction of approximate representations for mathematical functions. We want to consider the advantages of expanding a function defined over a given range of a real variable in terms of certain types of Fourier series in which one employs the so-called Chebyshev polynomials $T_n(x)$ and $T_n^*(x)$ which are defined by means of the relations

$$T_n(x) = \cos[n \cos^{-1}(x)] \quad , -1 < x < 1 \qquad (8\text{-}1a)$$

$$T_n^*(x) = \cos[n \cos^{-1}(2x - 1)] \quad , \quad 0 < x < 1 \qquad (8\text{-}1b)$$

In view of the fact that there exists such a voluminous literature upon the so-called Chebyshev approximation problem, it is almost regretable that the nomenclature "Chebyshev polynomials" has come to be associated with the method of trigonometrical interpoltation of empirical and analytical functions which has been developed since the publication in 1938 of a classic paper by

---

*I.P. Natanson, Constructive Theory of Functions, AEC Trans. TR-4503, 1961

E. Remez, General Computation Methods for Chebyshev Approximation, AEC Trans. TR-4491, 1960

N. I. Achieser, Theory of Approximation, New York, Ungar, 1956

Lanczos (Ref. 1). The "Chebyshev approximation problem" deals with the developing of a representation for a function which possesses the minimum maximum error. The reader who is not familiar with these so-called "best fit" procedures may find it particularly instructive to "thumb through" the small handbook prepared by Hastings (Ref. 2) and to read the chapter on these methods which appeared in a recent book by Ralston (Ref. 37).

The author wishes the reader to grasp the essential point that the Chebyshev polynomials arise only in those problems for which the form chosen for the approximation to the function is that of a polynomial. Since the Chebyshev approximation theory often deals with representations in the form of continued fractions, rational fractions, series of exponentials, etc., there is often no relevance between the papers on Chebyshev "best fit" and the class of polynomials problems which the author proposes as a means of representing certain functions which occur in diffraction theory. Even when the mode of representation which is chosen is that of a polynomial, the representation advocated by the author is that of a truncated Chebyshev series expansion. Although these truncated series may be closely related to the "best fit" polynomial, the amount of labour necessary to compute the "Chebyshev approximation" is far greater than that of the method of trigonometric interpolation which is employed when the truncated Chebyshev series expansion is used to represent the function. In a recent paper, Clenshaw (Ref. 3) has discussed this problem and his references will lead the reader to other sources in which the Chebyshev series expansions are compared with the "best fit" polynomials. The technical report by Murnaghan and Wrench (Ref. 4), which is summarized by the recent journal article (Ref. 5), affords an excellent example of the labour which is required to arrive at the "best fit" polynomial. For most practical problems, the obtaining of the "best fit" polynomial appears to the present author to be primarily of academic interest if one has

at his disposal the means of obtaining readily a truncated Chebyshev polynomial expansion.

8.2 Polynomial Expansions

Unless the reader has had experience with numerical analysis, he will probably be inclined to think of representations of functions in terms of Taylor series, asymptotic expansions in inverse powers of the variable, or in terms of Fourier series. The Taylor series and the Fourier series are the most attractive from the standpoint of our knowledge of what types of functions can be represented and of the rules for the determination of the coefficients in the expansions. Each of these types of expansions involve well-defined infinite processes. Within the range of convergence of the series, we can achieve an arbitrary degree of accuracy by considering a sufficient number of terms and by carrying each term to a sufficient number of significant figures. The asymptotic series, on the other hand, are far more difficult to deal with since the terms in the series may not display a behavior which permits one to evaluate the function for the values of the variable which are of interest.

We can work with the Taylor series with the knowledge that the series will converge within any circle about the expansion point which does not intersect a point at which the function possesses a singularity. Lanzcos (Ref. 1) has written a very interesting paper in which he demonstrates that the Taylor series is an extrapolating series which takes all its information about the function at a given point and then employs this set of information to represent the function in a finite region which surrounds the given point. The demands placed upon such a function are quite severe and we describe these conditions by saying that the function must be analytic in the neighborhood of the expansion point. The function must possess derivatives

of all orders at the point, and the value of the derivative must be independent of the direction in which we approach the point from a neighboring point in the complex plane.

Lanczos (Ref. 1) emphasizes the fact that the Fourier series is an interpolating series. If we consider the relations

$$f(x) = \sum_{n=-\infty}^{\infty} A_n(a,b)\exp\{i2\pi n[(x-a)/(b-a)]\} \tag{8-2}$$

$$A_n(a,b) = \int_a^b f(x)\exp\{-i2\pi n[(x-a)/(b-a)]\}[dx/(b-a)] \tag{8-3}$$

we see that $f(x)$ is represented on $a \leq x \leq b$ in terms of the values of the function over this range. The function $f(x)$ need not be defined anywhere except on the range $a \leq x \leq b$, and, in particular, it need not be analytic in the neighborhood of this range since the only requirement is that the integral be integrable. Therefore, the Fourier series permits us to work with a much broader class of functions than the exremely restricted class of functions which possess Taylor series.

Research in the diffraction of waves by obstacles which are larger than a wavelength requires the investigator to possess a certain amount of familiarity with asymptotic series. Since many of these series have the property that they converge in the beginning but that the terms increase in magnitude after a certain minimum, the practical use of such series is sufficiently difficult that even the subject of such series is avoided in many textbooks. However, the modern reader now has access to two very clear introductions in the form of the little monographs by Copson (Ref. 6) and Erdélyi (Ref. 7). Lanzcos (Ref. 1) discusses these types of series from the standpoint of interpolation and extrapolation and shows that they resemble the Taylor series in that they are extrapolating

series which involve information about the function at a single point (usually the point at infinity since the series are usually in inverse powers of the variable). Furthermore, Lanzcos remarks that, unlike the Taylor series, the value of the derivative need not be the same when infinity is approached along different rays in the complex plane. This phenomena greatly complicates the practical use of asymptotic expansions of functions for complex values of the variable since it is often found that different asymptotic expansions are required for different sectors in the complex plane as the variable is allowed to tend to infinity. With a number of problems that occur in diffraction theory, this phenomenon is known as the "Stokes' phenomenon" and it has been discussed in a number of texts which have appeared in recent years. The author particularly recommends to the reader who is unfamiliar with this phenomenon that he consult the monographs by Jeffreys (Ref. 8) and by Heading (Ref. 9).

The Taylor series and the asymptotic series have something in common in the sense that when they are employed to evaluate a function that only a finite number of terms are employed. The numerical problem then becomes that of evaluating a polynomial. Until the usage of electronic computers became widespread, very little thought was given to the fact that the classical Taylor series and the asymptotic series are very inefficient except for the cases in which the value of the function is obtained by use of only a few terms. Not only does the electronic computer afford us a motivation for seeking a more efficient polynomial representation, but the ease of evaluating polynomials on a computer has given a "new life" to the classic theorem of Weierstrass which asserts that: For any continuous function $f(x)$ defined upon the range $a < x < b$, there exists a sequence of algebraic polynomials of the form

$$P_n(x) = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots + c_n x^n$$

which converges uniformly to $f(x)$ on $a \leq x \leq b$. This theorem informs us that there exist polynomial representations for functions which may not possess a Taylor series over the range of interest.

The finite word-length of the electronic computer has changed some of the basic philosophy regarding what constitutes a "good" algorithm for the evaluation of a function. If we are to work with floating point numbers which consist of 8-digits (which is generally the case in FORTRAN compilers when one works in single-precision), we readily see that any term in the polynomial which is less than 10^{-8} times the sum of the polynomial will not be used in forming the sum. Also, since we seldom can find a single algorithm which is useful for an arbitrary value of the variable, we will find ourselves frequently writing a computer program which is to be used only for a certain range of the variables. When the values of the variable are taken over a real range, and the value of the variable which we will use in the program is likely to lie anywhere in the range with almost equal probability, we arrive at a situation where the conventional power series and asymptotic expansions become relatively inefficient as compared with a Fourier series for the function developed over this range.

The Chebyshev polynomials are nothing more than a specific type of Fourier series which are particularly useful when the function is to be computed for arbitrary values of x lying in the range $a \leq x \leq b$ and the computer to be employed has a finite word-size and hence the accuracy to which the function can be computed is limited by the computer.

Since the transformation

$$x = (2y - b - a)/(b - a) \qquad (8\text{-}4)$$

transforms the segment $a \leq y \leq b$ into the segment $-1 \leq x \leq 1$, we can simplify the discussion by restricting the range to the

x-axis between -1 and +1 when discussing the applications of the Chebyshev polynomials $T_n(x)$. In a similar manner, the transformation

$$x = (y - a)/(b - a) \tag{8-5}$$

transforms the segment $a < y < b$ into the segment $0 < x < 1$ which is the range over which the "T-star" polynomials $T_n^*(x)$ have been defined. However, the $T_n^*(x)$ polynomials will generally be most useful when the function is to be represented over the infinite range $a \leq y < \infty$. In this important class of problems we can define the variable x to be

$$x = y/a \tag{8-6}$$

and then we see that $x = 0$ corresponds to $y \to \infty$ and $x = 1$ corresponds to $y = a$.

If we now assume that a function $f(x)$ which is defined on $(-1,1)$ has been expanded in a series of Chebyshev polynomials with coefficients A_n

$$f(x) = \frac{1}{2} A_0 + \sum_{n=1}^{\infty} A_n T_n(x) = \frac{1}{2} A_0 + \sum_{n=1}^{\infty} A_n \cos(n \cos^{-1} x) \tag{8-7}$$

we can intuitively make some predictions about the behavior of the A_n if the function $f(x)$ behaves in a sufficiently smooth manner. Since Eq. (8-7) is essentially a resolution of the function $f(x)$ into its "spectral components," we can expect that after the first several values of A_n that, as n increases, the magnitude of A_n will diminish rather rapidly because they represent "small ripples" of successive "corrections." Since no A_n which is less than 10^{-8} times the largest of the A_n will contribute to the sum (when employing conventional FORTRAN compilers), we can discard all such "spectral terms" and arrive thereby at an efficient means of evaluating the function $f(x)$ for values of x in the range $(-1,1)$. From the conventional theory of Fourier series, we recognize the fact that we can use Eq. (8-3) in the form

$$A_n = \frac{2}{\pi}\int_0^{\pi} f(\cos\theta)\cos(n\theta)d\theta = \frac{2}{\pi}\int_{-1}^{1} \frac{f(x)T_n(x)}{\sqrt{1-x^2}}\,dx \qquad (8\text{-}8)$$

However, we shall show below that it is seldom necessary to resort to Eq. (8-8) to determine the A_n.

8.3 Properties of Chebyshev Polynomials

The reader who undertakes to employ the Chebyshev polynomials should arrange to have access to Refs. (10) and (11) where a rather complete summary of the properties of the Chebyshev polynomials will be found.

Explicit expressions for the first several Chebyshev polynomials are

$$T_0(x) = 1$$
$$T_1(x) = x$$
$$T_2(x) = 2x^2 - 1$$
$$T_3(x) = 4x^3 - 3x$$
$$T_4(x) = 8x^4 - 8x^2 + 1$$
$$T_5(x) = 16x^5 - 20x^3 + 5x$$

Further polynomials are readily obtained from the recursion formula

$$T_{n+1}(x) - 2x\,T_n(x) + T_{n-1}(x) = 0 \qquad (8\text{-}9)$$

The $T_n^*(x)$ polynomials are related to the $T_n(x)$ polynomials through the relationships

$$T_n^*(x^2) = T_n(2x^2 - 1) = T_{2n}(x) \qquad (8\text{-}10)$$

and therefore it is merely a matter of notational convenience to introduce the "T-star" polynomials. The first few $T_n^*(x)$ are

$$T_0^*(x) = 1$$
$$T_1^*(x) = 2x - 1$$
$$T_2^*(x) = 8x^2 - 8x + 1$$
$$T_3^*(x) = 32x^3 - 48x^2 + 18x - 1$$
$$T_4^*(x) = 128x^4 - 256x^3 + 160x^2 - 32x + 1$$
$$T_5^*(x) = 512x^5 - 1280x^4 + 1120x^3 - 400x^2 + 50x - 1$$

and the recursion formula is

$$T_{n+1}^*(x) = 2(2x - 1)T_n^*(x) - T_{n-1}^*(x) \qquad (8\text{-}11)$$

These relations can be employed when one wishes to convert a representation in terms of Chebyshev polynomials into an algebraic polynomial, i.e., when carrying out the operations indicated by the following formulae:

$$f(x) = \tfrac{1}{2}A_0 + A_1T_1(x) + A_2T_2(x) + A_3T_3(x) + \cdots + A_NT_N(x)$$
$$= B_0 + B_1x^1 + B_2x^2 + B_3x^3 + B_4x^4 + \cdots + B_Nx^N \qquad (8\text{-}12)$$

$$f(x) = \tfrac{1}{2}C_0 + C_1T_1^*(x) + C_2T_2^*(x) + C_3T_3^*(x) + \cdots + C_NT_N^*(x)$$
$$= D_0 + D_1x^1 + D_2x^2 + D_3x^3 + D_4x^4 + \cdots + D_Nx^N \qquad (8\text{-}13)$$

If one wishes to start with the coefficients B_n in Eq. (8-12) and arrive at the coefficients A_n, the following inverse relations will be found useful

$$x^0 = T_0(x)$$
$$x^1 = T_1(x)$$
$$2x^2 = T_0(x) + T_2(x)$$
$$4x^3 = 3T_1(x) + T_3(x)$$
$$8x^4 = 3T_0(x) + 4T_2(x) + T_4(x)$$

If one wishes to start with the coefficients D_n in Eq. (8-13) and arrive at the coefficients C_n, the following inverse relations will be found useful

$$x^0 = T_0^*(x)$$
$$2x^1 = T_0^*(x) + T_1^*(x)$$
$$8x^2 = 3T_0^*(x) + 4T_1^*(x) + T_2^*(x)$$
$$32x^3 = 10T_0^*(x) + 15T_1^*(x) + 6T_2^*(x) + T_3^*(x)$$
$$128x^4 = 35T_0^*(x) + 56T_1^*(x) + 28T_2^*(x) + 8T_3^*(x) + T_4^*(x)$$

Lanczos (Ref. 10) has listed both types of inverse relations for all terms up and including x^{12}. However, in actual practice one will seldom need these explicit representations because the necessary coefficients are readily generated by algorithms which will be discussed below.

8.4 Summing Series of Chebyshev Polynomials

Extensive tables of the Chebyshev polynomials are contained in a publication (Ref. 10) in the NBS Applied Mathematics Series. However, in practice one seldom has a need for these tables since Clenshaw (Ref. 11) has published an algorithm which permits one to sum a series of Chebyshev polynomials without actually computing the values of $T_n(x)$ or $T_n^*(x)$ for the particular value of x and the range of values of n. Before we present the algorithm, let us consider what might be necessary if the algorithm were not known.

Let us introduce a set of FORTRAN variables which are related to the mathematical variables as follows:

$$X = x$$
$$SUM = f(x)$$
$$A0 = A_0$$
$$A(I) = A_n$$

where I = n, for n = 1,2,3,...,N. In order to evaluate the sum in Eq. (8-12) we could employ the FORTRAN program designated as Program 8-1.

Program 8-1

```
      SUBRØUTINE CHEBYS(X,SUM,AO,A,N)
      DIMENSIØN A(N),T(N)
      IF(ABS(X).GT.(1.0)) GØ TØ 10
      T(1) = 1.0
      T(2) = X
      DØ 1  I = 3,N
      T(I) = 2.0*X*T(I-1) - T(I-2)
    1 CØNTINUE
      SUM = 0.5*AO
      DØ 2 I = 2,N
      SUM = SUM + A(I)*T(I)
    2 CØNTINUE
      RETURN
   10 WRITE(6,11) X
      CALL EXIT
   11 FØRMAT(37H1SUBRØUTINE CHEBYS CANNØT HANDLE X = ,E15.8)
      END
```

In the above program we have first set up an array of numbers denoted by T(N) which correspond to the values of the Chebyshev polynomials $T_n(x)$. This has been done by programming the recursion formula in Eq. (8-9). Then, in the second DØ-loop we have formed the sum which appears in Eq. (8-12).

Clenshaw's algorithm permits us to sum the series in Eq. (8-12) by letting

$$b_{N+2} = b_{N+1} = 0$$

and then recurring backwards with the recurrence formula

$$b_r = 2xb_{r+1} - b_{r+2} + A_r \qquad (8\text{-}14)$$

until we have b_2, b_1, and b_0. We then determine f(x) by means of the relation

$$f(x) = \frac{1}{2}(b_0 - b_2)$$
$$= xb_1 - b_2 + \frac{1}{2}A_0 \qquad (8\text{-}15)$$

We observe that the only difference between the recurrence relation in Eq. (8-14) and that in Eq. (8-9) is the presence of the term A_r in Eq. (8-14).

We can carry out the operations indicated in Eqs. (8-14) and (8-15) by means of the FORTRAN program which is designated as Program 8-2.

Program 8-2

```
      SUBRØUTINE CLENSW(X,SUM,AO,A,N)
      DIMENSIØN A(N)
      IF(ABS(X).GT.(1.0)) GØ TØ 10
      B2 = 0.0
      B1 = 0.0
      DØ 1 I = 1,N
      M = N + 1 - I
      B0 = 2.0*X*B1 - B2 + A(M)
      B2 = B1
      B1 = B0
    1 CØNTINUE
      SUM = X*B1 - B2 + 0.5*AO
      RETURN
   10 WRITE(6,11) X
      CALL EXIT
   11 FØRMAT(37H1SUBRØUTINE CLENSW CANNØT HANDLE X = ,E15.8)
      END
```

A comparison of Program 8-1 with Program 8-2 reveals that we now have only one DØ-loop, and that it is no longer necessary to set up the array T(I), I = 1 to N.

Since $T_n^*(x) = T_n(2x - 1)$, we observe that the algorithm in Eq. (8-14) can be employed to sum the series in Eq. (8-13) if we merely replace the recurrence formula of Eq. (8-14) by

$$b_r = 2(2x - 1)b_{r+1} - b_{r+2} + C_r \qquad (8\text{-}16)$$

Therefore, we can sum a series of "T-star" polynomials as given in Eq. (8-13) by means of Program 8-3. We observe that

$$C0 = C_0$$

$$C(I) = C_n$$

for I = n, with n ranging from 1 to N.

Program 8-3

```
      SUBRØUTINE TNSTAR(X,SUM,C0,C,N)
      DIMENSIØN C(N)
      IF((X.LT.(0.0)).ØR.(X.GT.(1.0))) GØ TØ 10
      Z = 2.0*X - 1.0
      CALL CLENSW(Z,SUM,C0,C,N)
      RETURN
   10 WRITE(6,11) X
      CALL EXIT
   11 FØRMAT(37H1SUBRØUTINE TNSTAR CANNØT HANDLE X = ,E15.8)
      END
```

The property of the Chebyshev polynomials

$$T_{2n}(x) = T_n(2x^2 - 1)$$

permits us to sum a series of even polynomials of the form

$$f(x) = \tfrac{1}{2}E_0 + E_1T_2(x) + E_2T_4(x) + E_3T_6(x) + \cdots + E_NT_{2N}(x) \quad (8\text{-}17)$$

by replacing the recurrence formula of Eq. (8-14) by

$$b_r = 2(2x^2 - 1)b_{r+1} - b_{r+2} + E_r \quad (8\text{-}18)$$

If we then let

$$E0 = E_0$$

$$E(I) = E_n$$

for I = n, and n ranging from 1 to N, we can evaluate the sum in Eq. (8-17) by means of Program 8-4.

Program 8-4

```
      SUBRØUTINE TNEVEN(X,SUM,EO,E,N)
      DIMENSION E(N)
      IF(ABS(X).GT.(1.0)) GØ TØ 10
      Z = 2.0*X*X - 1.0
      CALL CLENSW(Z,SUM,EO,E,N)
      RETURN
   10 WRITE(6,11) X
      CALL EXIT
   11 FØRMAT(37H1SUBRØUTINE TNEVEN CANNØT HANDLE X = ,E15.8)
      END
```

If $f(x)$ is expressed in the form of a odd series of the form

$$f(x) = F_1T_1(x) + F_2T_3(x) + F_3T_5(x) + \cdots + F_NT_{2N-1}(x) \qquad (8\text{-}19)$$

it will often be convenient to use the property described by Eq. (8-9) to express $f(x)$ in the form

$$f(x) = x[\tfrac{1}{2}E_0 + E_1T_2(x) + E_2T_4(x) + \cdots + E_{N-1}T_{2N-2}(x)] \qquad (8\text{-}20)$$

where

$$2F_1 = E_0\ ,\ 2F_2 = E_1 + E_2,\ 2F_3 = E_2 + E_3,\ \cdots,\ 2F_N = E_{N-1}$$

Clenshaw (Ref. 11) has pointed out that the form in Eq. (8-20) has the desirable property of yielding the correct behavior of $f(x)/x$ near $x = 0$. However, if one finds an occasion where the form in Eq. (8-19) is preferred, the recurrence relation in Eq. (8-14) has to be replaced by

$$b_r = 2(2x^2 - 1)b_{r+1} - b_{r+2} + F_r \qquad (8\text{-}21)$$

and the value of $f(x)$ is given by

$$f(x) = x(b_0 - b_1) \qquad (8\text{-}22)$$

The form of Eq. (8-22) reveals that the Clenshaw algorithm yields the correct behavior of $f(x)/x$ for x near zero. Since many practical applications lead to indeterminate forms (such as the

familiar form sin(αx)/x which occurs in many diffraction problems) in which the ratio f(x)/x occurs, it will be convenient to replace Eq. (8-22) by the alternative form

$$f(x) = x\,g(x) = x(b_0 - b_1) \qquad (8\text{-}23)$$

and prepare a FORTRAN program in which

SUM1 = f(x)
SUM2 = g(x)
F(I) = F_n

where I = n, and n ranges through the values 1,2,3,...,N. The series given in Eq. (8-19) can be summed by Program 8-5.

Program 8-5

```
      SUBRØUTINE TNØDDS(X,SUM1,SUM2,F0,F,N)
      DIMENSIØN F(N)
      IF(ABS(X).GT.(1.0)) GØ TØ 10
      Z = 2.0*(2.0*X*X - 1.0)
      B2 = 0.0
      B1 = 0.0
      DØ 1 I = 1,N
      M = N + 1 - I
      B0 = Z*B1 - B2 + F(M)
      B2 = B1
      B1 = B0
    1 CØNTINUE
      SUM2 = B1 - B2
      SUM1 = X*SUM2
      RETURN
   10 WRITE(6,11) X
      CALL EXIT
   11 FORMAT(37H1SUBRØUTINE TNØDDS CANNØT HANDLE X = ,E15.8)
      END
```

The reader will observe that each of the programs given above has been provided with a CALL EXIT which will cause the computer program which employs these subroutines to exit from the electronic computer when the variable x lies outside the allowed range. However, in each case, the computer will write a message to indicate

that the appropriate subroutine could not handle the value of x which caused the program to be terminated by the CALL EXIT statement.

8.5 Further Remarks on the Clenshaw Algorithm

The technique which we have employed in the discussion above to sum the Chebyshev series belongs to a wider class of algorithms which was introduced by Clenshaw (Ref. 12). Let f(x) be represented in the form

$$f(x) = \sum_{n=0}^{N} A_n p_n(x) \tag{8-24}$$

where the functions denoted by $p_n(x)$ satisfy a linear recurrence relation of the form

$$p_{n+1}(x) + \alpha_n p_n(x) + \beta_n p_{n-1}(x) = 0 \tag{8-25}$$

where α_n and β_n may be functions of x as well as of n. The Clenshaw algorithm asserts that if one defines

$$b_{N+2} = b_{N+1} = 0$$

and then constructs the sequence $b_N, b_{N-1}, b_{N-2}, \cdots, b_1, b_0$ according to the rule

$$b_r + \alpha_r b_{r+1} + \beta_{r+1} b_{r+2} = A_r , \quad r = N, N-1, \cdots, b_1, b_0 \tag{8-26}$$

that the sum of the series in Eq. (8-24) is given by

$$f(x) = b_0 p_0(x) + b_1 [p_1(x) + \alpha_0 p_0(x)] \tag{8-27}$$

A well-known example of this algorithm is the case in which

$$p_n(x) = x^n$$

for which Eq. (8-25) takes the simple form

$$p_{n+1}(x) - x p_n(x) = 0$$

Therefore, we see that for this case

$$\alpha_n = -x \quad , \quad \beta_n = 0$$

and Eq. (8-27) reduces to the simple form

$$f(x) = b_0$$

and the recurrence relation reduces to

$$b_r = xb_{r+1} + A_r$$

If we consider a simple case, such as

$$f(x) = A_0 + A_1x + A_2x^2 + A_3x^3$$

we see that the algorithm yields the successive results

$$b_3 = A_3$$
$$b_2 = xb_3 + A_2 = A_2 + A_3x$$
$$b_1 = xb_2 + A_1 = A_1 + A_2x + A_3x^2$$
$$b_0 = xb_1 + A_0 = A_0 + A_1x + A_2x^2 + A_3x^3$$

The reader who is experienced in the methods of numerical analysis will probably recognize this algorithm as being merely the so-called "nested-multiplication" method of evaluating a polynomial, i.e.,

$$f(x) = A_0 + A_1x + A_2x^2 + A_3x^3 = \{A_0 + x[A_1 + x(A_2 + xA_3)]\}$$

The Clenshaw algorithm provides an extension of the "nested multiplication" technique which enables one to evaluate the function f(x) and its derivatives when it is expressed in the form of a Taylor series

$$f(x) = \sum_{n=0}^{N} A_n \frac{x^n}{n!} \qquad (8\text{-}28a)$$

and

$$f^{(m)}(x) = \frac{d^m f(x)}{dx^m} = \sum_{n=m}^{N} A_n \, [n(n-1)\cdots(n-m+1)] \frac{x^n}{n!} \qquad (8\text{-}28b)$$

$$= \sum_{k=0}^{N-m} A_{m+k} \frac{x^k}{k!} \qquad (8\text{-}28c)$$

If we observe that the $p_n(x)$ in Eq. (8-28) are $x^n/x!$ we see that the recurrence relation is

$$p_{n+1}(x) = \frac{x}{n+1} p_n(x)$$

and hence α_n and β_n are defined by

$$\alpha_n = -\frac{x}{n+1} \quad , \quad \beta_n = 0$$

Therefore, if we let

$$b_{N-m+1} = 0$$

$$b_r = \frac{x}{r+1} b_{r+1} + A_{r+m} \quad , \; r = N-m, \; N-m-1, \ldots, 2, 1, 0$$

we will find that

$$f^{(m)}(x) = b_0$$

The summation of Fourier series of the form

$$f(\theta) = \sum_{n=0}^{N} A_n \cos(n\theta) \qquad (8\text{-}29a)$$

$$g(\theta) = \sum_{n=1}^{N} A_n \sin(n\theta) \qquad (8\text{-}29b)$$

can be accomplished readily by using the algorithm in the form

$$b_{N+2} = b_{N+1} = 0$$

$$b_r = 2\cos\theta\, b_{r+1} - b_{r+2} + A_r \ , \quad r = N, N-1, N-2, \ldots, 2, 1, 0$$

The algorithm asserts that

$$f(\theta) = b_0 - b_1 \cos\theta \tag{8-30a}$$

$$g(\theta) = b_1 \sin\theta \tag{8-30b}$$

Since many diffraction problems involving a circular cylinder can be solved by series of the form in Eq. (8-29), the use of this algorithm can be extremely helpful in making more efficient computer programs for the calculation of the scattering patterns, current distributions, and other quantities which can be expressed in a Fourier series of the form of Eq. (8-29).

One can also obtain more efficient programs for the summing of certain series of spherical harmonics such as occur in the theory of the diffraction of waves by a sphere. In the case of a series of Legendre functions $P_n(\cos\theta)$ such as

$$f(\theta) = \sum_{n=0}^{N} A_n P_n(\cos\theta) \tag{8-31}$$

we consider the recurrence formula for the polynomials, namely,

$$(n+1)P_{n+1}(\cos\theta) = (2n+1)\cos\theta\ P_n(\cos\theta) - nP_{n-1}(\cos\theta)$$

and find that the α_n and β_n of Eq. (8-25) are

$$\alpha_n = -\frac{2n+1}{n+1}\cos\theta \ , \qquad \beta_n = \frac{n}{n+1}$$

Therefore, if we set

$$b_{N+2} = b_{N+1} = 0$$

and recur backwards with the formula

$$b_r = \frac{2n+1}{n+1}\cos\theta\, b_{r+1} - \frac{r+1}{r+2} b_{r+2} + A_r \tag{8-32}$$

we will find* that

$$f(\theta) = b_0 \tag{8-33}$$

The result contained in Eq. (8-33) can be used as the basis for summing the many series of the form of Eq. (8-31) which occur in boundary value problems involving a sphere.

There are many other problems in applied mathematics which can be summed by means of the Clenshaw algorithm. For example, in his original paper (Ref. 12), Clenshaw observed that the finite Neumann series of Bessel functions

$$f(x) = \sum_{n=0}^{N} A_n J_{n+\nu}(x)$$

could be evaluated in this manner.

The author has encountered several cases in which the coefficients in series of the form described by Eq. (8-28) behaved in such a manner that the backward recursion led to incorrect results because of some unfavorable cancellation which took place during the recurrence process. Therefore, until such time as the theory of the propagation of errors in such processes is better understood, the author would like to advise the reader that it may be advisable to test the algorithm by comparing its results with the results obtained by computing the magnitude of the quantities $A_n p_n(x)$ in Eq. (8-24) and listing these for visual inspection and

---

*The author suggests that "skeptics" join him in observing that Eq. (8-33) is correct by considering the case of the simple function $f(\theta) = P_0(\cos\theta) + P_1(\cos\theta) + 2P_2(\cos\theta) = \cos\theta + 3\cos^2\theta$. We start with $b_4 = b_3 = 0$, and then compute $b_2 = 2; b_1 = (3/2)\cos\theta\ (2) + 1 = 1 + 3\cos\theta$; and $b_0 = \cos\theta(1 + 3\cos\theta) - (1/2)(2) + 1 = \cos\theta + 3\cos^2\theta$. Therefore, we do find that $f(\theta) = b_0$ as implied by Eq. (8-33).

comparing the sum of this "column" of figures with the results obtained from the algorithm.

8.6 Determination of Coefficients

We now want to take up the problem of the determination of the coefficients A_n in the truncated series of Chebyshev polynomials which is employed to represent a function f(x).

$$f(x) = \frac{1}{2} A_0 + A_1 T_1(x) + A_2 T_2(x) + A_3 T_3(x) + \cdots + A_N T_N(x)$$

$$= B_0 + B_1 x^1 + B_2 x^2 + B_3 x^3 + B_4 x^4 + \cdots + B_N x^N \qquad (8\text{-}34)$$

Let us first of all dispose of a related problem. Suppose we already have the A_n for $n = 0,1,2,\ldots,N$ and we would like to convert from the series of Chebyshev polynomials to the conventional algebraic polynomial, i.e., we know the A_n in Eq. (8-34) and we would like to know the B_n. This was one of the first problems which the author undertook under the present contract, but during the course of the work there have appeared several programs which will accomplish this task. Golden (Ref. 47, pp. 120-124) has presented a good treatment of this problem, and actual FORTRAN IV programs are constructed. Prager (Ref. 79) restricts his discussion to the "shifted" Chebyshev polynomials (which we have referred to as the "T star" polynomials) and he has given FORTRAN IV programs that permit one to convert from the C_n of Eq. (8-13) to the D_n, or vice versa. Arden has discussed this problem when it is in the form of Eq. (8-34) (which is the same as Eq.(8-12)), but his treatment (Ref. 48) seems (to the present author) to fail to correctly yield the B_n from the A_n if N is greater than 5. Since it would be convenient to have the ability to carry out these operations as a subroutine, we give in Program 8-6 a subroutine which will take a set of A(I) = A_n, for I = n (and n < 97) and convert them to a set of B(I) = B_n.

The algorithm which has been programmed in Program 8-6 has two features which may require modification if the reader wishes to compile this FORTRAN IV subroutine on the electronic computer which is available to him. The UNIVAC 1107 compiler permits one to use positive and negative integers, as well as zero, as a subscript. This compiler also permits one to write DØ-loops which proceed in "negative" increments. Both of these features have been employed in Program 8-6 (and in Program 8-7). The reader will observe that in order for the program to have access to

Program 8-6

```
      SUBRØUTINE CBS2PS(N,A,B)
      DIMENSIØN A(N),B(N)
      DIMENSIØN AB1(98),AAB1(99),AB2(98),AAB2(99)
      EQUIVALENCE (AB1(1),AAB1(2)),(AB2(1),AAB2(2))
      IF(N.GT.0) GØ TØ 10
      B(0) = A(0)
      RETURN
   10 IF(N.LE.97) GØ TØ 20
      WRITE(6,11) N
      CALL EXIT
   11 FØRMAT(37H1SUBRØUTINE CBS2PS CANNØT HANDLE N = ,I6)
   20 DØ 9 I=0,N
      AB1(I) = A(I)
    9 CØNTINUE
      NM1 = N-1
      DØ 8 I=NM1,1,-1
      AB2(I+2) = AB2(I+1)
      DØ 7 J=I,1,-1
      AB2(J) = 2.0*AB1(J+1) - AB2(J+2)
    7 CØNTINUE
      AB2(0) = AB1(1) - 0.5*AB2(2)
      B(N-I-1) = AB1(0) - 0.5*AB2(1)
      DØ 6 J=0,I
      AB1(J) = AB2(J)
    6 CØNTINUE
    8 CØNTINUE
      B(N-1) = AB1(0)
      B(N)   = AB1(1)
      RETURN
      END
```

the array AB1(0), AB1(1), AB1(2),... that it has been necessary to dimension AB1(98) and then introduce a second array AAB1(99). The EQUIVALENCE statement is then used in such a manner that a reference to AB1(0) is actually a reference to AAB1(1) and hence the compiler is able to work with zero as a subscript. Many compilers for FORTRAN IV will not permit a DØ-loop of the form

```
DØ 8 I = NM1,1,-1
```

since the allowed form is generally

```
DØ 8 I = N1,N2,N3
```

where N1,N2, and N3 are unsigned (i.e., positive) integers. Both of these features of the UNIVAC 1107 compiler (and certain other FORTRAN IV compilers) are very convenient because they permit us to write an algorithm which is closer to the mathematical language.

Since the Program 8-6 expects to find a value for A(0) and will give back a result B(0), the calling program must contain a "gimmick" such as:

```
DIMENSIØN A(N),B(N),AA(2),BB(2)
EQUIVALENCE (A(1),AA(2)),(B(1),BB(2))
```

Program 8-6 may be found useful in the case of a function which is to be evaluated many times because there are fewer steps involved in evaluating the series when expressed in the form of an algebraic polynomial. However, Clenshaw (Ref. 11, p. 13) has shown that the Chebyshev coefficients of ascending order usually decrease more rapidly than the coefficients of ascending powers. Therefore, the evaluation of the algebraic polynomial may be faster than that of the equivalent Chebyshev expansion, but the Chebyshev expansion may be subject to smaller errors. Clenshaw's example shows definitely some advantages to leaving the function represented as a Chebyshev series. However, Beasley (Ref. 27) has shown that a different rearrangement of the series

used in Clenshaw's example can lead to an algebraic polynomial which does not suffer the disadvantages of the algebraic polynomial which was considered by Clenshaw.

Let us turn now to the problem of obtaining the coefficients A_n in Eq. (8-34) when the coefficients B_n are assumed to be known. A program which will accomplish this is given as Program 8-7. Many of the remarks made in connection with the discussion of Program 8-6 apply equally well to this program since we have also employed the zero subscripts in A(0), B(0), etc., and backward steps are made in DØ-loops such as "DØ 7 J = I,1,-1."

Program 8-7

```
      SUBRØUTINE PS2CBS(N,A,B)
      DIMENSIØN A(N),B(N)
      DIMENSIØN AB1(98),AAB1(99),AB2(98),AAB2(99)
      EQUIVALENCE (AB1(1),AAB1(2)),(AB2(1),AAB2(99))
      IF(N.GT.0) GØ TØ 10
      A(0) = B(0)
      RETURN
   10 IF(N.LE.97) GØ TØ 20
      WRITE(6,11) N
      CALL EXIT
   11 FØRMAT(37H1SUBRØUTINE PS2CBS CANNØT HANDLE N = ,I6)
   20 DØ 2 I=0,N
      AB1(I) = B(I)
    2 CØNTINUE
      NMS1 = N-1
      DØ 3 I=NMS1,1,-1
      AB2(I+2) = 0.0
      AB2(I+1) = 0.0
      DØ 4 J=I,1,-1
      AB2(J) = 2.0*AB1(J+1) - AB2(J+2)
    4 CØNTINUE
      AB2(0) = AB1(1) - 0.5*AB2(2)
      A(N-I-1) = AB1(0) - 0.5*AB2(1)
      DØ 5 J=1,I
      AB1(J) = AB2(J)
    5 CØNTINUE
    3 CØNTINUE
      A(N-1) = AB1(0)
      A(N)   = AB1(1)
      RETURN
      END
```

Although Program 8-7 is quite limited in its usefulness (since one needs to already have a polynomial represenation for the function), most of the literature on the Chebyshev series has evolved around the concept of using the Chebyshev polynomials to "telescope" a power series. This concept has been quite clearly discussed by Prager (Ref. 79), Golden (Ref. 47), Arden (Ref. 48), Lanzcos (Ref. 1 and Ref. 36), Minnick (Ref. 44), Spielberg (Ref. 42), and other authors whose papers are listed in the references for this Section. Often times the polynomial which one starts with is the Taylor's series and then the process of "telescoping" permits one to obtain a power series containing fewer terms which can be used to compute the function to the same accuracy over the particular range of interest.

The telescoping process has attracted so much attention in the literature of the recent past that the important concepts associated with the idea of trigonometrical interpolation (which were stressed by Lanzcos (Ref. 1)) have not received sufficient emphasis. In Eq. (8-8) we showed that the ccefficients in the expansion in a series of Chebyshev polynomials could be obtained by a process of numerical integration.

The most promising approach to the determination of of the A_n in Eq. (8-34) is associated with an orthogonality property of the Fourier series which, as Lanzcos has emphasized, is peculiar to the Fourier series when one examines the Fourier series as being an example of expansions in terms of orthogonal functions. We refer the reader to the paper by Lanzcos (Ref. 1) and the discussion by Clenshaw (Ref. 11) for more details. the result which we want to employ is the following:

$$NA_r = f(x_0)+2f(x_1)T_1(x_r)+\cdots+2f(x_{N-1})T_{N-1}(x_r)+f(x_N)T_N(x_r) \quad (8\text{-}35)$$

where $r = 0,1,2,\ldots,N$, and the arguments x_j appearing in $f(x_j)$

are defined by

$$x_j = \cos(j\pi/N) \tag{8-36a}$$

and the arguments x_r of the Chebyshev polynomials $T_j(x_r)$ are defined by

$$x_r = \cos(r\pi/N) \tag{8-36b}$$

In the discussion above we have made no distinction in notations between the A_n which appear in the infinite series of Eq. (8-7) and the A_n which appear in the truncated series of Eq. (8-12) or Eq. (8-34). The reader will find this discussed by Clenshaw in his tables (Ref. 11) and in a recent paper (Ref. 3). Further study of the relationships are needed, but it will suffice for our purposes to point out that we are assured by certain aspects of the theory of Fourier series that as $N \to \infty$ that the A_n which are computed on the basis of Eq. (8-35) will converge to the A_n employed in Eq. (8-7) and defined by an integral in Eq. (8-8).

It is interesting to observe that the formula in Eq. (8-35) is itself a truncated Chebyshev series and hence it can be summed by use of the Clenshaw algorithm. All we need to do is to be able to compute the value of $f(x)$ at $N + 1$ points x_j defined by Eq. (8-36a). This aspect of the determination of the A_n emphasizes the interpolating nature of the truncated series of Chebyshev polynomials. An ALGOL 60 procedure for the calculation of these coefficients was given by Schwarz (Ref. 29) in his tutorial paper on this programming language. This is quite a sophisticated program in that the degree of approximation which the user feeds in as an input to the ALGOL procedure determines the degree N of the truncated Chebyshev series, and a random number generator is employed to check that the resulting series does indeed provide the desired interpolating polynomial.

In this discussion, we will be content with the much simpler program for the evaluation of Eq. (8-35) as given in Program 8-8.

Program 8-8

```
      SUBRØUTINE CLNSHW(FUNCTN,A,B,N,CØEF)
      DØUBLE PRECISIØN A,B, CØEF(N),BPA,BMA,DFLØAT,DCØS,PI,PIØN,
     A       ZERØ,ØNE,TWØ,X(100),T(100),E(100),F(100)
      DATA PI,ZERØ,ØNE,TWØ/3.14159265358979 3D0,0.0D0,1.0D0,2.0D0/
  199 FØRMAT(37H SUBRØUTINE CLNSHW CANNØT HANDLE A = ,D25.16,
     A       5H B = ,D25.16,5H N = ,I4)
      DFLØAT(KK) = DBLE(FLØAT(KK))
      IF(N.LT.2) GØ TØ 98
      IF(N.GT.98) GØ TØ 98
      IF(DABS(A-B).LT.(1.0D-8)) GØ TØ 98
      BPA = B + A
      BMA = B - A
      PIØN = PI/DFLØAT(N-1)
      DØ 1 I=1,N
      T(I) = DCØS(PIØN*DFLØAT(I-1))
      X(I) = (BPA + BMA*T(I))/TWØ
      CALL FUNCTN(X(I),E(I))
    1 CØNTINUE
      F(N+2) = ZERØ
      F(N+1) = ZERØ
      E(N)   = E(N)/TWØ
      DØ 2 K =1,N
      DØ 3 L =N,1,-1
      F(L) = TWØ*T(K)*F(L+1) - F(L+2) + E(L)
    3 CØNTINUE
      COEF(K) = (F(1) - F(3))/DFLØAT(N-1)
    2 CØNTINUE
      RETURN
   98 WRITE( ,199) A,B,N
      CALL EXIT
      END
```

There are a number of aspects of Program 8-8 which require discussion. In the argument list, FUNCTN refers to another SUBRØUTINE which it is assumed the reader will have provided. It must be of the form:

```
      SUBRØUTINE FUNCTN(ARG,VALUE)
      DØUBLE PRECISIØN ARG, VALUE
        .
        .
      VALUE = ...
      RETURN
      END
```

When the statement CALL FUNCTN(ARG,VALUE) appears the function that is served is that of obtaining a value for f(x) where the relation

between the mathematical quantities x and f(x) and the FORTRAN symbols are

$$x = \text{ARG}$$
$$f(x) = \text{VALUE}$$

The A and B which appear in Program 8-8 give the range of the variable x over which the function is to be approximated, i.e,

$$A < \text{ARG} < B$$

Therefore, if $A \neq -1$, $B \neq +1$, the user will have to arrange to use Eq. (8-4) to employ the coefficients obtained from the program. The coefficients A_n are contained in the array COEF(N), i.e.,

$$A_n = \text{COEF(I)}, \text{ with } n = (I-1), \text{ for } n = 0,1,2,\ldots,(N-1)$$

The appearance of the statement "DØ 3 L=N,1,-1" reveals that this program must be modified if the compiler employed does not permit a DØ-loop to proceed from "high" values to "low" values.

In Program 8-9 we give a test program which was used to test the algorithm in Program 8-8.

Program 8-9

```
      DØUBLE PRECISIØN COEF(100)
      EXTERNAL FUNC
  110 FØRMAT(10X,I4,D25.16)
      CALL CLNSHW(FUNC,-1.0D0,+1.0D0,18,COEF)
      WRITE(6,110) (I,COEF(I),I=1,18)
      CALL EXIT
      END
      SUBRØUTINE FUNC(ARG,VALUE)
      DØUBLE PRECISIØN ARG, VALUE, DEXP
      VALUE = DEXP(ARG)
      RETURN
      END
```

In Table 8-1 we present the output from the use of Programs 8-8 and 8-9. In the right hand column we present, for comparison, the Chebyshev coefficients for $f(x) = \exp(x)$ on $-1 < x < +1$ which are given by Clenshaw(Ref. 11).

Table 8-1

CHEBYSHEV COEFFICIENTS FOR EXP(X)

N	From Program 8-9	From Clenshaw (Ref.11)
1	.2532131755504014+01	.25321317555040166712 0+01
2	.1130318207984963+01	.11303182079849700544 2+01
3	.2714953395340757+00	.27149533953407656237 +00
4	.4433684984866365-01	.44336849848663804 95 -01
5	.5474240442093978-02	.54742404420937326 5 -02
6	.5429263119140001-03	.54292631191394375 -03
7	.4497732295421072-04	.44977322954295 15 -04
8	.3198436462465850-05	.31984364624019 9 -05
9	.1992124804986286-06	.19921248066728 -06
10	.1103677183631729-07	.1103677172552 -07
11	.5505895478103851-09	.55058960797 -09
12	.2497978294801374-10	.2497956617 -10
13	.1039116505259752-11	.103915223 -11
14	.4004639757059681-13	.3991263 -13
15	.1345329076898718-14	.142376 -14
16	.6530723674265623-16	.4741 -16
17	.0000000000000000	.148 -17
18	.0000000000000000	.4 -18

A study of the output from Program 8-9 with the more precise results printed alongside reveals that the initial results (N near 1) are accurate to about 15 decimal places, but that as N increases there is a growth of the error and by the time one reaches N = 14 that there is only one significant figure and that this is in the 14th decimal place. These results are quite encouraging and suggest that Eq. (8-35) can be quite useful for the determination of the coefficients in the Chebyshev series which is truncated as in Eq. (8-34).

8.7 The Future for Diffraction Theory

From the many references at the end of this section, the author has assembled a set of BLOCK DATA statements which contain (on punched cards) virtually all the Chebyshev coefficients which have been published to the present date. It was originally part

of the author's plan for this report that these would be contained in the Appendices. However, the compilation runs to many pages and the results are most useful when they can be read directly into an electronic computer. Therefore, the author plans to run further checks upon these listings (to insure their accuracy) and to then submit the data (in the form of a strip of magnetic tape) to the SHARE organization so that it will be unnecessary for programmers who have access to this tape to punch the data directly from the published tables.*

Before we discuss some possible applications of the Chebyshev series in diffraction theory, let us pause to point out a field which needs further investigation. The results in Eqs. (8-34) and (8-35) permit us to interpolate (using the values of the function at N+1 points) with efficiency and accuracy a function $f(x)$ defined on $-1 < x < +1$. In diffraction theory (and in many other fields) we meet the need to interpolate when the function is defined in the complex plane and therefore we find ourselves dealing with a function of two (or more) variables. The ideas advanced by Lanzcos could be extended to such problems by "brute force." For example, if $F(z) = F(x,y)$ is defined on $-1 < x < +1$, $-1 < y < +1$, we could write

$$F(x,y) = \frac{1}{2}f_0(y) + f_1(y)T_1(x) + f_2(y)T_2(x) + \cdots + \frac{1}{2}f_N(y)T_N(x)$$

where


$$f_r(y) = \frac{1}{2}A_0^r + A_1^r T_1(y) + A_2^r T_2(y) + \cdots + A_{M-1}^r T_{M-1}(y) + \frac{1}{2}A_M^r T_M(y)$$

*The questions related to the status of the "copyright" on these tables has also enterred the decision to omit these DATA BLOCKS from this report. However, since keypunching and verifying is relatively expensive, it is hoped that arrangements can be made whereby this file (consisting of several thousand IBM cards) can be shared with other scientists who employ the Chebyshev series.

where the coefficients A_p^r could be determined from a result derived from Eq. (8-35). We observe that this approach would require the knowledge of the value of the function F(x,y) at (N+1)(M+1) points in the domain of interest, i.e., the square $-1 < x < +1$, $-1 < y < +1$. It is quite conceiveable that as we push foward in the field of the constructive theory of functions that we will find means to interpolate over the rectangle which require many fewer points than the (N+1)(M+1) that the above approach requires. The interpolation problem bears some resemblance to the problem of choosing points for the evaluation of multiple integrals. This has begun to be a field which is attracting considerable interest. One author recently pointed out that the use of a Gauss 3-point formula for an n-dimensional integral would require 3^n points, whereas comparable accuracy can be obtained with about $2n^2$ points. Results such as this for the interpolation of functions of more than one variable are badly needed. However, the problem is sufficiently difficult that we may have to wait years for significant advances. In the meantime, the outstanding paper on interpolation of one-dimensional functions which was written in 1938 by Lanzcos (Ref. 1) has only now begun to be properly appreciated by people who face applied problems.

The outstanding advantages which will accrue to diffraction theorists with the further development of the Chebyshev series representations for special functions will be demonstrated by considering several types of problems. Let us start with the Airy functions. The behavior of the Airy functions Ai(x) and Bi(x) are depicted in Fig. 8-1. An essential point to be borne in mind when seeking representations in terms of truncated Chebyshev polynomials is that the coefficients in the expansion can be expected to decrease rapidly with increasing order only if the function which is to be represented is "smooth."

If we recall that we know Taylor series (See Eq. (B-5)) for these functions, we can recognize the possibility of using the terms in



s	a_s	$Ai'(-a_s)$
1	2·33810 741	+0·70121 082
2	4·08794 944	− ·80311 137
3	5·52055 983	+ ·86520 403
4	6·78670 809	− ·91085 074

Fig. 8-1 Behavior of the Airy Functions

these series in an algorithm (such as Program 8-7) to obtain efficient representations of the form

$$Ai(x) = \frac{1}{2} A_0 + A_1 T_1(x) + \cdots + A_N T_N(x) \qquad (8\text{-}37a)$$

$$Bi(x) = \frac{1}{2} B_0 + B_1 T_1(x) + \cdots + B_M T_M(x) \qquad (8\text{-}37b)$$

where N need not be the same as M, and where $-1 < x < +1$. It might even be worthwhile to expand the range to $-2 < x < +2$, and employ series of the form

$$g(x) = \frac{1}{2} C_0 + C_1 T_1\left(\frac{x}{2}\right) + \cdots + C_K T_K\left(\frac{x}{2}\right) \qquad (8\text{-}38)$$

The use of either Eq. (8-37) or Eq. (8-38) would be merely a conventional use of the Chebyshev polynomials to "telescope" a power

series. However, if one expected to compute the Airy functions many times on an electronic computer, it would be economically sound to seek these more efficient series.

Before leaving the subject of the Airy functions near the origin, the author would like to point out that it may prove to be advantageous to consider the form of Eq. (B-5) and express Ai(x) and Bi(x) in the form

$$\mathrm{Ai}(x) = \alpha f_1(\xi) - \beta x f_2(\xi) \quad , \quad \mathrm{Bi}(x) = \sqrt{3}\,[\alpha f_1(\xi) + \beta x f_2(\xi)] \qquad (8\text{-}39)$$

where

$$\xi = x^3/A^3$$

and $f_1(\xi)$ and $f_2(\xi)$ are expressed in the form of Chebyshev series

$$f_j(\xi) = \frac{1}{2} E_0 + E_1 T_1(\xi) + \cdots + E_N T_N(\xi) \quad , \quad j = 1,2$$

and the expansion would be employed for the range $-A < x < +A$. The author hopes that this alternative causes the reader to see that the employment of the Chebyshev representations is not purely "mechanical." One will often find that some preliminary analytical work will suggest an argument to be employed in the Chebyshev polynomial which is most appropriate for the problem under consideration.

Let us turn now to an aspect of the Chebyshev representation of the Airy functions which is more than just "financially economic" in the sense that we will consider representations which cannot be "telescoped" because they start from asymptotic series. Miller has shown (See Eq. (B-3)) that the Airy functions can be expressed in terms of certain auxiliary functions F(x) and $\chi(x)$ when x is large and negative. We recall that the forms are

$$\mathrm{Ai}(-\eta) = F(-\eta)\sin[\chi(-\eta)] \quad , \quad \mathrm{Bi}(-\eta) = F(-\eta)\cos[\chi(-\eta)] \qquad (8\text{-}40)$$

where asymptotic expansions for $F(-\eta)$ and $\chi(-\eta)$ are given in Eq.

(6-9). Suppose we seek a representation for $A < \eta < \infty$. Since this is a "one-sided" range, we would probably choose to use the shifted (or T-star) polynomials $T_n^*(x)$. An inspection of Eq. (6-9) suggests that we express $F(-\eta)$ and $\chi(-\eta)$ in the forms

$$F^2(-\eta) = \frac{f_1(\xi)}{\pi\sqrt{\eta}} \quad , \quad \chi(-\eta) - \frac{\pi}{4} = \frac{2}{3}\sqrt{\eta^3}\, f_2(\xi) \, , \quad \xi = A^3/\eta^3 \qquad (8\text{-}41)$$

where $f_1(\xi)$ and $f_2(\xi)$ are to be expressed in the form of series of shifted Chebyshev polynomials

$$f(x) = \frac{1}{2} A_0 + A_1 T_1^*(x) + A_2 T_2^*(x) + \cdots + A_N T_N^*(x) \qquad (8\text{-}42a)$$

where the counterpart of Eq. (8-35) is

$$NA_r = f(x_0) + 2f(x_1)T_1^*(x_r) + \cdots + 2f(x_{N-1})T_{N-1}^*(x_r) + f(x_N)T_N^*(x_r) \qquad (8\text{-}42b)$$

and the x_r and x_j ($j = 0,1,2,\ldots,N$) are given by

$$x_r = \cos\left(\frac{j\pi}{2N}\right) \quad , \quad x_j = \cos\left(\frac{r\pi}{2N}\right) \, , \quad \left.\begin{matrix} r \\ j \end{matrix}\right\} = 0,1,2,\ldots,N \qquad (8\text{-}42c)$$

In arriving at the choice of this form of expansion we have been guided by the asymptotic theory which led to Eq. (6-9), but unless we take A extremely large (in which case we can evaluate the asymtotic series in Eq. (6-9) by considering only a few decreasing terms) we cannot convert the series of inverse powers in Eq. (6-9) into series of Chebyshev polynomials by means of Program 8-7. If, however, we use Eq. (6-9) only for those values of η for which the asymptotic series yields the desired accuracy, and then by "brute force" (such as by numerical integration from large values of η to smaller values, or vice versa) obtain the values of $F(-\eta)$ and $\chi(-\eta)$ at $(N + 1)$ points such as those required in Eq. (8-35) and Program 8-8, we can determine a set of values of A_n for use in Eq. (8-42). In this way we will obtain a polynomial which

possesses many of the features of the asymptotic series, but which will possess the advantage of being able to be used down to very small values of the variable (η in Eq. (8-41)) which were completely inconceiveable with the asymptotic series. The procedures suggested above depend upon the analytical study of the asymptotic behavior to provide the motivation for which portion of the function will be sufficiently "smooth" to possess a rapidly decreasing set of coefficients A_n in the Chebyshev series. Furthermore, we have used the asymptotic theory to provide a motivation for the selection of a "argument" to employ in the Chebyshev polynomial.*

A similar set of relations can be developed for the computation of the Airy functions for large, positive values of the argument by using the relations in Eqs. (6-52a) and (6-52b). However, this would so closely parallel the result in Eq. (8-41) that we will choose instead to consider the asymptotic expansions

$$\mathrm{Ai}(x) \sim \frac{1}{2}\pi^{-1/2}x^{-1/4}e^{-\xi}\left(1 - \frac{3\cdot5}{1!\ 216}\frac{1}{\xi} + \frac{5\cdot7\cdot9\cdot11}{2!(216)^2}\frac{1}{\xi^2} - \frac{7\cdot9\cdot11\cdot13\cdot15\cdot17}{3!\,(216)^3}\frac{1}{\xi^3} + \cdots\right)$$

$$\mathrm{Bi}(x) \sim \pi^{-1/2}x^{-1/4}e^{\xi}\left(1 + \frac{3\cdot5}{1!\ 216}\frac{1}{\xi} + \frac{5\cdot7\cdot9\cdot11}{2!\,(216)^2}\frac{1}{\xi^2} + \frac{7\cdot9\cdot11\cdot13\cdot15\cdot17}{3!\,(216)^3}\frac{1}{\xi^3} + \cdots\right)$$

where

$$\xi = \frac{2}{3}\sqrt{x^3}$$

*This field is so new that there has not been accumulated (in the literature or in the experience of the present author) enough experience to be certain that the natural argument suggested by the asymptotic theory is the "best" argument to employ. The reader who undertakes work along the lines suggested above should enter this field with an "open mind" and be prepared to "experiment."

to suggest that representations be sought for Ai(x) and Bi(x) which are of the form

$$2\sqrt{\pi\sqrt{x}}\,\mathrm{Ai}(x) = \exp(-\xi) f_1(\eta)\ , \quad \sqrt{\pi\sqrt{x}}\,\mathrm{Bi}(x) = \exp(+\xi) f_2(\eta) \qquad (8\text{-}43)$$

where

$$\eta = \sqrt{(A/x)^3}\ , \quad A < x < \infty$$

and the functions $f_1(\eta)$ and $f_2(\eta)$ are series of the form given in Eq. (8-42)

We can look foward to finding many expansions similar to those above in the literature of the next several years. The reader's attention is directed to the papers of Wimp (Ref. 67) and of Luke and Wimp (Ref. 68) as being typical of some of the work which has been done to date.

The present author has begun work on the obtaining of Chebyshev polynomial representations for the roots $t_s(q)$ of the Airy function combination

$$w_1'(t_s) - q\, w_1(t_s) = 0$$

where arg q is held fixed (for certain values of arg q which occur in practical problems) and the Chebyshev expansions are with respect to the modulus of q.

Some more interesting applications, however, arise in the case of some of the diffraction functions. Consider the case of the asymptotic expansion for the current distribution function $g(\xi)$.

$$g(\xi) \xrightarrow[\xi \to -\infty]{} 2\exp\left(-i\frac{\xi^3}{3}\right)\left\{1 + i\frac{1}{4\xi^3} - \frac{1}{\xi^6} - i\frac{469}{64}\frac{1}{\xi^9} + \frac{5005}{64}\frac{1}{\xi^{12}} + i\frac{11\,22121}{1024}\frac{1}{\xi^{15}}\right.$$

$$\left. - \frac{24\,33368}{128}\frac{1}{\xi^{18}} - i\frac{16102\,89919}{4096}\frac{1}{\xi^{21}} + \ldots\right\} \qquad (8\text{-}44)$$

We know a wide variety of representations for $g(\xi)$. If we use

one of the following representations

$$g(\xi) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \frac{\exp(i\xi t)}{w_1'(t)}\, dt = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\exp(i\xi t)}{Bi'(t) + i\,Ai'(t)}\, dt = 2\sqrt{\pi}\, i \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s^o)}{t_s^o\, w_1(t_s^o)}$$

$$= \sum_{s=1}^{\infty} \frac{\exp\left(-\frac{\sqrt{3}-i}{2}\beta_s \xi\right)}{\beta_s\, Ai(-\beta_s)} = -\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\exp\left(\frac{\sqrt{3}-i}{2} p\xi\right)}{Ai'(p)}\, dp$$

$$= \frac{1}{\pi} \int_{0}^{\infty} \frac{\exp\left(-\frac{\sqrt{3}+i}{2}\xi t\right)}{Bi'(t) - i\,Ai'(t)}\, dt + \frac{1}{\pi} \int_{0}^{\infty} \frac{\exp(i\xi t)}{Bi'(t) + i\,Ai'(t)}\, dt = \sum_{n=o}^{\infty} g^{(n)}(0)\, \frac{\xi^n}{n!} \qquad (8\text{-}45)$$

or make use of the fact that $g(\xi)$ satisfies the integral equation

$$g(\xi) = 2\exp\left(-i\,\tfrac{1}{3}\xi^3\right) - \frac{1}{2}\,\frac{\exp\left(-i\frac{\pi}{4}\right)}{2\sqrt{\pi}} \int_{-\infty}^{\xi} g(x) \exp\left[-i\,\tfrac{1}{12}(\xi - x)^3\right] \sqrt{\xi - x}\, dx \qquad (8\text{-}46)$$

we can devise "brute force" means of evaluating $g(\xi)$ for all values of ξ. We can then use the interpolation properties of the Chebyshev polynomials to find a representation in terms of Chebyshev polynomials which will be more convenient (as well as more economical) to employ when further values of $g(\xi)$ are required for arbitrary values of ξ.

Let us start with the case in which ξ is large and negative. Although it might turn out to be convenient to find Chebyshev polynomial representations for the real and the imaginary parts of the quantity in braces in Eq. (8-44), the present author suggests that Eq. (8-44) be replaced by

$$g(\xi) = 2\,\exp(-i\tfrac{1}{3}\xi^3) M(\eta) \exp\left[-i\frac{P(\eta)}{4(-\xi)^3}\right] \qquad (8\text{-}47)$$

where $M(\eta)$ is an amplitude function and $P(\eta)$ is a "modified" (since it has been divided by $4(-\xi)^3$) phase function. Both $M(\eta)$ and $P(\eta)$ tend to unity as $\xi \to -\infty$. In order to find a representation for these functions for ξ in the range $-\infty < \xi \leq -A < 0$, it is proposed that one take η to be defined by

$$\eta = (-A/\xi)^3$$

and seek representations for $M(\eta)$ and $P(\eta)$ in the form of the series contained in Eq. (8-42).

We can draw many parallels between methods to be employed for the functions of which $g(\xi)$ is an example, and the methods which we have suggested for the representation of the Airy functions. The form for $g(\xi)$ in Eq. (8-47) is the analog of Eq. (8-40). Near the origin, we can use an analog of Eq. (8-37) and seek to represent $g(\xi)$ in the form of a Chebyshev series

$$g(\xi) = \frac{1}{2} A_0 + A_1 T_1(x) + A_2 T_2(x) + \cdots + A_N T_N(x) , \quad x = \xi/A$$

where $-A < \xi < +A$, and the result in Eq. (8-35) or the method of telescoping a power series can be used to determine the A_n.

For large positive values of ξ we know that

$$g(\xi) \xrightarrow[\xi \to \infty]{} \frac{\exp(-\frac{\sqrt{3} - i}{2} \beta_1 \xi)}{\beta_1 \mathrm{Ai}(-\beta_1)} = G_\infty(\xi) \qquad (8\text{-}48)$$

where β_1 is the root of $\mathrm{Ai}'(-\beta_1) = 0$ which has the value $\beta_1 = 1.101879\ldots$. For smaller values of ξ, but $\xi \geq 0$, we have to replace the approximation in Eq. (8-48) by a sum over a finite number of the roots β_s. If $g(\xi)$ is to be computed very frequently, it might be advantageous to express $g(\xi)$ in the form

$$g(\xi) = G_\infty(\xi) M(\eta) \exp[iP(\eta)] \qquad (8\text{-}49)$$

where $M(x)$ and $P(x)$ are expressed in the form of series of the type defined in Eq. (8-42) and the variable η is defined by

$\eta = (A/\xi)$ when the range of ξ is $0 < A < \xi < \infty$.

The present author believes that the development of methods such as those described above can transform diffraction theory from an interesting theoretical exercise into a practical tool for engineers and scientists who are concerned with the problems that arise in connection with the diffraction and scattering of waves by convex surfaces. Most of the theory leads to representations which permit the evaluation of the functions that occur in only limited regions. For example, it is characteristic of most of these problems that there is a "gap" between the largest values of a variable for which it is practical to employ a power series representation and the smallest values of the variable for which the asymptotic series can be employed. If this "gap" lies in the region for which engineering results are required one is forced to considering graphs of the functions and the use of "artistic license" in drawing the curves so as to close the "gap." However, the use of such an approach requires considerable manual manipulation of graphs and "French curves." If a "brute force" method is used to close the "gap" in order to determine the coefficients in the Chebyshev polynomial representations, one will then be able to handle the functions in a manner which will permit the use of an electronic computer to assist in the practical applications.*

---

*The author has been seeking to determine "brute force" methods to close the "gap" by employing methods of numerical analysis in order to obtain relatively accurate representations for the diffraction functions. However, in a practical problem when one is faced with the need to obtain values in the "gap" to employ in either Eq. (8-35) or Eq. (8-42), it should be borne in mind that an "educated guess" can sometimes be made by drawing a curve of the function and attempting to see if a "French curve" will provide an approximate means of bridging the "gap."

REFERENCES FOR SECTION 8

1. C. Lanzcos, "Trigonometrical Interpolation of Empirical and Analytical Functions," J. Math. Physics, Vol. 17, 1938, pp. 123-199. (The polynomials denoted by $T_n(x)$ in this paper are the same as those defined by $T_n^*(x)$ in later papers.)

2. C. Hastings, Approximations for Digital Computers, Princeton, Princeton Univ. Press, 1955

3. C. W. Clenshaw, "A Comparison of 'Best' Polynomial Approximations with Truncated Chebyshev Series Expansions," J. SIAM Numer. Anal., Vol. 1, 1964, pp. 26-37

4. David Taylor Model Basin, The Approximation of Differentiable Functions by Polynomials, by F. D. Murnaghan and J. W. Wrench, Report 1175, Washington, D. C., May, 1960

5. F. D. Murnaghan and J. W. Wrench, "The Determination of the Chebyshev Approximating Polynomial for a Differentiable Function," Math. Tables, Vol. 13, July-Aug.,1959, pp. 185-193

6. E. T. Copson, Asymptotic Expansions, Cambridge, Cambridge University Press, 1965. (This is an extension of the Admiralty Computing Service report entitled "The Asymptotic Expansion of a Function Defined by a Definite Integral or Contour Integral" which has received widespread distribution since it was first issued in 1943.)

7. A. Erdélyi, Asymptotic Expansions, New York, Dover, 1956

8. H. Jeffreys, Asymptotic Approximations, Oxford, Oxford Univ. Press, 1962

9. J. Heading, An Introduction to Phase-Integral Methods, London, Methuen, 1962

10. National Bureau of Standards Applied Mathematics Series, No. 9, Table of Chebyshev Polynomials, Washington, U.S. Government Printing Office, 1952. (This report contains an Introduction written by C. Lanzcos which is indispensable to anyone doing research involving Chebyshev polynomials.)

11. C. W. Clenshaw, Chebyshev Series for Mathematical Functions, National Physical Laboratory Mathematical Tables, Vol. 5, London, Her Majesty's Stationery Office, 1962

12. C. W. Clenshaw, "A Note on the Summation of Chebyshev Series," Math. Tables, Vol. 9, Oct., 1955, pp. 118-120

13. C. W. Clenshaw, "Polynomial Approximations to Elementary Functions," Math. Tables, Vol. 8, July, 1954, pp. 143-147

14. J.W. Clenshaw, G. F. Miller, and M. Woodger, "Algorithms for Special Functions I," Numerische Math., Vol. 4, Pt. 5, 1963, pp. 403-419. (Contains a collection of ALGOL procedures for the evaluation of several special functions.)

15. C. W. Clenshaw, "The Numerical Solution of Linear Differential Equations in Chebyshev Series," Proc. Cambridge Phil. Soc., Vol. 53, Jan., 1957, pp. 134-149

16. C. W. Clenshaw, "The Numerical Solution of Linear Differential Equations in Chebyshev Series," Symposium, Provisional International Computation Centre, Basel, Birkhäuser, 1960

17. C. W. Clenshaw and D. Elliott, "A Numerical Treatment of the Orr-Sommerfeld Equation in the Case of a Laminar Jet," Quart. J. Mech. Appl. Math., Vol. 13, Aug., 1960, pp. 300-313

18. C. W. Clenshaw and F. W. J. Olver, "The Use of Economized Polynomials in Mathematical Tables," Proc. Cambridge Phil. Soc., Vol. 51, Oct., 1955, pp. 614-628

19. C. W. Clenshaw and H. J. Norton, "The Solution of Nonlinear Ordinary Differential Equations in Chebyshev Series," Computer J., Vol. 6, April, 1963, pp. 88-92

20. C. W. Clenshaw, "Curve Fitting with a Digital Computer," Computer J., Vol. 2, Oct. 1959, pp. 170-172

21. C. W. Clenshaw and A. R. Curtis, "A Method for Numerical Integration on an Automatic Computer," Numerische Math., Vol. 2, 1960, pp. 197-205

22. J. P. Imhof, "On the Method of Numerical Integration of Clenshaw and Curtis," Numerische Math., Vol. 5, Pt. 2, 1963, pp. 138-141

23. S. Filippi, "Angenäherte Tschebyscheff-Approximation einer Stammfunktion--eine Modifikation des Verfahrens von Clenshaw und Curtis," Numerische Math., Vol. 6, Pt. 4, 1964, pp. 320-328

24. J. C. P. Miller, "Two Numerical Applications of Chebyshev Polynomials," Proc. Roy. Soc. (Edinburgh), Vol. 62, 1943-9, pp. 204-210

25. B. van der Pol and T. J. Weijers, "Tchebycheff Polynomials and Their Relation to Circular Functions, Bessel Functions, and Lissajous Figures," Physica, Vol. , pp. -

26. G. Shortley, "Use of Tschebyscheff-Polynomial Operators in the Numerical Solution of Boundary-Value Problems," J. Appl. Physics, Vol. 24, April, 1953, pp. 392-396

27. J. D. Beasley, "A Note on the Rearrangement of Chebyshev Series," Computer J., Vol. 8, Oct., 1965, pp. 278-279

28. G. A. Chisnall, "A Modified Chebyshev-Everett Interpolation Formula," Math. Tables, Vol. 10, Apr., 1956, pp. 66-73

29. H. R. Schwarz, "Am Introduction to ALGOL," Comm. Assoc. Comput. Mach., Vol. 5, Feb., 1962, pp. 82-95. (Contains an ALGOL procedure for evaluating Chebyshev coefficients for a function defined by another procedure.)

30. W. Gautschi, "Numerical Integration of Ordinary Differential Equations Based Upon Trigonometric Polynomials," Numerische Math., Vol. 3, Pt. 5, 1961, pp. 381-397

31. L. Fox, "Chebyshev Methods for Ordinary Differential Equations," Computer J., Vol. 4, Oct., 1962, pp. 318-331

32. L. Fox, "Chebyshev Solution of Ordinary Differential Equations," Numerical Solution of Ordinary and Partial Differential Equations, L. Fox, ed., Reading, Mass., Addison-Wesley, 1962

33. L. Fox, "Chebyshev Approximation," Numerical Solution of Ordinary and Partial Differential Equations, L. Fox, ed., Reading, Mass., Addison-Wesley, 1962

34. R. W. Hamming, "Chebyshev Polynomials," Numerical Methods for Scientists and Engineers, New York, McGraw-Hill, 1962, pp. 249-264

35. J. Todd, "The Chebyshev Theory," Introduction to the Constructive Theory of Functions, Basel, Birkhauser, 1963, pp. 24-35

36. C. Lanczos, "Power Expansions," Applied Analysis, Englewood Cliffs, N.J., Prentice-Hall, 1956, pp. 438-507

37. A. Ralston, "Functional Approximation-Minimum Maximum Error Techniques," A First Course in Numerical Analysis, New York, McGraw-Hill, 1965, pp. 271-317

38. A. Ralston, "On Economization of Rational Functions," J. Assoc. Comput. Mach., Vol. 10, July, 1963, pp. 278-282

39. K. G. Kogbetliantz, "Generation of Special Functions," Mathematical Methods for Digital Computers, A. Ralston and H. F. Wilf, eds., New York, John Wiley, 1960, pp. 7-35

40. H. J. Maehly, "Methods for Fitting Rational Approximations, Part I," J. Assoc. Comput. Mach., Vol. 7, April, 1960, pp. 150-162

41. H. J. Maehly,"Methods for Fitting Rational Approximations, Parts II and III," J. Assoc. Comput. Mach., Vol. 10, July, 1963, pp. 257-277. (Prepared posthumously by C. Witzgall.)

42. K. Spielberg, "Representations of Power Series in Terms of Polynomials, Rational Approximations and Continued Fractions," J. Assoc. Comput. Mach., Vol. 8, Oct., 1961, pp. 613-627

43. E. Stiefel, "Methods - Old and New - For Solving the Tchebycheff Approximation Problem," J. SIAM Numer. Anal., Vol. 1, 1964, pp. 164-176

44. R. C. Minnick, "Tchebysheff Approximations for Power Series," J. Assoc. Comput. Mach., Vol. 4, Oct., 1957, pp. 487-504

45. R. De Vogelaere, "Remarks on the Paper 'Tchebysheff Approximations for Power Series,'" J. Assoc. Comput. Mach., Vol. 6, Jan.,1959, pp. 111-114

46. G.N. Lance, "Evaluation of Functions," Numerical Methods for High Speed Computers, London, Iliffe, 1960, pp. 13-50

47. J. T. Golden, "Chebyshev Approximations," FORTRAN IV Programming and Computing, Englewood Cliffs, N.J., Prentice-Hall, 1965, pp. 120-124. (Presents FORTRAN IV coding for "economizing" a power series.)

48. B. W. Arden, "Chebyshev Economization," An Introduction to Digital Computing, Reading, Mass., Addison-Wesley, 1963, pp. 248-257. (The present author is of the opinion that Arden's algorithm presented in Fig. 15-13 is in error since it apparently will not convert a Chebyshev polynomial series of degree greater than 5 to a power series with the correct coefficients.)

49. H. C. Thacher, "Conversion of a Power Series to a Series of Chebyshev Polynomials," Comm. Assoc. Comput. Mach., Vol. 7, March, 1964, pp. 181-182

50. R. Barakat, "Evaluation of the Incomplete Gamma Function of Imaginary Argument by Chebyshev Polynomials," Math. Comp., Vol. 16, Jan., 1961, pp. 7-11

51. R. Barakat and A. Houston, "Chebyshev Expansion of the Sine and Cosine Integrals," J. Math. Physics, Vol. 42, Dec., 1963, pp. 331-333

52. D. Elliott, "The Numerical Solution of Integral Equations Using Chebyshev Polynomials," J. Australian Math. Soc., Vol. 1, 1960, pp. 344-356

53. D. Elliott, "A Chebyshev Series Method for the Numerical Solution of Fredholm Integral Equations," Computer J., Vol. 6, April, 1963, pp. 102-111

54. D. Elliott, "Truncation Errors in Two Chebyshev Series Approximations," Math. Comp., Vol. 19, April, 1965, pp. 234-248

55. D. Elliott and G. Szekeres, "Some Estimates of the Coefficients in the Chebyshev Series Expansion of a Function," Math. Comp., Vol. 19, Jan., 1965, pp. 25-32

56. F. Briones, "Calcul des Coefficients des Developpements de $\int_0^x f(y)\, y^{-N} dy$ et de f(x)/(x-b) en Polynomes de Techebycheff," Chiffres, Vol. 7, No. 3, 1964, pp. 177-184

57. A. J. W. Duijvestijn and A. J. Dekkers, "Chebyshev Approximations of Some Transcendental Functions for Use in Digital Computing," Philips Res. Rep., Vol. 16, April, 1961, pp. 145-174

58. F. J. Smith, "An Algorithm for Summing Orthogonal Polynomial Series and Their Derivatives With Applications to Curve-Fitting and Interpolation," Math. Comp., Vol. 19, Jan., 1965, pp. 33-36

59. G. E. Forsythe, "Generation and Use of Orthogonal Polynomials for Data Fitting with a Digital Computer," J. Soc. Indust. Appl. Math., Vol. 5, June, 1957, pp. 74-88

60. Z. Kopal, "Trigonometric Interpolation. Tchebysheff Polynomials and Their Use for Optimum-Interval Interpolation," Numerical Analysis, 2nd Ed., London, Chapman and Hall, 1961, pp. 523-542

61. R. E. Scraton, "The Solution of Linear Differential Equations in Chebyshev Series," Computer J., Vol. 8, April, 1965, pp. 57-61

62. W. J. Cody, "Chebyshev Approximations for the Complete Elliptic Integrals K and E," Math. Comp., Vol. 19, Jan., 1965, pp. 105-112

63. W. J. Cody, "Chebyshev Polynomial Expansions of Complete Elliptic Integrals," Math. Comp., Vol. 19, April, 1965, pp. 249-259

64. F. D. Burgoyne, "Polynomial Approximations to $I_0(x)$, $I_1(x)$ and Related Functions," Math. Comp., Vol. 17, Oct., 1963, pp. 497-498

65. J. Wimp, "Polynomial Approximations to Integral Transforms," Math. Comp., Vol. 15, April, 1961, pp. 174-178

66. J. F. Fields and J. Wimp, "Expansion of Hypergeometric Functions in Hypergeometric Functions," Math. Comp., Vol. 15, Oct., 1961, pp. 390-395

67. J. Wimp, "Polynomial Expansions of Bessel Functions and Some Associated Functions," Math. Comp., Vol. 16, Oct., 1961, pp. 446-458

68. Y.L. Luke and J. Wimp, "Jacobi Polynomial Expansions of a Generalized Hypergeometric Function over a Semi-Infinite Ray," Math. Comp., Vol. 17, Oct., 1963, pp. 395-404

69. A.M. Arthurs and R. Carroll, "Expansion of Spherical Bessel Functions in a Series of Chebyshev Polynomials," Math. Comp., Vol. 15, April, 1961, pp. 159-162

70. D. G. Hummer,"Expansion of Dawson's Function in a Series of Chebyshev Polynomials," Math. Comp., Vol. 18, April, 1964, pp. 317-319

71. E. L. Stiefel, "Approximations," An Introduction to Numerical Mathematics, New York, Academic Press, 1963, pp. 218-253

72. L. Lapidus, "Further Methods of Approximation," Digital Computations for Chemical Engineers, New York, McGraw-Hill, 1962, pp. 323-368

73. C. Lanzcos, "Global Integration by Chebyshev Polynomials," Linear Differential Operators, Princeton, N.J., Van Nostrand, 1961, pp. 536-540

74. C.E. Froberg," Approximation," Introduction to Numerical Analysis, Reading, Mass., Addison-Wesley, 1965, pp. 277-299

75. E. L. Stiefel, "Numerical Methods of Tchebycheff Approximation," On Numerical Approximation, R. E. Langer, ed., Madison, Wisc., Univ. Wisconsin Press, 1959, pp. 217-232

76. L. Fox, "Minimax Methods in Table Construction," On Numerical Approximation, R. E. Langer, ed., Madison, Wisc., Univ. Wisconsin, Press, 1959, pp. 233-244

77. J. C. P. Miller, "Extremal Approximation-A Summary," On Numerical Approximation, R. E. Langer, ed., Madison, Wisc., Univ. Wisconsin Press, 1959, pp. 329-340

78. L. Fox, "Chebyshev Polynomials," The Use and Construction of Mathematical Tables, National Physical Laboratory Mathematical Tables, Vol. 1, London, Her Majesty's Stationery Office, 1956, pp. 12-14

79. W. Prager, "Computing with Polynomials," Introduction to Basic FORTRAN Programming and Numerical Methods, New York, Blaisdell, 1965, pp. 61-79. (Program VII.5.1 performs the transformation of a Chebyshev polynomial expansion into a polynomial.)

80. Staff of the National Physical Laboratory, "Chebyshev Series," Modern Computing Methods, London, Her Majesty's Stationery Office, 1961, pp. 71-79, 88-90

81 G. Meinardus, Approximation von Funktionen und ihre numerische Behandlung, Berlin, Springer, 1964. (The 7-page list of references in this book provide a good starting point for a discovery of many papers which deal with the "best fit" type of minimum maximum approximation.)

Appendix A

FORTRAN AND NUMERICAL ANALYSIS

At the present time, numerical methods are assuming a role of increasing importance in applied science; the actual, rather than the theoretical, production of a solution is of paramount interest. Diffraction theory provides many interesting challenges to the numerical analyst. Although very few of the papers which have come to the attention of the present author have actually been written by people engaged in studies in diffraction theory, the author finds that almost every new issue of journals devoted to numerical analysis contains papers which are of direct interest to those who face problems in diffraction theory.

We are now at the threshold of an era in which the electronic digital computer is becoming as common an engineering tool as the slide rule. We are reaching the point where even relatively simple problems which are traditionally performed by persons operating manual-type desk calculators can be more economically (as well as more accurately---in the sense of the elimination of the chances of human error in transcribing numbers from computation sheets to the keyboard and from the registers to the computation sheets) executed on the electronic computer. The problem of setting up a set of computation sheets for the hand-computer is hardly more difficult than the devising of the FORTRAN program for the electronic computer. Much of the inertia which causes scientists and engineers to continue the traditional hand-computations is due to the fact that although the task might require only 30 seconds (perhaps a charge of three to five dollars for computer time), there is present the delays in getting the IBM cards punched by the "keypunch pool" and the "turn around" time in getting the problem onto and off the electronic computer.

We are also at the threshold of a new era in which even the use of the electronic computer is being revolutionized through the introduction of "time-sharing" concepts and other innovations. These new advances will soon make the use of the manually-operated desk machine less attractive.

The present author is a strong advocate of the "open shop" operation of computing facilities; that is, he advocates that the programming be done by the scientist or engineer who devises the theoretical formulae (or by one of their close assistants) instead of being "farmed out" to a "closed shop" facility where specialists in programming and numerical analysis translate the formulae into output in the form of tables or graphs. The "do-it-yourself" programmer will build up a background of experience in the particular types of calculations which his field of research is centered around and he will be able to draw upon his experience with previous computations to devise methods for the computation of new problems. Furthermore, since new problems are often only relatively small perturbations of problems which have already been considered, the engineer and scientist will begin to develop a nucleus of programs which often require but minor changes in order to undertake a new problem. Moreover, since these engineers and scientists have an understanding of the importance of the particular problem in their field, they are in a position to be able to recognize when a program that has been devised will be valuable to his colleagues. The reporting of these computing programs can become a vital part of the task of preparing the results of the theory and the calculations for inclusion in technical reports.

We are now able to undertake problems which would have been far too complex to have attempted to do by hand on manually-operated desk machines. However, we are also able, in a matter of seconds, to produce thousands of meaningless results because of the use of

an algorithm which is not appropriate for the problem under consideration for the range of values of the parameters which are of interest. A method of calculation (an algorithm) which may have worked well for a hand computation for a small value of a parameter may lead to mere computer noise when the electronic computer is used to evaluate the same physical phenomena for a large value of the parameter. While doing a problem by hand, the human operator would generally recognize when the results were fading away into the roundoff noise (such as when finding the difference between two numbers which are identical to all of the significant digits). However, when using the electronic computers which are available today, most of the intermediate results are "out-of-sight" and the user must be alert to the possibility that the results which he obtains may be meaningless in spite of the fact that a "correct" program has been written and has been used for a value of the parameter for which results are known from hand computations or from the results published by a previous author.

The experience of the present author has offered a good example of where "mathematically correct" procedures can lead to erroneous results when used with the finite word size of the modern electronic computer. In 1955 the author supervised the preparation of a set of data for the scattering properties of perfectly conducting spheres with diameters up to about seven wavelengths. Curves based upon these data were published in an appendix of a monograph on diffraction theory, namely

> R.W.P. King and T.T. Wu, The Scattering and Diffraction of Waves, Cambridge, Mass., Harvard University Press, 1959

and have been reproduced in the Encyclopedia of Physics and in J. van Bladel's Electromagnetic Fields (New York, McGraw-Hill, 1964). On three occasions the author has received long distance telephone calls from individuals who have programmed the exact

solution for the sphere and have had difficulty in obtaining results which correspond to those presented in the curves which were based upon the author's data. On each occasion, the author was able to resolve the problem on the telephone by inquiring as to how the spherical Bessel functions had been obtained. In each case, the investigators had made use of the property that the first two $j_n(x)$ functions are

$$j_0(x) = \frac{\sin(x)}{x} \quad , \quad j_1(x) = \frac{\sin(x)}{x^2} - \frac{\cos(x)}{x}$$

and the other values obtained from the recursion formula

$$j_{n+1}(x) = - j_{n-1}(x) + \frac{2n + 1}{x} j_n(x)$$

When x is larger than unity, and n is allowed to range through values that exceed the magnitude of x, the calculation (using the finite word size of the computer) deteriorates rapidly and eventually one even gets results which show the calculated values of $j_n(x)$ increasing with increasing values of n instead of possessing the correct property of decreasing with increasing values of n. The author was able to help his telephone callers by referring them to literature in which it is shown that the $j_n(x)$ functions should be calculated by "backward recurrence" by means of

$$j_{n-1}(x) = - j_{n+1}(x) + \frac{2n + 1}{x} j_n(x)$$

There is, by now, a rather extensive literature upon this subject. For example:

I. A. Stegun and M. Abramowitz, "Generation of Bessel Functions on High Speed Computers," Math. Tables, Vol. 11, Oct., 1957, pp. 255-257

J. B. Randels and R. F. Reeves, "Note on Empirical Bounds for Generating Bessel Functions," Comm. Assoc. Comput. Mach., Vol, 1, May, 1958, pp. 3-5

M. Goldstein and R. M. Thaler, "Recurrence Techniques for the Calculation of Bessel Functions," Math. Tables, Vol. 13, April, 1959, pp. 102-108

F. J. Corbato and J. L. Uretsky, "Generation of Bessel Functions in Digital Computers," J. Assoc. Comput. Mach., Vol. 6, July, 1959, pp. 366-375

W. Gautschi, "Recursive Computation of the Repeated Integrals of the Error Function," Math. Comp., Vol. 15, July, 1961, pp. 227-232

F. W. J. Olver, "Error Analysis of Miller's Recurrence Algorithm," Math. Comp., Vol. 18, Jan., 1964, pp. 65-74

The present author considers this problem of the computation of the Bessel functions to be a good example of why the individual research worker in diffraction theory cannot afford to "divorce himself" from the field of numerical analysis. The textbooks from which we learn our basic applied mathematics present us with devices such as the recurrence formulae for the Bessel functions, but we often fail to realize that these exact mathematical properties may have to be experimented with before they can be used on a calculating device which works with only a finite number of digits.

The author highly recommends that the reader acquaint himself with the paper

I. A. Stegun and M. Abramowitz, "Pitfalls in Computation," J. Soc. Indust. Appl. Math., Vol. 4, Dec., 1956, pp. 207-219

Among the related papers on this subject of difficulties associated with translating exact mathematical formulae into numerically meaningful results, the author recommends the following:

A. S. Householder, "Generation of Errors in Digital Computation," Bull. Amer. Math. Soc., Vol. 60, June, 1954, pp. 234-247

G. E. Forsythe, "Singularity and Near Singularity in Numerical Analysis," Amer. Math. Monthly, Vol. 65, April, 1958, pp. 229-240

Mathematics Research Center, The University of Wisconsin, Some Considerations in Practical Computation, by A. S. Chai, and H. J. Wertz, MRC Technical Summary Report No. 469, Contract No. DA-11-022-ORD-2059, Madison, Wisc., May, 1964

To date, the author knows of no book which has been written expressly for the engineer and scientist who is interested in the diffraction and the propagation of radio waves. However, such books are beginning to appear in other fields. For example:

R. S. Ledley, Use of Computers in Biology and Medicine, New York, McGraw-Hill, 1965

L. Lapidus, Digital Computation for Chemical Engineers, New York, McGraw-Hill, 1962

An examination of these books will reveal, however, that they are primarily concerned with "classical" numerical analysis and that they are primarily a collection of methods which are not very different from those to be found in books which are concerned with numerical analysis and which refrain from the use of "digital" or "computer" in their titles. However, the author does recommend the book by Ledley for its introductory material which will

provide the reader with some photographs and descriptions of some of the digital equipment which is currently available.

A different type of book which is beginning to appear is to be found in the efforts of certain authors to make available to other scientists the actual computer programs which have been developed in specific fields. Examples of books of this type are:

M. A. Melkanoff, D. S. Saxon, J. S. Nodvik, and D. G. Cantor, A FORTRAN Program for Elastic Scattering Analyses with the Nuclear Optical Model, Berkeley, University of California Press, 1961

P. A. D. de Maine and R. D. Seawright, Digital Computer Programs for Physical Chemistry, New York, Macmillan, 1963

F. Herman and S. Skillman, Atomic Structure Calculations, Englewood Cliffs, N.J., Prentice-Hall, 1963

The book by de Maine and Seawright is unique in that it presents the programs in both FORTRAN II and ALGOL 60. The present author has had no experience with the use of ALGOL 60 and, therefore, is in no position to pass judgement upon the relative merits of the use of these languages. However, since ALGOL 60 has been adopted as a "publication" language by several journals, the scientist who wishes to adapt for his useage the published algorithms of other scientists will often find the need to be able to read an ALGOL 60 procedure. The reader who wishes to acquaint himself with this programming language will probably find that one of the best sources for "self-study" is the following manual:

D. D. McCracken, A Guide to ALGOL Programming, New York, John Wiley, 1962

Other books which deal with ALGOL are:

E. W. Dijkstra, A Primer of ALGOL 60 Programming, New York, 1962

R. Wooldridge and J. F. Ractliffe, An Introduction to ALGOL Programming, Princeton, N.J., Van Nostrand, 1963

C. M. Reeves and M. Wells, A Course on Programming in ALGOL 60, Chicago, Educational Methods Inc., 1964

The reader who is not familiar with FORTRAN and who wishes to set out to learn this programming language will find himself confronted with the making of a choice between a number of texts if he wishes to begin a "self-study" program to acquaint himself with the features of the language.* The suggestion as to what text a particular person would find most helpful as a starting point is dependent upon the background, interests, and temperament of the individual. In the opinion of the present author, none of the texts presently available provide the mature scientists and engineers (assumed to have an education in electromagnetic theory and advanced calculus which includes some graduate courses) with an adequate "marriage" between the teaching of programming concepts and the use of these in numerical methods. One of the best efforts in this direction is the text

W. Prager, Introduction to Basic FORTRAN Programming and Numerical Methods, New York, Blaisdell, 1965

This text was written for a sophomore course for students of applied mathematics. It provides a readable account of the elements of FORTRAN (essentially FORTRAN II) and has the advantage of introducing at an early stage some flow charts and programs

*In the San Francisco area one of the local TV stations has been conducting a course in FORTRAN during the early morning hours. A least one course (of twenty lessons) has been taped and is available for classroom use in certain universities.

which will serve to answer the demands of the reader who starts his study and very soon becomes impatient and starts saying to himself: "Okay, so FORTRAN provides a set of precise rules for the writing of mathematical expressions. Now show me how you employ these rules in a typical scientific problem." However, after giving a few programs, the emphasis of the text shifts to the presentation (in a rather lucid manner) of a "minimum" knowledge of classical numerical methods. The present author is of the opinion that this text will be useful to many persons (engaged in electromagnetic studies) who work with "closed shop" programmers or with younger assistants who have been "college trained" in computer programming and numerical methods. The study of this text will help to open up the lines of communication between the scientist-engineer and his programming assistant.

For the person who wishes to persue his study of FORTRAN to the point of undertaking to do his own programming, the present author highly recommends either of the following texts:

J. T. Golden, FORTRAN IV Programming and Computing, Englewood Cliffs, N.J., Prentice-Hall, 1965

D. D. McCracken, A Guide to FORTRAN IV Programming, New York, John Wiley, 1965

Either of these texts will, if diligently studied (and if help is sought from colleagues who have had more experience with programming when obscure points arise), should provide the necessary background to undertake to write FORTRAN IV programs.

The reader who wishes to "shop around" for other texts will find himself faced with a "flood" of books (some of which are rather expensive) from which to choose. A partial list includes the following:

J. N. Haag, Comprehensive FORTRAN Programming, New York, Hayden, 1965

L. D. Harris, Numerical Methods Using FORTRAN, Columbus, Ohio, Charles E. Merrill, 1964

J. M. McCormick and M. G. Salvadori, Numerical Methods in FORTRAN, Englewood Cliffs, N.J., Prentice-Hall, 1964

D. D. McCracken, A Guide to FORTRAN Programming, New York, John Wuley, 1961

D. D. McCracken and W. S. Dorn, Numerical Methods and FORTRAN Programming, New York, John Wiley, 1964

E. I. Organick, A FORTRAN Primer, Reading, Mass., Addison-Wesley, 1963

R. H. Pennington, Introductory Computer Methods and Numerical Analysis, New York, Macmillan, 1965

S. V. Pollack, A Guide to FORTRAN IV, New York, Columbia University Press, 1965

The present author has also derived some benefits from sections in the book

B. W. Arden, An Introduction to Digital Computing, Reading, Mass., Addison-Wesley, 1963.

However, this book contains programs which are written in the MAD (Michigan Algorithm Decoder) language which resembles FORTRAN (FORmula TRANslation) in many respects, and yet has some of the features of ALGOL 60, while at the same time having features not found in either of the two more common languages.

Several of the titles above reveal that many writers are trying

teach numerical methods and computer programming with the same text. Although this is very desirable in that it affords the scientist-engineer the opportunity to see the programming rules applied to the type of problems which he is likely to encounter in his work, the result is generally that the book is an inferior text on numerical analysis.

The present author feels that the two books which he personally feels to be the most valuable books on numerical analysis are

C. Lanczos, Applied Analysis, Englewood Cliffs, N.J., Prentice-Hall, 1956

A. Ralston, A First Course in Numerical Analysis, New York, McGraw-Hill, 1965

In many respects, the book by Lanczos is "unorthodox" since its emphasis is somewhat different from a host of other books. However, Lanczoz has presented many "tricks of the trade" which he has found useful during his long and highly successful career as a physicist and a consultant in applied mathematics. The book by Ralston, although it is newly off the press, promises to become a classic. Ralston acknowledges his endebtedness to his teachers who wrote the excellent treatises:

F. B. Hildebrand, Introduction to Numerical Analysis, New York, McGraw-Hill, 1956

Z. Kopal, Numerical Analysis, 2nd ed., London, Chapman and Hall, 1961

and to his former colleague who wrote the very useful book:

R. W. Hamming, Numerical Methods for Scientists and Engineers, New York, McGraw-Hill, 1962

Two other books which the author has found quite useful are:

Staff of National Physical Laboratory, Modern Computing Methods, London, Her Majesty's Stationery Office, 1961

G. N. Lance, Numerical Methods for High Speed Computers, New York, Gordon and Breach, 1960

Three books have appeared which have been the result of the joint efforts of a number of authors and which contain much valuable reference material. These are:

L. Fox, ed., Numerical Solution of Ordinary and Partial Differential Equations, Reading, Mass., Addison-Wesley, 1962

A. Ralston and H. S. Wilf, eds., Mathematical Methods for Digital Computers, New York, John Wiley, 1960

J. Todd, ed., Survey of Numerical Analysis, New York, McGraw-Hill, 1962

The collection of papers edited by Ralston and Wilf is especially recommended to the reader who wishes to study some examples of the preparation of flow charts which can often be of great help in the explanation of the methods which underlie a computer program.

There are many other textbooks available which provide the reader with a collection of techniques for use in numerical analysis. Much of the material which is included is classical and can be found in some of the older texts, such as

E.T. Whittaker and G. Robinson, The Calculus of Observations; a Treatise on Numerical Mathematics, London, Blackie, 1924

Among the newer texts, the author has consulted the following:

P. J. Davis, Interpolation and Approximation, New York, Blaisdell, 1963

C. E. Fröberg, Introduction to Numerical Analysis, Reading, Mass., Addison-Wesley, 1965

D. R. Hartree, Numerical Analysis, Oxford, Oxford University Press, 1958

P. Henrici, Elements of Numerical Analysis, New York, John Wiley, 1964

J. G. Herriot, Methods of Mathematical Analysis and Computation, New York, John Wiley, 1963

A. S. Householder, Principles of Numerical Analysis, New York, McGraw-Hill, 1953

C. Lanczos, Linear Differential Operators, New York, Van Nostra nd, 1961

B. Noble, Numerical Methods: 1 and 2, New York, Interscience, 1964

R. G. Stanton, Numerical Methods for Science and Engineering, Englewood Cliffs, N.J., Prentice-Hall, 1961

E. L. Stiefel, An Introduction to Numerical Mathematics, New York, Academic Press, 1963

Since two of the most important problems which face the numerical analyst are those of evaluating integrals and of solving differential equations, the author wishes to direct attention to the two books:

V. I. Krylov (trans. by A. H. Stroud), Approximate Calculation of Integrals, New York, Macmillan, 1962

P. Henrici, Discrete Variable Methods in Ordinary Differential Equations, New York, John Wiley, 1962

This survey of some of the recent books which have appeared on FORTRAN programming and numerical methods serves to indicate that the scientist-engineer (whose formal education has included no courses in programming and numerical methods) who wishes to acquire skills which will permit him to make use of the electronic digital computer will find a wealth of source material available to him. The problem which the individual will have to resolve is that of choosing where he is going to start. The author feels that most of the above books are excellent and the choice of which books are to be singled out for use by any given individual will depend largely upon the background of that person. However, with apologies to all the authors whose works are slighted, the present author recommends to most of his colleagues in the field of electromagnetic theory that they consider as a basic library the three books:

W. Prager, Introduction to Basic FORTRAN Programming and Numerical Methods, New York, Blaisdell, 1965

J. T. Golden, FORTRAN IV Programming and Computing, Englewood Cliffs, N.J., Prentice-Hall, 1965

A. Ralston, A First Course in Numerical Analysis, New York, McGraw-Hill, 1965

For the use of any given computer, the potential user will have to consider the manuals on FORTRAN which have been prepared by the manufacturer. Although these manuals are generally written in such terse language that they do not constitute a good text

for the learning of FORTRAN, they constitute a vital reference once one has begun to acquire a familiarity with the language. A great deal of care has been taken in the preparation of most of these manuals.

The present author feels that one of the most difficult concepts to master in the FORTRAN language is that of the FORMAT statements for the input and output of the data. When one has mastered these concepts it is possible to devise a display of the output data which is essentially the format which one would like to have reproduced as a table in a technical report. One of the best ways to master FORMAT is to actually write programs and gradually attempt to be more sophisticated in the layout of the output data. In connection with the INPUT/OUTPUT statements, the author would like to direct attention to one manufacturer's manual in which he believes that an exceptionally good job has been done in illustrating the concepts associated with this important aspect of computer programming. The manual is:

> Honeywell Electronic Data Processing, AUTOMATH 1800 Language Manual, Wellesley Hills, Mass., 1964

The AUTOMATH 1800 language is a "dialect" of FORTRAN IV. However, most of the features of the FORMAT specifications are identical with those of the features of most of the FORTRAN IV compilers for the larger electronic computers. Through the use of skillfully-executed drawings and illustrations this manual succeeds (in the opinion of the present author) in clearing up a lot of points which one might be forced to learn to master from experience if one tries to follow some of the more terse descriptions given in some of the other manuals and texts. It is extremely important to master FORMAT because in many instances one can have a program which is working "perfectly" from a mathematical point of view but one will be forced to re-run the job because of a mistake in format specification.

Appendix B

TABLES OF THE AIRY FUNCTION ON THE REAL AXIS

In this appendix we will present some tables of the Airy functions which we have constructed for real values of the argument. The definitions which we use in this section are those employed by Miller*.

$$\mathrm{Ai}(x) = \frac{1}{\pi}\int_0^\infty \cos\left(\frac{t^3}{3} + xt\right) dt \qquad \text{(B-1)}$$

$$\mathrm{Bi}(x) = \frac{1}{\pi}\int_0^\infty \left[\sin\left(\frac{t^3}{3} + xt\right) + \exp\left(-\frac{t^3}{3} + xt\right)\right] dt \qquad \text{(B-2)}$$

We recall that Miller also defined the auxiliary functions defined by

$$\mathrm{Ai}(x) = F(x)\sin[\chi(x)] \quad , \quad \mathrm{Bi}(x) = F(x)\cos[\chi(x)] \qquad \text{(B-3)}$$

$$\mathrm{Ai}'(x) = G(x)\sin[\psi(x)] \quad , \quad \mathrm{Bi}'(x) = G(x)\cos[\psi(x)] \qquad \text{(B-4)}$$

For small values of x, one can evaluate these functions by means of the Taylor series

$$\mathrm{Ai}(x) = \alpha y_1 - \beta y_2 \qquad \mathrm{Bi}(x) = 3^{1/2}(\alpha y_1 + \beta y_2) \qquad \text{(B-5)}$$

$$y_1 = 1 + \frac{1}{3!}x^3 + \frac{1\cdot 4}{6!}x^6 + \frac{1\cdot 4\cdot 7}{9!}x^9 + \ldots \qquad y_2 = x + \frac{2}{4!}x^4 + \frac{2\cdot 5}{7!}x^7 + \frac{2\cdot 5\cdot 8}{10!}x^{10} + \ldots$$

---

* J.C.P. Miller, The Airy Integral, British Association Mathematical Tables, Part-Volume B, Cambridge, Cambridge University Press, 1946

where α and β are the constants

$$\alpha = 3^{-\frac{2}{3}}/(-\tfrac{1}{3})! = 0.355028053887817239256\ldots$$

$$\beta = 3^{-\frac{1}{3}}/(-\tfrac{2}{3})! = 0.258819403792806798405\ldots$$

The tables given by Miller include the following

Ai(x) and Ai'(x). x = - 20.00(0.01) + 2.00. 8D
Log_{10} Ai(x) and Ai'(x)/Ai(x). x = 0.0(0.1)25.0(1)75. 7-8D
Zeros and Turning-Values of Ai(x) and Ai'(x). The first 50 of each. 8D
Bi(x) and Reduced Derivatives. x = - 10.0(0.1) + 2.5. 7-8D
Zeros and Turning-Values of Bi(x) and Bi'(x). The first 20 of each. 8D
Log_{10} Bi(x) and Bi'(x)/Bi(x). x = 0.0(0.1)10.0. 7-8D
Auxiliary Functions. F(x), χ(x), G(x), -ψ(x). x = -80(1) - 30.0(0.1)+ 2.5. 8D

Values of Ai(z), Bi(z), Ai'(z), and Bi'(z) have been given by Woodward and Woodward* for complex values of z = x +iy for which the magnitudes of x and y are less than 2.4. If one makes use of the relationships between these Airy functions and the modified Hankel functions which were computed by the staff of the Harvard Computation Laboratory**, one can obtain values of these functions for complex values of z for which the modulus of z is less than 6.

We have programmed (using data from the Harvard tables**) Eq. (B-3) and Eq. (B-4) so as to be able to evaluate the Airy functions in

---

*P. M. Woodward and A. M. Woodward, "Four-Figure Tables of the Airy Function in the Complex Plane," Phil. Mag., Vol. 37, pp. 236-261

**Harvard University Computation Laboratory, Tables of the Modified Hankel Functions of Order One-Third and of Their Derivatives, Annals of the Computation Laboratory of Harvard University, Vol. II, Cambridge, Mass., Harvard University Press, 1945

the complex plane under the condition that the modulus of z be less than 6.1. The program is known as

SUBRØUTINE AIBIZ(Z,AI,BI,AIP,BIP)

and is listed in Program B-1. The relation of the FORTRAN variables to the mathematical definitions are as follows:

$$z = Z$$
$$Ai(z) = AI$$
$$Bi(z) = BI$$
$$Ai'(z) = AIP$$
$$Bi'(z) = BIP$$

The presence of the type declaration

CØMPLEX Z,AI,BI,AIP,BIP

reveals that the complex arithmetic features of FORTRAN IV are utilized. However, the program actually employs double precision arithmetic. For values of z which are somewhat smaller than the limit of 6.1 which has been placed on the modulus of z, the program can be easily modified* so as to obtain more than the eight

*For example, let z = ZR + iZI and add "DØUBLE PRECISION ZR,ZI" while retaining "CØMPLEX Z." Omit the statements defining AI, BI, AIP, and BIP. Then rename the subprogram as follows: "SUBRØUTINE AIBIZ(ZR,ZI,DAIR,DAII,DAIPR,DAIPI,DBIR,DBII,DBIPR,DBIPI)." Let the first executable statement be "Z = -CMPLX(ZR,ZI)." Replace the statements defining DRZ and DIZ with "DRZ = -ZR" and "DIZ = -ZI." The "CALL EXIT" which will occur when CABS(Z) is greater than 6.1 will have to be modified according to what accuracy the user wishes to achieve. Since the greatest loss of accuracy occurs when computing Ai(x) and Ai'(x) for positive values of x, a comparison of the output of these quantities with the tabulated values given in Block B-2 will provide the basis for a criterion for the number of significant figures available.

decimal digits which are available when employing the "CØMPLEX" features of FORTRAN IV. We have employed this feature in the argument list of the subprogram because we had in mind the needs of a user who is employing complex arithmetic in the calling program.

If the modulus of z exceeds 6.1, the program writes a message

AIBIZ CANNOT HANDLE Z = ...

(where the dots indicate that the value of z is printed) and then executes a CALL EXIT.

The NPREC subroutines (described in the Preface) have been employed to compute values of the Airy functions on the real axis which are accurate to from 15 to 21 significant figures. In Block B-2 we present the values of the Airy functions and their derivatives for the interval x = 0.1(0.1)10.0. These data have been arranged in the form of a BLOCK DATA subprogram with the data stored in a labeled CØMMØN block referred to by the FORTRAN name EQUADP for its relation to the EQUal-interval Airy function in Double Precision. If we let the mathematical variable x be related to the FORTRAN index I by means of the relationship

x = 0.1*FLØAT(I)

then the functions that are tabulated are

Ai(x) = EAIRYF(I)
Bi(x) = EBIRYF(I)
Ai'(x) = EAIRYD(I)
Bi'(x) = EBIRYD(I)

The values of Bi(x) and Bi'(x) are accurate to 20 figures throughout most of the range, but the values of Ai(x) and Ai'(x) deteriorate to about 15 or 16 figures in the middle of the range. The author has been unable to devise an algorithm which will achieve more accuracy for Ai(x) and Ai'(x) for x in the vicinity of 5 when doing the calculations with a "word length" of 21 decimal

digits. Since the need for higher accuracy has not arisen, the author has not undertaken the computation of the auxiliary functions $r(x)$, $\mu(x)$, $p(x)$, and $\lambda(x)$ in the representations

$$2\sqrt{\pi}\ \mathrm{Ai}(x) = \sqrt{r(x)}\ \exp[-\mu(x)] \tag{B-6a}$$

$$\sqrt{\pi}\ \mathrm{Bi}(x) = \sqrt{r(x)}\ \exp[+\mu(x)] \tag{B-6b}$$

$$-2\sqrt{\pi}\mathrm{Ai}'(x) = \sqrt{p(x)}\ \exp[-\lambda(x)] \tag{B-7a}$$

$$\sqrt{\pi}\mathrm{Bi}'(x) = \sqrt{p(x)}\ \exp[+\lambda(x)] \tag{B-7b}$$

where the differential equations for the functions have been given in Eqs. (6-38), (6-50), (6-53) and (6-54). Accurate values for these auxiliary functions can be obtained for small values of x from the Taylor series in Eq. (B-5) and for large values of x from the asymptotic expansions of Eqs. (6-52) and (6-55). If step-by-step integration is employed from the small and large values of x, it is quite possible that one can close the "gap" that occurs around $x = 5$ with very precise values (whose accuracy will be apparent from the degree of overlap between the values obtained by starting at the two edges of the "gap").

In Table B-1 we give the values of Ai(x) and Bi(x) for x = 10.0 (0.1)25.0. In Table B-2 we give the values of Ai'(x) and Bi'(x) over the same range. The reader should observe that for x greater than 25.0 one will soon reach the point where the values of the Airy functions will lie outside the range 10^{-38} to 10^{+38} which is employed by most FORTRAN compilers. The manner in which the Airy functions continue to grow with increasing x is displayed in Tables B-3 and B-4 where we present the values of these functions for (1/x) in the range 0.001(0.001)0.100. Since problems occur in which we require values (for large values of x) for products such as Ai(x)Bi'(x) or ratios such as Bi'(x)/[Bi'(x) - qBi(x)], it is not practical to use the Airy functions from Tables B-3 and B-4. These products and ratios are well within the range of the FORTRAN numbers and therefore it is convenient to employ

tables of the auxiliary functions $r(x)$, $\mu(x)$, $p(x)$ and $\lambda(x)$ which are defined in Eq. (B-7). In order to tabulate a function which will permit accurate and easy interpolation, it is convenient to consider the asymptotic expansions in Eqs. (6-52) and (6-55) and define auxiliary functions which have the property of approaching unity for x tending to infinity. Let us first introduce the quantity $\xi = \xi(x)$ which is defined by

$$\xi = \frac{2}{3}(\sqrt{x^3}) \tag{B-8}$$

We then define the auxiliary functions

$$Q(x) = \sqrt{x}\, r(x) = 2\pi\sqrt{x}\, Ai(x)Bi(x) \tag{B-9a}$$

$$M(x) = \xi^{-1}\mu(x) \tag{B-9b}$$

$$P(x) = \frac{-1}{\sqrt{x}}\, p(x) = -\frac{2\pi}{\sqrt{x}} Ai'(x)Bi'(x) \tag{B-10a}$$

$$\Lambda(x) = \xi^{-1}\lambda(x) \tag{B-10b}$$

Tables of $Q(x)$ and $M(x)$ are given in Table B-5 for $(1/x) = 0.001(0.001)0.100$. Tables over the same range are given for $P(x)$ and $\Lambda(x)$ in Table B-6.

Fox* has also considered the case of Airy functions for large, positive values of the argument. The auxiliary functions which he employs are suggested by the asymptotic expansions which are of the form described by Eq. (6-48). Let us supplement Eq. (6-48) by giving the explicit form for the asymptotic expansions for $Ai'(x)$ and $Bi'(x)$. Let ξ be as defined in Eq. (B-8) and let us express the asymptotic expansions in the form

---

*L. Fox, Tables of Weber Parabolic Cylinder Functions and Other Functions for Large Arguments, National Physical Laboratory Mathematical Tables, Vol. 4, London, Her Majesty's Stationery Office, 1960

$$2\sqrt{\pi}\ \mathrm{Ai}(x) = x^{-1/4}\exp(-\xi)L(-\xi)\ ,\quad \sqrt{\pi}\ \mathrm{Bi}(x) = x^{-1/4}\exp(\xi)L(\xi) \tag{B-11}$$

$$2\sqrt{\pi}\ \mathrm{Ai}'(x) = -x^{1/4}\exp(-\xi)K(-\xi)\ ,\quad \sqrt{\pi}\ \mathrm{Bi}'(x) = x^{1/4}\exp(\xi)K(\xi) \tag{B-12}$$

where $L(\xi)$ and $K(\xi)$ are the asymptotic series

$$L(\xi) = \sum_{s=0}^{\infty} \frac{u_s}{\xi^s} = 1 + \frac{3\cdot 5}{1!216}\frac{1}{\xi} + \frac{5\cdot 7\cdot 9\cdot 11}{2!(216)^2}\frac{1}{\xi^2} + \frac{7\cdot 9\cdot 11\cdot 13\cdot 15\cdot 17}{3!(216)^3}\frac{1}{\xi^3} + \cdots \tag{B-13}$$

$$M(\xi) = \sum_{s=0}^{\infty} \frac{v_s}{\xi^s} = 1 - \frac{3\cdot 7}{1!216}\frac{1}{\xi} - \frac{5\cdot 7\cdot 9\cdot 13}{2!(216)^2}\frac{1}{\xi^2} - \frac{7\cdot 9\cdot 11\cdot 13\cdot 15\cdot 19}{3!(216)^3}\frac{1}{\xi^3} - \cdots \tag{B-14}$$

in which $u_o = 1$ and

$$u_s = \frac{(2s+1)(2s+3)(2s+5)\ldots(6s-1)}{s!\,(216)^s}\ ,\quad v_s = -\frac{6s+1}{6s-1}\,u_s \tag{B-15}$$

The functions tabulated by Fox can be defined as follows:

$$R(z) = L(-\tfrac{1}{z}) \tag{B-16}$$

$$S(z) = L(\tfrac{1}{z}) \tag{B-17}$$

$$W(z) = -K(-\tfrac{1}{z}) \tag{B-18}$$

$$X(z) = K(\tfrac{1}{z}) \tag{B-19}$$

Fox showed that $R(z)$ and $S(z)$ satisfy the differential equation

$$y''(z) + 2\left(\frac{1}{z} \pm \frac{1}{z^2}\right)y'(z) + \frac{5}{36z^2}\,y(z) = 0 \tag{B-20}$$

where the + sign is used for $R(z)$ and the - sign for $S(z)$. He also showed that $W(z)$ and $X(z)$ satisfy the differential equation

$$y''(z) + 2\left(\frac{1}{z} \pm \frac{1}{z^2}\right)y'(z) - \frac{7}{36z^2}\,y(z) = 0 \tag{B-21}$$

where the + sign is used for $W(z)$ and the - sign is used for $X(z)$. These differential equations are very promising for the

+.79

purpose of developing a method of numerical integration which can be used to obtain the values of the Airy functions in the "gap" between the region where the Taylor series of Eq. (B-5) become impractical* and the smallest values of x for which the asymptotic series of Eq. (B-13) can be employed to compute the functions to the desired accuracy. A very promising approach would be that of employing the asymptotic series of Eqs. (B-13) and (B-14) for large values of ξ (i.e., for large values of x or small values of $z = \xi^{-1}$) and developing a program for the step-by-step integration of Eqs. (B-20) and (B-21) which would carry the computation to smaller values of x and ξ (or to larger values of z).

In Tables B-7 and B-8 we present our results for the functions R(z), S(z), W(z), and X(z). The tables are given as a function of (1/x), where $x = (\frac{2}{3}z)^{\frac{2}{3}}$, for (1/x) = 0.001(0.001)0.100. The tables are accurate to about 20 decimals.

Since Fox has tabulated the functions as a function of $z = \xi^{-1}$, there is only one point in common in the two sets of calculations, namely x = 25 where (1/x) = 0.04 and z = 0.012. The values given by Fox are

---

*The Taylor series for the Airy function Ai(x) presents the same numerical difficulties for large positive values of x as does the Taylor series for the more familiar function exp(-x). The terms in the series so completely cancel each other that when one employs a computer word length of 21 decimal digits the output has become "computer noise" by the time one reaches $x \approx 10$. The use of step-by-step integration starting from x = 0 and going towards x = 10 was found to yield essentially the same accuracy as that obtained with the Taylor series. However, step-by-step integration starting from large values of x and working towards smaller values is a more stable numerical process and one can preserve a relatively high degree of accuracy.

$$R(0.012) = 0.9991719494 \ , \ S(0.012) = 1.0008387474$$
$$-W(0.012) = 1.0011604193 \ , \ X(0.012) = 0.9988269392$$

The reader can readily see for himself that these agree with our results in Tables B-7 and B-8 for $(1/x) = 0.040$.

Fox computed these auxiliary functions to 10 decimals over the range $z = 0.000(0.001)0.050$. In terms of the argument x of the Airy functions Ai(x) and Bi(x), this range is $x_o < x < \infty$, where $x_o \approx 9.65$. This is essentially the same range as that of Tables B-7 and B-8, namely $10 < x < \infty$.

The tables which have been given as Tables B-5 through B-8 provide excellent examples of slowly varying functions which can be represented very efficiently by Chebyshev polynomial series of the form given in Eq. (8-42). A representation in terms of a series of the form $\Sigma a_n T_n^*(\eta)$ would require only a small number of coefficients a_n in order to compute these functions to high accuracy over the range employed in Tables B-5 through B-8. If x lies in the range $A < x < \infty$, then we would take η to be in the form $\eta = (A/x)^\nu$, where appropriate choices of ν might be $\nu = 3$ for Tables B-5 and B-6 and $\nu = 1.5$ for Tables B-7 and B-8.

Let us now direct our attention to negative values of x. In Tables B-9 and B-10 we present values for the Airy functions and their derivatives for the range $-10 < x < 0$ at increments of 0.1. However, we have followed the convention employed by Miller in The Airy Integral (see footnote on p. B-1) and we give tables of Ai(-x), Bi(-x), Ai'(-x) and Bi'(-x). Therefore, the arguments in the tables are positive and $x = 0(0.1)10.0$. When these calculations were compared with the results obtained from the asymptotic series the agreement was found to exist to 19 decimals. Therefore, we assume that throughout most of these tables the results are good to approximately 20 decimals.

Since the facilities to work with 21 decimal digits are not generally available (and since the NPREC subroutines are not especially suited for writing output on magnetic tape to be later reread into the computer), we formed the auxiliary functions $F(-x)$, $\chi(-x)$, $G(-x)$, and $\psi(-x)$ at the time of computing the Airy functions in the range $0 < x < 10.0$. The auxiliary functions are defined by means of the relations

$$Ai(-x) = F(-x)\sin[\chi(-x)] \quad , \quad Ai'(-x) = G(-x)\sin[\psi(-x)] \tag{B-22}$$

$$Bi(-x) = F(-x)\cos[\chi(-x)] \quad , \quad Bi'(-x) = G(-x)\cos[\psi(-x)] \tag{B-23}$$

or

$$F^2(-x) = Ai^2(-x) + Bi^2(-x) \quad , \quad \chi(-x) = \tan^{-1}[\, Ai(-x)/Bi(-x) \,] \tag{B-24}$$

$$G^2(-x) = Ai'^2(-x) + Bi'^2(-x) \quad , \quad \psi(-x) = \tan^{-1}[Ai'(-x)/Bi'(-x)] \tag{B-25}$$

In Tables B-11 and B-12 we present our output for these auxiliary functions for $x = 0.0(0.1)10.0$. Unfortunately, the computer program was written without adequate forethought and the angles defined as $\chi(-x)$ and $\psi(-x)$ which appear in these tables require a correction for those cases in which the angle lies in either the second or the third quadrants since the tabular values are in the range from $-\pi/2$ to $+\pi/2$. The reader who wishes to use these values can readily determine if a quadrant correction is required by consulting the corresponding values of $Ai(-x)$, $Bi(-x)$, $Ai'(-x)$ and $Bi'(-x)$ which are given in Tables B-9 and B-10.

For x between 10 and 25, we computed the auxiliary functions by means of the asymptotic series given in Eq. (6-9). We then used Eqs. (B-22) and (B-23) to compute the Airy functions. The Airy functions are given in Tables B-13 and B-14 and the auxiliary functions are given in Tables B-15 and B-16. In Tables B-15 and B-16 the angles $\chi(-x)$ and $\psi(-x)$ are given in radians, but the reader will observe that they are not confined to the range from

0 to 2π radians. The functions which we have tabulated (in radians) will have the behavior depicted in Fig. B-1 if we extend the tables back towards the origin. Since the quantities tabulated in the tables are $\chi(-x)$ and $\psi(-x)$, where $10 < x < 25$, the reader will observe that the tables represent the behavior of the functions plotted in Fig. B-1 in the region far to the left of the




Fig. B-1 The Auxiliary Functions $\chi(x)$ and $\psi(x)$

+ 41

portions of the curves which are depicted in the figure.

For large values of x, the auxiliary functions possess the asymptotic expansions

$$\left|F(-x)\right|^2 \sim \frac{1}{\pi x^{1/2}}\left(1 - \frac{1\cdot3\cdot5}{1!\,96}\frac{1}{x^3} + \frac{1\cdot3\cdot5\cdot7\cdot9\cdot11}{2!\,96^2}\frac{1}{x^6} - \frac{1\cdot3\cdot5\cdot7\cdot9\cdot11\cdot13\cdot15\cdot17}{3!\,96^3}\frac{1}{x^9} + \ldots\right) \tag{B-26}$$

$$\chi(-x) - \tfrac{1}{4}\pi \sim \tfrac{2}{3}x^{3/2}\left(1 - \frac{5}{32}\frac{1}{x^3} + \frac{1105}{6144}\frac{1}{x^6} - \frac{82825}{65536}\frac{1}{x^9} + \frac{1282031525}{58720256}\frac{1}{x^{12}} - \ldots\right) \tag{B-27}$$

$$\left|G(-x)\right|^2 \sim \frac{1}{\pi}x^{1/2}\left(1 + \frac{1\cdot3}{1!\,96}\frac{7}{x^3} - \frac{1\cdot3\cdot5\cdot7\cdot9}{2!\,96^2}\frac{13}{x^6} + \frac{1\cdot3\cdot5\cdot7\cdot9\cdot11\cdot13\cdot15}{3!\,96^3}\frac{19}{x^9} - \ldots\right) \tag{B-28}$$

$$\psi(-x) + \tfrac{1}{4}\pi \sim \tfrac{2}{3}x^{3/2}\left(1 + \frac{7}{32}\frac{1}{x^3} - \frac{1463}{6144}\frac{1}{x^6} + \frac{4\,95271}{3\,27680}\frac{1}{x^9} - \frac{206530429}{8388608}\frac{1}{x^{12}} + \ldots\right) \tag{B-29}$$

It would be easier to interpolate in the tables if the quantities which were tabulated were the slowly varying functions $f(x)$, $\theta(x)$, $g(x)$ and $\varphi(x)$ which are defined as follows:

$$f^2(x) = \pi\sqrt{\pi}\, F^2(-x) \tag{B-30}$$

$$\theta(x) = \tfrac{3}{2}x^{\frac{2}{3}}\,[\chi(-x) - \pi/4] \tag{B-31}$$

$$g^2(x) = \frac{\pi}{\sqrt{x}}\, G^2(-x) \tag{B-32}$$

$$\varphi(x) = \tfrac{3}{2}x^{\frac{2}{3}}\,[\psi(-x) + \pi/4] \tag{B-33}$$

There was not adequate time under the present contract period to

prepare tables of these function. The author plans to return to this problem and to compute the functions for $(1/x) = 0.000(0.001)(0.100)$. The functions $f(x)$, $\theta(x)$, $g(x)$ and $\varphi(x)$ should also be developed into series of the Chenbyshev polynomials $T_n^*(\eta)$ where $\eta = (A/x)^3$ and $A < x < \infty$.

The asymptotic expansions for $f(x)$ and $g(x)$, namely

$$f^2(x) = 1 - \frac{1\times3\times5}{96\,x^3} + \frac{1\times3\times5\times7\times9\times11}{21(96)^2x^6} - \frac{1\times3\times5\times7\times9\times11\times13\times15\times17}{31(96)^3\,x^9} + \cdots \quad \text{(B-34)}$$

$$g^2(x) = 1 + \frac{1\times3\times7}{96\,x^3} - \frac{1\times3\times5\times7\times9\times13}{21(96)^2x^6} + \frac{1\times3\times5\times7\times9\times11\times13\times15\times19}{31(96)^3\,x^9} + \cdots \quad \text{(B-35)}$$

can be readily extended to higher order terms. However, it is not so obvious as to how to develop the further terms in the expansions for $\theta(x)$ and $\varphi(x)$, namely

$$\theta(x) = 1 - \frac{5}{32\,x^3} + \frac{1105}{6144\,x^6} - \frac{82825}{65536\,x^9} + \frac{1282031525}{58720256\,x^{12}} + \cdots \quad \text{(B-37)}$$

$$\varphi(x) = 1 + \frac{7}{32\,x^3} - \frac{1463}{6144\,x^6} + \frac{495271}{327680x^9} - \frac{206530429}{8388608\,x^{12}} + \cdots \quad \text{(B-38)}$$

However, the properties

$$\pi F^2(-x)\,\frac{d\chi(-x)}{dx} = 1 \quad , \quad \pi G^2(-x)\,\frac{d\psi(-x)}{dx} = x \quad \text{(B-39)}$$

permit us to assume a series representation for $\theta(x)$ and $\varphi(x)$

$$\theta(x) = 1 + \sum_{n=1}^{\infty} A_n x^{-3n} \quad , \quad \varphi(x) = 1 + \sum_{n=1}^{\infty} B_n x^{-3n} \quad \text{(B-40)}$$

and to determine the relationship between the known coefficients in the expansions of $f^2(x)$ and $g^2(x)$ and the unknown coefficients in the expansions of $\theta(x)$ and $\varphi(x)$.

Block B-1

THE COEFFICIENTS FOR ASYMPTOTIC EXPANSION OF $\theta(x)$ AND $\varphi(x)$

```
      DOUBLE PRECISION CHI(18), PSI(18)
      DATA (CHI(I),PSI(I),I=1,18)/
     A -0.15625000000000000000D+00 ,+0.21875000000000000000D+00 ,
     B +0.17985026041666666667D+00 ,-0.23811848958333333333D+00 ,
     C -0.12638092041015625000D+01 ,+0.15114471435546875000D+01 ,
     D +0.21832866753850664411 3D+02 ,-0.24620345711708068847 7D+ 2 ,
     E -0.69682822536884082688 2D+03 ,+0.76086294204038050439 6D+ 3 ,
     F +0.35553981581391285131 3D+05 ,-0.38092128868188640229 7D+ 5 ,
     G -0.26532584171176254655 6D+07 ,+0.28082562244080195275 9D+ 7 ,
     H +0.27258321302415599342 3D+09 ,-0.28607946453262385636 6D+ 9 ,
     I -0.36894027011098001901 6D+11 ,+0.38480790905747179385 8D+11 ,
     J +0.63630204766133750347 9D+13 ,-0.66050619870875374085 1D+13 ,
     K -0.13622439180500760352 1D+16 ,+0.14087107203786266895 5D+16 ,
     L +0.35446828581078074793 6D+18 ,-0.36542984737250173263 8D+18 ,
     M -0.11017990287911060070 0D+21 ,+0.11329639450011994200 7D+21 ,
     N +0.40320823577747335533 8D+23 ,-0.41371744404875883016 5D+23 ,
     O -0.17159646692716793386 0D+26 ,+0.17574349642737663059 0D+26 ,
     P +0.84031035110193005527 5D+28 ,-0.85924160506996514557 4D+28 ,
     Q -0.46916652616593043608 9D+31 ,+0.47906551025465751515 7D+31 ,
     R +0.29625489083114947131 5D+34 ,-0.30213269421511401499 7D+34 /
```

In Block B-1 we present numerical values for the coefficients. The relationship between A_n and B_n and the CHI(I) and PSI(I) is

$$A_I = \text{CHI}(I) \quad , \quad B_I = \text{PSI}(I)$$




Fig. B-2 The Behavior of $\chi'(t)$ and $\psi'(t)$

The behavior of the derivatives of $\chi(t)$ and $\psi(t)$ is depicted in Fig. B -2.

In Program B-2 we present a FORTRAN subroutine which evaluates the Airy functions and their derivatives by using the classical expansions

$$\sqrt{\pi}\,Ai(x) = \frac{1}{2}\,x^{-1/4}\,e^{-\xi}\left(1 - \frac{3\cdot 5}{1!\,216}\,\frac{1}{\xi} + \frac{5\cdot 7\cdot 9\cdot 11}{2!\,(216)^2}\,\frac{1}{\xi^2} - \frac{7\cdot 9\cdot 11\cdot 13\cdot 15\cdot 17}{3!\,(216)^3}\,\frac{1}{\xi^3} + \cdots\right) \tag{B-41}$$

$$\sqrt{\pi}\,Bi(x) = x^{-1/4}\,e^{\xi}\left(1 + \frac{3\cdot 5}{1!\,216}\,\frac{1}{\xi} + \frac{5\cdot 7\cdot 9\cdot 11}{2!\,(216)^2}\,\frac{1}{\xi^2} + \frac{7\cdot 9\cdot 11\cdot 13\cdot 15\cdot 17}{3!\,(216)^3}\,\frac{1}{\xi^3} + \cdots\right) \tag{B-42}$$

where

$$\xi = \frac{2}{3}\,x^{3/2}$$

The reader will note that Program B-2 will set the Airy functions equal to the following

$$Ai(x) \approx 0 \quad , \quad Ai'(x) \approx 0$$

$$Bi(x) \approx 10^{+38} \quad Bi'(x) \approx 10^{+38}$$

when ξ becomes large enough that the FORTRAN subroutine cannot compute $\exp(\pm\xi)$. In Table B-17 we present a table of values of $\exp(\pm\xi)$. The reader will observe that the overflow and underflow occur for x between 25.5 and 26.0.

In Table B-18 we give tables of values of the sine and cosine of $(\xi + \pi/4)$ which will be helpful in considerations associated with the employment of the asymptotic expansions

$$\sqrt{\pi}\;Ai(-x) = x^{-1/4}[P(x)\sin(\xi + \pi/4) - Q(x)\cos(\xi + \pi/4)] \tag{B-43}$$

$$\sqrt{\pi}\;Bi(-x) = x^{-1/4}[P(x)\cos(\xi + \pi/4) + Q(x)\sin(\xi + \pi/4)] \tag{B-44}$$

where, if we let

$$\xi = \frac{2}{3} x^{3/2}$$

the functions $P(\xi)$ and $Q(\xi)$ are the asymptotic expansions

$$P(x) = 1 - \frac{5\times7\times9\times11}{2!(216)^2\xi^2} + \frac{9\times11\times13\times15\times17\times19\times21\times23}{4!(216)^4\xi^4} + \cdots \qquad \text{(B-43)}$$

$$Q(x) = \frac{3\times5}{1!(216)^1\xi^1} - \frac{7\times9\times11\times13\times15\times17}{3!(216)^3\xi^3} + \cdots \qquad \text{(B-44)}$$

The reader will observe that Eqs. (B-41) and (B-42) and Eqs. (B-43) and (B-44) involve the expressions

$$A_1(x) = \frac{3\times5}{1!(216)^1\xi^1}$$

$$A_2(x) = \frac{5\times7\times9\times11}{2!(216)^2\xi^2}$$

$$A_3(x) = \frac{7\times9\times11\times13\times15\times17}{3!(216)^3\xi^3}$$

In Table B-19 we present the values of $A_n(x)$ for n = 1(1)30 for x = 4.0(1.0)8.0. These tables have been retyped from computer output so as to present the numbers in "fixed point" format, i.e., as decimal digits which are not multiplied at powers of 10. This display of the behavior of the $A_n(x)$ provides a graphic picture of the asymptotic nature of series in the classical asymptotic expansions for the Airy functions. In Table B-20 we present the values of $A_n(x)$ for n = 1(1)14 for x = 1.0(1.0)20.0. These results have been reproduced from computer output and appear as "floating point" numbers. It is hoped that the availability of these tables which present the values of the successive terms in the series will attract the interest of people who have been interested in summation techniques for asymptotic series.

```
      BLOCK DATA
C     DOUBLE PRECISION EAIRYF(100),EBIRYF(100),EAIRYD(100),EBIRYD(100)
      COMMON/EQUADP/   EAIRYF(100),EBIRYF(100),EAIRYD(100),EBIRYD(100)
      DOUBLE PRECISION EAIRYF,EBIRYF,EAIRYD,EBIRYD
      DATA (EAIRYF(I),I= 1,20 )/
     A +0.32920312994353810016 7D+00, +0.30370315428638199488 8D+00,      BIRYF 51
     B +0.27880648195500492193 7D+00, +0.25474235429567634607 9D+00,      BIRYF 52
     C +0.23169360648083348976 6D+00, +0.20980006166637946834 6D+00,      BIRYF 53
     D +0.18916240039815007330 3D+00, +0.16984631744436485921 8D+00,      BIRYF 54
     E +0.15188680364054436109 4D+00, +0.13529241631288141552 5D+00,      BIRYF 55
     F +0.12004942735539766797 1D+00, +0.10612576226331254273 6D+00,      BIRYF 56
     G +0.93474665771502709861 8D-01, +0.82038049807610677630 0D-01,      BIRYF 57
     H +0.71749497008105409675 5D-01, +0.62536907968931960324 3D-01,      BIRYF 58
     I +0.54324792732919467751 1D-01, +0.47036216866845805227 1D-01,      BIRYF 59
     J +0.40594420031529496664 2D-01, +0.34924130423274379128 5D-01/      BIRYF 60
      DATA (EAIRYF(I),I=21,40 )/
     A +0.29952602115866526607 1D-01, +0.25610404421773219541 4D-01,      BIRYF 61
     B +0.21831993180622632369 0D-01, +0.18556093622975467356 4D-01,      BIRYF 62
     C +0.15725923380470490002 9D-01, +0.13289282529671484168 0D-01,      BIRYF 63
     D +0.11198535451065880947 1D-01, +0.94105069149239629606 6D-02,      BIRYF 64
     E +0.78863123041212308399 2D-02, +0.65911393574607191148 6D-02,      BIRYF 65
     F +0.54939963491868373173 6D-02, +0.45674392740398209069 1D-02,      BIRYF 66
     G +0.37872884268267532691 8D-02, +0.31323427799042077955 3D-02,      BIRYF 67
     H +0.25840987869896349475 4D-02, +0.21264786826381708314 3D-02,      BIRYF 68
     I +0.17455720006099791352 2D-02, +0.14293928921077907950 8D-02,      BIRYF 69
     J +0.11676548729914494880 5D-02, +0.95156385120480187348 6D-03/      BIRYF 70
      DATA (EAIRYF(I),I=41,60 )/
     A +0.77362966378159718693 4D-03, +0.62749586830916337476 7D-03,      BIRYF 71
     B +0.50778716815614929258 1D-03, +0.40997358638696215596 6D-03,      BIRYF 72
     C +0.33025032351430898362 1D-03, +0.26543212392445024288 6D-03,      BIRYF 73
     D +0.21286092135859752183 8D-03, +0.17032552328643494846 9D-03,      BIRYF 74
     E +0.13599211701506753694 0D-03, +0.10834442813607441733 7D-03,      BIRYF 75
     F +0.86132427064788441011 3D-04, +0.68328555925248100896 4D-04,      BIRYF 76
     G +0.54090531013400567112 3D-04, +0.42729861694116620324 5D-04,      BIRYF 77
     H +0.33685311908598144214 0D-04, +0.26500613296849970985 4D-04,      BIRYF 78
     I +0.20805817713260686112 7D-04, +0.16301750585877252927 4D-04,      BIRYF 79
     J +0.12747094509184487559 2D-04, +0.99476943602528895690 7D-05/      BIRYF 80
```

```
      DATA (EAIRYF(I),I=61,80 )/
     A +0.7747731032448427535420-05, +0.6022460719688198216460-05,     BIRYF 81
     B +0.4672260820574287162440-05, +0.3617762318851802992420-05,     BIRYF 82
     C +0.2795882343204913585140-05, +0.2156599952596919985460-05,     BIRYF 83
     D +0.1660343478187535373500-05, +0.1275879416876668148380-05,     BIRYF 84
     E +0.9786113339266037630050-06, +0.7492128863997167079880-06,     BIRYF 85
     F +0.5725322885877657257840-06, +0.4367166359142262286050-06,     BIRYF 86
     G +0.3325137824437757599820-06, +0.2527171939266752444640-06,     BIRYF 87
     H +0.1917256067513430751420-06, +0.1451946174801253705780-06,     BIRYF 88
     I +0.1097616850202778423170-06, +0.8282960066226890174600-07,     BIRYF 89
     J +0.6239640097283940477980-07, +0.4692207616099231625120-07/     BIRYF 90
      DATA (EAIRYF(I),I=81,100)/
     A +0.3522435623573567884570-07, +0.2639741834028278329010-07,     BIRYF 91
     B +0.1974861749667693136270-07, +0.1479355499471921868320-07,     BIRYF 92
     C +0.1099700975519550650850-07, +0.8185506151345623966920-08,     BIRYF 93
     D +0.6082608218774544496880-08, +0.4512440519153703554440-08,     BIRYF 94
     E +0.3342061042518703485810-08, +0.2471168430872489843080-08,     BIRYF 95
     F +0.1824228253564026075190-08, +0.1344621833707132745200-08,     BIRYF 96
     G +0.9892686613164267493950-09, +0.7267411770779201682370-09,     BIRYF 97
     H +0.5330263704617491638220-09, +0.3903233530415135479830-09,     BIRYF 98
     I +0.2853715931493100113700-09, +0.2083102822430873119120-09,     BIRYF 99
     J +0.1518195814104911539430-09, +0.1104753255289870539770-09/     BIRYF100
      DATA (EBIRYF(I),I= 1,20 )/
     A +0.6598616901941892336590+00, +0.7054642029186612354000+00,     BIRYF  1
     B +0.7524855850873156380040+00, +0.8017730000135972398350+00,     BIRYF  2
     C +0.8542770431031554932940+00, +0.9110633416949404818130+00,     BIRYF  3
     D +0.9733286558781659366820+00, +0.1042422171231560700410+01,     BIRYF  4
     E +0.1119872813134447244010+01, +0.1207423594952871259450+01,     BIRYF  5
     F +0.1307074247516306954960+01, +0.1421133675610348069500+01,     BIRYF  6
     G +0.1552284162344543746450+01, +0.1703659711538680882510+01,     BIRYF  7
     H +0.1878941503747895000940+01, +0.2082474171254111889800+01,     BIRYF  8
     I +0.2319407506938924947300+01, +0.2595869356743906290120+01,     BIRYF  9
     J +0.2919176863042108953160+01, +0.3298094999978214710350+01/     BIRYF 10
      DATA (EBIRYF(I),I=21,40 )/
     A +0.3743153564956175138160+01, +0.4267036581766644730020+01,     BIRYF 11
     B +0.4885061581835644686720+01, +0.5615770654120597364950+01,     BIRYF 12
     C +0.6481660738460578608210+01, +0.7510087698082272365320+01,     BIRYF 13
     D +0.8734387649988913889580+01, +0.1019527037763705519690+02,     BIRYF 14
```

AIRY FUNCTIONS ON POSITIVE AXIS

```
      E +0.11942554067750536993 2D+02, +0.14037328963730232032 1D+02,   BIRYF 15
      F +0.16554660922469495368 3D+02, +0.19586975733441185124 2D+02,   BIRYF 16
      G +0.23248303262941579479 7D+02, +0.27679609423659011366 3D+02,   BIRYF 17
      H +0.33055506754611479415 7D+02, +0.39592715074025013650 0D+02,   BIRYF 18
      I +0.47560747499589442857 0D+02, +0.57295430971052160129 1D+02,   BIRYF 19
      J +0.69216043201447079638 2D+02, +0.83847071408468139924 9D+02/   BIRYF 20
        DATA (EBIRYF(I),I=41,60 )/
     A +0.10184588752305113796 5D+03, +0.12403800986864210685 2D+03,   BIRYF 21
     B +0.15146210883707224410 4D+03, +0.18542754839855758803 5D+03,   BIRYF 22
     C +0.22758808183559971847 0D+03, +0.28003639880129145404 3D+03,   BIRYF 23
     D +0.34542563075723357214 9D+03, +0.42712576758084800319 7D+03,   BIRYF 24
     E +0.52942535802223195437 9D+03, +0.65779204417117118249 5D+03,   BIRYF 25
     F +0.81920965867996127747 5D+03, +0.10226151169136374873 0D+04,   BIRYF 26
     G +0.12794655954684410401 6D+04, +0.16044760782413450248 4D+04,   BIRYF 27
     H +0.20165800386595313945 3D+04, +0.25401828375814989053 5D+04,   BIRYF 28
     I +0.32067997242347116960 7D+04, +0.40571999739612470156 3D+04,   BIRYF 29
     J +0.51442181542807988145 3D+04, +0.65364461048098634543 2D+04/   BIRYF 30
        DATA (EBIRYF(I),I=61,80 )/
     A +0.83230894240154933337 88D+04, +0.10620366113617851116 8D+05,  BIRYF 31
     B +0.13579950691419307528 7D+05, +0.17400135568084821470 8D+05,   BIRYF 32
     C +0.22340607718396998160 0D+05, +0.28742042901040523044 2D+05,   BIRYF 33
     D +0.37052129373357501864 8D+05, +0.47860185574291982244 3D+05,   BIRYF 34
     E +0.61943282486771132753 7D+05, +0.80327790709430247013 8D+05,   BIRYF 35
     F +0.10437163938045399852 3D+06, +0.13587442849813917848 9D+06,   BIRYF 36
     G +0.17722505516442812635 0D+06, +0.23159994089891552781 8D+06,   BIRYF 37
     H +0.30322961511253340232 2D+06, +0.39775777780342409613 0D+06,   BIRYF 38
     I +0.52272566398783376136 1D+06, +0.68822643370722585050 8D+06,   BIRYF 39
     J +0.90779061606199380915 0D+06, +0.11995860041244599310 3D+07/   BIRYF 40
        DATA (EBIRYF(I),I=81,100)/
     A +0.15880461279294299545 3D+07, +0.21060837099317048127 8D+07,   BIRYF 41
     B +0.27981037510671466248 3D+07, +0.37241112472959810232 3D+07,   BIRYF 42
     C +0.49653195414713019815 1D+07, +0.66318188792060827961 5D+07,   BIRYF 43
     D +0.88730774696071810630 9D+07, +0.11892342454717134287 3D+08,   BIRYF 44
     E +0.15966418120232306481 9D+08, +0.21472868891435349095 0D+08,   BIRYF 45
     F +0.28927488903264973132 5D+08, +0.39035987736433834489 8D+08,   BIRYF 46
     G +0.52765389153991047572 0D+08, +0.71442803588713036963 8D+08,   BIRYF 47
     H +0.96892265580451092833 0D+08, +0.13162457704426055782 4D+09,   BIRYF 48
     I +0.17910106467773553106 8D+09, +0.24410055050457074447 8D+09,   BIRYF 49
     J +0.33323064825676906841 8D+09, +0.45564115354822514108 5D+09/   BIRYF 50
```


+45

```
      DATA (EAIRYD(I),I= 1,20 )/
     A -0.25713042190758618869 4D+00, -0.25240547028561953611 3D+00,      AIRYF 51
     B -0.24514636421905480343 5D+00, -0.23583203441920821727 5D+00,      AIRYF 52
     C -0.22491053266468389313 6D+00, -0.21279325938915852051 3D+00,      AIRYF 53
     D -0.19985119158228047517 4D+00, -0.18641286380727170902 3D+00,      AIRYF 54
     E -0.17276384346163467386 6D+00, -0.15914744129679321278 9D+00,      AIRYF 55
     F -0.14576640734501625283 5D+00, -0.13278537855722617420 7D+00,      AIRYF 56
     G -0.12033386559018358247 3D+00, -0.10850959048013916365 8D+00,      AIRYF 57
     H -0.97382012842301319219 8D-01, -0.86995908448104131813 0D-01,      AIRYF 58
     I -0.77374889525325028085 8D-01, -0.68524780118610934566 0D-01,      AIRYF 59
     J -0.60436781785756540153 5D-01, -0.53090384433653631715 2D-01/      AIRYF 60
      DATA (EAIRYD(I),I=21,40 )/
     A -0.46455994032674599469 7D-01, -0.40497263244453135269 0D-01,      AIRYF 61
     B -0.35173122720818063130 3D-01, -0.30439520128972592688 2D-01,      AIRYF 62
     C -0.26250881035903230356 7D-01, -0.22561310886108747630 4D-01,      AIRYF 63
     D -0.19325560692377637544 7D-01, -0.16499780994915139731 5D-01,      AIRYF 64
     E -0.14042089387786424733 4D-01, -0.11912976705951318514 1D-01,      AIRYF 65
     F -0.10075576032316879519 1D-01, -0.84958172185685955856 2D-02,      AIRYF 66
     G -0.71424877858847379925 0D-02, -0.59872190251137937938 1D-02,      AIRYF 67
     H -0.50044139679525826459 2D-02, -0.41711317444193810236 5D-02,      AIRYF 68
     I -0.34669407490276282004 4D-02, -0.28737510609047910600 0D-02,      AIRYF 69
     J -0.23756347375622522185 2D-02, -0.19586409502041788995 6D-02/      AIRYF 70
      DATA (EAIRYD(I),I=41,60 )/
     A -0.16106114612269867785 5D-02, -0.13210006638876865555 9D-02,      AIRYF 71
     B -0.10807033052246403308 3D-02, -0.88189208649176807236 9D-03,      AIRYF 72
     C -0.71786656755750888609 3D-03, -0.58291417781033317107 2D-03,      AIRYF 73
     D -0.47218363998626423999 7D-03, -0.38157072868873844049 1D-03,      AIRYF 74
     E -0.30761599633764974329 7D-03, -0.24741389086846247597 1D-03,      AIRYF 75
     F -0.19853254788180524130 8D-03, -0.15894345264594751644 4D-03,      AIRYF 76
     G -0.12696012333659671728 3D-03, -0.10118495655699361226 9D-03,      AIRYF 77
     H -0.80463391305565143370 2D-04, -0.63844581246177234678 7D-04,      AIRYF 78
     I -0.50547811684537210944 4D-04, -0.39933952437098958392 1D-04,      AIRYF 79
     J -0.31481297117112764236 3D-04, -0.24765200397034954751 3D-04/      AIRYF 80
      DATA (EAIRYD(I),I=61,80 )/
     A -0.19440985375102954124 8D-04, -0.15229651696941566672 3D-04,      AIRYF 81
     B -0.11905970459957270811 4D-04, -0.92886034448629829034 4D-05,      AIRYF 82
     C -0.72319314666017925589 7D-05, -0.56193194434578581449 D-05,      AIRYF 83
     D -0.43575841632977717417 7D-05, -0.33724647753763918519 9D-05,      AIRYF 84
```

```
E -0.26049260870862621552 7D-05, -0.20081508947387919909 3D-05,      AIRYF 85
F -0.15451003667897689471 0D-05, -0.11865410717176396516 2D-05,      AIRYF 86
G -0.90945403888334594451 4D-06, -0.69575554020805930728 0D-06,      AIRYF 87
H -0.53127139597205446841 7D-06, -0.40491682045077797699 4D-06,      AIRYF 88
I -0.30804239001035928623 4D-06, -0.23391386990032383902 1D-06,      AIRYF 89
J -0.17729958329430352741 7D-06, -0.13414392979067865741 3D-06/      AIRYF 90
 DATA (EAIRYD(I),I=81,100)/
A -0.10130972032660834050 5D-06, -0.76375329841861791263 6D-07,      AIRYF 91
B -0.57475397363380090271 1D-07, -0.43176027504858731478 9D-07,      AIRYF 92
C -0.32377254404476022552 0D-07, -0.24236992122415316101 1D-07,      AIRYF 93
D -0.18111876046176122803 3D-07, -0.13511349359955728625 9D-07,      AIRYF 94
E -0.10062109921836922694 8D-07, -0.74806413896589463957 1D-08,      AIRYF 95
F -0.55520373443859135251 8D-08, -0.41137124428079200209 3D-08,      AIRYF 96
G -0.30428998756186507745 7D-08, -0.22470755570506690492 5D-08,      AIRYF 97
H -0.16566394593740665525 7D-08, -0.12193337781681122595 9D-08,      AIRYF 98
I -0.89599458489931542399 5D-09, -0.65732504194716696680 9D-09,      AIRYF 99
J -0.48144951964682449351 9D-09, -0.35206336767389239548 7D-09/      AIRYF100
 DATA (EBIRYD(I),I= 1,20 )/
A +0.45151263114964650127 3D+00, +0.46178928436215093039 2D+00,      AIRYF  1
B +0.48004902875244802216 1D+00, +0.50728167605062244342 3D+00,      AIRYF  2
C +0.54457256414059230182 6D+00, +0.59314447863428570749 2D+00,      AIRYF  3
D +0.65440591917214003029 1D+00, +0.73000690161525176703 2D+00,      AIRYF  4
E +0.82190389030720902608 0D+00, +0.93243593339277563296 9D+00,      AIRYF  5
F +0.10644146482419911093 8D+01, +0.12212313987048950137 8D+01,      AIRYF  6
G +0.14069858538563174731 6D+01, +0.16266411607804522590 3D+01,      AIRYF  7
H +0.18862122548481654887 3D+01, +0.21929954372537653714 2D+01,      AIRYF  8
I +0.25558493569004380161 7D+01, +0.29855400508465990729 0D+01,      AIRYF  9
J +0.34951656628916143290 3D+01, +0.41006820499328898894 8D+01/      AIRYF 10
 DATA (EBIRYD(I),I=21,40 )/
A +0.48215499257174953555 5D+01, +0.56815417695852521672 4D+01,      AIRYF 11
B +0.67097408127238260494 9D+01, +0.79417858797353189804 7D+01,      AIRYF 12
C +0.94214233173343017557 7D+01, +0.11202445467843920930 1D+02,      AIRYF 13
D +0.13351116152330930918 2D+02, +0.15949210720073756977 8D+02,      AIRYF 14
E +0.19097832882836592643 9D+02, +0.22922214966382170185 8D+02,      AIRYF 15
F +0.27577765207770731447 1D+02, +0.33257698980545098963 5D+02,      AIRYF 16
G +0.40202685120884542480 2D+02, +0.48713060081951407077 7D+02,      AIRYF 17
H +0.59164319581360987037 1D+02, +0.72026800334887057303 9D+02,      AIRYF 18
I +0.87890727262833410890 6D+02, +0.10749814238907310360 3D+03,      AIRYF 19
J +0.13178367436912177763 0D+03, +0.16192668350461340184 7D+03/      AIRYF 20
```

```
 DATA (EBIRYD(I),I=41,60 )/
A +0.19941806743320975343 2D+03, +0.24614599171178559262 0D+03,      AIRYF 21
B +0.30450608885753505724 4D+03, +0.37754334370077789968 5D+03,      AIRYF 22
C +0.46913507732796639796 9D+03, +0.58422732232556571145 4D+03,      AIRYF 23
D +0.72914066855334527957 0D+03, +0.91196664368974605428 9D+03,      AIRYF 24
E +0.11430822653649948566 3D+04, +0.14358190802179825187 9D+04,      AIRYF 25
F +0.18073344813513677844 5D+04, +0.22797482935833352424 0D+04,      AIRYF 26
G +0.28816277721431462519 6D+04, +0.36499309318307159566 2D+04,      AIRYF 27
H +0.46325537331390424206 7D+04, +0.58916740862081323352 5D+04,      AIRYF 28
I +0.75081489113299978487 2D+04, +0.95873050303703078842 6D+04,      AIRYF 29
J +0.12266577761776943280 6D+05, +0.15725602621930476840 7D+05/      AIRYF 30
 DATA (EBIRYD(I),I=61,80 )/
A +0.20199568849304028408 5D+05, +0.25996916653612601523 3D+05,      AIRYF 31
B +0.33522827688550219097 3D+05, +0.43310453588538678801 8D+05,      AIRYF 32
C +0.56062495842522860753 5D+05, +0.72706653568706398278 6D+05,      AIRYF 33
D +0.94469678154312250938 2D+05, +0.12297643030844547499 3D+06,      AIRYF 34
E +0.16038258323096269439 7D+06, +0.20955267087397131952 7D+06,      AIRYF 35
F +0.27429933826435646503 1D+06, +0.35970531748976032441 1D+06,      AIRYF 36
G +0.47255738639870335073 1D+06, +0.62193213587641413747 0D+06,      AIRYF 37
H +0.81998783535879962100 8D+06, +0.10830365079310609034 3D+07,      AIRYF 38
I +0.14329975282373650337 2D+07, +0.18993714378644851420 6D+07,      AIRYF 39
J +0.25219241139567816834 1D+07, +0.33543423127445388769 2D+07/      AIRYF 40
 DATA (EBIRYD(I),I=81,100)/
A +0.44692194243083495314 2D+07, +0.59648654324238731605 3D+07,      AIRYF 41
B +0.79746220844948902620 5D+07, +0.10679623386635035452 1D+08,      AIRYF 42
C +0.14326301030662058335 4D+08, +0.19250433795750532407 8D+08,      AIRYF 43
D +0.25910234760755175571 7D+08, +0.34931938027946960148 9D+08,      AIRYF 44
E +0.47172696726445880962 6D+08, +0.63807489780908213859 4D+08,      AIRYF 45
F +0.86449372333449287978 4D+08, +0.11731609833731916670 2D+09,      AIRYF 46
G +0.15946132356019694092 2D+09, +0.21709587056397884317 8D+09,      AIRYF 47
H +0.29603476386800503866 8D+09, +0.40432014156182941256 5D+09,      AIRYF 48
I +0.55309044723402311177 7D+09, +0.75779495800325827182 3D+09,      AIRYF 49
J +0.10398932020639825052 2D+10, +0.14292361344828657763 8D+10/      AIRYF 50
 END
```

```
      SUBROUTINE AIBIZ(Z,AI,BI,AIP,BIP)
      COMPLEX           Z,AI,BI,AIP,BIP
      DOUBLE PRECISION AZERO,A(22),BZERO,B(22),CZERO,C(22),DZERO,D(22)
      DOUBLE PRECISION DRZ,DIZ,D2RZ,D2IZ,D3RZ,D3IZ,FR,FI,FPR,FPI,GR,GI,
     A       GPR,GPI,AREAL(23),AIMAG(23),ZERO,TWO,CA,CB,
     B       DAIR,DAII,DAIPR,DAIPI,DBIR,DBII,DBIPR,DBIPI
      DATA AZERO,BZERO,(A(M),B(M),M=1,11) /
     A             0.93043671693   D  0 ,        0.67829872514   D  0 ,
     B            31.01455723097   D  0 ,       11.30497875240   D  0 ,
     C           206.76371487316   D  0 ,       53.83323215431   D  0 ,
     D           574.34365242545   D  0 ,      119.62940478735   D  0 ,
     E           870.21765519008   D  0 ,      153.37103177865   D  0 ,
     F           828.77871922864   D  0 ,      127.80919314888   D  0 ,
     G           541.68543740434   D  0 ,       74.74221821572   D  0 ,
     H           257.94544638302   D  0 ,       32.35593862152   D  0 ,
     I            93.45849506631   D  0 ,       10.78531287384   D  0 ,
     J            26.62635187074   D  0 ,        2.85325737403   D  0 ,
     K             6.121000430056  D  0 ,        0.613603736351  D  0 ,
     L             1.159280384480  D  0 ,        0.109376780098  D  0 /
      DATA (A(M),B(M),M=12,22)                 /
     A             0.184012759441  D  0 ,        0.16422939955   D- 1 ,
     B             0.24833030964   D- 1 ,        0.2105505122    D- 2 ,
     C             0.2884208010    D- 2 ,        0.233167788     D- 3 ,
     D             0.291334142     D- 3 ,        0.22528289      D- 4 ,
     E             0.25827495      D- 4 ,        0.1915671       D- 5 ,
     F             0.2025686       D- 5 ,        0.144470        D- 6 ,
     G             0.141557        D- 6 ,        0.9729          D- 8 ,
     H             0.8870          D- 8 ,        0.589           D- 9 ,
     I             0.501           D- 9 ,        0.32            D-10 ,
     J             0.26            D-10 ,        0.2             D-11 ,
     K             0.1             D-11 ,        0.0             D  0 /
      DATA CZERO,DZERO,(C(M),D(M),M=1,11) /
     A             0.46521835846   D  0 ,        0.67829872514   D  0 ,
     B             6.20291144619   D  0 ,       45.21991500962   D  0 ,
     C            25.84546435915   D  0 ,      376.83262508015   D  0 ,
     D            52.21305931140   D  0 ,     1196.29404787350   D  0 ,
     E            62.15840394215   D  0 ,     1993.82341312250   D  0 ,
     F            48.75168936639   D  0 ,     2044.94709038206   D  0 ,
     G            27.08427187022   D  0 ,     1420.10214609865   D  0 ,
```

Program B-1 (Cont'd)

SUBROUTINE AIBIZ(Z,AI,BI,AIP,BIP)

```
H  11.215019407960  D 0 ,    711.830649673510  D 0 ,
I     3.594557502550  D 0 ,    269.632821846030  D 0 ,
J     0.918150064510  D 0 ,     79.891206472900  D 0 ,
K     0.191281263439  D 0 ,     19.021715826880  D 0 ,
L     0.331222966990  D- 1 ,    3.718810523339  D 0 /
   DATA (C(M),D(M),M=12,22)
   /
A     0.484244103800  D- 2 ,    0.607648778323  D 0 ,
B     0.605683682000  D- 3 ,    0.842202048960  D- 1 ,
C     0.655501820000  D- 4 ,    0.100262148690  D- 1 ,
D     0.619859900000  D- 5 ,    0.103630127800  D- 2 ,
E     0.516550000000  D- 6 ,    0.938678690000  D- 4 ,
F     0.382200000000  D- 7 ,    0.751243500000  D- 5 ,
G     0.252800000000  D- 8 ,    0.535074000000  D- 6 ,
H     0.150000000000  D- 9 ,    0.341350000000  D- 7 ,
I     0.800000000000  D-11 ,    0.196200000000  D- 8 ,
J     0.000000000000  D 0 ,     0.102000000000  D- 9 ,
K     0.000000000000  D 0 ,     0.500000000000  D-11 /
   DATA ZERO,TWO,CA,CB/0.0D0,2.0D0,0.66090107608300,0.38157141418400/
110 FORMAT(25H AIBIZ CANNOT HANDLE Z = ,(E15.8,E15.8))
    IF(CABS(Z).GT.(6.1)) GO TO 99
    M=3+INT(20.*CABS(Z)/6.1)
    DRZ=DBLE(REAL(-Z))
    DIZ=DBLE(REAL((0.,1.)*Z))
    D2RZ=DRZ*DRZ-DIZ*DIZ
    D2IZ=TWO*DRZ*DIZ
    D3RZ=DRZ*D2RZ-DIZ*D2IZ
    D3IZ=DRZ*D2IZ+DIZ*D2RZ
    D3RZ = D3RZ/(-200.0D0)
    D3IZ = D3IZ/(-200.0D0)
    DO 39  I=1,23
    AREAL(I)= ZERO
    AIMAG(I)= ZERO
39  CONTINUE
    IF(M.GT.22)  M=22
    DO 1 N=M,1,-1
    AREAL(N)=D3RZ*AREAL(N+1) - D3IZ*AIMAG(N+1) + A(N)
    AIMAG(N)=D3RZ*AIMAG(N+1) + D3IZ*AREAL(N+1)
1   CONTINUE
```

Program B-1 (Cont'd)

SUBROUTINE AIBIZ(Z,AI,BI,AIP,BIP)

```
      FR = D3RZ*AREAL(1)-D3IZ*AIMAG(1)+AZERO
      FI = D3RZ*AIMAG(1)+D3IZ*AREAL(1)
      DO 2 N=M,1,-1
      AREAL(N)=D3RZ*AREAL(N+1) - D3IZ*AIMAG(N+1)  + B(N)
      AIMAG(N)=D3RZ*AIMAG(N+1)  + D3IZ*AREAL(N+1)
    2 CONTINUE
      GPR= D3RZ*AREAL(1)-D3IZ*AIMAG(1)+BZERO
      GPI= D3RZ*AIMAG(1)+D3IZ*AREAL(1)
      GR = DRZ*GPR-DIZ*GPI
      GI = DRZ*GPI+DIZ*GPR
      DO 3 N=M,1,-1
      AREAL(N)=D3RZ*AREAL(N+1)-D3IZ*AIMAG(N+1)+C(N)
      AIMAG(N)=D3RZ*AIMAG(N+1)+D3IZ*AREAL(N+1)
    3 CONTINUE
      GPR = D3RZ*AREAL(1)-D3IZ*AIMAG(1)+CZERO
      GPI = D3RZ*AIMAG(1)+D3IZ*AREAL(1)
      FPR =-D2RZ*GPR + D2IZ*GPI
      FPI =-D2RZ*GPI - D2IZ*GPR
      DO 4 N=M,1,-1
      AREAL(N)=D3RZ*AREAL(N+1)-D3IZ*AIMAG(N+1)+D(N)
      AIMAG(N)=D3RZ*AIMAG(N+1)+D3IZ*AREAL(N+1)
    4 CONTINUE
      GPR = D3RZ*AREAL(1)-D3IZ*AIMAG(1)+DZERO
      GPI = D3RZ*AIMAG(1)+D3IZ*AREAL(1)
      WRITE(6,1000) FR,FI,GR,GI,FPR,FPI,GPR,GPI
C-----NOW CONSTRUCT MILLER AIRY FUNCTIONS
      DAIR   = CB*(FR + GR)
      DAII   = CB*(FI + GI)
      DAIPR =-CB*(FPR + GPR)
      DAIPI =-CB*(FPI + GPI)
      DBIR   = CA*(FR - GR)
      DBII   = CA*(FI - GI)
      DBIPR =-CA*(FPR - GPR)
      DBIPI =-CA*(FPI - GPI)
      AI     = CMPLX(SNGL(DAIR),SNGL(DAII))
      AIP    = CMPLX(SNGL(DAIPR),SNGL(DAIPI))
      BI     = CMPLX(SNGL(DBIR),SNGL(DBII))
      BIP    = CMPLX(SNGL(DBIPR),SNGL(DBIPI))
      RETURN
   99 WRITE(6,110)  Z
      CALL EXIT
      END
```

```
      SUBROUTINE ASYPOS(X,AL,AM,AI,AIP,BL,BM,BI,BIP,ERROR,FLOVER)
C-----THIS SUBROUTINE EVALUATES THE CLASSICAL ASYMPTOTIC EXPANSION OF
C-----THE AIRY FUNCTION. THE LOGICAL VARIABLE AUNDER IS TRUE WHEN THE
C-----ARGUMENT X IS SO LARGE THAT AI AND ADP ARE LESS THAN 1.0E-38.
C-----IF THE TERMS IN THE ASYMPTOTIC EXPANSIONS DO NOT DECEASE TO
C-----1.0D-17, THE VALUE OF THE LAST TERM EMPLOYED WILL BE FOUND STORED
C-----IN ERROR.
      LOGICAL FLOVER
      DOUBLE PRECISION  X,AL,AM,AI,AIP,BL,BM,BI,BIP,ERROR
      DOUBLE PRECISION DFLOAT,DSQRT,DEXP,Z,XI,D72XI,RL,RM,SL,SM,TL,TM,W
      DATA             ZERO,ONE,D1P5,D5,D7,D203,D10M17,D88,PI,D72
     1 1.0D0,1.5D0,5.0D0,7.0D0,0.66666666666666667D0,1.0D-18,88.0D0,
     2 3.14159265358979320D0,72.0D0/
  110 FORMAT(6H    I =,I2,8H    TL = ,D22.16,8H    TM = ,D22.16)
  111 FORMAT(6H    I =,I2,8H    SL = ,D22.16,8H    SM = ,D22.16)
  120 FORMAT(11H0  FOR X = ,D22.16,59H SUMMING 40 TERMS DID NOT LEAD TO
     A ATERMS AS SMALL AS 1.0D-17//8H    TL = ,D22.16,8H    TM = ,D22.16/
     B                            8H    AL = ,D22.16,8H    AM = ,D22.16)
  121 FORMAT(11H0  FOR X = ,D22.16,59H SUMMING 40 TERMS DID NOT LEAD TO
     A ATERMS AS SMALL AS 1.0D-17//8H    SL = ,D22.16,8H    SM = ,D22.16/
     B                            8H    BL = ,D22.16,8H    BM = ,D22.16)
  199 FORMAT(11H0  FOR X = ,D22.16,33H TERMS STARTED TO GROW AT INDEX =,
     A  I3,41H AND PROGRAM LEFT THE DO-LOOP WITH VALUES//
     B                            8H    RL = ,D22.16,8H    RM = ,D22.16/
     C                            8H    TL = ,D22.16,8H    TM = ,D22.16)
  998 FORMAT(69H1  PROGRAM EXIT BECAUSE SUBROUTINE ASYPOS(X,...) WILL NO
     AT ACCEPT X = ,D22.16)
      DFLOAT(N) = DBLE(FLOAT(N))
      IF(X.LT.D10M17) GO TO 999
C-----X MUST BE POSITIVE
      FLOVER = .FALSE.
      Z      = DSQRT(X)
      XI     = D203*(Z**3)
      D72XI  =-D72*XI
      AL     = ONE
      AM     = ONE
```

```
      BL      = ONE
      BM      = ONE
      TL      = D5/D72XI
      TM      =-D7/D72XI
      SL      =-TL
      SM      =-TM
      AL      = AL + TL
      AM      = AM + TM
      BL      = BL + SL
      BM      = BM + SM
      ERROR   = DABS(TL)
      DO 10 K=1,40
      INDEX   = K
      RL      = DFLOAT((6*K+1)*(6*K+5))/(D72XI*DFLOAT(K+1))
      RM      = DFLOAT((6*K-1)*(6*K+7))/(D72XI*DFLOAT(K+1))
      IF(DABS(RL).GT.ONE) GO TO 99
C-----TRANSFER OUT WHEN TERMS START TO INCREASE
      TL      = RL*TL
      TM      = RM*TM
      SL      = -RL*SL
      SM      = -RM*SM
      AL      = AL + TL
      AM      = AM + TM
      BL      = BL + SL
      BM      = BM + SM
      IF(DABS(TL).LT.D10M17) GO TO 89
C-----TRANSFER OUT WHEN TERMS ARE SMALLER THAN 1.0D-17
      ERROR   = DABS(TL)
      WRITE(6,110) INDEX,TL,TM
      WRITE(6,111) INDEX,SL,SM
   10 CONTINUE
      WRITE(6,120) X,TL,TM,AL,AM
      WRITE(6,121) X,SL,SM,BL,BM
C-----OUTPUT A MESSAGE THAT COMPLETION OF SUM OF 40 TERMS DID NOT
C-----INCLUDE SUFFICIENT TERMS TO HAVE DABS(TL) GO BELOW 1.0E-17.
   99 WRITE(6,199) X,INDEX,RL,RM,TL,TM
```

```
   89 IF(XI.GT.86.0D0) GO TO 20
      XI   = DEXP(XI)
      W    = DSQRT(Z*PI)
      Z    = 2.0D0*W*XI
      CALL OVERFL(JOVER)
      IF(JOVER.EQ.1) GO TO 20
      W    = XI/W
      AI   = AL/Z
      AIP  =-(AM/Z)*DSQRT(X)
      BI   = BL*W
      BIP  = BM*W*DSQRT(X)
      RETURN
   20 AI   = ZERO
      AIP  = ZERO
      BI   = 1.0D+38
      BIP  = 1.0D+38
C-----WHEN AIRY FUNCTIONS UNDERFLOW THEY ARE SET EQUAL TO ZERO
      FLOVER = .TRUE.
      RETURN
  999 WRITE(6,998) X
      CALL EXIT
      END
```

Table B-1

THE AIRY FUNCTIONS ($10 < x < 25$)

x	Ai(x)	Bi(x)
10.0	.1104753255289868593546 -9	.4556411535482251406835E 9
10.1	.8026504702909595103846-10	.6246208078751005586706E 9
10.2	.5822567393145971366406-10	.8559919521429397243416 9
10.3	.4217281922626017618806-10	.1176064627812018572156 10
10.4	.3049888186594001301986-10	.1618378355925988532496 10
10.5	.2202274519283401643786-10	.2236554441136695228826 10
10.6	.1587809078467610204956-10	.3079116097411633706416 10
10.7	.1143055704006619006122E-10	.4257124992535575608136 10
10.8	.8216403415359878791036-11	.5894955352414365464516 10
10.9	.5897178542355832097356-11	.8175515560123620780946 10
11.0	.4226275864960359592786-11	.1135578253043047628106 11
11.1	.3024302943390258477106-11	.1579730840830296144186 11
11.2	.2160973464206401724706-11	.2206949657963227378916 11
11.3	.1541817736556407183006-11	.3071107495362083891456 11
11.4	.1098446317760130698176-11	.4291753611130391608196 11
11.5	.7814290183962854347616-12	.6006568015889603654826 11
11.6	.5550956150241364982696-12	.8419124146271474906786 11
11.7	.3937460927991327741416-12	.1181825636207017335756 12
11.8	.2788929386938675794166-12	.1661433640266314817876 12
11.9	.1972577843025202799846-12	.2339121892419696753236 12
12.0	.1393184688875360839166-12	.3298072258290741761556 12
12.1	.9825703489721669599696-13	.4656956415799357210546 12
12.2	.6919927013665690639616-13	.6585308813868049516966 12
12.3	.4866589575637091911936-13	.9325655972826153889886 12
12.4	.3417713851204040939626-13	.1322541314728766546846 13
12.5	.2396827826078049936816-13	.1878291935622051866976 13
12.6	.1678535402849066328506-13	.2671396401305799377976 13
12.7	.1173867394773565622306-13	.3804805113201950027166 13
12.8	.8197940328162533909746-14	.5426787378459237553196 13
12.9	.5717286166520787109466-14	.7751165486331977631856 13
13.0	.3981776078833335363686-14	.1108670671905940474526 14
13.1	.2769285274543039419246-14	.1587987282606835661786 14
13.2	.1923378383265192324286-14	.2277707673685720484986 14
13.3	.1334043904436132248136-14	.3271546334039509215746 14
13.4	.9240299231256377377636-15	.4705545112262510632466 14
13.5	.6391673876741866651826-15	.6777469026570790720766 14
13.6	.4415277017994101159846-15	.9775066832787969634006 14
13.7	.3045917781195526435726-15	.1411783489044016659646 15
13.8	.2098443778378903728606-15	.2041780754334598085516 15
13.9	.1443768268407858795476-15	.2956926569962840911246 15
14.0	.9920205491192377268286-16	.4288053617865341494506 15

Table B-1 (Cont'd)

THE AIRY FUNCTIONS ($10 < x < 25$)

x	Ai(x)	Bi(x)
14.0	.9920205491192377268288-16	.6226814667522772416538 15
14.1	.6807205687849341421468-16	.6226814667522772416538 15
14.2	.4664918869921745149148-16	.9054314867754664954338 15
14.3	.3192627866595625152678-16	.1318338838918478707238 16
14.4	.2182144145304546435478-16	.1923[illegible]09844194420959028 16
14.5	.1489537454965927195848-16	.2806124832CC504044172E 16
14.6	.1015439812427434527998-16	.4102148199942648738708 16
14.7	.6913430483931700947898-17	.6004671459575673172808 16
14.8	.4700797454945289694808-17	.8801135154033906120388 16
14.9	.3192198248746920689928-17	.1291688158385227532938 17
15.0	.2164962520737992299478-17	.1898209956749358968038 17
15.1	.1466408391804858967658-17	.2793164C928367285103E 17
15.2	.9919854874426999211118-18	.4115402921684211767988 17
15.3	.6701982780894072132938-18	.6071417160935860208458 17
15.4	.4522209409365553027538-18	.8968662880532094818808 17
15.5	.3047538152456012684268-18	.1326549227800920276868 18
15.6	.2051164470567301669458-18	.1964604969518255824938 18
15.7	.1378816228734705248358-18	.2913274919751699110578 18
15.8	.9256963564546363576908-19	.4325539212594780857768 18
15.9	.6207C990076353263458608-19	.6430574783789611089508 18
16.0	.4156888289170243945908-19	.9572123906049186526158 18
16.1	.2780413696180730236288-19	.1426638856618374484178 19
16.2	.1857433775509541683208-19	.2128949011402162701928 19
16.3	.1239315243034588919098-19	.3180974736290147066288 19
16.4	.8258789304858416992758-20	.4758797419299714626318 19
16.5	.5496911172967060764278-20	.7128112715195081312688 19
16.6	.3654183640230173357348-20	.1069031654140924404858 20
16.7	.2426235220630674717528-20	.1605253309941089863648 20
16.8	.1608969730614833049908-20	.2413414617572006749568 20
16.9	.1065704552753245682098-20	.3632904882346292749568 20
17.0	.7050197298388614543168-21	.5475303811330586981818 20
17.1	.4658463916423829588858-21	.8262147069178404931198 20
17.2	.3074416599652194659208-21	.1248264262423132459228 21
17.3	.2026575086929258866508-21	.1888197835392386640598 21
17.4	.1334271347613304927368-21	.2859658924801236123978 21
17.5	.8774220823294709737128-22	.4336159183543006998558 21
17.6	.5763115854826131186o3E-22	.6582924931758761992848 21
17.7	.3780871941448598017258-22	.1000584510513478612968 22
17.8	.2477501007475754212988-22	.1522678902673337313238 22
17.9	.1621527729533390286918-22	.2319963088817650458978 22
18.0	.1060046682524795565698-22	.3538918250356568669798 22

Table B-1 (Cont'd)

THE AIRY FUNCTIONS ($10 < x < 25$)

x	Ai(x)	Bi(x)
18.0	.106004668252479556569E-22	.353891825035656866979E 22
18.1	.692176842161265116230E-23	.54C474526324040891267E 22
18.2	.451442021548144010205E-23	.826406488524846671C57E 22
18.3	.294090577881388059609E-23	.126509912416052574242E 23
18.4	.191362047645285272726E-23	.193894866042865782519E 23
18.5	.124373376697194045779E-23	.29752C9591110722317346 23
18.6	.807415896027368370647E-24	.457C6389412462818654 3E 23
18.7	.523560345022757982345E-24	.702979994427667652976E 23
18.8	.339IC726236531551257 4E-24	.108246617442C08372776E 24
18.9	.219386367536543900949E-24	.166874373649704369106E 24
19.0	.141770437779335271884E-24	.257553555223445854590E 24
19.1	.915C956799708753936 5E-25	.397966656714 C2327415E 24
19.2	.5900C186828420946736 2E-25	.615638430900988868981E 24
19.3	.379968497345214649698E-25	.953461869446631873350E 24
19.4	.244427673798422745413E-25	.147835314814324184241E 25
19.5	.157059056151781837797E-25	.229482196854864866637E 25
19.6	.1008C6063153014384290E-25	.356627195427772272298E 25
19.7	.646282826485243200913E-26	.554847051820414897862E 25
19.8	.413877756396473509949E-26	.864219768 5C5250258610E 25
19.9	.264750186892370283040E-26	.134761529914774766915E 26
20.0	.169167286867054031329E-26	.21C3765C4965110381481E 26
20.1	.107972611148515300070E-26	.328788681557529046869E 26
20.2	.688381387028602595842E-27	.514426774414172293774E 26
20.3	.438393496762 4C279288CE-27	.805779334 1CC291031133E 26
20.4	.278881510801988761086E-27	.126355258865 5C5297776E 27
20.5	.177213635431494210350E-27	.198359929862883354653E 27
20.6	.112465745214810135618E-27	.311743C623218662116 9E 27
20.7	.713217397495334314911E-28	.49C4793C0961905262920E 27
20.8	.451722354958226535600E-28	.772546135 3C9276695631E 27
20.9	.285790363311651692469E-28	.121816703817634209032E 28
21.0	.180613844247038337893E-28	.192294462486380835187E 28
21.1	.114020531328895726437E-28	.303880666246597862256E 28
21.2	.719026264127932 4C0624E-29	.48C7447735C5085303756E 28
21.3	.452936387392152928570E-29	.761377682662431374138E 28
21.4	.285010932756958239155E-29	.120714281033582144510E 29
21.5	.179150801726944136460E-29	.191597119702290587854E 29
21.6	.112488983696C01255802E-29	.304431788340486276995E 29
21.7	.705563806817C01060242E-30	.484239946720569689753E 29
21.8	.442077811985405880216E-30	.771C80497483682132865E 29
21.9	.276693089647270999206E-30	.122915378863818866915E 30
22.0	.172996024035369828514E-30	.196145814091862286920E 30

Table B-1 (Cont'd)

THE AIRY FUNCTIONS ($10 < x < 25$)

x	Ai(x)	Bi(x)
22.0	.17299602403536982851[illegible]E-30	.19614581409186228692[illegible]E 30
22.1	.10804711642474469036[illegible]E-30	.31334C911729386914638E 30
22.2	.67410961052849720394[illegible]E-31	.5C1C941468119295771[illegible]3E 30
22.3	.42013528533642463[illegible]5698E-31	.80220370531853044337[illegible]E 30
22.4	.26157131312813773276[illegible]E-31	.12856183106117536995[illegible]E 31
22.5	.16268008568516215603[illegible]E-31	.20625307844122523084[illegible]E 31
22.6	.10107017069554396017[illegible]E-31	.33124458954810973594[illegible]E 31
22.7	.62727348205772493470[illegible]E-32	.53254471362464699362[illegible]E 31
22.8	.38889930625606735672[illegible]E-32	.85707981410548220626[illegible]E 31
22.9	.24086001140253398630[illegible]E-32	.13808399397799464617[illegible]E 32
23.0	.14901860957672060912[illegible]E-32	.22270054796228608941[illegible]E 32
23.1	.92101255694955599597[illegible]E-33	.35954569827C528538825E 32
23.2	.56864444347153485328[illegible]E-33	.58108648573206848848[illegible]E 32
23.3	.35072539246137599091[illegible]E-33	.94011358261754828752[illegible]E 32
23.4	.21609550500447974569[illegible]E-33	.15225507139836513366[illegible]E 33
23.5	.13300786289394327606[illegible]E-33	.24683919148919015836[illegible]E 33
23.6	.81782973411361752393[illegible]E-34	.40059580569517966270[illegible]E 33
23.7	.50234653323415700611[illegible]E-34	.65080015550553884972[illegible]E 33
23.8	.30824765615505017332[illegible]E-34	.10583684985681477538[illegible]E 34
23.9	.18895262910264662000[illegible]E-34	.17229521405783458231[illegible]E 34
24.0	.11570810853985424654[illegible]E-34	.28077318368107314534[illegible]E 34
24.1	.70783699869893122141[illegible]E-35	.45801868968272610901[illegible]E 34
24.2	.43257582028565731027[illegible]E-35	.74791974360481371845[illegible]E 34
24.3	.26408980480389827980[illegible]E-35	.12225597389543507790[illegible]4E 35
24.4	.16106543154802954600[illegible]E-35	.20004493062891869641[illegible]5E 35
24.5	.98133037974629947798[illegible]E-36	.32766228915633382400[illegible]E 35
24.6	.59729823155596202928[illegible]E-36	.53723699162809420704[illegible]8E 35
24.7	.36316773524105446214[illegible]E-36	.88174901889136705362[illegible]9E 35
24.8	.22061546467231486275[illegible]E-36	.14486477253904681188[illegible]3E 36
24.9	.13387714163375527874[illegible]E-36	.23824205322378942031[illegible]9E 36
25.0	.81160268246913866848[illegible]7E-37	.39220307780413817732[illegible]5E 36

Table B-2

DERIVATIVE OF THE AIRY FUNCTION ($10 < x < 25$)

x	Ai′(x)	Bi′(x)
10.0	-.35206336767389236353 3E -9	.14292361344828657769 2E 10
10.1	-.25703607274884638753 9E -9	.19674071623825106230 3E 10
10.2	-.18735878902127557984 0E -9	.27124163612814786622 8E 10
10.3	-.13635289499567024049 2E -9	.37453049707728368328 6E 10
10.4	-.99075995195010803676 1E-10	.51794502061654251681 5E 10
10.5	-.71876967814515670894 4E-10	.71736922452832991822 0E 10
10.6	-.52063296282141648576 5E-10	.99508785193257181177 7E 10
10.7	-.37652820868652853445 6E-10	.13824096312236110430 2E 11
10.8	-.27188835349391645979 6E-10	.19233831121236351300 3E 11
10.9	-.19602621700717091448 1E-10	.26800671934077690764 5E 11
11.0	-.14111441246628517334 3E-10	.37400168196926977018 8E 11
11.1	-.10142975815598381671 5E-10	.52269290479748818908 7E 11
11.2	-.72794625245019053037 6E-11	.73158038850450559872 3E 11
11.3	-.52164638717962404102 8E-11	.10254563135155427989 9E 12
11.4	-.37324936170788681002 4E-11	.14394918898724550154 5E 12
11.5	-.26666799675045314054 2E-11	.20236507276638385748 8E 12
11.6	-.19023683505348665444 7E-11	.28490070674762372989 6E 12
11.7	-.13551055027635776603 0E-11	.40168028783807018232 3E 12
11.8	-.96384954042280151215 3E-12	.56714480425065589871 0E 12
11.9	-.68455104418886800275 7E-12	.80192045829278418796 9E 12
12.0	-.48547365549853084610 5E-12	.11355075024433707428 7E 13
12.1	-.34378860722501380411 1E-12	.16101547369077681490 4E 13
12.2	-.24310057456061155369 0E-12	.22864497015690094467 8E 13
12.3	-.17165318121916935085 0E-12	.32513946831844889012 3E 13
12.4	-.12102963821607193514 3E-12	.46300888905784713726 6E 13
12.5	-.85213465646738564426 8E-13	.66026486813642953926 7E 13
12.6	-.59910633493315866042 4E-13	.94287457506803277C823E 13
12.7	-.42061167984049883696 0E-13	.13483238385828256565 3E 14
12.8	-.29487841939274785788 1E-13	.19307976320097349248 1E 14
12.9	-.20643882490681672869 0E-13	.27687215946102551585 7E 14
13.0	-.14432080573972626042 0E-13	.39757544969908345410 8E 14
13.1	-.10075302908577197242 1E-13	.57168309370941144714 2E 14
13.2	-.70239477570717307676 8E-14	.82316038540744184664 8E 14
13.3	-.48899091759095196266 1E-14	.11868742945786722086 4E 15
13.4	-.33995310287076281793 3E-14	.17136178824921227053 7E 15
13.5	-.23601425439243112847 3E-14	.24774797864941903868 2E 15
13.6	-.16362917805603644216 8E-14	.35866667121725397162 0E 15
13.7	-.11328927909353160006 8E-14	.51994165256713809640 0E 15
13.8	-.78329374580342010496 9E-15	.75474253014743001188 3E 15
13.9	-.54084210346759040973 1E-15	.10970378769209520759 5E 16
14.0	-.37293101100179006789 8E-15	.15966914115880027889 2E 16

Table B-2 (Cont'd)

DERIVATIVE OF THE AIRY FUNCTION ($10 < x < 25$)

x	Ai'(x)	Bi'(x)
14.0	-.37293101100179006789 8E-15	.15969914115880027889 2E 16
14.1	-.25680343780402770184 6E-15	.23269921922250179105 9E 16
14.2	-.17659956607020601461 5E-15	.33957949308473008247 7E 16
14.3	-.12128218326719278872 4E-15	.49620212606471855345 9E 16
14.4	-.83181173764816682769 6E-16	.72601314436393178615 1E 16
14.5	-.56973882061857806098 6E-16	.10636460360636523139 1E 17
14.6	-.38971879748295069941 6E-16	.15603234961526904369 4E 17
14.7	-.26622797656396853281 0E-16	.22918973914706352265 6E 17
14.8	-.18162884692409363646 7E-16	.33708298376494263964 1E 17
14.9	-.12375043585074923167 7E-16	.49640686840978901002 7E 17
15.0	-.84205679540177727648 9E-17	.73197492034070104973 2E 17
15.1	-.57223011302340032621 0E-17	.10807127578903909611 7E 18
15.2	-.38836190865703871960 2E-17	.15976374133915111633 9E 18
15.3	-.26323342842520720371 7E-17	.23648209178981062418 7E 18
15.4	-.17819102628187293878 3E-17	.35048434853487195862 1E 18
15.5	-.12046832044534437417 6E-17	.52010088403587950189 9E 18
15.6	-.81339961504917980691 1E-18	.77277563971730403868 4E 18
15.7	-.54850548550771709320 5E-18	.11496467437264554336 4E 19
15.8	-.36940749326960159550 5E-18	.17124538227039954471 0E 19
15.9	-.24847341459447657902 6E-18	.25539660082021048749 3E 19
16.0	-.16691886768381809551 7E-18	.38137435071218626576 0E 19
16.1	-.11199121614129490799 5E-18	.57019883715307105891 2E 19
16.2	-.75044221881281201842 6E-19	.85356778954178150229 7E 19
16.3	-.50223508206133003653 2E-19	.12793369266079108086 3E 20
16.4	-.33570284877685947070 5E-19	.19198419440174920468 9E 20
16.5	-.22411085425252972568 8E-19	.28845498537888041726 8E 20
16.6	-.14942811301127750977 2E-19	.43393140292806888580 9E 20
16.7	-.99509592582740908946 0E-20	.65357135190807018898 1E 20
16.8	-.66185437122113038319 4E-20	.98558090774208264708 3E 20
16.9	-.43966958135242079810 2E-20	.14880494683320821645 9E 21
17.0	-.29171482192933137922 0E-20	.22494002910657269280 5E 21
17.1	-.19331264721812734168 5E-20	.34043868722439385687 9E 21
17.2	-.12794794641938736880 2E-20	.51586059363581868788 7E 21
17.3	-.84582247463981999719 7E-21	.78261037862651348076 2E 21
17.4	-.55846958593876467679 7E-21	.11887134719749202212 7E 22
17.5	-.36829496287900966898 3E-21	.18076970117807844095 8E 22
17.6	-.24258810762983384801 4E-21	.27522604090025685368 7E 22
17.7	-.15959608155819316173 7E-21	.41953422780366137094 3E 22
17.8	-.10487100587489765219 1E-21	.64026217368755104264 1E 22
17.9	-.68828867455032806477 4E-22	.97827160987644170067 0E 22
18.0	-.45120018606819418867 8E-22	.14964796503287850692 3E 23

Table B-2 (Cont'd)

DERIVATIVE OF THE AIRY FUNCTION ($10 < x < 25$)

x	Ai'(x)	Bi'(x)
18.0	-.45120018606819418867 8E-22	.14964796503287850692 3E 23
18.1	-.29542880426488718901 4E-22	.22918730225199886076 9E 23
18.2	-.19320696319236526215 7E-22	.35141256374270517818 E 23
18.3	-.12620615097354644929 7E-22	.53944807163278436547 5E 23
18.4	-.82343155520978735785 8E-23	.82906060550209160860 8E 23
18.5	-.53661788234147277077 8E-23	.12756332855677990941 3E 24
18.6	-.34929697986106265165 5E-23	.19650201233732411917 0E 24
18.7	-.22710043486971815384 7E-23	.30304591486518825144 1E 24
18.8	-.14748095012585425797 6E-23	.46789545347539204189 8E 24
18.9	-.95664340594635767627 6E-24	.72324738675452147815 3E 24
19.0	-.61981458271300150556 5E-24	.11192350063395887816 0E 25
19.1	-.40111822638376606413 6E-24	.17340067290834801324 5E 25
19.2	-.25928848928775746575 0E-24	.26895156781359414706 9E 25
19.3	-.16741548998076847580 0E-24	.41762830666520199183 1E 25
19.4	-.10797194222506161638 7E-24	.64922773685233428732 6E 25
19.5	-.69555322364636242547 5E-25	.10104021269463098594 7E 26
19.6	-.44756415255379278952 5E-25	.15742737363291075199 5E 26
19.7	-.28766496909027958593 3E-25	.24555785744008457301 8E 26
19.8	-.18468290283814512567 4E-25	.38345445773147984859 4E 26
19.9	-.11843381000807503940 5E-25	.59945820527435787035 3E 26
20.0	-.75863916257483549553 2E-26	.93818393361339643555 9E 26
20.1	-.48540764173800191552 9E-26	.14699408113831285753 1E 27
20.2	-.31023514034732898071 6E-26	.23056495514802672687 2E 27
20.3	-.19805673852988694918 4E-26	.36204884531652782816 6E 27
20.4	-.12630010992037822208 4E-26	.56914155007993862826 6E 27
20.5	-.80451567937554895820 3E-27	.89567719810491614958 4E 27
20.6	-.51189769462191681922 9E-27	.14111061891774011578 8E 28
20.7	-.32535006762689870719 E-27	.22255824521902328682 6E 28
20.8	-.20655650487392371723 4E-27	.35140048943366627062 4E 28
20.9	-.13099305495395139946 1E-27	.55543672840977525635 4E 28
21.0	-.82981302583004457425 2E-28	.87889960528431639204 0E 28
21.1	-.52509267836882000786 8E-28	.13922437598712822970 5E 29
21.2	-.33190711479645123276 4E-28	.22078091354393049578 5E 29
21.3	-.20956713886167784386 0E-28	.35049103594683008365 5E 29
21.4	-.13217732523843468987 2E-28	.55700662706918190776 0E 29
21.5	-.83275837516556894649 1E-29	.88615724051337808188 9E 29
21.6	-.52409552103970105749 5E-29	.14113253070316347408 8E 30
21.7	-.32948252273316654480 8E-29	.22501336575970430928 1E 30
21.8	-.20691211055876686022 2E-29	.35913133103359480064 4E 30
21.9	-.12979923210521480227 6E-29	.57380062683074017254 0E 30
22.0	-.81337740384475402708 0E-30	.91776278690818259747 9E 30

Table B-2 (Cont'd)

DERIVATIVE OF THE AIRY FUNCTION ($10 < x < 25$)

x	Ai'(x)	Bi'(x)
22.0	-.8133774038447540270 80E-30	.9177627869081825974 79E 30
22.1	-.5091513814570804396 50E-30	.1469469370062620749 43E 31
22.2	-.3183740491628665495 73E-30	.2355321911624826582 38E 31
22.3	-.1988682112743385163 67E-30	.3779189127195225467 37E 31
22.4	-.1240883756640589611 03E-30	.6070222166076143401 86E 31
22.5	-.7734565180555525404 49E-31	.9760389394046413871 62E 31
22.6	-.4815937242698697765 36E-31	.1571032972105910461 03E 32
22.7	-.2995483126331075991 25E-31	.2531384162693390558 19E 32
22.8	-.1861208832320103983 08E-31	.4083047504270874706 43E 32
22.9	-.1155224841207865908 26E-31	.6592701928500502945 97E 32
23.0	-.7162788572866303807 74E-32	.1065599743705194806 68E 33
23.1	-.4436525158286744929 55E-32	.1724151656700924838 86E 33
23.2	-.2745048151903463985 42E-32	.2792585940677542004 15E 33
23.3	-.1696696558050527717 03E-32	.4527791303873281900 77E 33
23.4	-.1047626726464527566 87E-32	.7348760176376941726 17E 33
23.5	-.6461868523316313292 18E-33	.1193956827424054106 72E 34
23.6	-.3981619871879011887 89E-33	.1941820619926214464 61E 34
23.7	-.2450826508873351273 94E-33	.3161364682904613830 36E 34
23.8	-.1507014388807029044 04E-33	.5152098561413822265 06E 34
23.9	-.9257106035293633551 80E-34	.8404983849513808698 74E 34
24.0	-.5680506160122678291 81E-34	.1372561596919796144 15E 35
24.1	-.3482200157885005862 08E-34	.2243717284105895650 28E 35
24.2	-.2132437254729110482 47E-34	.3671511126023557142 87E 35
24.3	-.1304534461099634758 61E-34	.6013967683448903568 98E 35
24.4	-.7972463494423577379 78E-35	.9860886596766367794 42E 35
24.5	-.4867300156198381658 84E-35	.1618484644343298733 00E 36
24.6	-.2968542273959660809 57E-35	.2659120782786928049 20E 36
24.7	-.1808667657561200447 06E-35	.4373242197258069961 13E 36
24.8	-.1100868979879259541 1E-35	.7199529373614427891 72E 36
24.9	-.6693830621864682549 57E-36	.1186421289825409424 90E 37
25.0	-.4066089337243281003 32E-36	.1957073508323330897 99E 37

Table B-3

THE AIRY FUNCTIONS ($0.001 < \frac{1}{x} < 0.100$)

(1/x)	Ai(x)	Bi(x)
0.001	.9306933063179554850 86E-9157	.5407711839194947217 73E 9155
0.002	.5441298591207528694 06E-3238	.1308074777327973248 89E 3237
0.003	.6284959438456742745 04E-1763	.1387305814297621569 05E 1762
0.004	.2423540281225583456 45E-1145	.4153362995256493047 17E 1144
0.005	.9153624308452684222 05E-820	.1229453361044710141 37E 819
0.006	.8438860408561284345 02E-624	.1460871338124800040 21E 623
0.007	.3536757930056225126 02E-495	.3764990103370007880 20E 494
0.008	.1977615996743770057 72E-405	.7198188136930865714 32E 404
0.009	.6882515587854900686 74E-340	.2193785856124747154 65E 339
0.010	.2634482152088184467 72E-290	.6041223996670201449 04E 289
0.011	.1002234034702566998 00E-251	.1665510663976726612 17E 251
0.012	.5217473910100956477 71E-221	.3341570173945931063 80E 220
0.013	.4412989072529058245 63E-196	.4112056798869610349 69E 195
0.014	.1596335043580413801 02E-175	.1179669329062711270 16E 175
0.015	.2479435686287269118 40E-158	.7061639815050477762 09E 157
0.016	.8770565519900959950 18E-144	.2295370541452945730 65E 143
0.017	.2426064265282311144 48E-131	.8553475677940503446 34E 130
0.018	.1329645206807488123 08E-120	.1605909279513916260 29E 120
0.019	.2944228246595371060 49E-111	.7451194887644830998 84E 110
0.020	.4584941724074823469 67E-103	.4909099699444219338 08E 102
0.021	.7773230709276491532 51E-96	.2967077223268002783 56E 95
0.022	.2033259830638090952 09E-89	.1161018964941738873 93E 89
0.023	.1087012518145611918 39E-83	.2220497711046715504 71E 83
0.024	.1493487650490034604 52E-78	.1650916297457943451 71E 78
0.025	.6365742658552914899 75E-74	.3953139302438593539 63E 73
0.026	.9838359243493339752 96E-70	.2608466849877495585 34E 69
0.027	.6280372368090100734 94E-66	.4164069364529677962 62E 65
0.028	.1847539096024221540 57E-62	.1441474618501625568 49E 62
0.029	.2747831744710070162 60E-59	.9803490923301602813 65E 58
0.030	.2235801350309873066 77E-56	.1232961106967074390 05E 56
0.031	.1064754960520526313 9E-53	.2631802700913084972 14E 53
0.032	.3145397562248238252 37E-51	.9051525907126628085 89E 50
0.033	.6060836350401680558 54E-49	.4770314174985777396 06E 48
0.034	.7957492415467714084 35E-47	.3687958490888511027 85E 46
0.035	.7395081747456895603 84E-45	.4026373736101710147 64E 44
0.036	.5029409729121682661 67E-43	.6004232915227383955 40E 42
0.037	.2577705578359161779 55E-41	.1187657238831848101 74E 41
0.038	.1021713248959822927 46E-39	.3036593297631266514 87E 39
0.039	.3204448922427185330 67E-38	.9808511549393338213 67E 37
0.040	.8116026824691386684 87E-37	.3922030778041381773 25E 36

Table B-3 (Cont'd)

THE AIRY FUNCTIONS ($0.001 < \frac{1}{x} < 0.100$)

(1/x)	Ai(x)	Bi(x)
0.040	.81160268246913866848 7E-37	.39220307780413817 7325E 36
0.041	.16903349773031311782 2E-35	.19365313887560780408 6E 35
0.042	.29422278871824573951 9E-34	.11085964872672168137 3E 34
0.043	.43426730848820675o539E-33	.75998044762676613550 1E 32
0.044	.55065338056794053424 7E-32	.60628097771062644016 7E 31
0.045	.60693014697048055390 7E-31	.55628053563620872546 4E 30
0.046	.58768146629960775894 0E-30	.58084955401975883466 4E 29
0.047	.50472325974961185729 2E-29	.68363263884932644709 6E 28
0.048	.38783722366488311615 7E-28	.89908089371453720254 5E 27
0.049	.26875856832382015678 3E-27	.13108827226411639301 8E 27
0.050	.16916728686705403132 9E-26	.21037650496511038148 1E 26
0.051	.97355835151233548608 4E-26	.36919196345183798426 0E 25
0.052	.51535206386756710286 2E-25	.70425076922147543992 5E 24
0.053	.25230584347718423032 2E-24	.14522482617089541989 5E 24
0.054	.11482036418815261854 8E-23	.32211339316815612052 0E 23
0.055	.48796406386298457848 3E-23	.76493570577340381074 9E 22
0.056	.19448333110611106986 8E-22	.19366167558603117271 4E 22
0.057	.72980298291130348656 0E-22	.52067233223832984205 9E 21
0.058	.25877906568393746141 8E-21	.14812154625905582334 7E 21
0.059	.86997171990867775212 5E-21	.44438045472904910959 4E 20
0.060	.27814894654297473448 5E-20	.14016287788856544837 46E 20
0.061	.84818599661438540164 2E-20	.46345691460254334097 0E 19
0.062	.24734395364384114515 1E-19	.16022521542827647023 7E 19
0.063	.69148218005379122693 8E-19	.57773196451880410891 7E 18
0.064	.18574966030135299225 6E-18	.21677035840091138294 3E 18
0.065	.48047573526233513837 1E-18	.84454735029663015266 7E 17
0.066	.11991603475696082113 0E-17	.34098428888581227896 0E 17
0.067	.28930335628570772408 5E-17	.14240475535220287703 6E 17
0.068	.67585710817487390553 9E-17	.61410277930146948585 3E 16
0.069	.15313951191623517047 3E-16	.27301074275482236700 1E 16
0.070	.33706305827494480015 7E-16	.12493418799590003928 8E 16
0.071	.72168128513567927248 8E-16	.58766277416168686543 2E 15
0.072	.15051176531107936115 2E-15	.28375354236337750289 3E 15
0.073	.30614760786123307141 2E-15	.14046791169091478753 7E 15
0.074	.60804875225738850685 6E-15	.71207399783639927543 2E 14
0.075	.11805221820522025617 9E-14	.36923705813759235077 8E 14
0.076	.22428015533457376038 9E-14	.19564372102205437131 0E 14
0.077	.41736387529548471224 4E-14	.10582338951981994827 8E 14
0.078	.76146164490234622996 4E-14	.58378328630072983847 5E 13
0.079	.13632309130186781129 6E-13	.32816916123665511469 5E 13
0.080	.23968278260780499368 1E-13	.18782919356220518669 7E 13

Table B-3 (Cont'd)

THE AIRY FUNCTIONS ($0.001 < \frac{1}{x} < 0.100$)

(1/x)	Ai(x)	Bi(x)
0.080	.23968278260780499368 1E-13	.18782919356220518669 7E 13
0.081	.41417918024876274100 8E-13	.10937309650421835577 2E 13
0.082	.70395227259232082775 2E-13	.64747250510867894711 6E 12
0.083	.11776166555487056858 5E-12	.38939835978437933971 4E 12
0.084	.19402448908498408544 8E-12	.23776259436736067617 8E 12
0.085	.31504461367894286457 4E-12	.14729883102432882239 0E 12
0.086	.50443748745398340669 9E-12	.92534836703140523854 8E 11
0.087	.79690412384C931864865E-12	.58913994988826675330 0E 11
0.088	.12427948456930252489 6E-11	.37993419262490346627 7E 11
0.089	.19142869222579940934 0E-11	.24805961241291447720 4E 11
0.090	.29136505932362936630 0E-11	.16389039953441143821 5E 11
0.091	.43841873097949159047 6E-11	.10352241225205918751 7E 11
0.092	.65245505518868816368 7E-11	.73997395182226368551 9E 10
0.093	.96073042387386831147 0E-11	.50525986417070864419 1E 10
0.094	.14002737167751875901 9E-10	.34851995590906447288 0E 10
0.095	.20209134365870218122 1E-10	.24276863753850022257 9E 10
0.096	.28890907566741250278 4E-10	.17070835987551715221 2E 10
0.097	.40926206619449957965 6E-10	.12113416240466738262 2E 10
0.098	.57465870368458024102 1E-10	.86713609269089501292 7E 9
0.099	.80005827769078545057 9E-10	.62601130664973833851 2E 9
0.100	.11047532552898685935 4E -9	.45564115354822514083 5E 9

+55

Table B-4

DERIVATIVE OF THE AIRY FUNCTIONS ($0.001 < \frac{1}{x} < 0.100$)

(1/x)	Ai'(x)	Bi'(x)
0.001	-.29431339179103355694 7E-9155	.17100551146246152919 7E 9157
0.002	-.12167385585536581794 5E-3236	.29248787143627118140 3E 3238
0.003	-.11475184848386729516 5E-1761	.25322105368168540728 2E 1763
0.004	-.38321959606420465251 8E-1144	.65666281052399218463 6E 1145
0.005	-.12946323592218823028 0E-818	.17385559018472874985 3E 820
0.006	-.10895787402736215856 4E-622	.18857575934395081313 7E 624
0.007	-.42278529685766730443 7E-494	.44993647707552895427 1E 495
0.008	-.22114372475605720519 0E-404	.80463787123599175195 9E 405
0.009	-.72563561704402489567 2E-339	.23119594691136902435 3E 340
0.010	-.26351403616044099047 9E-289	.60397127453106029754 1E 290
0.011	-.95586807372670387432 1E-251	.15875437638309473166 0E 252
0.012	-.47644442149850358152 3E-220	.30494189732992795069 0E 221
0.013	-.38718803509164449067 9E-195	.36051747320248292017 5E 196
0.014	-.13497074943035330847 5E-174	.99658923983385266003 4E 175
0.015	-.20253794842141834410 9E-157	.64160505201550562926 2E 158
0.016	-.69372446546794929587 2E-143	.18137304331266745753 9E 144
0.017	-.18617360058201443040 3E-130	.65565760979540122697 7E 131
0.018	-.99165646328667080367 8E-120	.11962503510052386884 3E 121
0.019	-.21373638960075771925 6E-110	.54021197921061259929 0E 111
0.020	-.32443318198287992837 6E-102	.34687987795459767376 1E 103
0.021	-.53681120166702596253 6E-95	.20459171117597914328 5E 96
0.022	-.13719395214478207753 1E-88	.78211895731242093570 0E 89
0.023	-.71737850545463266563 6E-83	.14628732475266392360 8E 84
0.024	-.96493615479342041186 6E-78	.10646690231160086472 5E 79
0.025	-.40300179776006780325 2E-73	.24977079681706968810 3E 74
0.026	-.61078696694558781201 8E-69	.16160024927678842096 7E 70
0.027	-.38263404801476112807 9E-65	.25313533203081795033 5E 66
0.028	-.11054053087986536782 6E-61	.86043384384589757329 5E 62
0.029	-.16155684343639989772 3E-58	.57848680366469773889 7E 59
0.030	-.12925119586807573348 1E-55	.71092267936702106650 1E 56
0.031	-.60556257328606774990 3E-53	.14927168267098359276 3E 54
0.032	-.17608380830590861878 9E-50	.50526894539699806496 3E 51
0.033	-.33413622259452125015 1E-48	.26220211385739486499 1E 49
0.034	-.43222937932856851146 9E-46	.19969291447175601426 2E 47
0.035	-.39592819355925760304 9E-44	.21486496514458973563 4E 45
0.036	-.26552390115662440373 1E-42	.31590814833278409975 9E 43
0.037	-.13424592017799798180 4E-40	.61633007522092069419 3E 41
0.038	-.52509402068000385733 3E-39	.15546412234372643595 5E 40
0.039	-.16257452556622606732 1E-37	.49571230144035192483 3E 38
0.040	-.40660893372432810033 2E-36	.19570735083233308979 9E 37

Table B-4 (Cont'd)

DERIVATIVE OF THE AIRY FUNCTIONS ($0.001 < \frac{1}{x} < 0.100$)

(1/x)	Ai'(x)	Bi'(x)
0.040	-.4066089337243281003 32E-36	.1957073508323330897 99E 37
0.041	-.8365206560309995014 90E-35	.9396042767250602864 15E 35
0.042	-.1438733135021021274 59E-33	.5397693255363286106 77E 34
0.043	-.2098866588957736665 11E-32	.3656735217380572252 31E 33
0.044	-.2631159710408983165 21E-31	.2883623048787266608 61E 32
0.045	-.2867883948778790662 03E-30	.2616035553010649144 60E 31
0.046	-.2746796231993884572 99E-29	.2701503427524654061 74E 30
0.047	-.2334008087615885657 66E-28	.3145277581605668595 37E 29
0.048	-.1774850562058588653 01E-27	.4092863024576799410 42E 28
0.049	-.1217397780193827761 26E-26	.5905795506722959320 92E 27
0.050	-.7586391625748354955 32E-26	.9381839336133964355 59E 26
0.051	-.4323313860659652773 38E-25	.1630067809019576711 20E 26
0.052	-.2266619087103967958 43E-24	.3079119571781972263 10E 25
0.053	-.1099264563377160997 08E-23	.6288771550622668418 45E 24
0.054	-.4956463163569071515 08E-23	.1381772044815899728 39E 24
0.055	-.2087342192226579383 47E-22	.3251092460124100640 63E 23
0.056	-.8245427242095785359 02E-22	.8156357653836406057 97E 22
0.057	-.3067121307471676036 16E-21	.2173372395781950574 47E 22
0.058	-.1078241787798392987 45E-20	.6128743254976836764 82E 21
0.059	-.3594336430150746624 21E-20	.1822870753459433742 14E 21
0.060	-.1139673046498057506 79E-19	.5700903566542330819 80E 20
0.061	-.3447020419108459900 34E-19	.1869346369964630892 85E 20
0.062	-.9971548896915655668 07E-19	.6409719212133567813 91E 19
0.063	-.2765716622104355674 80E-18	.2292547819994330715 39E 19
0.064	-.7371826259857567280 25E-18	.8533557686445932783 77E 18
0.065	-.1892309684751347400 29E-17	.3298717532062540533 15E 18
0.066	-.4687306922920829907 27E-17	.1321592101630269224 05E 18
0.067	-.1122470995415373747 05E-16	.5477457321138906849 67E 17
0.068	-.2603159188872139165 95E-16	.2344417998748619484 75E 17
0.069	-.5856046715798151367 25E-16	.1034569836697351435 42E 17
0.070	-.1279810758341796189 74E-15	.4699944547845370242 08E 16
0.071	-.2721084100146867839 22E-15	.2194901078140800635 22E 16
0.072	-.5636016379300547701 83E-15	.1052315907743040327 21E 16
0.073	-.1138623402575360084 67E-14	.5172989451983895947 40E 15
0.074	-.2246343130218858290 49E-14	.2604291716267379634 28E 15
0.075	-.4332516530842191009 80E-14	.1341248159271972172 45E 15
0.076	-.8177568260717941158 13E-14	.7059068494211716404 21E 14
0.077	-.1512005247092788000 52E-13	.3792958023801241722 24E 14
0.078	-.2741122682521209525 32E-13	.2078735369901919640 63E 14
0.079	-.4876719943955520918 56E-13	.1160999030413177261 55E 14
0.080	-.8521346564673856442 68E-13	.6602648681364295392 67E 13

Table B-4 (Cont'd)

DERIVATIVE OF THE AIRY FUNCTIONS ($0.001 < \frac{1}{x} < 0.100$)

(1/x)	Ai'(x)	Bi'(x)
0.080	-.852134656467385644268E-13	.660264868136429539267E 13
0.081	-.146354755583667727203E-12	.382050005569360244412E 13
0.082	-.247253428802173484712E-12	.224759682439674223376E 13
0.083	-.411164986569351596034E-12	.134341607744334080985E 13
0.084	-.673462750454590197180E-12	.815286957702759259337E 12
0.085	-.100718876731495257755E-11	.502050328074181791250E 12
0.086	-.173079812775502557116E-11	.313519054390867627011E 12
0.087	-.271882280270428414382E-11	.198434376832673242852E 12
0.088	-.421637245777368617479E-11	.127225728507594103672E 12
0.089	-.645861697402928647589E-11	.825882901959071354759E 11
0.090	-.977666173457181014304E-11	.542548879304808233160E 11
0.091	-.146315347787940395985E-10	.360526995364261165995E 11
0.092	-.216583566590634897483E-10	.242229234423285103867E 11
0.093	-.317231603948524590943E-10	.164484734880062291083E 11
0.094	-.459952704428099147495E-10	.112840216897124317157E 11
0.095	-.660386350814962365078E-10	.781769217261657434957E 10
0.096	-.939260337825592019538E-10	.546782042248172793551E 10
0.097	-.132380646401629340827E -9	.385942680805962674046E 10
0.098	-.184950069381856421738E -9	.274829224136809255301E 10
0.099	-.256218489240527352727E -9	.197378390237140312213E 10
0.100	-.352063367673892363533E -9	.142923613448286577692E 10

Table B-5

THE AUXILIARY FUNCTIONS Q(x) AND M(x)

1/x	Q(x)	M(x)
0.001	.1000000000156250000018E 1	.1000000000156250000056E 1
0.002	.1000000001250000001151E 1	.1000000001250000003609E 1
0.003	.1000000004218750013111E 1	.1000000004218750041113E 1
0.004	.1000000010000000073666E 1	.1000000010000000231000E 1
0.005	.1000000019531252810l6E 1	.1000000019531258811l96E 1
0.006	.1000000033750008391l0E 1	.1000000033750026312l0E 1
0.007	.1000000053593771159l5E 1	.1000000053593816350l5E 1
0.008	.1000000080000047146l3E 1	.1000000080000147840l7E 1
0.009	.1000000113906345580l9E 1	.1000000113906549716l5E 1
0.010	.1000000156250179851l2E 1	.1000000156250563971l3E 1
0.011	.1000000207969068618l8E 1	.1000000207969749113[illegible] 1
0.012	.1000000270000537036l2E 1	.1000000270001684023l9E 1
0.013	.1000000343282118116l5E 1	.1000000343283972219l2E 1
0.014	.1000000428751354215l2E 1	.1000000428754246527l2E 1
0.015	.1000000527345798655l5E 1	.1000000527350174161l1E 1
0.016	.1000000640003017473l1E 1	.1000000640009462206l3E 1
0.017	.1000000767660591297l5E 1	.1000000767669863510l2E 1
0.018	.1000000911256117358l5E 1	.1000000911269182986l1E 1
0.019	.1000001071727211621l1E 1	.1000001071745284317l8E 1
0.020	.1000001250011511063l2E 1	.1000001250036097074l6E 1
0.021	.1000001447046676063l7E 1	.1000001447079624234l1E 1
0.022	.1000001663770302931l9E 1	.1000001663813950118l8E 1
0.023	.1000001901120376569l7E 1	.1000001901177248732l3E 1
0.024	.1000002160034473259l3E 1	.1000002160107792514l8E 1
0.025	.1000002441450163577l3E 1	.1000002441543961501l7E 1
0.026	.1000002746305565446l9E 1	.1000002746424252897l1E 1
0.027	.1000003075538437316l1E 1	.1000003075687291060l8E 1
0.028	.1000003430086681470l8E 1	.1000003430271837895l8E 1
0.029	.1000003810888247471l7E 1	.1000003811116803668l3E 1
0.030	.1000004218881135727C0E 1	.1000004219161258225l3E 1
0.031	.1000004655003401199l9E 1	.1000004655344442629l4E 1
0.032	.1000005120193157238l9E 1	.1000005120605781217l0E 1
0.033	.1000005615388579542l5E 1	.1000005615884894069l4E 1
0.034	.1000006141527910263l7E 1	.1000006142121609894l6E 1
0.035	.1000006699549462233l7E 1	.1000006700255979340l0E 1
0.036	.1000007290391623326C1E 1	.1000007291228288719l4E 1
0.037	.1000007914992850953l1E 1	.1000007915979074159l0E 1
0.038	.1000008574291726692l6E 1	.1000008575449136172l4E 1
0.039	.1000009269226861045l1E 1	.1000009270579554654l1E 1
0.040	.1000010000736990833l7E 1	.1000010002311704309l6E 1

Table B-5 (Cont'd)

THE AUXILIARY FUNCTIONS Q(x) AND M(x)

1/x	Q(x)		M(x)	
0.041	.1C0001076976C971726761	1	.100001077158727049547E	1
0.042	.1C0001157723771840300E	1	.100001157934826551341E	1
0.043	.1C0001242410628484637E	1	.100001242653704530862E	1
0.044	.1C0001331130583227746E	1	.100001331409632662665E	1
0.045	.1C0001423977564221988E	1	.100001424296920459017E	1
0.046	.1C0001521045512220230E	1	.100001521409917071965E	1
0.047	.1C0001622428381159664E	1	.100001622843013139332E	1
0.048	.1000017282201387592 74E	1	.100001728690642674998E	1
0.049	.1C0001838514767131C25E	1	.100001839047285003697E	1
0.050	.1000019534062634048 05E	1	.100001954007466740617E	1
0.051	.1C0002072988640367171E	1	.100002073665763816092E	1
0.052	.1C0002197355927113984E	1	.100002198116803545646E	1
0.053	.1C0002326602169716955E	1	.100002327455266745739E	1
0.054	.1C0002460821431904192E	1	.100002461775889895509E	1
0.055	.1C0002600107795754790E	1	.100002601173467344850E	1
0.056	.1C0002744555362407559E	1	.100002745742853569192E	1
0.057	.1000028942582527839 24E	1	.100002895578965471325E	1
0.058	.100C0304931060832510CE	1	.100003050776784730674E	1
0.059	.1C0003209806591743 6C1E	1	.100003211431360200396E	1
0.060	.1000033758403877991 66E	1	.100003377637810352735E	1
0.061	.1C0003547506204029187E	1	.100003549491325773059E	1
0.062	.1C0003724898271643716E	1	.100003727087171703020E	1
0.063	.1C0003908110846235145E	1	.100003910520690633315E	1
0.064	.1C0004097238208652650E	1	.100004099887304946525E	1
0.065	.1C0004292374665831492E	1	.100004295282519610546E	1
0.066	.1C0C044936145516477 6E	1	.100004496801924923122E	1
0.067	.1C0C047010522277852 61E	1	.100004704541199308041E	1
0.068	.1C0004914782084624848E	1	.100004918596112163548E	1
0.069	.1000513489854213933 1E	1	.100005139062526763575E	1
0.070	.1C0005361496050811034E	1	.100005366036403212376E	1
0.071	.1C0005594669C92561968E	1	.100005599613801453231E	1
0.072	.1C0005834512181700096E	1	.100005839890884331853E	1
0.073	.1C0006081119865881372E	1	.100006086963920715183E	1
0.074	.1C0006334586727087645E	1	.100006340929288666294E	1
0.075	.1000065950073826205 98E	1	.100006601883478676125E	1
0.076	.1C0006862476486111837E	1	.100006869923096952806E	1
0.077	.1000071370887285492 92E	1	.100007145144868769375E	1
0.078	.1C0C074189388393200 64E	1	.100007427645641870690E	1
0.079	.1000077081215872698 89E	1	.100007717522389940402E	1
0.080	.1000080047317817793 64E	1	.100008014872216128844E	1

Table B-5 (Cont'd)

THE AUXILIARY FUNCTIONS Q(x) AND M(x)

1/x	Q(x)	M(x)
0.081	.1000083088642738571126 1	.1000083197923566427816 1
0.082	.1000086206139572500526 1	.1000086323801843979386 1
0.083	.1000089400757695709516 1	.1000089527332127352986 1
0.084	.1000092673446934434366 1	.1000092809490992021896 1
0.085	.1000096025157576646706 1	.1000096171256493992066 1
0.086	.1000099456840393858716 1	.1000099613608208940656 1
0.087	.1000102969446603108776 1	.1000103137527272035236 1
0.088	.1000106563927979129796 1	.1000106743996418445296 1
0.089	.1000110241236766702246 1	.1000110434000024558376 1
0.090	.1000114002325743194116 1	.1000114208524149913246 1
0.091	.1000117848148221290216 1	.1000118068556579863506 1
0.092	.1000121779658061912996 1	.1000122015086868984846 1
0.093	.1000125797809687337606 1	.1000126049106385240346 1
0.094	.1000129903558094503396 1	.1000130171608354918266 1
0.095	.1000134097858868524836 1	.1000134383587908357586 1
0.096	.1000138381668196404216 1	.1000138686042126476876 1
0.097	.1000142755942880949216 1	.1000143079970088123106 1
0.098	.1000147221640354898036 1	.1000147566372918257136 1
0.099	.1000151779718695255136 1	.1000152146253836993556 1
0.100	.1000156431136637840716 1	.1000156820618209513336 1

Table B-6

THE AUXILIARY FUNCTIONS P(x) AND Λ(x)

1/x	P(x)	Λ(x)
0.001	.99999999978124999763E +0	.99999999978124999335E +0
0.002	.99999999824999984762E +0	.99999999824999957345E +0
0.003	.99999999409374982641 2E +0	.99999999409374951411 9E +0
0.004	.99999998599999902466 8E +0	.99999998599999726999 9E +0
0.005	.99999997265624627939 8E +0	.99999997265623958586 5E +0
0.006	.99999995274998889033 0E +0	.99999995274996890352 2E +0
0.007	.99999992496872198553 8E +0	.99999992496867158619 1E +0
0.008	.99999988799993757844 4E +0	.99999988799982527902 7E +0
0.009	.99999984053112345348 8E +0	.99999984053089578968 8E +0
0.010	.99999978124976180501E +0	.99999978124933348884 1E +0
0.011	.99999970884332815500 9E +0	.99999970884256923056 9E +0
0.012	.99999962199928897420 3E +0	.99999962199800979256 6E +0
0.013	.99999951940510063150 4E +0	.99999951940303283600 6E +0
0.014	.99999939974820704703 2E +0	.99999939974498138493 5E +0
0.015	.99999926171603762347 3E +0	.99999926171115782500 9E +0
0.016	.99999910399600493079 5E +0	.99999910398881742138 1E +0
0.017	.99999892527550221927 5E +0	.99999892526516135550 2E +0
0.018	.99999872424190076075 8E +0	.99999872422732928052 0E +0
0.019	.99999849958254701809 7E +0	.99999849956239139501 5E +0
0.020	.99999824998475964270 7E +0	.99999824995734003466 3E +0
0.021	.99999797413582630011 8E +0	.99999797409°080781451E +0
0.022	.99999767072300032347 2E +0	.99999767067442308997 9E +0
0.023	.99999733843349719482 3E +0	.99999733837007043034 7E +0
0.024	.99999697595449085415 1E +0	.99999697587260994707 0E +0
0.025	.99999658197310983594 9E +0	.99999658186850163341 9E +0
0.026	.99999615517643323325 4E +0	.99999615504406702050 6E +0
0.027	.99999569425148648894 3E +0	.99999569408547738036 6E +0
0.028	.99999519788523701417 1E +0	.99999519767874144228 5E +0
0.029	.99999466476458963373 5E +0	.99999466450969262145 7E +0
0.030	.99999409357638185818 6E +0	.99999409326397575906 9E +0
0.031	.99999348300737898248 1E +0	.99999348262703337280 0E +0
0.032	.99999283174426901093 5E +0	.99999283128409141663 7E +0
0.033	.99999213847365740826 4E +0	.99999213792014454887 4E +0
0.034	.99999140188206167641 1E +0	.99999140121994090702 8E +0
0.035	.99999062065590575694 3E +0	.99999061986796638833 4E +0
0.036	.99998979348151425866 8E +0	.99998979254842843446 0E +0
0.037	.99998891904510651017 9E +0	.99998891794523931884 5E +0
0.038	.99998799603279043702 9E +0	.99998799474199893514 9E +0
0.039	.99998702313055626311 9E +0	.99998702162197708505 0E +0
0.040	.99998599902427003598 7E +0	.99998599726809526363 5E +0

Table B-6 (Cont'd)

THE AUXILIARY FUNCTIONS P(x) AND Λ(x)

1/x	P(x)	Λ(x)
0.041	.9999849223996669755 74E +0	.9999849203629079404 43E +0
0.042	.9999837919423446460 42E +0	.9999837895885833341 84E +0
0.043	.9999826063377559502 53E +0	.9999826036268836789 80E +0
0.044	.9999813642712019463 90E +0	.9999813611591449799 42E +0
0.045	.9999800644278244862 90E +0	.9999800608662562556 65E +0
0.046	.9999787054925986749 16E +0	.9999787014286382652 16E +0
0.047	.9999772861503251505 09E +0	.9999772815262217170 01E +0
0.048	.9999758050856221847 93E +0	.9999757998384249567 75E +0
0.049	.9999742609829176026 81E +0	.9999742550441311319 41E +0
0.050	.9999726575264405208 69E +0	.9999726458216648291 40E +0
0.051	.9999709784002129046 28E +0	.9999709708487681819 79E +0
0.052	.9999692372880409422 21E +0	.9999692288025764456 38E +0
0.053	.9999674278735062361 00E +0	.9999674183595930348 87E +0
0.054	.9999655488399568103 02E +0	.9999655381956640219 59E +0
0.055	.9999635988704979331 50E +0	.9999635869859520904 83E +0
0.056	.9999615766479827545 65E +0	.9999615634049099416 52E +0
0.057	.9999594808550027570 79E +0	.9999594661262531484 28E +0
0.058	.9999573101738780197 24E +0	.9999572938229324536 28E +0
0.059	.9999550632866472939 18E +0	.9999550451671055074 27E +0
0.060	.9999527388750578903 58E +0	.9999527188301080396 42E +0
0.061	.9999503356205553759 88E +0	.9999503134824244620 24E +0
0.062	.9999478522042730800 26E +0	.9999478277936578955 42E +0
0.063	.9999452873070214080 03E +0	.9999452604324996174 57E +0
0.064	.9999426396092769626 94E +0	.9999426100666979227 84E +0
0.065	.9999399077911714708 80E +0	.9999398753630263945 20E +0
0.066	.9999370905324805146 83E +0	.9999370549872515767 97E +0
0.067	.9999341865126120663 33E +0	.9999341476041000448 81E +0
0.068	.9999311944105948250 75E +0	.9999311518772248657 35E +0
0.069	.9999281129050663549 08E +0	.9999280664691714426 00E +0
0.070	.9999249406742610218 08E +0	.9999248900413427367 63E +0
0.071	.9999216763959977290 58E +0	.9999216212539638595 38E +0
0.072	.9999183187476674492 12E +0	.9999182587660460270 92E +0
0.073	.9999148664062205511 84E +0	.9999148012353498705 29E +0
0.074	.9999113180481539209 60E +0	.9999112473183480934 06E +0
0.075	.9999076723494978742 59E +0	.9999075956701874684 81E +0
0.076	.9999039279858028595 56E +0	.9999038449446501652 58E +0
0.077	.9999000836321259496 94E +0	.9998999937941143995 43E +0
0.078	.9998961379630171203 39E +0	.9998960408695143959 29E +0
0.079	.9998920896525053134 28E +0	.9998919848202996537 70E +0
0.080	.9998879373740842837 52E +0	.9998878242943935069 08E +0

Table B-6 (Cont'd)

THE AUXILIARY FUNCTIONS P(x) AND Λ(x)

1/x	P(x)	Λ(x)
0.081	.99988367980069822663 9E +0	.99988355793815096697 7E +0
0.082	.99987931560472718481 7E +0	.99987918439631583983 8E +0
0.083	.99987484345797223226 8E +0	.99987470231197710424 4E +0
0.084	.99987026203164043296 1E +0	.99987011032652454147 6E +0
0.085	.99986556999632957226 3E +0	.99986540707960360425 0E +0
0.086	.99986076602201265862 6E +0	.99986059120906951281 0E +0
0.087	.99985584877802219329 2E +0	.99985566135094056567 2E +0
0.088	.99985031603034205460 E +0	.99985061613935065196 3E +0
0.089	.99984566915505205041 9E +0	.99984545420650095191 7E +0
0.090	.99984040411138968045 E +0	.99984017418261081157 2E +0
0.091	.99983502046865639889 4E +0	.99983477469586777710 8E +0
0.092	.99982951689273705517 4E +0	.99982925437237677387 9E +0
0.093	.99982389204877774364 0E +0	.99982361183610841452 8E +0
0.094	.99981814460116693746 4E +0	.99981784570884642002 2E +0
0.095	.99981227321351809402 8E +0	.99981195461013413686 2E +0
0.096	.99980627654865171539 6E +0	.99980593715722013307 2E +0
0.097	.99980015326857714824 2E +0	.99979979196500285493 5E +0
0.098	.99979390203447411981 5E +0	.99979351764597432571 4E +0
0.099	.99978752150667400643 9E +0	.99978711281016286693 7E +0
0.100	.99978101034464083095 3E +0	.99978057606507482204 4E +0

Table B-7

THE AUXILIARY FUNCTIONS R(z) AND S(z) FOR $0.001 < \frac{1}{x} < 0.100$

$$z = \frac{3}{2} x^{-3/2}$$

1/x	R(z)	S(z)
0.001	.9999967C60443169741S4E +0	.1C0000329412278372087E 1
0.002	.9999906837184C4807046E +0	.1000009317618400785S6E 1
0.003	.9999828859253686759S3E +0	.1000017118586350499S2E 1
0.004	.9999736530313125820S2E +0	.1C3002635766313425321E 1
0.005	.9999631819599442953C0E +0	.1C3003683892765163C54E 1
0.006	.9999516057421886341S8E +0	.1C0004843035158859894E 1
0.007	.9999390221717768453S9E +0	.1C0006103514383002080E 1
0.008	.9999255071316273771276 +0	.1C3007457842408119226E 1
0.009	.9999111217693533325S8E +0	.1CC008900004736307109E 1
0.010	.9998959167557449530Z9E +0	.1C0010425034553319610E 1
0.011	.9998799349952748257626 +0	.1000120287416783552l6E 1
0.012	.9998632134452231362686 +0	.1000137C753065198110 4E 1
0.013	.9998457843884690266466 +0	.1000154582734574137 25E 1
0.014	.9998276763544857548456 +0	.1000172782174214913 46F 1
0.015	.9998089148045334612566 +0	.1000191649166959229 76E 1
0.016	.9997895226534484894706 +0	.1000211161800789239 81E 1
0.017	.9997695206749735254C4E +0	.1C0023130030482812003E 1
0.018	.9997489278220668028366 +0	.1000252046729037383 93E 1
0.019	.9997277614838364962266 +0	.1CC0273384687289431 26E 1
0.020	.9997060376943676368206 +0	.1CC0295299148547995 89E 1
0.021	.9996837713044347169766 +0	.1CC0317776265158328 27E 1
0.022	.9996609761242072621336 +0	.1000340803230174833 21E 1
0.023	.9996376650428675530956 +0	.1CC0364368157701980 81E 1
0.024	.9996138501298789208136 +0	.1CC0388459981685068 99E 1
0.025	.9995895427212332992196 +0	.1CC0413068369647152 42E 1
0.026	.9995647534935061980346 +0	.1000438183648648976 75E 1
0.027	.9995394925278242281226 +0	.1000463796741331786 12E 1
0.028	.9995137693654493788836 +0	.1000489899110343805 15E 1
0.029	.9994875930563417713146 +0	.1000516482709788672 75E 1
0.030	.9994609722018016460596 +0	.1000543539942595069 06E 1
0.031	.9994339149920874997536 +0	.1000571063622910623 29E 1
0.032	.9994064292397465810746 +0	.1000599046942783892 05E 1
0.033	.9993785224092662599646 +0	.1000627483442525895 58E 1
0.034	.9993502016435524270226 +0	.1000656366984245056 33E 1
0.035	.9993214737876595003366 +0	.1000685691728131963 45E 1
0.036	.9992923454101214788876 +0	.1000715452111137525 80E 1
0.037	.9992628228222065917506 +0	.1000745642827742965 20E 1
0.038	.9992329120953169099746 +0	.1000776258812565285 45E 1
0.039	.9992026190767868802496 +0	.1000807295224579255 12E 1
0.040	.9991719494042476691206 +0	.1000838747432768029 75E 1

Table B-7 (Cont'd)

THE AUXILIARY FUNCTIONS R(z) AND S(z) FOR $0.001 < \frac{1}{x} < 0.100$

$$z = \frac{3}{2} x^{-3/2}$$

1/x	R(z)	S(z)
0.041	.99914C9085187261276826E +0	.1000870611003040584266E 1
0.042	.9991C95016766173850356E +0	.1000902881686275970C0E 1
0.043	.9990777339606525770136E +0	.1000935555407372879466E 1
0.044	.9990456102899675873316E +0	.1000968628255198647576E 1
0.045	.9990131354293653626016E +0	.1001002096473345129806E 1
0.046	.9989803139978529579876E +0	.1001035956451610302616E 1
0.047	.9989471504765247302006E +0	.1001070204718134171676E 1
0.048	.9989136492158546681036E +0	.1001104837932126000486E 1
0.049	.9988798144424535949166E +0	.1001139852877127127586E 1
0.050	.9988456502653406947716E +0	.1001175246454759922816E 1
0.051	.9988111606817733250436E +0	.1001211015678918923676E 1
0.052	.9987763495826743291206E +0	.1001247157670364940326E 1
0.053	.9987412207576918787806E +0	.1001283669651687103246E 1
0.054	.9987057778999242277C76E +0	.1001320548942601475236E 1
0.055	.9986700246103305458906E +0	.1001357792955555806138E 1
0.056	.9986339644018741644966E +0	.1001395399191630391236E 1
0.057	.9985976007033806072953E +0	.1001433365236668334836E 1
0.058	.9985609368632042801786E +0	.1001471688757683044936E 1
0.059	.9985239761525867037886E +0	.1001510367499462865356E 1
0.060	.9984867217689214881806E +0	.1001549399281385809016E 1
0.061	.9984491768387491025306E +0	.1001588781994423758826E 1
0.062	.9984113444206101840986E +0	.1001628513598320930846E 1
0.063	.9983732275077318333486E +0	.1001668592118934397596E 1
0.064	.9983348290305639690C5E +0	.1001709015645725069616E 1
0.065	.9982961518591763362C2E +0	.1001749782329388544036E 1
0.066	.9982571988055258709716E +0	.1001790890379616127466E 1
0.067	.9982179726256033012376 +0	.1001832338062977153486E 1
0.068	.9981784760214671430166 +0	.1001874123700914444876E 1
0.069	.9981387116431725802176 +0	.1001916245667845437236E 1
0.070	.9980986820906021200286 +0	.1001958702389362078286E 1
0.071	.9980583899152043721836 +0	.1002001492340523161106E 1
0.072	.9980178376216468031136 +0	.1002044614044233247146E 1
0.073	.9979770276693878748106 +0	.1002088066069702775746E 1
0.074	.9979359624741735598596 +0	.1002131847033098437651E 1
0.075	.9978946444094628558586 +0	.1002175955585580768386E 1
0.076	.9978530758077865854746 +0	.1002220390433119967066E 1
0.077	.9978112589620433448126 +0	.1002265150314093842996E 1
0.078	.9977691961267370143496 +0	.1002310234008656347016E 1
0.079	.9977268895191570858466 +0	.1002355640335477989506E 1
0.080	.9976843413205086076696 +0	.1002401368150653395496E 1

Table B-7 (Cont'd)

THE AUXILIARY FUNCTIONS R(z) AND S(z) FOR $0.001 < \frac{1}{x} < 0.100$

$$z = \frac{3}{2} x^{-3/2}$$

1/x	R(z)	S(z)
0.081	.9976415536769910980 51E +0	.1002447416346658979 50E 1
0.082	.9975985287008313621 86E +0	.1002493783851357983 62E 1
0.083	.9975552684712720643 34E +0	.1002540469627050314 26E 1
0.084	.9975117750355185660 80E +0	.1002587472669564780 32E 1
0.085	.9974680504096462797 54E +0	.1002634792007391494 46E 1
0.086	.9974240965794706359 59E +0	.1002682426700852340 55E 1
0.087	.9973799155013816359 46E +0	.1002730375841307611 43E 1
0.088	.9973355091031448279 61E +0	.1002778638550396800 45E 1
0.089	.9972908792846704352 91E +0	.1002827213979312146 95E 1
0.090	.9972460279187522550 24E +0	.1002876101308103014 04E 1
0.091	.9972009568517778482 49E +0	.1002925299745009780 88E 1
0.092	.9971556679044114486 57E +0	.1002974808525825772 89E 1
0.093	.9971101628722509313 08E +0	.1003024626913285911 72E 1
0.094	.9970644435264601053 90E +0	.1003074754196480836 15E 1
0.095	.9970185116143775165 83E +0	.1003125189690295323 63E 1
0.096	.9969723688601028321 65E +0	.1003175932734869905 45E 1
0.097	.9969260169650622076 43E +0	.1003226982695084643 49E 1
0.098	.9968794576085525859 71E +0	.1003278338960064084 77E 1
0.099	.9968326924482676132 11E +0	.1003330000942702477 86E 1
0.100	.9967857231208043091 65E +0	.1003381968079208381 53E 1

+61

Table B-8

THE AUXILIARY FUNCTIONS W(z) AND X(z) FOR $0.001 < \frac{1}{x} < 0.100$

$$z = \frac{3}{2} x^{-3/2}$$

1/x	-W(z)	X(z)
0.001	.10000046115561842615 3E 1	.99999538824633309895 2E +0
0.002	.10000130429400406991 1E 1	.99998695548009814917 1E +0
0.003	.10000239601965108013 4E 1	.99997603447145748614 3E +0
0.004	.10000368869222432252 3E 1	.99996310043886528715 6E +0
0.005	.10000515475331251893 6E 1	.99994842778153503616 3E +0
0.006	.10000677558948056167 6E 1	.99993220144891478207 2E +0
0.007	.10000853752048543465 5E 1	.99991455705852587134 5E +0
0.008	.10001042993358148788 2E 1	.99989559955290769071 0E +0
0.009	.10001244427894809735 4E 1	.99987541324533809865 6E +0
0.010	.10001457347350096532 1E 1	.99985406778171699647 1E +0
0.011	.10001681152142021157 5E 1	.99983162193528153189 4E +0
0.012	.10001915325947187318 5E 1	.99980812615338676121 5E +0
0.013	.10002159417881609941 8E 1	.99978362433941890221 9E +0
0.014	.10002413029606691432 3E 1	.99975815514219245327 1E +0
0.015	.10002675805736435151 6E 1	.99973175291522313263 9E +0
0.016	.10002947426532311397 2E 1	.99970444844723511017 1E +0
0.017	.10003227602228555919 3E 1	.99967626952962452710 6E +0
0.018	.10003516068547764934 8E 1	.99964724140489290802 1E +0
0.019	.10003812583103740666 6E 1	.99961738712635506158 6E +0
0.020	.10004116922477844829 5E 1	.99958672785049572784 9E +0
0.021	.10004428879814885024 3E 1	.99955528307737247530 0E +0
0.022	.10004748262825522419 2E 1	.99952307085036600956 6E +0
0.023	.10005074892110861621 0E 1	.99949010792371180282 7E +0
0.024	.10005408599745335040 2E 1	.99945640990420181312 7E +0
0.025	.10005749228068829664 5E 1	.99942199137196150143 5E +0
0.026	.10006096628649932205 3E 1	.99938686598411454949 5E +0
0.027	.10006450661390330375 0E 1	.99935104656433150176 7E +0
0.028	.10006811193746581654 0E 1	.99931454518064119497 4E +0
0.029	.10007178100050185138 2E 1	.99927737321341141530 0E +0
0.030	.10007551260910545810 3E 1	.99923954141503984789 6E +0
0.031	.10007930562688274515 1E 1	.99920105996261098988 0E +0
0.032	.10008315897028516393 3E 1	.99916193850454975018 1E +0
0.033	.10008707160445788689 3E 1	.99912218620212364359 9E +0
0.034	.10009104253953241623 8E 1	.99908181176650220886 7E +0
0.035	.10009507082730412189 7E 1	.99904082349196666344 0E +0
0.036	.10009915555824480680 6E 1	.99899922928576881287 5E +0
0.037	.10010329585880808217 1E 1	.99895703669506138663 1E +0
0.038	.10010749088899166141 1E 1	.99891425293125871195 5E +0
0.039	.10011173984012591643 7E 1	.99887088489213428740 0E +0
0.040	.10011604193286239470 4E 1	.99882693918191827005 9E +0

Table B-8 (Cont'd)

THE AUXILIARY FUNCTIONS W(z) AND X(z) FOR $0.001 < \frac{1}{x} < 0.100$

$$z = \frac{3}{2} x^{-3/2}$$

1/x	-W(z)	X(z)
0.041	.10012039641533963890 8E 1	.99878242212962145583 9E +0
0.042	.10012480256150671192 9E 1	.99873733980578172492 2E +0
0.043	.10012925966958741328 2E 1	.99869169803780308753 2E +0
0.044	.10013376706067036454 8E 1	.99864550242403555343 4E +0
0.045	.10013832407741200578 2E 1	.99859875834672540266 3E +0
0.046	.10014293008284113897 0E 1	.99855147098394949444 0E +0
0.047	.10014758445925502169 8E 1	.99850364532063358016 5E +0
0.048	.10015228660719819107 4E 1	.99845528615874281744 8E +0
0.049	.10015703594451621468 8E 1	.99840639812672251444 2E +0
0.050	.10016183190547744640 4E 1	.99835698568825832380 3E +0
0.051	.10016667393995663027 8E 1	.99830705315041745008 4E +0
0.052	.10017156151267486381 4E 1	.99825660467122575551 1E +0
0.053	.10017649410249101564 2E 1	.99820564426672980976 8E +0
0.054	.10018147120174020370 3E 1	.99815417581758781960 2E +0
0.055	.10018649231561539097 7E 1	.99810220307522886408 1E +0
0.056	.10019155696158855109 9E 1	.99804972966761590855 7E +0
0.057	.10019666466886820717 3E 1	.99799675910464455998 5E +0
0.058	.10020181497789045667 8E 1	.99794329478320643065 4E +0
0.059	.10020700743984087048 2E 1	.99788933999194322604 3E +0
0.060	.10021224161620489891 8E 1	.99783489791571522242 4E +0
0.061	.10021751707834463563 5E 1	.99777997163980562280 5E +0
0.062	.10022283340709998475 1E 1	.99772456415388033085 9E +0
0.063	.10022819019241245088 6E 1	.99766867835572094209 1E +0
0.064	.10023358703296992811 3E 1	.99761231705474718669 1E +0
0.065	.10023902353587100353 7E 1	.99755548297534366138 0E +0
0.066	.10024449931630741800 8E 1	.99749817876000441967 8E +0
0.067	.10025001399726343942 7E 1	.99744040697230786016 8E +0
0.068	.10025556720923100724 0E 1	.99738217009973331974 9E +0
0.069	.10026115858993959915 1E 1	.99732347055632985823 0E +0
0.070	.10026678778409985520 7E 1	.99726431068524687107 5E +0
0.071	.10027245444316007028 5E 1	.99720469276113541831 3E +0
0.072	.10027815822507473553 7E 1	.99714461899242845383 3E +0
0.073	.10028389879408437192 8E 1	.99708409152350751738 5E +0
0.074	.10028967582050595623 5E 1	.99702311243676287795 5E +0
0.075	.10029548898053329241 2E 1	.99696168375455359135 9E +0
0.076	.10030133795604672840 6E 1	.99689980744107346318 6E +0
0.077	.10030722243443166249 2E 1	.99683748540412846779 4E +0
0.078	.10031314210840532289 2E 1	.99677471949683077692 8E +0
0.079	.10031909667585134098 8E 1	.99671151151921418568 5E +0
0.080	.10032508583966167189 8E 1	.99664786321977538884 0E +0

Table B-8 (Cont'd)

THE AUXILIARY FUNCTIONS W(z) AND X(z) FOR $0.001 < \frac{1}{x} < 0.100$

$$z = \frac{3}{2} x^{-3/2}$$

1/x	-W(z)	X(z)
0.081	.10033110930758544733 2E 1	.99658377629694524817 1E +0
0.082	.10033716679208437332 6E 1	.99651925240049391507 1E +0
0.083	.10034325801019431219 9E 1	.99645429313287340409 4E +0
0.084	.10034938268339271162 0E 1	.99638890005050097781 4E +0
0.085	.10035554053747156597 0E 1	.99632307466498647979 6E +0
0.086	.10036173130241561569 5E 1	.99625681844430654688 5E +0
0.087	.10036795471228550912 6E 1	.99619013281392844402 6E +0
0.088	.10037421050510566885 5E 1	.99612301915788608789 5E +0
0.089	.10038049842275662082 5E 1	.99605547881981066492 9E +0
0.090	.10038681821087155930 1E 1	.99598751310391809833 1E +0
0.091	.10039316961873693496 8E 1	.99591912327595547759 3E +0
0.092	.10039955239919686603 3E 1	.99585031056410843717 7E +0
0.093	.10040596630856118447 3E 1	.99578107615987134785 2E +0
0.094	.10041241110651694066 5E 1	.99571142121888207253 5E +0
0.095	.10041888655604319976 7E 1	.99564134686172293682 1E +0
0.096	.10042539242332897340 9E 1	.99557085417468946154 6E +0
0.097	.10043192847769413868 0E 1	.99549994421052832011 6E +0
0.098	.10043849449151320531 8E 1	.99542861798914589338 1E +0
0.099	.10044509024014179945 3E 1	.99535687649828871947 0E +0
0.100	.10045171550184573987 8E 1	.99528472069419705933 6E +0

Table B-9

THE AIRY FUNCTIONS Ai(-x) AND Bi(-x) FOR 0 < x < 10

x	Ai(-x)	Bi(-x)
0.0	.35502805388781723925 6E +0	.61492662744600073514 7E +0
0.1	.38084866812012151177 9E +0	.56999904300295486043 6E +0
0.2	.40628418744480141036 2E +0	.52450903281848551282 4E +0
0.3	.43090309528558085827 1E +0	.47797784010989294675 7E +0
0.4	.45422561388866739839 0E +0	.43002093994850337519 1E +0
0.5	.47572809161053958879 8E +0	.38035265975105385017 0E +0
0.6	.49484952543114968829 6E +0	.32879184076086943820 6E +0
0.7	.51100397575010149392 E +0	.27526801198787969865 3E +0
0.8	.52357394970577400835 2E +0	.21982751089758181041 6E +0
0.9	.53195994561097391354 8E +0	.16263894892596624932 6E +0
1.0	.53556088329235211880 1E +0	.10399738949694461188 7E +0
1.1	.53381051043050215980 9E +0	.44326586740826603010 4E -1
1.2	.52619437480212007386 7E +0	-.15821370184632083689 7E -1
1.3	.51227200604103093087 9E +0	-.75769644150340703534 4E -1
1.4	.49170018106129072860 3E +0	-.13472406095279475205 0E +0
1.5	.46425657774886940647 9E +0	-.19178486115704122001 7E +0
1.6	.42986297676913517919 0E +0	-.24596320021137348593 7E +0
1.7	.38860703739632873707 E +0	-.29620265761049573531 9E +0
1.8	.34076155912421395331 3E +0	-.34140583183013496392 7E +0
1.9	.28680005627949084406 7E +0	-.38046587745364882174 2E +0
2.0	.22740742820168557599 4E +0	-.41230258795639848808 8E +0
2.1	.16348451299924273734 2E +0	-.43590234823067674541 8E +0
2.2	.96145378007669001804 1E -1	-.45036098416820731241 2E +0
2.3	.26706333057356845660 4E -1	-.45492823439436498165 9E +0
2.4	-.43334140440309514204 9E -1	-.44905227628210751389 8E +0
2.5	-.11232506769296608919 0E +0	-.43242247184070529303 0E +0
2.6	-.17850242893640334380 1E +0	-.40500827813003355803 2E +0
2.7	-.24003810974245725446 7E +0	-.36709211182100781894 4E +0
2.8	-.29509759299920866541 3E +0	-.31929388893831209992 4E +0
2.9	-.34190509567298304096 5E +0	-.26258499816469701493 3E +0
3.0	-.37881429367765807435 0E +0	-.19828962637492654321 1E +0
3.1	-.40438222239097832428 8E +0	-.12807165128975454031 0E +0
3.2	-.41744342056415136517 9E +0	-.53905755630539283651 2E -1
3.3	-.41718093737455012877 1E +0	.21967999989777454377 3E -1
3.4	-.40319048424589972740 2E +0	.97106190548660719253 2E -1
3.5	-.37553382314043191193 2E +0	.16893983748105861185 1E +0
3.6	-.33477747747482189978 6E +0	.23486630624776703497 5E +0
3.7	-.28201306184193150173 8E +0	.29235261007145199472 1E +0
3.8	-.21885547561885578381 6E +0	.33904647075666408299 1E +0
3.9	-.14741930564074419 5021E +0	.37289057831939561931 8E +0
4.0	-.70265532949289515095 7E -1	.39223470570699289558 E +0

Table B-9 (Cont'd)

THE AIRY FUNCTIONS Ai(-x) AND Bi(-x) FOR 0 < x < 10

x	Ai(-x)	Bi(-x)
4.1	.96769795187143328313 8E -2	.39593974C7401389214 62E +0
4.2	.89210763239450579C145E -1	.38346736127C944661885E +0
4.3	.16499783927277230 4379E +0	.3549490626 1C930446158E +C
4.4	.23370325807316312 9975E +0	.3112285990063957812 49E +0
4.5	.29215278105539546 6887E +C	.2538726576969326367 90E +0
4.6	.33749597548946 28631 87E +0	.1851457579472126770 30E +0
4.7	.36736748287733133 0869E +0	.1079469455485536552 93E +0
4.8	.38003667664279293 8416E +0	.2570779336508269253 14E -1
4.9	.37453635470583879 9451E +0	-.5774655314043531203 49E -1
5.0	.35076100902411431 9779E +0	-.1383691349016005763 75E +0
5.1	.30952599628731751 5927E +0	-.2120891315690364759 36E +0
5.2	.25258C33810474473 4695E +0	-.2750270441896407740 21E +0
5.3	.18256793106833949 9387E +0	-.3237160767487924039 92E +0
5.4	.10293459611013465 2261E +0	-.3553170800 8C467545504E +0
5.5	.17761541276574975 5965E -1	-.3678134539157119910 96E +0
5.6	-.68330699686167863 9290E -1	-.3601722254377133442 44E +0
5.7	-.15062015797853529 3745E +0	-.3324582531879016843 19E +0
5.8	-.22435191633495887 5935E +0	-.2858902110493242468 50E +0
5.9	-.28512277955517990 3034E +0	-.2228296929478136933 66E +0
6.0	-.32914517362982310 5240E +0	-.1466983766 7C557037848E +0
6.1	-.35351167612C964874673E +0	-.6182254919628C89021 04E -1
6.2	-.35642107366896143 1057E +0	.2679080896533582627 69E -1
6.3	-.33734764921613506 2331E +0	.1137370090081979768 96E +0
6.4	-.29713762213662779 6395E +0	.1935413606157801751 19E +0
6.5	-.23802030199711580 3576E +0	.2610126576364839518 42E +0
6.6	-.16352646272772955 2083E +0	.3115999458112095482 96E +0
6.7	-.78312471801255919 1819E -1	.3417277386067502601 27E +0
6.8	.12104524277365038 1496E -1	.3490841790 4C394786340E +0
6.9	.10168739773376451 5792E +0	.3328378429834287330 87E +0
7.0	.18428083525050563 7301E +0	.2937620718544140201 10E +0
7.1	.25403632856197836 4411E +0	.2342508783298568275 00E +0
7.2	.30585152336862657 1561E +0	.1582173900904976929 15E +0
7.3	.33577037051514730 8967E +0	.7087411376989631221 07E -1
7.4	.34132375223233864 1094E +0	-.2159651857188359989 81E -1
7.5	.32177571638064787 5261E +0	-.1124634950764908064 08E +0
7.6	.27825C23488019733 0050E +0	-.1949337564738764709 51E +C
7.7	.21372C37378919298 0439E +0	-.2626700688053946763 88E +0
7.8	.13285154462606732 6912E +0	-.3103005661474120823 15E +0
7.9	.41701883617387043 3440E -1	-.3338785630030469105 49E +0
8.0	-.52705050356386202 6505E -1	-.3312515807511378599 63E +0

Table B-9 (Cont'd)

THE AIRY FUNCTIONS Ai(-x) AND Bi(-x) FOR 0 < x < 10

x	Ai(-x)	Bi(-x)
8.1	-.142908147093581424377E +0	-.302303309060702033165E +0
8.2	-.221599454803604415167E +0	-.249040191297943596885E +0
8.3	-.282231759958830619293E +0	-.175505563096446039117E +0
8.4	-.319592189726197978383E +0	-.875179820004110091821E -1
8.5	-.330290237630203879022E +0	.775443644765840446080E -2
8.6	-.313112452617262463685E +0	.102356470012673830991E +0
8.7	-.269204540700509324562E +0	.188203626247790888932E +0
8.8	-.202054447376745762278E +0	.257782401703263694214E +0
8.9	-.117266306371752132940E +0	.304832413364962959881E +0
9.0	-.221337215473414036634E -1	.324947323455244917922E +0
9.1	.749598372735548019466E -1	.316034712393298914417E +0
9.2	.165268004651479631477E +0	.278584254357115239686E +0
9.3	.240473796853185974428E +0	.215708345763898821874E +0
9.4	.293477556112067757598E +0	.132938761140427214287E +0
9.5	.319103247713128201379E +0	.377854324894665022643E -1
9.6	.314651583311693258632E +0	-.609129273601140532029E -1
9.7	.280237501916297438275E +0	-.153794208777252970928E +0
9.8	.218867432663286421209E +0	-.231863308817480920680E +0
9.9	.136235026447979751373E +0	-.287383557724355170119E +0
10.0	.402412384864431906390E -1	-.314679829643838633171E +0

Table B-10

AIRY FUNCTION DERIVATIVES Ai'(-x) AND Bi'(-x) FOR 0 < x < 10

x	Ai'(-x)	Bi'(-x)
0.0	-.25881940379280679840E +0	.44828835735382635791E +0
0.1	-.25695811232364617461E +0	.45121336229346124209E +0
0.2	-.25103267400554781265E +0	.45938529458683409658E +0
0.3	-.24054512725815460873E +0	.47188021630064791999E +0
0.4	-.22503140930241503157E +0	.48773486404914753388E +0
0.5	-.20408167033954738614E +0	.50593371362384716657E +0
0.6	-.17736259869656603299E +0	.52540115229915356590E +0
0.7	-.14464128564332102740E +0	.54499912006918190665E +0
0.8	-.10580999118796886752E +0	.56353059459633115436E +0
0.9	-.60910998002198836248E -1	.57974926126528144469E +0
1.0	-.10160567116645209356E -1	.59237562642279235081E +0
1.1	.46029154710607865615E -1	.60011969648991087315E +0
1.2	.10703156927228079368E +0	.60171015743746440877E +0
1.3	.17199180675377403307E +0	.59592974940901122716E +0
1.4	.23981911993629659397E +0	.58165624392193851793E +0
1.5	.30918696720241042021E +0	.55790810302189735413E +0
1.6	.37854219195188082704E +0	.52389354144640984792E +0
1.7	.44612455463607505269E +0	.47906133847344781739E +0
1.8	.50999762771970664839E +0	.42315137373868928948E +0
1.9	.56809171681229921580E +0	.35624251263424769866E +0
2.0	.61825902074169104141E +0	.27879516692116952268E +0
2.1	.65834069281434206450E +0	.19168563232842986824E +0
2.2	.68624482490900170982E +0	.96229185938564530770E -1
2.3	.70003366287657600235E +0	-.58110593070515435988E -2
2.4	.69801760154444186707E +0	-.11223237096351617945E +0
2.5	.67895273426479436337E +0	-.22042015487462958769E +0
2.6	.64163798711106179488E +0	-.32739716870710052908E +0
2.7	.58600720014433144410E +0	-.42989534308201506299E +0
2.8	.51221098154347935003E +0	-.52445039743268310177E +0
2.9	.42118281160364836685E +0	-.60751828797109686723E +0
3.0	.31458376921659881364E +0	-.67561122268525853767E +0
3.1	.19482044600397890368E +0	-.72544957150816480299E +0
3.2	.65031146995263151366E -1	-.75412455310841361057E +0
3.3	-.70963617177836128662E -1	-.75926511504794454976E +0
3.4	-.20874904975273324321E +0	-.73920162510551817070E +0
3.5	-.34344343345404814630E +0	-.69311628490728880174E +0
3.6	-.46986396630325112664E +0	-.62117282664232020667E +0
3.7	-.58272780365295797804E +0	-.52461361490968349106E +0
3.8	-.67688257140887363406E +0	-.40581572062896559895E +0
3.9	-.74755808553547747198E +0	-.26829836289214567898E +0
4.0	-.79062857536858138030E +0	-.11667056743834089366E +0

Table B-10 (Cont'd)

AIRY FUNCTION DERIVATIVES Ai'(-x) AND Bi'(-x) FOR 0 < x < 10

x	Ai'(-x)	Bi'(-x)
4.1	-.80287253541821499018 9E +0	.43478717931659297743 1E -1
4.2	-.78221560786245196399 8E +0	.20575691122112262390 0E +0
4.3	-.72794081383881804657 0E +0	.36320468095665562906 8E +0
4.4	-.64085018328756365186 9E +0	.50858932154467567508 3E +0
4.5	-.52336253231574770069 7E +0	.63474476777366370975 4E +0
4.6	-.37953391433584533571 5E +0	.73494444367130626644 8E +0
4.7	-.21499018008957243134 1E +0	.80328925850719965966 0E +0
4.8	-.36765104312346077472 5E -1	.83508975839556434895 3E +0
4.9	.14695742731095607499 0E +0	.82721903336068038276 4E +0
5.0	.32719281855444313684 4E +0	.77841177300189924607 4E +0
5.1	.49458599838493705510 0E +0	.68948512842205131003 2E +0
5.2	.63990516690128384335 6E +0	.56345897957517367729 4E +0
5.3	.75457541994701104103 6E +0	.40555694088331552818 2E +0
5.4	.83122307200334625868 0E +0	.22307496421783366878 0E +0
5.5	.86419721777139839077 9E +0	.25111583073630925961 6E -1
5.6	.85003256004893156261 6E +0	-.17783759557946047209 7E +0
5.7	.78781722463519971378 1E +0	-.37440903400372219022 6E +0
5.8	.67943151996650276407 2E +0	-.55300203169101126131 3E +0
5.9	.52962857256300237831 4E +0	-.70247952231549159994 6E +0
6.0	.34593548728134289487 5E +0	-.81289878510506700045 2E +0
6.1	.13836393725271684847 2E +0	-.87622530153251547901 6E +0
6.2	-.81068556196304158032 7E -1	-.88697896206682812795 0E +0
6.3	-.29899160898473956420 8E +0	-.84276109680897025718 9E +0
6.4	-.50147985025496865641 8E +0	-.74461386630573286775 8E +0
6.5	-.67495249251320217304 6E +0	-.59717066629162201692 0E +0
6.6	-.80711924947739184717 9E +0	-.40856733931004221565 7E +0
6.7	-.88790796525555354459 3E +0	-.19009877716374965990 5E +0
6.8	-.91030400515880440593 5E +0	.44376784534242448915 7E -1
6.9	-.87103105868638765790 4E +0	.27926390596966473839 5E +0
7.0	-.77100816841012654770 2E +0	.49824459005811348880 5E +0
7.1	-.61552878754022817213 0E +0	.68542057764269389711 1E +0
7.2	-.41412428115703515910 5E +0	.82650634027200478553 0E +0
7.3	-.18009580448329322433 7E +0	.90998427043632467365 1E +0
7.4	.70276323643265995235 42E -1	.92812809007040669312 0E +0
7.5	.31880950669855459626 2E +0	.87780228154576092235 2E +0
7.6	.54671881905734806952 0E +0	.76095509188391053863 6E +0
7.7	.73605241760464206731 6E +0	.58474044748990395682 0E +0
7.8	.87115540424658935835 3E +0	.36122930432440057268 7E +0
7.9	.94004299802628012645 4E +0	.10670215481213814453 4E +0
8.0	.93556093819830655101 4E +0	-.15945049781298138945 1E +0

Table B-10 (Cont'd)

AIRY FUNCTION DERIVATIVES Ai'(-x) and Bi'(-x) FOR 0 < x < 10

x	Ai'(-x)	Bi'(-x)
8.1	.856218586328624562390E +0	-.416156639540127581404E +0
8.2	.706598697862805303377E +0	-.642322930908422933926E +0
8.3	.497276790253209576825E +0	-.818600440748798464895E +0
8.4	.244220894145284124529E +0	-.929109583748750229015E +0
8.5	-.323133482846391359413E -1	-.962969165120174798144E +0
8.6	-.309330272415632307481E +0	-.915479180196183452706E +0
8.7	-.562976849501854659673E +0	-.788826225075567710548E +0
8.8	-.770613009748042222000E +0	-.592213709228808931614E +0
8.9	-.912892757425249831881E +0	-.341364753721780747904E +0
9.0	-.975663980926331594730E +0	-.574005138436692543430E -1
9.1	-.951496815451917697499E +0	.234843786584969132257E +0
9.2	-.840671073803800818187E +0	.508944015545784449208E +0
9.3	-.651492407895600378317E +0	.739280283029872949861E +0
9.4	-.399862366389844822776E +0	.903485370676841537059E +0
9.5	-.108095318811871239011E +0	.984714070002119703933E +0
9.6	.196950442321259114068E +0	.973499179547113071310E +0
9.7	.486286291239268209056E +0	.868983876598210580717E +0
9.8	.731544863628323622557E +0	.679367742131133091187E +0
9.9	.907813331537150439126E +0	.421472089183833376124E +0
10.0	.996265044132790055937E +0	.119414113399909238093E +0

Table B-11

THE AUXILIARY FUNCTIONS F(-x) AND $\chi(-x)$ FOR $0 < x < 10$

x	F(-x)	$\chi(-x)$
0.0	.71005610777563447851 2E +0	.5235987755982988730 77E +0
0.1	.68552506667C180C80129E +0	.589033441900681614322E +0
0.2	.66345803671058627151 0E +0	.659059510580702280188E +0
0.3	.64353732849214952368 6E +0	.733650221730043295678E +0
0.4	.62549079408670369410 95E +0	.812764547904918292785E +0
0.5	.60908567782136109507 4E +0	.896350013442875280481E +0
0.6	.59412130694863560070 1E +0	.984345226848848781909E +0
0.7	.58042560655572210897 9E +0	.1076682099843916491 32E 1
0.8	.56785016981412519466 9E +0	.1173287750435715031 78E 1
0.9	.55626685272643500509 47E +0	.1274086103801057351 95E 1
1.0	.54556476859770152531 8E +0	.1378999215067597813 26E 1
1.1	.53564774557367049400 3E +0	.1487948343593070370 18E 1
1.2	.52643217590484833697 8E +0	-.1540737843238036163 88E 1
1.3	.51784519612329251551 0E +0	-.1423951983937848152 24E 1
1.4	.50982314644915685829 8E +0	-.1303363429082602261 06E 1
1.5	.50231026562486019158 2E +0	-.1179047241357311592 51E 1
1.6	.49525758414601869186 3E +0	-.1051076569882337579 19E 1
1.7	.48862198465631098624 7E +0	-.9195224570281577448 73E +0
1.8	.48236540317936521541 6E +0	-.7844537140981945779 01E +0
1.9	.47645414909358700889 7E +0	-.6459368514584304381 61E +0
2.0	.47085832522835198548 2E +0	-.5040360511706437806 40E +0
2.1	.46555133249049742164 5E +0	-.3588131723391825371 58E +0
2.2	.46050944591093225440 9E +0	-.2103277812194461323 54E +0
2.3	.45571145111193415842 2E +0	-.5863719968395807866 13E -1
2.4	.45113833195799565854 8E +0	.9620343306978824608 79E -1
2.5	.44677300162952435992 7E +0	.2541410896233404234 06E +0
2.6	.44260007059426724028 1E +0	.4151247774381380594 09E +0
2.7	.43860564598502292966 7E +0	.5791054584328735163 96E +0
2.8	.43477715775702590272 9E +0	.7460359669480052357 82E +0
2.9	.43110320772212495944 3E +0	.9158709275517183097 47E +0
3.0	.42757343816286219619 7E +0	.1088566673777727229 00E 1
3.1	.42417841723731198936 0E +0	.1264081168596777272 89E 1
3.2	.42090953881136783825 8E +0	.1442373927196252345 22E 1
3.3	.41775893471266318174 8E +0	-.1518186711125667699 18E 1
3.4	.41471939770080193425 6E +0	-.1334453040154904772 30E 1
3.5	.41178431370147611624 1E +0	-.1148053976741653312 23E 1
3.6	.40894760206519707267 8E +0	-.9590245117010296685 19E +0
3.7	.40620366279127953713 0E +0	-.7673984254739253779 30E +0
3.8	.40354732980981384569 0E +0	-.5732083470480919109 50E +0
3.9	.40097382954315016047 2E +0	-.3764858098015735744 56E +0
4.0	.39847874407765344636 0E +0	-.1772613043646245208 49E +0

Table B-11 (Cont'd)

THE AUXILIARY FUNCTIONS F(-x) AND $\chi(-x)$ FOR $0 < x < 10$

x	F(-x)	$\chi(-x)$
4.1	.396057778369296382848E +0	.24435671387510140357 4E -1
4.2	.393707730985646885583E +0	.22857656508817762119 8E +0
4.3	.391424467954153791204E +0	.43513372525059252162 9E +0
4.4	.389204899344157008727E +0	.64408035912564233923 5E +0
4.5	.387045958259268376954E +0	.85539049291513515513 7E +0
4.6	.384944781958960945037E +0	.10690389342113674570 7E 1
4.7	.382898694864436350938E +0	.12850012365372258367 3E 1
4.8	.380905193235010317135E +0	.15032536658653335186 8E 1
4.9	.378961931328120366931E +0	-.14178194845900982553 7E 1
5.0	.377066708879254078016E +0	-.11950553098811575088 6E 1
5.1	.375217459758157524567E +0	-.97006824308639981642 6E +0
5.2	.373412241675063848442E +0	-.74287946762744265834 4E +0
5.3	.371649226825785307056E +0	-.51350959608329055823 6E +0
5.4	.369926693377574382492E +0	-.28197869405254120662 1E +0
5.5	.368243917709197207595E +0	-.48306302736841235081 8E -1
5.6	.366596667328490950681E +0	.18748853967596673761 4E +0
5.7	.364986194399500362183E +0	.42538727775547259299 4E +0
5.8	.363410229818858493547E +0	.66537181963023768376 3E +0
5.9	.361867477787778881352E +0	.90742451878279986276 4E +0
6.0	.360356710831896444993E +0	.11515281567778717723 6E 1
6.1	.358876765226359340483E +0	.13976659268673850220 0E 1
6.2	.357426536738124267050E +0	-.14957712351700752429 7E 1
6.3	.356004977001422578536E +0	-.12456140514658953170 2E 1
6.4	.354611089445912374368E +0	-.99347083766699515545 9E +0
6.5	.353243926500171540619E +0	-.73935690195742403446 9E +0
6.6	.351902586295969483520E +0	-.48328721138303265717 0E +0
6.7	.350586209901225201809E +0	-.22527640397797786378 5E +0
6.8	.349293978711754666645E +0	.34661199682076420991 4E -1
6.9	.348025112033864465506E +0	.29651158546727678208 8E +0
7.0	.346778864841590233409E +0	.56026103519075197153 5E +0
7.1	.345554525693032724621E +0	.82589611658695338176 8E +0
7.2	.344351414798833436348E +0	.10934036737443194377 7E 1
7.3	.343168882211874670130E +0	.13627708179671325370 1E 1
7.4	.342006306158802590470E +0	-.15076077345467611422 5E 1
7.5	.340863091471970915822E +0	-.12345590566956612707 6E -1
7.6	.339738668131700251165E +0	-.35968793999872046813 3E +0
7.7	.338632489904355071462E +0	-.68300628787916640259 5E +0
7.8	.337544033069666956045E +0	-.40452577657112093056 7E +0
7.9	.336472795230487629845E +0	-.12425786212832343387 9E +0
8.0	.335418294198746428144E +0	.15778621285942263006 4E +0

Table B-11 (Cont'd)

THE AUXILIARY FUNCTIONS F(-x) AND $\chi(-x)$ FOR $0 < x < 10$

x	F(-x)	$\chi(-x)$
8.1	.33438006695192080018 7E +0	.44159541282702442498 0E +0
8.2	.33335766865431144582 9E +0	.72715890305794490681 1E +0
8.3	.33235067174185098711 5E +0	.10144660437578117002 9E 1
8.4	.33135866505557145311 1E +0	.13035063843652705543 9E 1
8.5	.33038125303721559097 95E +0	-.15473229955014936685 5E 1
8.6	.32941805496580303354 5E +0	-.12548468769349060859 6E 1
8.7	.32846870424226163268 8E +0	-.96066782832217491023 5E +0
8.8	.32753284771550002693 2E +0	-.66479559583747937092 9E +0
8.9	.32661014504754943919 0E +0	-.36723975169836563956 5E +0
9.0	.32570026811512269481 8E +0	-.68009748656361160551 0E -1
9.1	.32480290044514566435 7E +0	.23238517568012076327 6E +0
9.2	.32391773668200291145 7E +0	.53543589468276749144 3E +0
9.3	.32304448208441060364 7E +0	.83963343869285698949 3E +0
9.4	.32218285204998642199 4E +0	.11454680811562674234 8E 1
9.5	.32133257166572907229 40E +0	.14529338348974013991 1E 1
9.6	.32049337528275678037 6E +0	-.13795732050640180532 1E 1
9.7	.31966500611375666365 5E +0	-.10688752506204364622 5E 1
9.8	.31884721585174417433 1E +0	-.75657324546074339623 1E +0
9.9	.31803976430878315899 0E +0	-.44267535394077840448 8E +0
10.0	.31724241907348723917 2E +0	-.12718961779199818179 7E +0

Table B-12

THE AUXILIARY FUNCTIONS G(-x) AND $\psi(-x)$ FOR $0 < x < 10$

x	G(-x)	$\psi(-x)$
0.0	.5176388075856135968 10E +0	-.5235987755982988730 77E + 0
0.1	.5192503922012017849 80E +0	-.5176778438555394761 25E +0
0.2	.5235000219771535815 4E +0	-.5001162462038540683 69E +0
0.3	.5296535629858332807 05E +0	-.4714241937498935010 67E +0
0.4	.5371447037639591861 83E +0	-.4322776368403107911 71E +0
0.5	.5455439952467598111 10E +0	-.3834135891833329856 47E +0
0.6	.5545303077863981705 18E +0	-.3255637049540378017 80E +0
0.7	.5638662451226667833 63E +0	-.2594169083193005525 85E +0
0.8	.5733781346382966015 07E +0	-.1856016564017732963 86E +0
0.9	.5829402676220467056 37E +0	-.1046803373954442325 36E +0
1.0	.5924627582421763453 96E +0	-.1715055520345212861 78E -1
1.1	.6018823250424586268 03E +0	.7655007969627040303 08E -1
1.2	.6111553569955067060 59E +0	.1760376935864642719 48E +0
1.3	.6202527289912672912 61E +0	.2809755844621653943 87E +0
1.4	.6291559396369054780 01E +0	.3910679567946635677 34E +0
1.5	.6378542404854861891 14E +0	.5060542383254060356 01E +0
1.6	.6463425050675498803 17E +0	.6257040763561973060 08E +0
1.7	.6546196485511320555 84E +0	.7498130120679433767 67E +0
1.8	.6626874567823570921 90E +0	.8781987828606582862 82E +0
1.9	.6705497196470278882 28E +0	.1010698182743929442 19E 1
2.0	.6782115907495812845 33E +0	.1147164406356873993 15E 1
2.1	.6856791155171781113 47E +0	.1287464805635202273 79E 1
2.2	.6929588847404911072 91E +0	.1431478995138668736 21E 1
2.3	.7000577815942522133 66E +0	-.1562495403362766063 56E 1
2.4	.7069827983465704682 49E +0	-.1411373503203899006 07E 1
2.5	.7137409050164794952 46E +0	-.1256840226126908923 77E 1
2.6	.7203389567289560023 58E +0	-.1098980668444422256 30E 1
2.7	.7267836298545819387 42E +0	-.9378735919380261942 76E +0
2.8	.7330813795077827521 09E +0	-.7735921412188909828 68E +0
2.9	.7392384128342404623 40E +0	-.6062044604268881675 81E +0
3.0	.7452606739073196956 82E +0	-.4357742252535003348 46E +0
3.1	.7511538370950182249 37E +0	-.2623611035462756377 49E +0
3.2	.7569233065415514985 64E +0	-.8602115550449356559 95E -1
3.3	.7625742199966822624 94E +0	.9319281737520491138 62E -1
3.4	.7681114556698839720 95E +0	.2752309669682239872 82E +0
3.5	.7735396411215052525 07E +0	.4600461304266676863 54E +0
3.6	.7788631634562214308 85E +0	.6475935939746654881 67E +0
3.7	.7840861802753610769 24E +0	.8378308841049258855 33E +0
3.8	.7892126309892815072 16E +0	.1030717582394807205 88E 1
3.9	.7942462482001874371 80E +0	.1226215160747195248 69E 1
4.0	.7991905689481933685 65E +0	.1424286834349478414 84E 1

Table B-12 (Cont'd)

THE AUXILIARY FUNCTIONS G(-x) AND ψ(-x) FOR 0 < x < 10

x	G(-x)	ψ(-x)
4.1	.804048945675481671192E +0	-.151669522353918070788E 1
4.2	.808824556809993349956E +0	-.131357938539370748616E 1
4.3	.813520416904915027116E +0	-.110799059734451453425E 1
4.4	.818139386296111178574E +0	-.899959860854939646721E +0
4.5	.822684180258739585962E +0	-.689516973236561002359E +0
4.6	.827157377658214842839E +0	-.476690606528444655191E +0
4.7	.831561429100696942054E +0	-.261508379250849194564E +0
4.8	.835898664595332228468E +0	-.439969218317589530947E -1
4.9	.840171300745290503232E +0	.175818063599578120140E +0
5.0	.844381447487781477634E +0	.397911748446898110853E +0
5.1	.848531114405121591294E +0	.622260127379910565288E +0
5.2	.852622216629883459693E +0	.848839972102365533434E +0
5.3	.856656580367442898789E +0	.107762878874780C91706E 1
5.4	.860635948059031479432E +0	.130860477854412135033E 1
5.5	.864561983207849384927E +0	.154174680143015464324E 1
5.6	.868436274890002612295E +0	-.136455831124604539644E 1
5.7	.872260341971082121080E +0	-.112714517365717285347E 1
5.8	.876035637048161681424E +0	-.8876257961230'8307746E +0
5.9	.879763550135901422812E +0	-.6460189978344. 656487E +0
6.0	.883445412114338873721E +0	-.402343085697844260908E +0
6.1	.887082497954852809149E +0	-.156615877919021551990E +0
6.2	.890676029739714650430E +0	.911452739427772037477E -1
6.3	.894227179489609303784E +0	.340923464323467164194E +0
6.4	.897737071812519953795E +0	.592702214291307838158E +0
6.5	.901206786386434187105E +0	.846465453369565681894E +0
6.6	.904637360287444389125E +0	.110219750245156439426E 1
6.7	.908029790173984451370E +0	.135988305772384984232E 1
6.8	.911385034337167058519E +0	-.152208547806873814761E 1
6.9	.914704014626459944950E +0	-.126053739554673339458E 1
7.0	.917987618259263681754E +0	-.997079613718492100342E +0
7.1	.921236699522325546220E +0	-.731726078363095322299E +0
7.2	.924452081372341367564E +0	-.464490427673114399531E +0
7.3	.927634556942557341738E +0	-.195386OC3666475301416E +0
7.4	.930784890961684025294E +0	.755741369988520503742E -1
7.5	.933903821090972437148E +0	.348377212767067968128E +0
7.6	.936992059184874858238E +0	.623010707446694083374E +0
7.7	.940050292480318058296E +0	.899462360958682585056E +0
7.8	.943079184719251940367E +0	.117772016053921019430E 1
7.9	.946079377208799746445E +0	.145777233236833661661E 1
8.0	.949051489823024909124E +0	-.140198531999185832399E 1

Table B-12 (Cont'd)

THE AUXILIARY FUNCTIONS G(-x) AND ψ(-x) FOR 0 < x < 10

x	G(-x)	ψ(-x)
8.1	.95199612195004239496 3E +0	-.11183788088354661728 9E 1
8.2	.95491385338793712466 8E +0	-.83301189109689047991 4E +0
8.3	.95780524519270706352 1E +0	-.54589546090611159280 7E +0
8.4	.96067084048122226183 5E +0	-.25704020978147804965 1E +0
8.5	.96351116519198202078 7E +0	.33543367060635156807 9E -1
8.6	.96632672880625910299 6E +0	.32584496467269381438 1E +0
8.7	.96911802503204124433 8E +0	.61985446260376080301 6E +0
8.8	.97188553245301499848 0E +0	.91556191937358591326 5E +0
8.9	.97462971514468409164 8E +0	.12129575671794665221 0E 1
9.0	.97735102325957300222 5E +0	.15120318068232296214 9E 1
9.1	.98004989358333550741 8E +0	-.13288174507432667987 0E 1
9.2	.98272675006346662760 4E +0	-.10264141747263874665 2E 1
9.3	.98538200431220398699 3E +0	-.72236014050994613439 4E +0
9.4	.98801605608510039846 1E +0	-.41666431959919829128 3E +0
9.5	.99062929373665282740 8E +0	-.10933553734554693534 7E +0
9.6	.99322209465428320479 9E +0	.19961752307831493587 1E +0
9.7	.99579482567188330187 7E +0	.51018631685507214998 9E +0
9.8	.99834784346405855012 4E +0	.82236243384379104057 6E +0
9.9	.10008814949221338337 5E 1	.11361375950382348618 7E 1
10.0	.10033961175129174780 5E 1	.14515036491558851405 1E 1

Table B-13

THE AIRY FUNCTIONS Ai(-x) AND Bi(-x) FOR x = 10.0(0.1)25.0

x	Ai(-x)	Bi(-x)
10.0	.40241238486443190645OE -1	-.31467982964383863317 5E +0
10.1	-.59726811133454156364 4E -1	-.31076751228793916085 9E +0
10.2	-.15369678260708316794 6E +0	-.27573422910068368247 4E +0
10.3	-.23210801885482926713 8E +0	-.21282251839031565963 6E +0
10.4	-.28680545563082962100 3E +0	-.12818997344572347184 6E +0
10.5	-.31192603505 1C50600856E +0	-.30356123264021013166 9E -1
10.6	-.30457846499736678868 2E +0	.70620643405552847192 0E -1
10.7	-.26523367267346853213 7E +0	.16416118648516124796 4E +0
10.8	-.19776999205637179785 6E +0	.24027830578046136380 5E +0
10.9	-.10915179222515884141 2E +0	.29066745657553457493 1E +0
11.0	-.87595892557023813240 6E -2	.30965476742678188633 2E +0
11.1	.92570433646458226252 8E -1	.29489078683381124146 3E +0
11.2	.18370367405125353981 4E +0	.24770078655861551015 0E +0
11.3	.25443982511504728832 0E +0	.17303649043586083824 0E +0
11.4	.29668451242348505916 8E +0	.79016665218781160262 8E -1
11.5	.30542297004359265639 7E +0	-.23909272355945757578 3E -1
11.6	.27937454868332368680 0E +0	-.12409483526516101907 5E +0
11.7	.22123780856502705139 7E +0	-.21000838045383247243 6E +0
11.8	.13747955388507191353 8E +0	-.27157718924346775962 0E +0
11.9	.37673024339358523878 4E -1	-.30140609137784571662 6E +0
12.0	-.66555175054373129541 6E -1	-.29571991207807305671 4E +0
12.1	-.16285945350511373106 3E +0	-.25490469243486525528 6E +0
12.2	-.23964109627913476978 0E +0	-.18356517661700472991 9E +0
12.3	-.28747208025644136331 4E +0	-.90071313508555151972 7E -1
12.4	-.30029984453501425785 4E +0	.14371598200523129879 0E -1
12.5	-.27627456138116024821 9E +0	.11703336725739277661 7E +0
12.6	-.21808009991802631440 6E +0	.20520168053060346951 8E +0
12.7	-.13270691889389718572 6E +0	.26777225448472205711 0E +0
12.8	-.30673323846291911787 7E -1	.29668644672943310886 8E +0
12.9	.75227278581536191579 3E -1	.28802641802419088410 4E +0
13.0	.17151043937053704464 1E +0	.24261322909262719932 7E +0
13.1	.24572340714791971080 E +0	.16601077672138687180 7E +0
13.2	.28808667803239288202 4E +0	.67911972051572192234 8E -1
13.3	.29283500740797452870 6E +0	-.39035799213856985728 8E -1
13.4	.25906846994341417055 0E +0	-.14083154455696338936 6E +0
13.5	.19098124329622029266 4E +0	-.22395010358002279149 1E +0
13.6	.97413890384212864185 4E -1	-.27716252780599944500 7E +0
13.7	-.92354469683709487193 2E -2	-.29310029616882818361 1E +0
13.8	-.11461607446263517171 6E +0	-.26934032193708026538 4E +0
13.9	-.20434433220527378206 8E +0	-.20884394738523514060 7E +0
14.0	-.26598348278407779839 6E +0	-.11966555279762452310 0E +0

Table B-13 (Cont'd)

THE AIRY FUNCTIONS Ai(-x) AND Bi(-x) FOR x = 10.0(0.1)25.0

x	Ai(-x)	Bi(-x)
14.1	-.290810085149966542400E +0	-.139449179819057303158E -1
14.2	-.275111233707217394091E +0	.937010056063833635907E -1
14.3	-.220812567684753037941E +0	.188181084943215964558E +0
14.4	-.135323956376930263935E +0	.256056757239035563509E +0
14.5	-.305974189395514228158E -1	.287492243517527755240E +0
14.6	.784944011048661970258E -1	.277740807135961818392E +0
14.7	.176247385514908148444E +0	.227935185937784278396E +0
14.8	.248393528798947757662E +0	.145042705101175547154E +0
14.9	.284219508153073098728E +0	.409630871062547004781E -1
15.0	.278217490870828929532E +0	-.691265945310100611765E -1
15.1	.231006381313208420753E +0	-.168958580952192226378E +0
15.2	.149360880167607559296E +0	-.243582556265014671221E +0
15.3	.453142726642754678562E -1	-.281639100357291913367E +0
15.4	-.655593766657786769160E -1	-.277148644512263496030E +0
15.5	-.166447954090419767435E +0	-.230526530754712210696E +0
15.6	-.241853776309456335635E +0	-.148642777434945160149E +0
15.7	-.280008778231409792267E +0	-.438863503404062751514E -1
15.8	-.274771784867787423562E +0	.676523425806300522417E -1
15.9	-.226693950298588365530E +0	.168624065964169058925E +0
16.0	-.143057931669099697712E +0	.243123151428227216730E +0
16.1	-.368522003634394557182E -1	.279229116187111048782E +0
16.2	.751891724478489316167E -1	.270976772753752360411E +0
16.3	.175195586927892081738E +0	.219420661690464175667E +0
16.4	.247016437066724377416E +0	.132593822101696703640E +0
16.5	.278868480560550838314E +0	.243335984326956926555E -1
16.6	.265334889965864CC6144E +0	-.878681881139096494492E -1
16.7	.208367580336450845224E +0	-.185666818653931467286E +0
16.8	.117096950227064733015E +0	-.252874291645393359750E +0
16.9	.644492332728703505772E -2	-.278182521778200078603E +0
17.0	-.105262300290952390249E +0	-.257135921002343182134E +0
17.1	-.199309661928061248022E +0	-.192999160312283731216E +0
17.2	-.259744953337002906108E +0	-.963407131723165460050E -1
17.3	-.276134329617757533339E +0	.166345106620348942834E -1
17.4	-.245438411334952268887E +0	.126754403954158017658E +0
17.5	-.172660590662226267760E +0	.215120245578695335030E +0
17.6	-.701200288287906168874E -1	.266374079965369105099E +0
17.7	.445633965857197437876E -1	.271424902922468651251E +0
17.8	.151456073406667976582E +0	.229141108518447431837E +0
17.9	.231771307055489357663E +0	.146682144363863919525E +0
18.0	.271204540804414221595E +0	.383724885083839979915E -1

Table B-13 (Cont'd)

THE AIRY FUNCTIONS Ai(-x) AND Bi(-x) FOR x = 10.0(0.1)25.0

x	Ai(-x)	Bi(-x)
18.1	.26254622975241249499 5E +0	-.76722385474784147689 9E -1
18.2	.20708491826324285806 9E +0	-.17812044269450548583 7E +0
18.3	.11451480335836944917 0E +0	-.24757486160386942083 7E +0
18.4	.13234763638499023668 8E -2	-.27240184116085796201 7E +0
18.5	-.11208853977554047611 3E +0	-.24787069072043292520 6E +0
18.6	-.20506448027031100812 1E +0	-.17819401041305928427 0E +0
18.7	-.26047345019757374070 4E +0	-.75898311962065983569 8E -1
18.8	-.26792290176798836669 8E +0	.40352128765281317594 7E -1
18.9	-.22578996428749362666 6E +0	.14911548634577605836 5E +0
19.0	-.14166127688042265628 4E +0	.23012109009458831472 3E +0
19.1	-.31056856107879188249 6E -1	.26808155260491100834 2E +0
19.2	.85366443974465093724 6E -1	.25564609869517024572 0E +0
19.3	.18564166484965181741 0E +0	.19491310236499858577 3E +0
19.4	.25064664026586608860 6E +0	.97176400585324683600 7E -1
19.5	.26780027210258394575 1E +0	-.19091642257576624214 0E -1
19.6	.23357738324529758287 3E +0	-.13167759554280297720 8E +0
19.7	.15432579636976751633 2E +0	-.21885666862504631517 4E +0
19.8	.45183771258220759061 5E -1	-.26361300179118279158 7E +0
19.9	-.72738820111013022684 0E -1	-.25702629238684711614 4E +0
20.0	-.17640612707798468969 5E +0	-.20013930932265134918 6E +0
20.1	-.24536173925299817280 6E +0	-.10390009445385105850 3E +0
20.2	-.26581264230882624837 8E +0	.12857578511441220767 3E -1
20.3	-.23346817960876582377 7E +0	.12704193771341239525 9E +0
20.4	-.15453541737774759094 0E +0	.21585296293514630075 4E +0
20.5	-.44625680397011909639 8E -1	.26136213786923029214 6E +0
20.6	.74257771105699970962 3E -1	.25419791055362635141 4E +0
20.7	.17808610491682873076 4E +0	.19556724781879920465 8E +0
20.8	.24566765630448155807 9E +0	.97160953360554275495 0E -1
20.9	.26302274861374406909 6E +0	-.21088632996163868951 7E -1
21.0	.22635849367898896615 9E +0	-.13498730557827747612 0E +0
21.1	.14299516277795202989 7E +0	-.22101436210668334135 8E +0
21.2	.30003254695201258109 9E -1	-.26121095119355668991 1E +0
21.3	-.89218132530860659168 2E -1	-.24700005240800777388 5E +0
21.4	-.18975240330993084782 5E +0	-.18111215041707953699 4E +0
21.5	-.25038504298749525105 5E +0	-.77166876875313811352 1E -1
21.6	-.25813673053613818228 8E +0	.43055191387411079917 3E -1
21.7	-.21113063664897696011 3E +0	.15412397794381714160 4E +0
21.8	-.11914346967509305814 0E +0	.23233229025669939586 7E +0
21.9	-.16691709824184591091 1E -2	.26079659134132007266 3E +0
22.0	.11614415376051412929 2E +0	.23318112230442141606 6E +0

Table B-13 (Cont'd)

THE AIRY FUNCTIONS Ai(-x) AND Bi(-x) FOR x = 10.0(0.1)25.0

22.1	.20885298273289002315 5E +0	.15520836067172141391 9E +0
22.2	.25623726068416421984 0E +0	.43577662594042324884 30E -1
22.3	.24777817952575062748 0E +0	-.77529934887337660528 2E -1
22.4	.18508979044193268680 2E +0	-.18164821345723122979 1E +0
22.5	.81723500904036645619 3E -1	-.24581693455638800940 9E +0
22.6	-.39669365019314130121 0E -1	-.25569986940908184444 1E +0
22.7	-.15224473541320950148 3E +0	-.20887793332279489106 8E +0
22.8	-.23089319146455619514 7E +0	-.11554274327791784388 3E +0
22.9	-.25788171470767799650 0E +0	.36214680931358179592 4E -2
23.0	-.22693405337408287510 9E +0	.12195196890032855675 1E +0
23.1	-.14479389046039034381 8E +0	.21274931453866224030 4E +0
23.2	-.29861942686336177877 7E -1	.25532906376097741686 4E +0
23.3	.91845035223295394183 6E -1	.23980659170195122288 1E +0
23.4	.19254559087057344985 7E +0	.16949296042857731853 3E +0
23.5	.24904892692121109686 0E +0	.60300560228815318713 1E -1
23.6	.24815825583741328959 6E +0	-.62767463895516741495 7E -1
23.7	.18985000758036468919 6E +0	-.17129179813305858476 5E +0
23.8	.87442563738137360253 7E -1	-.24000043209073741156 2E +0
23.9	-.35347643155887028945 5E -1	-.25270599728356920511 1E +0
24.0	-.14983659008188653339 6E +0	-.20621104602689683959 1E +0
24.1	-.22906183229077721646 8E +0	-.11121935238440398431 1E +0
24.2	-.25417414164992341885 1E +0	.10023886009865782649 0E -1
24.3	-.21900764764881496224 2E +0	.12886953299622837124 2E +0
24.4	-.13170165747682098451 2E +0	.21701136686263882776 6E +0
24.5	-.12926044703241092501 7E -1	.25325983212568291536 4E +0
24.6	.10893192964526881472 9E +0	.22871516898466549979 8E +0
24.7	.20451908164387973001 5E +0	.14905939385319442061 9E +0
24.8	.25060928039252086254 4E +0	.33354119027650651168 7E -1
24.9	.23581960584548666047 57E +0	-.90433138158859617925 3E -1
25.0	.16352657883042946943 3E +0	-.19214681569037802382 4E +0

Table B-14

THE AIRY DERIVATIVES Ai'(-x) AND Bi'(-x) FOR x = 10.0(0.1)25.0

x	Ai'(-x)	Bi'(-x)
10.0	.99626504413279005589 2E +0	.11941411339990923818 4E +0
10.1	.98630683706012814809 5E +0	-.19752877902133186015 4E +0
10.2	.87698998000019466195 6E +0	-.49769246188428512101 3E +0
10.3	.67749282954091277893 5E +0	-.75018586978857374913 0E +0
10.4	.40656884037507168565 4E +0	-.92812682641186776958 9E +0
10.5	.90957487390681672806 5E -1	-.10116140816303775186 5E 1
10.6	-.23713206714159534241 7E +0	-.99010107964385990582 0E +0
10.7	-.54324590031593679610 1E +0	-.86387973414224358717 3E +0
10.8	-.79430738718621806417 4E +0	-.64446103069798284709 6E +0
10.9	-.96226006822773245908 E +0	-.35274773790404555739 8E +0
11.0	-.10273278736645794214 6E 1	-.22022995314464466625 3E -1
11.1	-.98050614982505289903 8E +0	.31508609193607221748 4E +0
11.2	-.82495955004224843419 4E +0	.62038365507859871608 0E +0
11.3	-.57610761992286277470 0E +0	.85922966430325768850 9E +0
11.4	-.26031689811010133445 2E +0	.10035593383790820954 0E 1
11.5	.87724154321784443111 5E -1	.10353264046930834408 9E 1
11.6	.42871139792683259104 7E +0	.94893760770695526378 0E +0
11.7	.72313366159037479788 9E +0	.75233866282229031482 9E +0
11.8	.93589786065535150147 9E +0	.46655210869869128816 5E +0
11.9	.10406290259592336172 8E 1	.12364175631678618702 6E +0
12.0	.10231104533679707298 5E 1	-.23673219783112331651 9E +0
12.1	.88340231823096755046 0E +0	-.57182121120131837294 3E +0
12.2	.63631251174038143469 3E +0	-.84086190019696732146 6E +0
12.3	.31007878814201665186 9E +0	-.10101178597436740115 8E 1
12.4	-.56663225663481390090 7E -1	-.10572617696936247002 8E 1
12.5	-.41933133041950516453 2E +0	-.97451653616717407210 2E +0
12.6	-.73277547091736961993 0E +0	-.77009836368870831516 0E +0
12.7	-.95694539101927640950 7E +0	-.46769574585194737509 0E +0
12.8	-.10621354773225733819 3E 1	-.10395630703946273409 3E +0
12.9	-.10331100184723800808 7E 1	.27578969397116391581 1E +0
13.0	-.87151967787995336664 6E +0	.62309724881928773364 2E +0
13.1	-.59621238207884556652 5E +0	.89259793395238440150 0E +0
13.2	-.24129921481379101677 9E +0	.10480275683464319996 1E 1
13.3	.14787197201463607509 2E +0	.10672821816670515536 4E 1
13.4	.52039301825738595019 8E +0	.94578137470553936217 6E +0
13.5	.82643275142525423805 1E +0	.69760874733408185728 4E +0
13.6	.10239783333201546158 8E 1	.35417395574824247196 8E +0
13.7	.10847644707773194267 9E 1	-.39532307075829851982 2E -1
13.8	.99853865825849945960 6E +0	-.43068271822482143234 5E +0
13.9	.77499796688398813275 6E +0	-.76565006660932960516 5E +0
14.0	.44302487700284364111 9E +0	-.99741181894933352408 9E +0

Table B-14 (Cont'd)

THE AIRY DERIVATIVES Ai'(-x) AND Bi'(-x) FOR x = 10.0(0.1)25.0

x	Ai'(-x)	Bi'(-x)
14.1	.4721154863556667658 85E -1	-.1092298861803436621 78E 1
14.2	-.3579536148211646294 13E +0	-.1035105941248726136 70E 1
14.3	-.7155102899955546591 17E +0	-.8317659880057362184 97E +0
14.4	-.9740665007723178672 65E +0	-.5091011119196131960 60E +0
14.5	-.1095321272880539215 04E 1	-.1115622228670333133 30E +0
14.6	-.1059955678469324704 35E 1	.3046961334913821072 92E +0
14.7	-.8709631196073223636 92E +0	.6796511915408309639 24E +0
14.8	-.5538226675711727537 13E +0	.9580843328875273855 00E +0
14.9	-.1533596645695470169 05E +0	.1097840900904636024 12E 1
15.0	.2723742043086420208 13E +0	.1076429753084374786 72E 1
15.1	.6604047286625017303 64E +0	.8949053233926250200 04E +0
15.2	.9521577832523489463 77E +0	.5783365716285292311 22E +0
15.3	.1102425630013205230 32E 1	.1726547305308247644 25E +0
15.4	.1086592300799084918 79E 1	-.2617825817252033970 75E +0
15.5	.9049379354302121993 87E +0	-.6590509566800734121 79E +0
15.6	.5832411648483673162 61E +0	-.9576666656218781995 72E +0
15.7	.1694412301027278264 46E +0	-.1110228511266440422 21E 1
15.8	-.2732700098353935395 71E +0	-.1091169277099032513 33E 1
15.9	-.6759748137663457337 44E +0	-.9013220878987427967 55E +0
16.0	-.9747644416212727180 65E +0	-.5684556059761353725 13E +0
16.1	-.1121015416635294229 16E 1	-.1435393846886623696 16E +0
16.2	-.1089540451198865617 24E 1	.3068225142291062131 65E +0
16.3	-.8832181769900260907 47E +0	.7107117910615042690 88E +0
16.4	-.5332183646697741173 92E +0	.1002396553596652831 10E 1
16.5	-.9462257996353213983 54E -1	.1133177108022710437 37E 1
16.6	.3620096253898709592 14E +0	.1079770385107840746 10E 1
16.7	.7618832999302872410 91E +0	.8487569775231355614 74E +0
16.8	.1038252466682624296 28E 1	.4762082104974518352 41E +0
16.9	.1143729800198708975 00E 1	.2238135728103492420 70E -1
17.0	.1058684576644660077 31E 1	-.4378020657909875096 81E +0
17.1	.7952045342413724567 95E +0	-.8270350629473295748 50E +0
17.2	.3957903023289157380 55E +0	-.1078670297887255294 83E 1
17.3	-.7318011263045013382 56E -1	-.1148327233556689487 47E 1
17.4	-.5322759987626111307 94E +0	-.1022014271756370472 70E 1
17.5	-.9024049204808416901 39E +0	-.7192395068395728134 53E +0
17.6	-.1118529827836791741 47E 1	-.2903953794665570366 89E +0
17.7	-.1141324531893842888 30E 1	.1913226227061095623 98E +0
17.8	-.9646473245242933319 49E +0	.6422293059966337839 63E +0
17.9	-.6173689563028825295 33E +0	.9826621194101303087 01E +0
18.0	-.1590389152049680154 08E +0	.1151187094108641798 77E 1

Table B-14 (Cont'd)

THE AIRY DERIVATIVES Ai'(-x) AND Bi'(-x) FOR x = 10.0(0.1)25.0

x	Ai'(-x)	Bi'(-x)
18.1	.33004279631029815655 5E +0	.11159490495000316001 63E 1
18.2	.76275159369174277768 1E +0	.88103100978871261408 6E +0
18.3	.10606792615372507975 9E 1	.48650797249558900716 2E +0
18.4	.11685209936692642261 4E 1	.19766678418310605686 5E -2
18.5	.10646439622797084435 5E 1	-.48547203836493565871 8E +0
18.6	.76577289959799539668 1E +0	-.88681248892578422222 8E +0
18.7	.32473691126077585187 9E +0	-.11274197142341149841 3E 1
18.8	-.17852907557958681039 7E +0	-.11611768008061627755 0E 1
18.9	-.65126782852747812434 3E +0	-.97965278434569550725 5E +0
19.0	-.10049611250051395937 8E 1	-.61447375395607405630 0E +0
19.1	-.11720441755041667589 3E 1	-.13222406926272946144 4E +0
19.2	-.11190984017222779704 2E 1	.37739355451102809038 7E +0
19.3	-.85390189666690688123 8E +0	.81809877385061736935 6E +0
19.4	-.42479723269402106260 9E +0	.11052596589291124545 5E 1
19.5	.87741088343757135920 0E -1	.11823541560575811492 1E 1
19.6	.58595259082647081691 9E +0	.10324332543293874452 7E 1
19.7	.97336641774272132321 6E +0	.68220710420223902307 0E +0
19.8	.11735978413886146362 7E 1	.19773108952387510241 4E +0
19.9	.11456883090128122293 6E 1	-.32771864942117713113 2E +0
20.0	.89286285673647123810 8E +0	-.79142903383953647971 0E +0
20.1	.46277308619950873339 0E +0	-.11013441608282842177 2E 1
20.2	-.61078109738297581370 1E -1	-.11945427682999161354 89E 1
20.3	-.57527966186227385017 1E +0	-.10503583127457746151 9E 1
20.4	-.97684078847900855170 3E +0	-.69534809235365174772 3E +0
20.5	-.11839330197051474971 4E 1	-.19886802802168598022 6E +0
20.6	-.11528532455466198291 7E 1	.34012601798836457298 1E +0
20.7	-.88764234575269905353 8E +0	.81261879271883039100 4E +0
20.8	-.44017738739689700416 8E +0	.11216040227739315832 7E 1
20.9	.99557418317793456547 2E -1	.12022167587914782100 5E 1
21.0	.62129444990892707317 1E +0	.10357155970136586022 2E 1
21.1	.10169346110626482792 9E 1	.65423703849929066603 4E +0
21.2	.12030790320491278565 8E 1	.13506760857723428415 5E +0
21.3	.11389241064531652161 0E 1	-.41466427453319116992 3E +0
21.4	.83562406644433628721 7E +0	-.87992674483851427401 4E +0
21.5	.35490253223925327914 0E +0	-.11619031340872722587 2E 1
21.6	-.20309296438529421680 0E +0	-.11992314270417519015 9E 1
21.7	-.72040284745129832565 4E +0	-.98175488362289309609 8E +0
21.8	-.10861529409902343654 1E 1	-.55363072784734502627 8E +0
21.9	-.12204983771696351628 9E 1	-.48345438675823459312 8E -2
22.0	-.10924127512708336725 8E 1	.54742191289194569928 7E +0

Table B-14 (Cont'd)

THE AIRY DERIVATIVES Ai'(-x) AND Bi'(-x) FOR x = 10.0(0.1)25.0

x	Ai'(-x)	Bi'(-x)
22.1	-.72729256534439474385 9E +0	.98360098428992021601 9E +0
22.2	-.20244195596469776585 4E +0	.12078178564127246348 7E 1
22.3	.36890135323481691070 6E +0	.11692272049223617449 3E 1
22.4	.86179376875745081246 3E +0	.87398979346284210827 4E +0
22.5	.11669360550027303556 6E 1	.38492284314438991452 3E +0
22.6	.12151609668289614312 3E 1	-.19141686918645025559 7E +0
22.7	.99352554448858690147 3E +0	-.72767260849400730844 7E +0
22.8	.54918478503141453143 5E +0	-.11037812242891088036 7E 1
22.9	-.20145461627674067885 5E -1	-.12340422445151733974 6E 1
23.0	-.58733509004493979480 7E +0	-.10870259966754336821 1E 1
23.1	-.10241054007409109539 3E 1	-.69362155987026516691 5E +0
23.2	-.12301647764183670595 8E 1	-.14108477781938338718 3E +0
23.3	-.11565770876666464215 9E 1	.44591497679011812905 3E +0
23.4	-.81785059526086790833 2E +0	.93323335420096508998 0E +0
23.5	-.28967206432050399306 5E +0	.12079654473588682764 4E 1
23.6	.30755551158443454797 0E +0	.12048980829009690928 0E 1
23.7	.83590607485603680949 4E +0	.92244416412372866205 7E +0
23.8	.11717799733772984028 3E 1	.42407478318014115341 0E +0
23.9	.12350640508438819421 2E 1	-.17545140078652670708 6E +0
24.0	.10086744076771970092 7E 1	-.73620256184307328736 7E +0
24.1	.54362536577734614968 5E +0	-.11256708395442330459 1E 1
24.2	-.51937134972497351159 4E -1	-.12502816856202535443 0E 1
24.3	-.63752268565667070778 6E +0	-.10782848815735957076 5E 1
24.4	-.10733179188282119077 66E 1	-.64834185975044122718 0E +0
24.5	-.12537174187587190878 7E 1	-.61397217333928337311 1E -1
24.6	-.11332952894489004690 2E 1	.54261466532762564772 2E +0
24.7	-.73874946811693732595 2E +0	.10179604591586547891 4E 1
24.8	-.16357772635071010669 9E +0	.12483731437990010468 6E 1
24.9	.45363254245185146429 8E +0	.11758414691803483676 9E 1
25.0	.96237885138769741044 5E +0	.81571971575460585732 2E +0

Table B-15

THE AUXILIARY FUNCTIONS F(-x) AND $\chi(-x)$ FOR x = 10.0(0.1)25.0

x	F(-x)	$\chi(-x)$
10.0	.3172424190734872391 76E +0	.2186395895733655448 74E 2
10.1	.3164549551860512749 27E +0	.2218102461599423499 40E 2
10.2	.3156771548299263595 88E +0	.2249966239174110678 27E 2
10.3	.3149088070389744208 17E +0	.2281986459304353902 83E 2
10.4	.3141497074192861112 90E +0	.2314162363953297887 26E 2
10.5	.3133996578847680966 17E +0	.2346493205937420253 94E 2
10.6	.3126584664057035269 20E +0	.2378978248671913401 15E 2
10.7	.3119259467695378109 73E +0	.2411616765924272899 04E 2
10.8	.3112019183531911598 56E +0	.2444408041575759305 26E 2
10.9	.3104862059062450597 83E +0	.2477351369390416445 89E 2
11.0	.3097786393443921706 40E +0	.2510446052791344459 96E 2
11.1	.3090790535525784000 71E +0	.2543691404643940310 80E 2
11.2	.3083872881973023240 92E +0	.2577086747045832079 45E 2
11.3	.3077031875475709272 91E +0	.2610631411123246223 07E 2
11.4	.3070266003040420612 89E +0	.2644324736833559149 04E 2
11.5	.3063573794359131953 25E +0	.2678166072773795968 68E 2
11.6	.3056953820251432055 20E +0	.2712154775994850190 40E 2
11.7	.3050404691176192146 33E +0	.2746290211821208429 44E 2
11.8	.3043925055809040323 39E +0	.2780571753675973983 57E 2
11.9	.3037513599682216760 86E +0	.2814998782910992386 10E 2
12.0	.3031169043883588987 90E +0	.2849570688641890824 62E 2
12.1	.3024890143811797229 55E +0	.2884286867587851641 18E 2
12.2	.3018675687984677834 37E +0	.2919146723915948027 00E 2
12.3	.3012524496898279067 77E +0	.2954149669089877521 16E 2
12.4	.3006435421933938901 24E +0	.2989295121722936039 74E 2
12.5	.3000407344311039665 34E +0	.3024582507435081921 49E 2
12.6	.2994439174083190287 96E +0	.3060011258713945898 59E 2
12.7	.2988529849175713983 11E +0	.3095580814779649004 56E 2
12.8	.2982678334462438306 31E +0	.3131290621453296236 70E 2
12.9	.2976883620879896024 06E +0	.3167140131029019309 16E 2
13.0	.2971144724577149782 26E +0	.3203128802149447086 73E 2
13.1	.2965460686099551589 30E +0	.3239256099684487287 55E 2
13.2	.2959830569604840102 35E +0	.3275521494613307803 88E 2
13.3	.2954253462110065036 97E +0	.3311924463909410523 16E 2
13.4	.2948728472767909092 52E +0	.3348464490428694851 13E 2
13.5	.2943254732171053956 95E +0	.3385141062800412256 35E 2
13.6	.2937831391683308551 30E +0	.3421953675320917079 07E 2
13.7	.2932457622796285002 69E +0	.3458901827850122592 31E 2
13.8	.2927132616510471177 22E +0	.3495985025710574872 84E 2
13.9	.2921855582739608225 54E +0	.3533202779589060448 25E 2
14.0	.2916625749737337707 72E +0	.3570554605440666939 47E 2

Table B-15 (Cont'd)

THE AUXILIARY FUNCTIONS F(-x) AND $\chi(-x)$ FOR x = 10.0(0.1)25.0

x	F(-x)	$\chi(-x)$
14.1	.291144236354513583034E +0	.360804002439521902410E 2
14.2	.290630468746060199807E +0	.364565856266601501424E 2
14.3	.290121200152521607838E +0	.368340975146079217640E 2
14.4	.289616360203072292703E +0	.372129312689485163145E 2
14.5	.289115880104334462204E +0	.375930822990627626320E 2
14.6	.288619692594506030177E +0	.379745460617317754181E 2
14.7	.288127731899123077938E +0	.383573180603290953677E 2
14.8	.287639933688388031791E +0	.387413938440319066418E 2
14.9	.287156235035998122645E +0	.391267690070507588131E 2
15.0	.286676574379411839734E +0	.395134391878772412064E 2
15.1	.286200891481494065713E +0	.399014000685490774521E 2
15.2	.285729127393483395255E +0	.402906473739321271326E 2
15.3	.285261224419227803452E +0	.406811768710187996739E 2
15.4	.284797126080637352563E +0	.410729843682424031242E 2
15.5	.284336777084305014675E +0	.414660657148069672663E 2
15.6	.283880123289248951148E +0	.418604168000320965874E 2
15.7	.283427111675731734923E +0	.422560335527124240694E 2
15.8	.282977690315114035561E +0	.426529119404912515833E 2
15.9	.282531808340702215800E +0	.430510479692479768647E 2
16.0	.282089415919551118409E +0	.434504376824989207114E 2
16.1	.281650464225185058858E +0	.438510771608111811284E 2
16.2	.281214905411201688123E +0	.442529625212291537282E 2
16.3	.280782692585724955671E +0	.446560899167133697836E 2
16.4	.280353779786674890120E +0	.450604555355913149307E 2
16.5	.279928121957823328428E +0	.454660556010199026933E 2
16.6	.279505674925606068004E +0	.458728863704592877061E 2
16.7	.279086395376663193588E +0	.462809441351577138482E 2
16.8	.278670240836080545720E +0	.466902252196471023969E 2
16.9	.278257169646306453637E +0	.471007259812490948712E 2
17.0	.277847140946718955535E +0	.475124428095912743964E 2
17.1	.277440114653819776414E +0	.479253721261332982698E 2
17.2	.277036051442032330971E +0	.483395103837026828991E 2
17.3	.276634912725081968936E +0	.487548540660399904765E 2
17.4	.276236660637937585255E +0	.491713996873531746409E 2
17.5	.275841258019294580082E +0	.495891437918808499609E 2
17.6	.275448668394579975819E +0	.500080829534642573932E 2
17.7	.275058855959461282526E +0	.504282137751277049250E 2
17.8	.274671785563841450908E +0	.508495328886672693904E 2
17.9	.274287422696322965609E +0	.512720369542475520067E 2
18.0	.273905733469124812522E +0	.516957226600062864940E 2

Table B-15 (Cont'd)

THE AUXILIARY FUNCTIONS F(-x) AND $\chi(-x)$ FOR x = 10.0(0.1)25.0

x	F(-x)	$\chi(-x)$
18.1	.27352668460343670381 2E +0	.52120586721666604715 5E 2
18.2	.27315024341519556500 9E +0	.52546625882156770664 3E 2
18.3	.27277637780126988131 2E +0	.52973836911237199282 1E 2
18.4	.27240505622603806648 8E +0	.53402216605134582058 3E 2
18.5	.27203624770834755885 6E +0	.53831761786182946636 3E 2
18.6	.27166992180884186603 1E +0	.54262469302471482754 4E 2
18.7	.27130604861764327452 8E +0	.54694336027498971746 9E 2
18.8	.27094459874237941306 2E +0	.55127358859834661601 4E 2
18.9	.27058554329654231069 4E +0	.55561534722785434136 7E 2
19.0	.27022885388816902350 0E +0	.55996860564069115299 6E 2
19.1	.26987450260883331748 1E +0	.56433333355493783865 8E 2
19.2	.26952246202293829172 4E +0	.56870950092642937954 0E 2
19.3	.26917270515730020508 3E +0	.57309707794566382767 1E 2
19.4	.26882520549101413302 8E +0	.57749603503476706835 6E 2
19.5	.26847993694559242916 1E +0	.58190634284451217775 5E 2
19.6	.26813687387536729927 7E +0	.58632797225139212178 6E 2
19.7	.26779599105814911522 9E +0	.59076089435474457757 6E 2
19.8	.26745726368613240196 6E +0	.59520508047392769238 6E 2
19.9	.26712066735704172461 4E +0	.59966050214554562767 5E 2
20.0	.26678617806550998381 3E +0	.60412713112072276755 6E 2
20.1	.26645377219468189739 4E +0	.60860493936242550161 3E 2
20.2	.26612342650803570535 0E +0	.61309389904283052159 0E 2
20.3	.26579511814141638341 6E +0	.61759398254073860014 1E 2
20.4	.26546882459527388887 7E +0	.62210516243903284776 7E 2
20.5	.26514452372710019104 5E +0	.62662741152218047075 2E 2
20.6	.26482219374405905841 5E +0	.63116070277377079075 5E 2
20.7	.26450181319580278542 4E +0	.63570500937413261864 9E 2
20.8	.26418336096747024425 7E +0	.64026030469789802634 0E 2
20.9	.26386681627286084171 2E +0	.64482656231173173031 7E 2
21.0	.26355215864777914804 3E +0	.64940375597200514099 6E 2
21.1	.26323936794354514434 0E +0	.65399185962254630010 7E 2
21.2	.26292842432066520765 6E +0	.65859084739242087721 6E 2
21.3	.26261930824265911905 3E +0	.66320069359374972360 1E 2
21.4	.26231200047003853934 2E +0	.66782137271956221402 7E 2
21.5	.26200648205443255072 8E +0	.67245285944168462638 8E 2
21.6	.26170273433285601023 8E +0	.67709512860866282841 0E 2
21.7	.26140073892211660278 9E +0	.68174815524371855903 0E 2
21.8	.26110047771335661838 8E +0	.68641191454273861016 8E 2
21.9	.26080193286672560951 3E +0	.69108638187229623181 1E 2
22.0	.26050508680618021126 0E +0	.69577153276770410056 7E 2

Table B-15 (Cont'd)

THE AUXILIARY FUNCTIONS F(-x) AND $\chi(-x)$ FOR x = 10.0(0.1)25.0

x	F(-x)	$\chi(-x)$
22.1	.26020992221440752871 2E +0	.70046734293109820794 1E 2
22.2	.25991642202786861334 5E +0	.70517378822955204074 0E 2
22.3	.25962456943195866327 5E +0	.70989084469322044152 3E 2
22.4	.25933434785628069099 0E +0	.71461848851351255170 3E 2
22.5	.25904574097002950709 3E +0	.71935669604129325509 6E 2
22.6	.25875873267748296960 5E +0	.72410544378511255322 3E 2
22.7	.25847330711359754574 4E +0	.72886470840946231800 2E 2
22.8	.25818944863970532696 3E +0	.73363446673305988050 9E 2
22.9	.25790714183930972847 5E +0	.73841469572715792758 1E 2
23.0	.25762637151397719175 9E +0	.74320537251388019078 1E 2
23.1	.25734712267932229263 5E +0	.74800647436458242429 6E 2
23.2	.25706938056108373862 1E +0	.75281797869823818050 6E 2
23.3	.25679313059128881757 7E +0	.75763986307984890342 5E 2
23.4	.25651835840450393511 3E +0	.76247210521887787151 3E 2
23.5	.25624504983416895110 8E +0	.76731468296770753240 8E 2
23.6	.25597319090901309599 9E +0	.77216757432011978270 2E 2
23.7	.25570276784955031536 2E +0	.77703075740979875622 7E 2
23.8	.25543376706465195681 3E +0	.78190421050885569464 7E 2
23.9	.25516617514819477654 3E +0	.78678791202637548363 5E 2
24.0	.25489997887578230383 2E +0	.79168184050698444770 3E 2
24.1	.25463516520153766C940E +0	.79658597462943900597 2E 2
24.2	.25437172125496599270 6E +0	.80150029320523480024 4E 2
24.3	.25410963433788471527 8E +0	.80642477517723591544 0E 2
24.4	.25384889192141984654 0E +0	.81135939961832382122 2E 2
24.5	.25358948164306673226 2E +0	.81630414573006567171 8E 2
24.6	.25333139130381353162 4E +0	.82125899284140160869 2E 2
24.7	.25307460886532587380 3E +0	.82622392040735072104 8E 2
24.8	.25281912244719114377 3E +0	.83119890800773532163 9E 2
24.9	.25256492032422090024 5E +0	.83618393534592320951 0E 2
25.0	.25231199092380997226 6E +0	.84117898224758759327 9E 2

Table B-16

THE AUXILIARY FUNCTIONS G(-x) AND $\psi(-x)$ FOR x = 10.0(0.1)25.0

x	G(-x)	$\psi(-x)$
10.0	.10033961175129174780 2E 1	.20301059570694644571 1E 2
10.1	.10058920396211575198 3E 1	.20618008490890211495 9E 2
10.2	.10083695808765665215 6E 1	.20936532371587778320 6E 2
10.3	.10108290524662373529 1E 1	.21256623425426061747 5E 2
10.4	.10132707574332222127 2E 1	.21578273980311584348 9E 2
10.5	.10156949909620004251 5E 1	.21901476476574848052 3E 2
10.6	.10181020406515169625 9E 1	.22226223464224871208 4E 2
10.7	.10204921867764329918 6E 1	.22552507600297707250 4E 2
10.8	.10228657025371917936 7E 1	.22880321646294801823 4E 2
10.9	.10252228542994679711 4E 1	.23209658465707268740 6E 2
11.0	.10275639018235347587 3E 1	.23540511021622374367 1E 2
11.1	.10298890984840532998 0E 1	.23872872374408716066 5E 2
11.2	.10321986914807588250 5E 1	.24206735679476764149 6E 2
11.3	.10344929220404915912 7E 1	.24542094185111609207 4E 2
11.4	.10367720256109950988 9E 1	.24878941230374918598 4E 2
11.5	.10390362320468803756 4E 1	.25217270243073257946 9E 2
11.6	.10412857657881328791 9E 1	.25557074737790076470 3E 2
11.7	.10435208460315177308 9E 1	.25898348313978789420 9E 2
11.8	.10457416868952194482 7E 1	.26241084654114517486 0E 2
11.9	.10479484975770340073 1E 1	.26585277521902162182 0E 2
12.0	.10501414825064138549 0E 1	.26930920760538608566 0E 2
12.1	.10523208414906503293 3E 1	.27278008291026952489 6E 2
12.2	.10544867698554627638 9E 1	.27626534110540749492 5E 2
12.3	.10566394585802492776 8E 1	.27976492290836376714 7E 2
12.4	.10587790944282408397 3E 1	.28327876976711688237 5E 2
12.5	.10609058600717875706 1E 1	.28680682384509228392 8E 2
12.6	.10630199342129943667 0E 1	.29034902800662347107 6E 2
12.7	.10651214916999117494 5E 1	.29390532580282636572 2E 2
12.8	.10672107036384773085 7E 1	.29747566145787179709 5E 2
12.9	.10692877375003931832 2E 1	.30105997985564168318 3E 2
13.0	.10713527572271156695 4E 1	.30465822652675512605 3E 2
13.1	.10734059233301242205 1E 1	.30827034763595124325 2E 2
13.2	.10754473929876287808 6E 1	.31189628996981613113 4E 2
13.3	.10774773201378o654461 E 1	.31553600092484190005 2E 2
13.4	.10794958555691318068 1E 1	.31918942849580623785 3E 2
13.5	.10815031470066755758 8E 1	.32285652126446144828 8E 2
13.6	.10834993391966049951 4E 1	.32653722838852237664 4E 2
13.7	.10854845739869063655 8E 1	.33023149959094307747 6E 2
13.8	.10874589904057096470 5E 1	.33393928514947249987 9E 2
13.9	.10894227247369067193 7E 1	.33766053588647986582 5E 2
14.0	.10913759105932303899 4E 1	.34139520315904079774 0E 2

Table B-16 (Cont'd)

THE AUXILIARY FUNCTIONS G(-x) AND ψ(-x) FOR x = 10.0(0.1)25.0

x	G(-x)		ψ(-x)	
14.1	.109331867898689612256E	1	.345143238849275613696E	2
14.2	.109525115839790371568E	1	.348904595354931553561E	2
14.3	.109717347484009166263E	1	.352679225580201028046E	2
14.4	.109908575192503266559E	1	.356467082926768295549E	2
14.5	.110098811092385473662E	1	.360268121285077270283E	2
14.6	.110288067082706848919E	1	.364082295025813449704E	2
14.7	.110476354840247759101E	1	.367909558991593220666E	2
14.8	.110663685825124590153E	1	.371749868488854062969E	2
14.9	.110850071286219154565E	1	.375603179279939416137E	2
15.0	.111035522266437506782E	1	.379469447575372211521E	2
15.1	.111220049607804586002E	1	.383348630026311297373E	2
15.2	.111403663956400825273E	1	.387240683717185200172E	2
15.3	.111586375767146599346E	1	.391145566158497871593E	2
15.4	.111768195308440130321E	1	.395063235279801267437E	2
15.5	.111949132666654229204E	1	.398993649422829793517E	2
15.6	.112129197750497022228E	1	.402936767334791833723E	2
15.7	.112308400295241592674E	1	.406892548161813748116E	2
15.8	.112486749866829261271E	1	.410860951442531894284E	2
15.9	.112664255865851030527E	1	.414841937101828383267E	2
16.0	.112840927531411530059E	1	.418835465444706433223E	2
16.1	.113016773944879620541E	1	.422841497150301329131E	2
16.2	.113191804033529642930E	1	.426859993266023136175E	2
16.3	.113366026574077136637E	1	.430890915201827448022E	2
16.4	.113539450196112694867E	1	.434934224724610579176E	2
16.5	.113712083385437471149E	1	.438989883952725733639E	2
16.6	.113883934487303757604E	1	.443057855350616799848E	2
16.7	.114055011709563752564E	1	.447138101723566535133E	2
16.8	.114225323125729842284E	1	.451230586212556011564E	2
16.9	.114394876677949178491E	1	.455335272289232299519E	2
17.0	.114563680179895551988E	1	.459452123750981465328E	2
17.1	.114731741319581282310E	1	.463581104716104056009E	2
17.2	.114899067662091784149E	1	.467722179619090336357E	2
17.3	.115065666652245362729E	1	.471875313205992632919E	2
17.4	.115231545617180692321E	1	.476040470529892224834E	2
17.5	.115396711768874338320E	1	.480217616946458303916E	2
17.6	.115561172206590593651E	1	.484406718109596605457E	2
17.7	.115724933919265814487E	1	.488607739967185387581E	2
17.8	.115888003787829358156E	1	.492820648756896510221E	2
17.9	.116050388587463147527E	1	.497045411002099435428E	2
18.0	.116212094989801810896E	1	.501281993507846038860E	2

Table B-16 (Cont'd)

THE AUXILIARY FUNCTIONS G(-x) AND ψ(-x) FOR x = 10.0(0.1)25.0

x	G(-x)	ψ(-x)
18.1	.116373129565075274389E 1	.505530363356934187516E 2
18.2	.116533498784195614778E 1	.509790487906048102232E 2
18.3	.116693209020789914564E 1	.514062334781973583964E 2
18.4	.116852266553180797714E 1	.518345871877886241696E 2
18.5	.117010677566316263735E 1	.522641067349710916125E 2
18.6	.117168448153650379484E 1	.526947889612550547970E 2
18.7	.117325584318976332265E 1	.531266307337182792017E 2
18.8	.117482091978213294163E 1	.535596289446622728956E 2
18.9	.117637976961148496169E 1	.539937805112750075773E 2
19.0	.117793245013135861307E 1	.544290823752999342688E 2
19.1	.117947901796752498622E 1	.548655315027111430253E 2
19.2	.118101952893414314437E 1	.553031248833945204057E 2
19.3	.118255403804951953627E 1	.557418595308347627023E 2
19.4	.118408259955148241741E 1	.561817324818081070264E 2
19.5	.118560526691238258536E 1	.566227407960806463081E 2
19.6	.118712209285373134790E 1	.570648815561120980834E 2
19.7	.118863312936048627053E 1	.575081518667649006598E 2
19.8	.119013842769499489330E 1	.579525488550185138064E 2
19.9	.119163803841060626229E 1	.583980696696888045747E 2
20.0	.119313201136495979141E 1	.588447114811524021978E 2
20.1	.119462039573296065218E 1	.592924714810759092442E 2
20.2	.119610324001945058348E 1	.597413468821498593283E 2
20.3	.119758059207158271928E 1	.601913349178273146848E 2
20.4	.119905249909090874911E 1	.606424328420669998760E 2
20.5	.120051900764518645418E 1	.610946379290808706770E 2
20.6	.120198016367991539915E 1	.615479474730860199648E 2
20.7	.120343601252960830778E 1	.620023587880608250794E 2
20.8	.120488659892880540711E 1	.624578692075052436600E 2
20.9	.120633196702283879092E 1	.629144760842051674945E 2
21.0	.120777216037835362741E 1	.633721767900007462841E 2
21.1	.120920722199359281913E 1	.638309687155585955730E 2
21.2	.121063719430845151307E 1	.642908492701478053635E 2
21.3	.121206211921430765770E 1	.647518158814196680926E 2
21.4	.121348203806363460845E 1	.652138659951910468030E 2
21.5	.121489699167940159606E 1	.656769970752313063686E 2
21.6	.121630702036426769087E 1	.661412066030527326449E 2
21.7	.121771216390957472169E 1	.666064920777043663321E 2
21.8	.121911246160414443989E 1	.670728510155691802308E 2
21.9	.122050795224288505619E 1	.675402809501645303590E 2
22.0	.122189867413521212178E 1	.680087794319458131944E 2

Table B-16 (Cont'd)

THE AUXILIARY FUNCTIONS G(-x) AND ψ(-x) FOR x = 10.0(0.1)25.0

x	G(-x)		ψ(-x)	
22.1	.122328466511328857329E	1	.684783440281132629844E	2
22.2	.122466596254008861577E	1	.689489723274218247463E	2
22.3	.122604260331728997647E	1	.694206619149940401842E	2
22.4	.122741462389299892615E	1	.698934104221358852944E	2
22.5	.122878206026931233349E	1	.703672154761554999842E	2
22.6	.123014494800972089066E	1	.708420747251847514566E	2
22.7	.123150332246357526O5E	1	.713179858330035745801E	2
22.8	.123285721768709490114E	1	.717949464788670338268E	2
22.9	.123420666862249577438E	1	.722729543573350527162E	2
23.0	.123555170893261990400E	1	.727520071781047580235E	2
23.1	.123689237209369105480E	1	.732321026658453872577E	2
23.2	.123822869118462757055E	1	.737132385600357091755E	2
23.3	.123956069889343987372E	1	.741954126148039082848E	2
23.4	.124088842752349815723E	1	.746786225987698854543E	2
23.5	.124221190899967343997E	1	.751628662948899278993E	2
23.6	.124353117487435506674E	1	.756481415003037028876E	2
23.7	.124484625633334764613E	1	.761344460261835306057E	2
23.8	.124615718420165033525E	1	.766217776975858926595E	2
23.9	.124746398894912129780E	1	.771101343533051336906E	2
24.0	.124876670069603008336E	1	.775995138457293145838E	2
24.1	.125006534921850059827E	1	.780899140406981766927E	2
24.2	.125135996395384726477E	1	.785813328173631774538E	2
24.3	.125265057400580689279E	1	.790737680680495586509E	2
24.4	.125393720814966871910E	1	.795672176981204094891E	2
24.5	.125521989483730500125E	1	.800616796258426874860E	2
24.6	.125649866220210448814E	1	.805571517822551610373E	2
24.7	.125777353806381102586E	1	.810536321110382382998E	2
24.8	.125904454993326949609E	1	.815511185683856478720E	2
24.9	.126031172501708122497E	1	.820496091228779374756E	2
25.0	.126157509022217094241E	1	.825491017553577576259E	2

Table B-17

THE FUNCTIONS $\exp(\pm\xi)$ WITH $\xi = \frac{2}{3} x^{3/2}$

x	ξ	$\exp(-\xi)$	$\exp(+\xi)$
5.0	0.74535593E 01	0.57937576E-03	0.17259955E 04
5.5	0.85990947E 01	0.18427251E-03	0.54267451E 04
6.0	0.97979584E 01	0.55564916E-04	0.17996967E 05
6.5	0.11047874E 02	0.15920949E-04	0.62810324E 05
7.0	0.12346838E 02	0.43434658E-05	0.23023088E 06
7.5	0.13693062E 02	0.11302602E-05	0.88475205E 06
8.0	0.15084942E 02	0.28099122E-06	0.35588299E 07
8.5	0.16521028E 02	0.66835672E-07	0.14962070E 08
9.0	0.17999999E 02	0.15229988E-07	0.65659930E 08
9.5	0.19520643E 02	0.33288345E-08	0.30040544E 09
10.0	0.21081848E 02	0.69866552E-09	0.14313000E 10
10.5	0.22682589E 02	0.14095370E-09	0.70945282E 10
11.0	0.24321912E 02	0.27360721E-10	0.36548743E 11
11.5	0.25998929E 02	0.51145606E-11	0.19552021E 12
12.0	0.27712811E 02	0.92146545E-12	0.10852279E 13
12.5	0.29462780E 02	0.16013157E-12	0.62486647E 13
13.0	0.31248108E 02	0.26860816E-13	0.37228949E 14
13.5	0.33068108E 02	0.43521416E-14	0.22977193E 15
14.0	0.34922130E 02	0.68157127E-15	0.14671979E 16
14.5	0.36809564E 02	0.10323075E-15	0.96870354E 16
15.0	0.38729829E 02	0.15130341E-16	0.66092360E 17
15.5	0.40682369E 02	0.21471936E-17	0.46572418E 18
16.0	0.42666659E 02	0.29519239E-18	0.33876210E 19
16.5	0.44682205E 02	0.39333694E-19	0.25423495E 20
17.0	0.46728522E 02	0.50823000E-20	0.19676130E 21
17.5	0.48805162E 02	0.63707001E-21	0.15696862E 22
18.0	0.50911681E 02	0.77506386E-22	0.12902162E 23
18.5	0.53047666E 02	0.91556933E-23	0.10922165E 24
19.0	0.55212715E 02	0.10505636E-23	0.95186996E 24
19.5	0.57406438E 02	0.11713866E-24	0.85368905E 25
20.0	0.59628470E 02	0.12696499E-25	0.78761863E 26
20.5	0.61878453E 02	0.13382249E-26	0.74725857E 27
21.0	0.64156047E 02	0.13720888E-27	0.72881578E 28
21.5	0.66460923E 02	0.13689499E-28	0.73048689E 29
22.0	0.68792757E 02	0.13294885E-29	0.75216895E 30
22.5	0.71151241E 02	0.12572113E-30	0.79541118E 31
23.0	0.73536075E 02	0.11579449E-31	0.86359887E 32
23.5	0.75946978E 02	0.10390730E-32	0.96239626E 33
24.0	0.78383660E 02	0.90867511E-34	0.11005033E 35
24.5	0.80845866E 02	0.77461553E-35	0.12909630E 36
25.0	0.83333325E 02	0.64386735E-36	0.15531149E 37
25.5	0.85845779E 02	0.52197713E-37	0.19157927E 38
26.0	0.88382991E 02	10^{-38}	10^{+38}

Table B-18

THE FUNCTIONS $\cos(\xi + \pi/4)$ AND $\sin(\xi + \pi/4)$ WITH $\xi = \frac{2}{3} x^{3/2}$

x	ξ	$\cos(\xi + \pi/4)$	$\sin(\xi + \pi/4)$
5.0	0.74535593E 01	-0.37553661E-00	0.92680752E 00
5.5	0.85990947E 01	-0.99918865E 00	0.40274188E-01
6.0	0.97979584E 01	-0.40064228E-00	-0.91623458E 00
6.5	0.11047874E 02	0.74310495E 00	-0.66917483E 00
7.0	0.12346838E 02	0.84412497E 00	0.53614651E 00
7.5	0.13693062E 02	-0.33470652E-00	0.94232239E 00
8.0	0.15084942E 02	-0.98684579E 00	-0.16166458E-00
8.5	0.16521028E 02	0.27663564E-01	-0.99961729E 00
9.0	0.17999999E 02	0.99794253E 00	-0.64114621E-01
9.5	0.19520643E 02	0.11406205E-00	0.99347363E 00
10.0	0.21081848E 02	-0.99233399E 00	0.12358493E-00
10.5	0.22682589E 02	-0.93819257E-01	-0.99558925E 00
11.0	0.24321912E 02	0.99967665E 00	-0.25427952E-01
11.5	0.25998929E 02	-0.80702017E-01	0.99673826E 00
12.0	0.27712811E 02	-0.97504447E 00	-0.22200969E-00
12.5	0.29462780E 02	0.39222410E-00	-0.91986965E 00
13.0	0.31248108E 02	0.81528299E 00	0.57906298E 00
13.5	0.33068108E 02	-0.76225052E 00	0.64728188E 00
14.0	0.34922130E 02	-0.40847914E-00	-0.91276778E 00
14.5	0.36809564E 02	0.99458133E 00	-0.10396106E-00
15.0	0.38729829E 02	-0.24286562E-00	0.97005986E 00
15.5	0.40682369E 02	-0.80975439E 00	-0.58676922E 00
16.0	0.42666659E 02	0.86268566E 00	-0.50574021E 00
16.5	0.44682205E 02	0.85386481E-01	0.99634793E 00
17.0	0.46728522E 02	-0.92489740E 00	-0.38021707E-00
17.5	0.48805162E 02	0.78075548E 00	-0.62483643E 00
18.0	0.50911681E 02	0.13875034E-00	0.99032743E 00
18.5	0.53047666E 02	-0.91063064E 00	-0.41322157E-00
19.0	0.55212715E 02	0.85223433E 00	-0.52315996E 00
19.5	0.57406438E 02	-0.72309267E-01	0.99738223E 00
20.0	0.59628470E 02	-0.74942156E 00	-0.66209325E 00
20.5	0.61878453E 02	0.98592072E 00	-0.16721262E-00
21.0	0.64156047E 02	-0.51310262E 00	0.85832710E 00
21.5	0.66460923E 02	-0.29353403E-00	-0.95594871E 00
22.0	0.68792757E 02	0.89466443E 00	0.44673906E-00
22.5	0.71151241E 02	-0.94923777E 00	0.31455915E-00
23.0	0.73536075E 02	0.47419146E-00	-0.88042159E 00
23.5	0.75946978E 02	0.23444170E-00	0.97213025E 00
24.0	0.78383660E 02	-0.80847432E 00	-0.58853170E 00
24.5	0.80845866E 02	0.99874297E 00	-0.50124244E-01
25.0	0.83333325E 02	-0.76207855E 00	0.64748432E 00

Table B-19

THE COEFFICIENTS $A_n(x)$ IN THE CLASSICAL ASYMPTOTIC EXPANSIONS

n	$A_n(4.0)$	$A_n(5.0)$	$A_n(6.0)$
1	0.01302 08333 33333	0.00931 69499 06249 123	0.00708 76439 31664 288
2	0.00130 54741 75347	0.00066 84027 77777 778	0.00038 68071 63065 844
3	0.00025 04425 28430	0.00009 17514 67913 566	0.00004 03921 91245 878
4	0.00007 12522 03727	0.00001 86783 37693 705	0.00000 62553 37501 375
5	0.00002 69051 29011	0.00000 50467 28962 103	0.00000 12857 32541 582
6	0.00001 26701 67091	0.00000 17005 61040 314	0.00000 03295 80122 467
7	0.00000 71505 37008	0.00000 06867 26050 060	0.00000 01012 46597 827
8	0.00000 47041 78155	0.00000 03232 68661 295	0.00000 00362 56731 648
9	0.00000 35349 39430	0.00000 01738 18821 499	0.00000 00148 30301 469
10	0.00000 29872 07930	0.00000 01051 03035 298	0.00000 00068 21762 064
11	0.00000 28040 43428	0.00000 00705 94281 263	0.00000 00034 85611 350
12	0.00000 28947 12367	0.00000 00521 46502 152	0.00000 00019 58676 684
13	0.00000 32594 50764	0.00000 00420 14497 292	0.00000 00012 00507 489
14	0.00000 39754 87102	0.00000 00366 67396 572	0.00000 00007 97028 606
15	0.00000 52212 77765	0.00000 00344 58907 563	0.00000 00005 69801 344
16	0.00000 73466 70944	0.00000 00346 93672 628	0.00000 00004 36415 705
17	0.00001 10256 33462	0.00000 00372 56171 056	0.00000 00003 56513 882
18	0.00001 75800 79049	0.00000 00425 06022 946	0.00000 00003 09426 018
19	0.00002 96784 31147	0.00000 00513 45876 150	0.00000 00002 84341 373
20	0.00005 28840 27375	0.00000 00654 67254 927	0.00000 00002 75795 229
21	0.00009 91903 41525	0.00000 00878 62584 873	0.00000 00002 81575 410
22	0.00019 53396 91281	0.00000 01238 11238 451	0.00000 00003 01841 498
23	0.00040 29986 99596	0.00000 01827 71382 114	0.00000 00003 38965 273
24	0.00086 91845 86792	0.00000 02820 66524 968	0.00000 00003 97948 334
25	0.00195 61180 20588	0.00000 04542 23465 271	0.00000 00004 87498 001
26	0.00458 56312 37166	0.00000 07619 18677 348	0.00000 00006 22071 098
27	0.01117 96875 75410	0.00000 13291 50442 553	0.00000 00008 25531 549
28	0.02830 37830 62708	0.00000 24078 19573 349	0.00000 00011 37657 689
29	0.07431 01387 75537	0.00000 45233 74094 742	0.00000 00016 25841 869
30	0.20206 29424 60391	0.00000 88010 82409 911	0.00000 00024 06467 581

Table B-19 (Cont'd)

THE COEFFICIENTS $A_n(x)$ IN THE CLASSICAL ASYMPTOTIC EXPANSIONS

n	$A_n(7.0)$	$A_n(8.0)$
1	0.00562 44713 24542 071	0.00460 35597 73349 918
2	0.00024 35870 18140 590	0.00016 31842 71918 403
3	0.00002 01853 76792 154	0.00001 10681 00634 412
4	0.00000 24806 75793 795	0.00000 11133 15683 228
5	0.00000 04046 22206 163	0.00000 01486 31243 536
6	0.00000 00823 07593 516	0.00000 00247 46420 099
7	0.00000 00200 64999 234	0.00000 00049 37688 679
8	0.00000 00057 01999 523	0.00000 00011 48480 995
9	0.00000 00018 50837 624	0.00000 00003 05124 468
10	0.00000 00006 75607 906	0.00000 00000 91162 351
11	0.00000 00002 73940 934	0.00000 00000 30254 488
12	0.00000 00001 22157 614	0.00000 00000 11042 451
13	0.00000 00000 59415 872	0.00000 00000 04396 018
14	0.00000 00000 31303 387	0.00000 00000 01895 660
15	0.00000 00000 17759 090	0.00000 00000 00880 242
16	0.00000 00000 10793 876	0.00000 00000 00437 896
17	0.00000 00000 06997 345	0.00000 00000 00232 348
18	0.00000 00000 04819 406	0.00000 00000 00130 982
19	0.00000 00000 03514 443	0.00000 00000 00078 178
20	0.00000 00000 02705 098	0.00000 00000 00049 252
21	0.00000 00000 02191 648	0.00000 00000 00032 661
22	0.00000 00000 01864 382	0.00000 00000 00022 741
23	0.00000 00000 01661 464	0.00000 00000 00016 587
24	0.00000 00000 01547 898	0.00000 00000 00012 648
25	0.00000 00000 01504 763	0.00000 00000 00010 064
26	0.00000 00000 01523 755	0.00000 00000 00008 341
27	0.00000 00000 01604 681	0.00000 00000 00007 190
28	0.00000 00000 01754 876	0.00000 00000 00006 436
29	0.00000 00000 01990 183	0.00000 00000 00005 974
30	0.00000 00000 02337 623	0.00000 00000 00005 743

Table B-20

THE COEFFICIENTS $A_n(x)$ IN THE CLASSICAL ASYMPTOTIC EXPANSIONS

n	$A_n(1.0)$	n	$A_n(3.0)$
1	.10416666666666666+00	1	.2004688434686201-01
2	.8355034722222217-01	2	.3094457304526749-02
3	.1282265745563271+00	3	.9139709547822118-03
4	.2918490264641404+00	4	.4003416000879845-03
5	.8816272674437575+00	5	.2327424509298436-03
6	.3321408281862767+01	6	.1687450227029808-03
7	.1499576298686259+02	7	.1466207352735687-03
8	.7892301301158699+02	8	.1485075728285682-03
9	.4744515388679998+03	9	.1718125647581276-03
10	.3207490090999999+04	10	.2235354993240953-03
11	.2408654959999999+05	11	.3230530821478668-03
12	.1989231199999999+06	12	.5134553428329405-03
13	.1791901999999999+07	13	.8901225166930805-03
14	.1748437699999999+08	14	.1671490117081857-02

n	$A_n(2.0)$	n	$A_n(4.0)$
1	.3682847818679937-01	1	.1302083333333333-01
2	.1044379340277778-01	2	.1305474175347221-02
3	.5666867524818843-02	3	.2504425284303265-03
4	.4560141038502203-02	4	.7125220372659678-04
5	.4870348588191003-02	5	.2690512901134514-04
6	.6487125550513236-02	6	.1267016709084612-04
7	.1035508368855157-01	7	.7150537007743164-05
8	.1926831372353205-01	8	.4704178155159174-05
9	.4095311285131311-01	9	.3534939429670571-05
10	.9788482943725630-01	10	.2987207929603755-05
11	.2598840722281831+00	11	.2804043423384427-05
12	.7588314819335980+00	12	.2894712379202246-05
13	.2416736708491792+01	13	.3259450750192627-05
14	.8337200641632135+01	14	.3975487061325111-05

Table B-20 (Cont'd)

n	$A_n(5.0)$
1	.9316949906249129-02
2	.6684027777777783-03
3	.9175146791356576-04
4	.1867833769370503-04
5	.5046728962103385-05
6	.1700561040313743-05
7	.6867260500602920-06
8	.3232686612954618-06
9	.1738188214990007-06
10	.1051030353018885-06
11	.7059428114303642-07
12	.5214650236928036-07
13	.4201449710962982-07
14	.3666739619430430-07

n	$A_n(7.0)$
1	.5624471324542075-02
2	.2435870181405896-03
3	.2018537679215353-04
4	.2480675793794599-05
5	.4046222061628177-06
6	.8230759351605851-07
7	.2006499923361713-07
8	.5701999522550561-08
9	.1850837624446468-08
10	.6756079061392759-09
11	.2739409335901954-09
12	.1221576141683263-09
13	.5941587181038829-10
14	.3130338639011568-10

n	$A_n(6.0)$
1	.7087643931664294-02
2	.3868071630658438-03
3	.4039219124587786-04
4	.6255337501374764-05
5	.1285732541581796-05
6	.3295801224667600-06
7	.1012465978266584-06
8	.3625673164759974-07
9	.1483030146893619-07
10	.6821762064333982-08
11	.3485611344451478-08
12	.1958676692325371-08
13	.1200507483750372-08
14	.7970285973939246-09

n	$A_n(8.0)$
1	.4603559773349922-02
2	.1631842719184028-03
3	.1106810063441180-04
4	.1113315683228076-05
5	.1486312435361024-06
6	.2474642009930891-07
7	.4937688679004468-08
8	.1148480994911912-08
9	.3051244679936253-09
10	.9116235136730252-10
11	.3025448790613831-10
12	.1104245139771371-10
13	.4396018463906861-11
14	.1895659952795571-11

Table B-20 (Cont'd)

n	$A_n(9.0)$	n	$A_n(11.0)$
1	.3858024691358024-02	1	.2855221066077309-02
2	.1146095297972869-03	2	.6277261248852160-04
3	.6514584898456896-05	3	.2640650700435702-05
4	.5491654322194565-06	4	.1647411669505825-06
5	.6144212011714596-07	5	.1364080105949640-07
6	.8573135330131605-08	6	.1408601341980650-08
7	.1433580940896128-08	7	.1743192599377849-09
8	.2794431984520721-09	8	.2514730417204718-10
9	.6221827729592102-10	9	.4143721988720650-11
10	.1557857328952611-10	10	.7678476262157784-12
11	.4332845327303985-11	11	.1580502862083735-12
12	.1325317987590837-11	12	.3577807164816551-13
13	.4421659748825643-12	13	.8833989802828937-14
14	.1597928946225705-12	14	.2362673846383553-14

n	$A_n(10.0)$	n	$A_n(12.0)$
1	.3294039229342062-02	1	.2505860543357751-02
2	.8355034722222220-04	2	.4835089538323045-04
3	.4054880321593885-05	3	.1785099521059007-05
4	.2918490264641404-06	4	.9773964845898059-07
5	.2787950212432688-07	5	.7102735929255482-08
6	.3321408281862769-08	6	.6437111766928895-09
7	.4742076629053547-09	7	.6991421474150121-10
8	.7892301301158706-10	8	.8851741124902261-11
9	.1500347502194787-10	9	.1280103361294624-11
10	.3207490091000004-11	10	.2081836567484729-12
11	.7616835771061771-12	11	.3760832852542712-13
12	.1989231200000002-12	12	.7471758622456988-14
13	.5666491663811048-13	13	.1619123425722283-14
14	.1748437700000002-13	14	.3800528513879401-15

Table B-20 (Cont'd)

n	$A_n(13.0)$
1	.2222356555389540-02
2	.3802928867647803-04
3	.1245182347741390-05
4	.6046417549651142-07
5	.3896815746645977-08
6	.3132074570459569-09
7	.3016917042901795-10
8	.3387532176858264-11
9	.4344668082341731-12
10	.6266355634582820-13
11	.1003943514868797-13
12	.1768906675025272-14
13	.3399530048116087-15
14	.7076846306335666-16

n	$A_n(15.0)$
1	.1793047845466397-02
2	.2475565843621399-04
3	.6539843789930920-06
4	.2562186240563100-07
5	.1332295530314341-08
6	.8639745162392619-10
7	.6714448506525714-11
8	.6082870183058153-12
9	.6294480063492231-13
10	.7324811241851952-14
11	.9468230395801812-15
12	.1345992373915983-15
13	.2087058885185549-16
14	.3505368842018450-17

n	$A_n(14.0)$
1	.1988550907086491-02
2	.3044837726757369-04
3	.8920760506585824-06
4	.3876055927804059-07
5	.2235243014034292-08
6	.1607570185860516-09
7	.1385556349862650-10
8	.1392089727185193-11
9	.1597582806544551-12
10	.2061791705747298-13
11	.2955711239413460-14
12	.4659943167431874-15
13	.8013413594672413-16
14	.1492661780839711-16

n	$A_n(16.0)$
1	.1627604166666666-02
2	.2039803398980033-04
3	.4891455633404814-06
4	.1739555755043866-07
5	.8210793765669292-09
6	.4833285175646258-10
7	.3409641746398528-11
8	.2803908678984149-12
9	.2633735112600454-13
10	.2782054179910341-14
11	.3264336175499981-15
12	.4212360915265520-16
13	.5928906375979980-17
14	.9039211048105041-18

Table B-20 (Cont'd)

n	$A_n(17.0)$
1	.1486125153408904-02
2	.1700597338127871-04
3	.3723556040551098-06
4	.1209106958799956-07
5	.5210964367439643-09
6	.2800799187888995-10
7	.1804076266354483-11
8	.1354617245418725-12
9	.1161800130671895-13
10	.1120550677967813-14
11	.1200514442213222-15
12	.1414505620569984-16
13	.1817859387797468-17
14	.2530595976950671-18

n	$A_n(19.0)$
1	.1257761698851765-02
2	.1218112658145827-04
3	.2257287290885441-06
4	.6203497952650543-08
5	.2262731415887266-09
6	.1029294982338471-10
7	.5611200117439291-12
8	.3565828447263459-13
9	.2588321660295779-14
10	.2112814209833057-15
11	.1915755918869904-16
12	.1910383001496502-17
13	.2077873171033199-18
14	.2448074004256393-19

n	$A_n(18.0)$
1	.1364017710622199-02
2	.1432619122466088-04
3	.2879066973946474-06
4	.8580709878429038-08
5	.3394229670727538-09
6	.1674440494166337-10
7	.9899363327025878-12
8	.6822343712208841-13
9	.5370479222374401-14
10	.4754203274391557-15
11	.4674963856154057-16
12	.5055686903346446-17
13	.5963488755215175-18
14	.7619518977287890-19

n	$A_n(20.0)$
1	.1164618738281141-02
2	.1044379340277778-04
3	.1792020857686831-06
4	.4560141038502205-08
5	.1540139453766902-09
6	.6487125550513241-11
7	.3274564981748065-12
8	.1926831372353206-13
9	.1295051138840621-14
10	.9788482943725640-16
11	.8218255958407700-17
12	.7588314819335988-18
13	.7642392503772465-19
14	.8337200641632148-20

Appendix C

TABLES TO FACILITATE HAND COMPUTATION OF THE ROOT t_1 FOR THE CASE OF A REAL EARTH

In this section we present some tables which permit one to rather easily compute (with a desk calculator) the numerical value of the first root t_1 of the equation

$$w_1'(t_1) - q\, w_1(t_1) = 0 \qquad \text{(C-1)}$$

when the impedance parameter q has a phase which is approximately 45°. This case occurs very frequently in the case of the propagation of radio waves around the earth's surface.

Let us first review the theoretical solution. We consider a sphere of radius a. We assume that the time dependance is of the form $\exp(-i\omega t)$ and that the propagation constants are of the form

$$k^2 = \omega^2 \varepsilon_o \mu_o \quad , \quad r > a \qquad \text{(C-2a)}$$

$$k_1^2 = \omega^2 \varepsilon_1 \mu_o + i\omega \mu_o \sigma \quad , \quad r < a \qquad \text{(C-2b)}$$

For the propagation of vertically polarized waves, the theory leads to an expression for the impedance parameter q which is of the form

$$q = i(ka/2)^{\frac{1}{3}}(k/k_1)\sqrt{1 - (k/k_1)^2} \qquad \text{(C-3a)}$$

Therefore, if $\omega\varepsilon_1 \ll \sigma$, and $\sigma \to \infty$, we see that

$$q \approx \sqrt{i}\,(ka/2)^{\frac{1}{3}}\sqrt{\omega\varepsilon_o/\sigma} \qquad \text{(C-3b)}$$

Eq. (C-3b) shows us that in the limiting case $\arg(q) = 45^\circ$; but if we retain the full form given in Eq. (C-3a) we find that the actual value of $\arg(q)$ is approximately 45°.

The theory of the propagation of waves around a convex surface has been developed (See Section 1) in terms of the diffraction function

$$D(x,y,y_o,q) = \frac{1}{2\pi}\int_{-\infty}^{\infty} \exp(ixt)\left|w_1(t-y_>)v(t-y_<) - \frac{v'(t)-qv(t)}{w_1'(t)-qw_1(t)}\, w_1(t-y_o)w_1(t-y)\right| dt$$

$$= -i\sum_{s=1}^{\infty} \frac{\exp(ixt_s)}{t_s-q^2}\,\frac{w_1(t_s-y)}{w_1(t_s)}\,\frac{w_1(t_s-y_o)}{w_1(t_s)} \qquad (C\text{-}4)$$

Fig. 2-2 reveals that the root t_1 has the smallest imaginary part when $30^\circ < \arg(q) < 210^\circ$. Therefore, when $x > \sqrt{y} + \sqrt{y_o}$ and $x \to \infty$, we find that we need only retain the first term in Eq. (C-4). Since the factor $\exp(ixt_1)$ then determines the manner in which the wave continues to propagate (when y and y_o are fixed), the ability to determine the numerical value of t_1 <u>accurately</u> will be required in many problems which involve the propagation of the ground wave to great distances.

Let us assume that we know the value of t_1 for some value q_o of the impedance parameter, and let us denote this value by $t_1(q_o)$. If we want to know the value of t_1 for some neighboring value of q, we can seek to express $t_1(q)$ in the form

$$t_1(q) \approx t_1(q_o) + A_1(q-q_c) + A_2(q-q_o)^2 + A_3(q-q_o)^3 + A_4(q-q_o)^4 + \ldots + A_N(q-q_o)^N \qquad (C\text{-}5)$$

We observe that the coefficients A_n will be functions of q_o. The coefficients can be determined by observing that t(q) satisfies the Ricatti differential equation

$$[t(q) - q^2]\frac{dt(q)}{dq} = 1 \qquad (C\text{-}6)$$

If Eq. (C-5) is inserted in Eq. (C-6), we find that the first several values of A_n are given by

$$A_1 = (t_o - q_o^2)^{-1} \tag{C-7a}$$

$$A_2 = -(1/2)[1 - 2q_o(t_o - q_o^2)](t_o - q_o^2)^{-3} \tag{C-7b}$$

and further terms can be obtained by means of the recurrence relation

$$nA_n(t_o - q_o^2) + (n-1)A_{n-1}(A_1 - 2q_o) + (n-2)A_{n-2}(A_2 - 1) + (n-3)A_{n-3}A_3 + \cdots + 2A_2A_{n-2} + A_1A_{n-1} = 0 \tag{C-8}$$

In Section 2 we showed that the roots for which the phase of q is zero (i.e., $q = Q$, where Q is real and positive) arise in a class of problems which are of practical importance. Therefore, in the tables that follow, we give the values of

$$t_o = t_1(q_o)$$

for the cases in which $q_o = Q\exp(i145^\circ)$ and $q_o = Q$. Table C-1 contains the data for the case $q_o = Q\exp(i145^\circ)$ for $Q = 0.00(0.05)4.00$, i.e., $Q = 0.00, 0.05, 0.10, 0.15, \ldots, 3.95$, and 4.00. The data for the case $q_o = Q$ is given in Table C-2 for $Q = 0.00(0.05)2.00$.

Table C-1

VALUES OF q_o, t_o, AND A_n

$q_o = 0.00\exp(i145^\circ)$

$t_o = 0.50939649 + i0.88230059$

n	Real A_n	Imag A_n
1	0.49077678E-00	-0.85005043E 00
2	-0.52716219E 00	0.
3	0.67202630E-01	0.11639834E-00
4	-0.35801253E-02	0.62010126E-02
5	0.98774254E-02	-0.23255247E-07
6	0.10064451E-02	0.17432125E-02
7	0.67314184E-03	-0.11658314E-02
8	0.56667980E-03	0.82246385E-07

$q_o = 0.05\exp(i145^\circ)$

$t_o = 0.55678567 + i0.87077604$

n	Real A_n	Imag A_n
1	0.45262659E-00	-0.88682587E 00
2	-0.53248200E 00	0.19428691E-01
3	0.65814348E-01	0.11701491E-00
4	-0.18961618E-02	0.79905694E-02
5	0.97795711E-02	0.62152158E-03
6	0.14634134E-02	0.16615978E-02
7	0.84904945E-03	-0.10119360E-02
8	0.60454502E-03	-0.74738907E-04

$q_o = 0.10\exp(i145^\circ)$

$t_o = 0.60407773 + i0.86158898$

n	Real A_n	Imag A_n
1	0.41272052E-00	-0.92261416E 00
2	-0.53804180E 00	0.38800108E-01
3	0.64594664E-01	0.11812087E-00
4	-0.34130665E-03	0.98899703E-02
5	0.97879910E-02	0.13325536E-02
6	0.19306374E-02	0.16650086E-02
7	0.10569446E-02	-0.86151452E-03
8	0.67524043E-03	-0.14223865E-03

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$$q_o = 0.15\exp(i145^\circ)$$

$$t_o = 0.65117579 + i0.85471159$$

n	Real A_n	Imag A_n
1	0.37104204E-00	-0.95743659E 00
2	-0.54389867E 00	0.58160997E-01
3	0.62907858E-01	0.11971661E-00
4	0.10900478E-02	0.11934237E-01
5	0.98866076E-02	0.21550881E-02
6	0.24161400E-02	0.17634061E-02
7	0.13074082E-02	-0.69951822E-03
8	0.78248135E-03	-0.16964135E-03

$$q_o = 0.20\exp(i145^\circ)$$

$$t_o = 0.69798303 + i0.85011455$$

n	Real A_n	Imag A_n
1	0.32756723E-00	-0.99131552E 00
2	-0.55011404E 00	0.77554256E-01
3	0.61312683E-01	0.12180702E-00
4	0.23884770E-02	0.14159878E-01
5	0.10058614E-01	0.31152634E-02
6	0.24251252E-02	0.19718236E-02
7	0.15990656E-02	-0.50853314E-03
8	0.93126450E-03	-0.18103705E-03

$$q_o = 0.25\exp(i145^\circ)$$

$$t_o = 0.74440248 + i0.84776678$$

n	Real A_n	Imag A_n
1	0.28226469E-00	-0.10242750E 01
2	-0.55675475E 00	0.97018810E-01
3	0.59561034E-01	0.12440058E-00
4	0.35361777E-02	0.16605039E-01
5	0.10283797E-01	0.42436861E-02
6	0.34591538E-02	0.23111593E-02
7	0.19377197E-02	-0.26731351E-03
8	0.11271954E-02	-0.15615424E-03

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.30\exp(i145^\circ)$

$t_o = 0.79033678 + i0.84763502$

n	Real A_n	Imag A_n
1	0.23509548E-00	-0.10563412E 01
2	-0.56369424E 00	0.11658908E-00
3	0.57596387E-01	0.12750845E-00
4	0.45135615E-02	0.19309778E-01
5	0.10536160E-01	0.55760220E-02
6	0.40143593E-02	0.28089224E-02
7	0.23231572E-02	0.50682307E-04
8	0.13759191E-02	-0.78781296E-04

$q_o = 0.35\exp(i145^\circ)$

$t_o = 0.83568808 + i0.84968355$

n	Real A_n	Imag A_n
1	0.18601296E-00	-0.10875440E 01
2	-0.57161380E 00	0.13629427E-00
3	0.55351876E-01	0.13114310E-00
4	0.52770384E-02	0.22315605E-01
5	0.10782291E-01	0.71528966E-02
6	0.45797289E-02	0.35000101E-02
7	0.27589741E-02	0.47896879E-03
8	0.16813800E-02	0.74375224E-04

$q_o = 0.40\exp(i145^\circ)$

$t_o = 0.88035785 + i0.85387373$

n	Real A_n	Imag A_n
1	0.13496263E-00	-0.11179169E 01
2	-0.58000366E 00	0.15615704E-00
3	0.52748542E-01	0.13531664E-00
4	0.57765931E-02	0.25664920E-01
5	0.10976752E-01	0.90198285E-02
6	0.51334415E-02	0.44271453E-02
7	0.32294668E-02	0.10594568E-02
8	0.20436674E-02	0.33615415E-03

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.45\exp(i145^\circ)$

$t_o = 0.92424677 + i0.86016357$

n	Real A_n	Imag A_n
1	0.81881676E-01	-0.11474991E 01
2	-0.58916447E 00	0.17619231E-00
3	0.49692807E-01	0.14003857E-00
4	0.57422909E-02	0.29398875E-01
5	0.11058241E-01	0.11225963E-01
6	0.56383629E-02	0.56409825E-02
7	0.37150677E-02	0.18441891E-02
8	0.24541350E-02	0.75138412E-03

$q_o = 0.50\exp(i145^\circ)$

$t_o = 0.96725465 + i0.86850721$

n	Real A_n	Imag A_n
1	0.26679860E-01	-0.11763366E 01
2	-0.59920627E 00	0.19640484E-00
3	0.46074087E-01	0.14551258E-00
4	0.56827610E-02	0.33555016E-01
5	0.10943777E-01	0.13621948E-01
6	0.56353474E-02	0.71987110E-02
7	0.41752204E-02	0.26950638E-02
8	0.28593176E-02	0.13779397E-02

$q_o = 0.55\exp(i145^\circ)$

$t_o = 1.0092804 + i0.87885439$

n	Real A_n	Imag A_n
1	-0.36002451E-01	-0.12044838E 01
2	-0.61025987E 00	0.21678644E-00
3	0.41702462E-01	0.15113252E-00
4	0.48808205E-02	0.36162646E-01
5	0.10522406E-01	0.16855521E-01
6	0.62365749E-02	0.91607014E-02
7	0.43412576E-02	0.42827047E-02
8	0.33004466E-02	0.22875719E-02

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.60\exp(i145^\circ)$

$t_o = 1.0502221 + i\ 0.89114979$

n	Real A_n	Imag A_n
1	-0.90292547E-01	-0.12320057E 01
2	-0.62245751E 00	0.23731278E-00
3	0.36606105E-01	0.15747670E-00
4	0.33889300E-02	0.43235994E-01
5	0.96477230E-02	0.20364205E-01
6	0.61130758E-02	0.11583455E-01
7	0.47035185E-02	0.60801339E-02
8	0.35994176E-02	0.35616199E-02

$q_o = 0.65\exp(i145^\circ)$

$t_o = 1.0899773 + i0.90533239$

n	Real A_n	Imag A_n
1	-0.15228532E-00	-0.12589797E 01
2	-0.63595340E 00	0.25793859E-00
3	0.30429444E-01	0.16430058E-00
4	0.10251596E-02	0.48764682E-01
5	0.81305329E-02	0.24363540E-01
6	0.54857755E-02	0.14507200E-01
7	0.44963928E-02	0.83514688E-02
8	0.36369886E-02	0.52810881E-02

$q_o = 0.70\exp(i145^\circ)$

$t_o = 1.1284428 + i0.92133459$

n	Real A_n	Imag A_n
1	-0.21674313E-00	-0.12854989E 01
2	-0.65091322E 00	0.27859244E-00
3	0.23032676E-01	0.17152733E-00
4	-0.24298231E-02	0.54700182E-01
5	0.57319522E-02	0.28829713E-01
6	0.41132031E-02	0.17934771E-01
7	0.36816528E-02	0.11128113E-01
8	0.31917967E-02	0.75018863E-02

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.75\exp(i145^\circ)$

$t_o = 1.1655153 + i0.93908155$

n	Real A_n	Imag A_n
1	-0.28377413E-00	-0.13116743E 01
2	-0.66751487E 00	0.29917005E-00
3	0.14192763E-01	0.17903575E-00
4	-0.72326457E-02	0.60937604E-01
5	0.21608219E-02	0.33674400E-01
6	0.16846272E-02	0.21799766E-01
7	0.19352160E-02	0.14370657E-01
8	0.17286288E-02	0.10211703E-01

$q_o = 0.80\exp(i145^\circ)$

$t_o = 1.2010921 + i0.95849018$

n	Real A_n	Imag A_n
1	-0.35349092E-00	-0.13576389E 01
2	-0.68594492E 00	0.31952658E-00
3	0.30683625E-02	0.18664596E-00
4	-0.13672118E-01	0.67292419E-01
5	-0.29219291E-02	0.38710555E-01
6	-0.21777098E-02	0.25920190E-01
7	-0.11544094E-02	0.17910162E-01
8	-0.56899035E-03	0.13257854E-01

$q_o = 0.85\exp(i145^\circ)$

$t_o = 1.2350714 + i0.97946823$

n	Real A_n	Imag A_n
1	-0.42600704E-00	-0.13635510E 01
2	-0.70639335E 00	0.33946810E-00
3	-0.87910885E-02	0.19410221E-00
4	-0.22055050E-01	0.73472234E-01
5	-0.98845738E-02	0.43609564E-01
6	-0.78935507E-02	0.29937501E-01
7	-0.60601790E-02	0.21465411E-01
8	-0.48798396E-02	0.16240927E-01

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$$q_o = 0.90\exp(i145^\circ)$$

$$t_o = 1.2673537 + i1.0019134$$

n	Real A_n	Imag A_n
1	-0.50143421E 00	-0.13895977E 01
2	-0.72904442E 00	0.35874235E-00
3	-0.23428267E-01	0.20105492E-00
4	-0.32678955E-01	0.79045778E-01
5	-0.19084015E-01	0.47852303E-01
6	-0.15877599E-01	0.33244967E-01
7	-0.13259773E-01	0.24044226E-01
8	-0.11446172E-01	0.18384216E-01

$$q_o = 0.95\exp(i145^\circ)$$

$$t_o = 1.2978426 + i1.0257123$$

n	Real A_n	Imag A_n
1	-0.57967604E 00	-0.14159986E 01
2	-0.75406393E 00	0.37702996E-00
3	-0.40452259E-01	0.20704291E-00
4	-0.45786110E-01	0.83412436E-01
5	-0.30791914E-01	0.50682299E-01
6	-0.26409216E-01	0.34919862E-01
7	-0.22978704E-01	0.24850582E-01
8	-0.20557209E-01	0.18410957E-01

$$q_o = 1.00\exp(i145^\circ)$$

$$t_o = 1.3264471 + i1.0507400$$

n	Real A_n	Imag A_n
1	-0.66142152E 00	-0.14430096E 01
2	-0.78158081E 00	0.39393754E-00
3	-0.60000297E-01	0.21148045E-00
4	-0.61494213E-01	0.85782430E-01
5	-0.45079466E-01	0.51078147E-01
6	-0.39472765E-01	0.33690391E-01
7	-0.35126768E-01	0.22249578E-01
8	-0.31879149E-01	0.14515670E-01

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.05\exp(i145^o)$

$t_o = 1.3530825 + i1.0768585$

n	Real A_n	Imag A_n
1	-0.74613634E 00	-0.14709239E 01
2	-0.81166378E 00	0.40899394E-00
3	-0.82086617E-01	0.21365441E-00
4	-0.79701199E-01	0.85179453E-01
5	-0.61657120E-01	0.47772381E-01
6	-0.54497832E-01	0.27987981E-01
7	-0.48729718E-01	0.14381056E-01
8	-0.43959461E-01	0.45776843E-02

$q_o = 1.10\exp(i145^o)$

$t_o = 1.3776740 + i1.1039177$

n	Real A_n	Imag A_n
1	-0.83405215E 00	-0.15000723E 01
2	-0.84429192E 00	0.42165411E-00
3	-0.10654067E-00	0.21273929E-00
4	-0.99968848E-01	0.80485356E-01
5	-0.79682529E-01	0.39355525E-01
6	-0.70071500E-01	0.16157454E-01
7	-0.61626483E-01	-0.55768762E-03
8	-0.53708897E-01	-0.13193293E-01

$q_o = 1.15\exp(i145^o)$

$t_o = 1.4001585 + i1.1317546$

n	Real A_n	Imag A_n
1	-0.92515410E 00	-0.15308213E 01
2	-0.87932145E 00	0.43131225E-00
3	-0.13293539E-00	0.20784044E-00
4	-0.12140156E-00	0.70547698E-01
5	-0.97578061E-01	0.24505138E-01
6	-0.83701950E-01	-0.31087198E-02
7	-0.70198639E-01	-0.23558985E-01
8	-0.56226173E-01	-0.39048227E-01

Table C-1 (Cont'd)

VALUES OF t_o, q_o, AND A_n

$q_o = 1.20\exp(145^\circ)$

$t_o = 1.4204878 + i1.1601938$

n	Real A_n	Imag A_n
1	-0.10193676E 01	-0.15635658E 01
2	-0.91645177E 00	0.43732899E-00
3	-0.16052481E-00	0.19807306E-00
4	-0.14255226E-00	0.54366705E-01
5	-0.11293600E-00	0.23640746E-02
6	-0.91802556E-01	-0.30091969E-01
7	-0.69592178E-01	-0.53707222E-01
8	-0.45527863E-01	-0.70082074E-01

$q_o = 1.25\exp(145^\circ)$

$t_o = 1.4386325 + i1.1890502$

n	Real A_n	Imag A_n
1	-0.11165450E 01	-0.15987185E 01
2	-0.95519596E 00	0.43907478E-00
3	-0.18820219E-00	0.18268245E-00
4	-0.16140678E-00	0.31361737E-01
5	-0.12262517E-00	-0.26959103E-01
6	-0.90114665E-01	-0.63235075E-01
7	-0.54779880E-01	-0.87042430E-01
8	-0.16715474E-01	-0.98916637E-01

$q_o = 1.30\exp(145^\circ)$

$t_o = 1.4545836 + i1.2181302$

n	Real A_n	Imag A_n
1	-0.12164553E 01	-0.16366923E 01
2	-0.99486415E 00	0.43599030E-00
3	-0.21450788E-00	0.16119861E-00
4	-0.17550460E-00	0.16844275E-02
5	-0.12321142E-00	-0.61880367E-01
6	-0.74737786E-01	-0.98498941E-01
7	-0.22621813E-01	-0.11606693E-00
8	0.30495062E-01	-0.11402132E-00

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.35\exp(i145^o)$

$t_o = 1.4683571 + i1.2472358$

n	Real A_n	Imag A_n
1	-0.13187756E 01	-0.16778798E 01
2	-0.10345690E 01	0.42765906E-00
3	-0.23770989E-00	0.13360475E-00
4	-0.18224201E-00	-0.33499251E-01
5	-0.11173595E-00	-0.99034086E-01
6	-0.45705740E-01	-0.12936444E-00
7	0.25519520E-01	-0.13078818E-00
8	0.88272503E-01	-0.10302761E-00

$q_o = 1.40\exp(i145^o)$

$t_o = 1.4799945 + i1.2761691$

n	Real A_n	Imag A_n
1	-0.14230932E 01	-0.17226270E 01
2	-0.10732608E 01	0.41388478E-00
3	-0.25597150E-00	0.10048096E-00
4	-0.17935414E-00	-0.71871694E-01
5	-0.86737260E-01	-0.13343360E-00
6	0.13580442E-02	-0.14794017E-00
7	0.81885668E-01	-0.12182394E-00
8	0.13995598E-00	-0.59056594E-01

$q_o = 1.45\exp(i145^o)$

$t_o = 1.4895648 + i1.3047369$

n	Real A_n	Imag A_n
1	-0.15289126E 01	-0.17712054E 01
2	-0.11097967E 01	0.39475746E-00
3	-0.26759717E-00	0.63073153E-01
4	-0.16549239E-00	-0.11009146E-00
5	-0.49195597E-01	-0.15928116E-00
6	0.54357100E-01	-0.14723893E-00
7	0.13288063E-00	-0.84913188E-01
8	0.16474293E-00	0.12669551E-01

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.50\exp(i145^o)$

$t_o = 1.4971633 + i1.3327571$

n	Real A_n	Imag A_n
1	-0.16356749E 01	-0.18237878E 01
2	-0.11430400E 01	0.37069287E-00
3	-0.27131661E-00	0.23242419E-01
4	-0.14073069E-00	-0.14424573E-00
5	-0.29466183E-02	-0.17136026E-00
6	0.10547882E-00	-0.12397455E-00
7	0.16306426E-00	-0.24660904E-01
8	0.14741138E-00	0.92863109E-01

$q_o = 1.55\exp(i145^o)$

$t_o = 1.5029100 + i1.3600639$

n	Real A_n	Imag A_n
1	-0.17427856E 01	-0.18804292E 01
2	-0.11719757E 01	0.34242941E-00
3	-0.26653893E-00	-0.16725001E-01
4	-0.10679856E-00	-0.17056987E-00
5	0.45805302E-01	-0.16660023E-00
6	0.14381885E-00	-0.80574372E-01
7	0.16147916E-00	0.45217568E-01
8	0.88044179E-01	0.15450577E-00

$q_o = 1.60\exp(i145^o)$

$t_o = 1.5069460 + i1.3865130$

n	Real A_n	Imag A_n
1	-0.18496485E 01	-0.19410583E 01
2	-0.11958211E 01	0.31097712E-00
3	-0.25350136E-00	-0.54421844E-01
4	-0.66899695E-01	-0.18624154E-00
5	0.89887080E-01	-0.14517745E-00
6	0.16101225E-00	-0.25134958E-01
7	0.12704954E-00	0.10636009E-00
8	0.46215098E-02	0.17554051E-00

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.65\exp(i145^\circ)$

$t_o = 1.5094280 + i1.4119853$

n	Real A_n	Imag A_n
1	-0.19557007E 01	-0.20054810E 01
2	-0.12141093E 01	0.27752244E-00
3	-0.23326014E-00	-0.87670457E-01
4	-0.25119858E-01	-0.18998946E-00
5	0.12301879E-00	-0.11061875E-00
6	0.15418245E-00	0.31000870E-01
7	0.69618856E-01	0.14309153E-00
8	-0.74772467E-01	0.15021475E-00

$q_o = 1.70\exp(i145^\circ)$

$t_o = 1.5105226 + i1.4363894$

n	Real A_n	Imag A_n
1	-0.20604455E 01	-0.20733957E 01
2	-0.12267265E 01	0.24330847E-00
3	-0.20751636E-00	-0.11483032E-00
4	0.14399479E-01	-0.18230417E-00
5	0.14140411E-00	-0.65806430E-01
6	0.12672086E-00	0.77010261E-01
7	0.57148401E-02	0.14857285E-00
8	-0.12565487E-00	0.91309742E-01

$q_o = 1.75\exp(i145^\circ)$

$t_o = 1.5104005 + i1.4596621$

n	Real A_n	Imag A_n
1	-0.21634762E 01	-0.21444190E 01
2	-0.12338984E 01	0.20951215E-00
3	-0.17833686E-00	-0.13498454E-00
4	0.48251616E-01	-0.16519434E-00
5	0.14443447E-00	-0.26305988E-01
6	0.80574337E-01	0.10588993E-00
7	-0.48447195E-01	0.12640835E-00
8	-0.13778460E-00	0.22142768E-01

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.80\exp(145^\circ)$

$t_o = 1.5092304 + i1.4817674$

n	Real A_n	Imag A_n
1	-0.22644915E 01	-0.22181155E 01
2	-0.12361307E 01	0.17714363E-00
3	-0.14782319E-00	-0.14797209E-00
4	0.74281961E-01	-0.14160773E-00
5	0.13433974E-00	0.11285232E-01
6	0.45226732E-01	0.11587257E-00
7	-0.82717918E-01	0.87439504E-01
8	-0.11672077E-00	-0.35463447E-01

$q_o = 1.85\exp(145^\circ)$

$t_o = 1.5071735 + i1.5026938$

n	Real A_n	Imag A_n
1	-0.23632956E 01	-0.22940300E 01
2	-0.12341227E 01	0.14698199E-00
3	-0.11783593E-00	-0.15428083E-00
4	0.91723242E-01	-0.11475278E-00
5	0.11509898E-00	0.40386754E-01
6	0.45425406E-02	0.10972737E-00
7	-0.94947357E-01	0.44297522E-01
8	-0.77406165E-01	-0.69537286E-01

$q_o = 1.90\exp(145^\circ)$

$t_o = 1.5043806 + i1.5224510$

n	Real A_n	Imag A_n
1	-0.24597927E 01	-0.23717146E 01
2	-0.12286731E 01	0.11955151E-00
3	-0.89826740E-01	-0.15485409E-00
4	0.10100264E-00	-0.87495010E-01
5	0.91154896E-01	0.59676795E-01
6	-0.24694620E-01	0.92771936E-01
7	-0.89262489E-01	0.68570913E-02
8	-0.35528116E-01	-0.79280217E-01

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.95\exp(i145^o)$

$t_o = 1.5009886 + i1.5410661$

n	Real A_n	Imag A_n
1	-0.25537726E 01	-0.24507305E 01
2	-0.12205941E 01	0.95133723E-01
3	-0.64762509E-01	-0.15087279E-00
4	0.10355550E-00	-0.62020496E-01
5	0.66394028E-01	0.69689599E-01
6	-0.43041542E-01	0.70744311E-01
7	-0.72666605E-01	-0.19745617E-01
8	-0.17330296E-02	-0.71207707E-01

$q_o = 2.00\exp(i145^o)$

$t_o = 1.4971197 + i1.5585788$

n	Real A_n	Imag A_n
1	-0.26458955E 01	-0.25307620E 01
2	-0.12106464E 01	0.73803559E-01
3	-0.43150991E-01	-0.14356328E-00
4	0.10040439E-00	-0.39712039E-01
5	0.45603485E-01	0.72107168E-01
6	-0.51524753E-01	0.48328004E-01
7	-0.51968273E-01	-0.34721857E-01
8	0.19796682E-01	-0.54013134E-01

$q_o = 2.05\exp(i145^o)$

$t_o = 1.4928805 + i1.5750388$

n	Real A_n	Imag A_n
1	-0.27356638E 01	-0.26114253E 01
2	-0.11994978E 01	0.55476008E-01
3	-0.25122084E-01	-0.13405856E-00
4	0.93811707E-01	-0.21220735E-01
5	0.24573738E-01	0.69057563E-01
6	-0.52466226E-01	0.28546988E-01
7	-0.32178292E-01	-0.40031955E-01
8	0.29761702E-01	-0.34945822E-01

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$$q_o = 2.10\exp(i145^o)$$

$$t_o = 1.4883628 + i1.5905017$$

n	Real A_n	Imag A_n
1	-0.28234249E 01	-0.26924697E 01
2	-0.11877027E 01	0.39957666E-01
3	-0.10531931E-01	-0.12331774E-00
4	0.85052217E-01	-0.66337522E-02
5	0.92893421E-02	0.62596819E-01
6	-0.48484058E-01	0.12833690E-01
7	-0.15935236E-01	-0.38701627E-01
8	0.31260084E-01	-0.18419635E-01

$$q_o = 2.15\exp(i145^o)$$

$$t_o = 1.4836442 + i1.6050261$$

n	Real A_n	Imag A_n
1	-0.29093366E 01	-0.27736759E 01
2	-0.11756996E 01	0.26988509E-01
3	0.93543349E-03	-0.11209558E-00
4	0.75302979E-01	0.43331879E-02
5	-0.17636636E-02	0.54423767E-01
6	-0.41871601E-01	0.14392892E-02
7	-0.41272982E-02	-0.33647855E-01
8	0.27792244E-01	-0.62236485E-02

$$q_o = 2.20\exp(i145^o)$$

$$t_o = 1.4787898 + i1.6186719$$

n	Real A_n	Imag A_n
1	-0.29935668E 01	-0.28548717E 01
2	-0.11638186E 01	0.16278045E-01
3	0.96834677E-02	-0.10094688E-00
4	0.65421522E-01	0.12165319E-01
5	-0.92962272E-02	0.45780734E-01
6	-0.34336335E-01	-0.60884871E-02
7	0.35238722E-02	-0.27113503E-01
8	0.22189391E-01	0.15994097E-02

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 2.25\exp(i145^\circ)$

$t_o = 1.4738531 + i1.6314986$

n	Real A_n	Imag A_n
1	-0.30762823E 01	-0.29359249E 01
2	-0.11522961E 01	0.75314543E-02
3	0.16140974E-01	-0.90251160E-01
4	0.55977318E-01	0.17420936E-01
5	-0.13981827E-01	0.37478443E-01
6	-0.26983559E-01	-0.10512557E-01
7	0.78323085E-02	-0.20551095E-01
8	0.16302145E-01	0.58631638E-02

$q_o = 2.30\exp(i145^\circ)$

$t_o = 1.4688782 + i1.6435635$

n	Real A_n	Imag A_n
1	-0.31576431E 01	-0.30167385E 01
2	-0.11412913E 01	0.46611596E-03
3	0.20721167E-01	-0.80245519E-01
4	0.47307265E-01	0.20647744E-01
5	-0.16502746E-01	0.29977524E-01
6	-0.20421964E-01	-0.12645830E-01
7	0.97333679E-02	-0.14743836E-01
8	0.11112744E-01	0.76129670E-02

$q_o = 2.35\exp(i145^\circ)$

$t_o = 1.4639002 + i1.6549218$

n	Real A_n	Imag A_n
1	-0.32377992E 01	-0.30972430E 01
2	-0.11309019E 01	-0.51776654E-02
3	0.23798249E-01	-0.71038545E-01
4	0.39575886E-01	0.22336286E-01
5	-0.17466106E-01	0.23484930E-01
6	-0.14908195E-01	-0.13210613E-01
7	0.10061622E-01	-0.10003487E-01
8	0.69951870E-02	0.77914037E-02

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 2.40\exp(i145^o)$

$t_o = 1.45589476 + i1.6656250$

n	Real A_n	Imag A_n
1	-0.33168883E 01	-0.31773922E 01
2	-0.11211799E 01	-0.96317171E-02
3	0.25696705E-01	-0.62740476E-01
4	0.32829142E-01	0.22899927E-01
5	-0.17369481E-01	0.18040799E-01
6	-0.10475364E-01	-0.12783258E-01
7	0.94661792E-02	-0.63559423E-02
8	0.39676845E-02	0.71139441E-02

$q_o = 2.45\exp(i145^o)$

$t_o = 1.4540429 + i1.6757223$

n	Real A_n	Imag A_n
1	-0.33950352E 01	-0.32571570E 01
2	-0.11121429E 01	-0.13098456E-01
3	0.26688969E-01	-0.55288094E-01
4	0.27037656E-01	0.22671692E-01
5	-0.16576942E-01	0.13586625E-01
6	-0.70342459E-02	-0.11789446E-01
7	0.84051536E-02	-0.36807614E-02
8	0.18790521E-02	0.60625041E-02

$q_o = 2.50\exp(i145^o)$

$t_o = 1.4492032 + i1.6852585$

n	Real A_n	Imag A_n
1	-0.34723518E 01	-0.33365223E 01
2	-0.11037847E 01	-0.15751884E-01
3	0.26998235E-01	-0.48663935E-01
4	0.22128706E-01	0.21910591E-01
5	-0.15430962E-01	0.10013570E-01
6	-0.44409981E-02	-0.10524012E-01
7	0.71761011E-02	-0.18016712E-02
8	0.52418395E-03	0.49292143E-02

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 2.55\exp(i145^o)$

$t_o = 1.4444420 + i1.6942755$

n	Real A_n	Imag A_n
1	-0.35489374E 01	-0.34154832E 01
2	-0.10960830E 01	-0.17739509E-01
3	0.26803680E-01	-0.42810526E-01
4	0.18008585E-01	0.20812096E-01
5	-0.14070429E-01	0.71941244E-02
6	-0.25387698E-02	-0.91789643E-02
7	0.59563193E-02	-0.53809933E-03
8	-0.29505823E-03	0.38711811E-02

$q_o = 2.60\exp(i145^o)$

$t_o = 1.4397693 + i1.7028119$

n	Real A_n	Imag A_n
1	-0.36248801E 01	-0.34940425E 01
2	-0.10890052E 01	-0.19185089E-01
3	0.26246816E-01	-0.37660400E-01
4	0.14577132E-01	0.19519807E-01
5	-0.12649424E-01	0.50010202E-02
6	-0.11802283E-02	-0.78709296E-02
7	0.48402283E-02	0.26992033E-03
8	-0.74419327E-03	0.29585057E-02

$q_o = 2.65\exp(i145^o)$

$t_o = 1.4351924 + i1.7109028$

n	Real A_n	Imag A_n
1	-0.37002569E 01	-0.35722084E 01
2	-0.10825128E 01	-0.20191938E-01
3	0.25437778E-01	-0.33142857E-01
4	0.11736647E-01	0.18136401E-01
5	-0.11253928E-01	0.33174241E-02
6	-0.23781255E-03	-0.66642665E-02
7	0.36690072E-02	0.75289337E-03
8	-0.95010809E-03	0.22100975E-02

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 2.70\exp(i145^\circ)$

$t_o = 1.4307162 + 1.7185812$

n	Real A_n	Imag A_n
1	-0.37751360E 01	-0.36499930E 01
2	-0.10765644E 01	-0.20845375E-01
3	0.24461174E-01	-0.29188336E-01
4	0.93968058E-02	0.16733225E-01
5	-0.99355357E-02	0.20414288E-02
6	0.39336304E-03	-0.55888654E-02
7	0.30519950E-02	0.10116339E-02
8	-0.10044345E-02	0.16177900E-02

$q_o = 2.75\exp(i145^\circ)$

$t_o = 1.4263443 + i1.7258766$

n	Real A_n	Imag A_n
1	-0.38495766E 01	-0.37274108E 01
2	-0.10711180E 01	-0.21215896E-01
3	0.23581160E-01	-0.25731035E-01
4	0.74770088E-02	0.15358130E-01
5	-0.87220672E-02	0.10871763E-02
6	0.79646157E-03	-0.46530651E-02
7	0.23811133E-02	0.11209490E-02
8	-0.97028439E-03	0.11612948E-02

$q_o = 2.80\exp(i145^\circ)$

$t_o = 1.4220782 + i1.7328160$

n	Real A_n	Imag A_n
1	-0.39236308E 01	-0.38044778E 01
2	-0.10661326E 01	-0.21361263E-01
3	0.22245742E-01	-0.22710320E-01
4	0.59070366E-02	0.14041748E-01
5	-0.76254664E-02	0.38407662E-03
6	0.10356723E-02	-0.38525677E-02
7	0.18400040E-02	0.11346001E-02
8	-0.88934766E-03	0.81675809E-03

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 2.85\exp(i145^\circ)$

$t_o = 1.4179189 + i1.7394242$

n	Real A_n	Imag A_n
1	-0.39973436E 01	-0.38812114E 01
2	-0.10615686E 01	-0.21328687E-01
3	0.21090315E-01	-0.20071374E-01
4	0.46267830E-02	0.12802324E-01
5	-0.66474937E-02	-0.12485612E-03
6	0.11595015E-02	-0.31763460E-02
7	0.14095004E-02	0.10901563E-02
8	-0.78803023E-03	0.56122144E-03

$q_o = 2.90\exp(i145^\circ)$

$t_o = 1.4138661 + i1.7457250$

n	Real A_n	Imag A_n
1	-0.40707552E 01	-0.39576285E 01
2	-0.10573889E 01	-0.21156660E-01
3	0.19940514E-01	-0.17765286E-01
4	0.35854494E-02	0.11649373E-01
5	-0.57837112E-02	-0.48498742E-03
6	0.12039574E-02	-0.26104024E-02
7	0.10706633E-02	0.10131820E-02
8	-0.68226423E-03	0.37459977E-03

$q_o = 2.95\exp(i145^\circ)$

$t_o = 1.4099192 + i1.7517384$

n	Real A_n	Imag A_n
1	-0.41438998E 01	-0.40337463E 01
2	-0.10535590E 01	-0.20876610E-01
3	0.18814492E-01	-0.15748823E-01
4	0.27404917E-02	0.10586388E-01
5	-0.50261766E-02	-0.73205028E-03
6	0.11953606E-02	-0.21400315E-02
7	0.80628451E-03	0.92059250E-03
8	-0.58105485E-03	0.24028598E-03

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 3.00\exp(i145^\circ)$

$t_o = 1.4060774 + i1.7574821$

n	Real A_n	Imag A_n
1	-0.42168079E 01	-0.41695823E 01
2	-0.10500468E 01	-0.20513897E-01
3	0.17724701E-01	-0.13984065E-01
4	0.20565450E-02	0.96128322E-02
5	-0.43652582E-02	-0.69392930E-03
6	0.11526068E-02	-0.17511155E-02
7	0.60149744E-03	0.82321673E-03
8	-0.48897856E-03	0.14504905E-03

$q_o = 3.05\exp(i145^\circ)$

$t_o = 1.4023372 + i1.7629775$

n	Real A_n	Imag A_n
1	-0.42895061E 01	-0.41851520E 01
2	-0.10468232E 01	-0.20089266E-01
3	0.16679364E-01	-0.12437939E-01
4	0.15043601E-02	0.87256134E-02
5	-0.37908032E-02	-0.99224318E-03
6	0.10895353E-02	-0.14307850E-02
7	0.44587375E-03	0.72770145E-03
8	-0.40789157E-03	0.78618832E-04

$q_o = 3.10\exp(i145^\circ)$

t_o 1.3986976 + i1.7682386

n	Real A_n	Imag A_n
1	-0.43620173E 01	-0.42604715E 01
2	-0.10436616E 01	-0.19619147E-01
3	0.15683445E-01	-0.11081672E-01
4	0.10598423E-02	0.79200251E-02
5	-0.32928073E-02	-0.10436270E-02
6	0.10147535E-02	-0.11676928E-02
7	0.32325066E-03	0.63786249E-03
8	-0.33604566E-03	0.33175654E-04

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 3.15\exp(145^o)$

$t_o = 1.3951562 + i1.7732802$

n	Real A_n	Imag A_n
1	-0.44343621E 01	-0.43355558E 01
2	-0.10411378E 01	-0.19117019E-01
3	0.14739561E-01	-0.98903269E-02
4	0.10319847E-03	0.71905120E-02
5	-0.28618536E-02	-0.10608169E-02
6	0.93535078E-03	-0.95205159E-03
7	0.23145616E-03	0.55565957E-03
8	-0.27881856E-03	0.28548413E-05

$q_o = 3.20\exp(145^o)$

$t_o = 1.3917100 + i1.7781165$

n	Real A_n	Imag A_n
1	-0.45065583E 01	-0.44104189E 01
2	-0.10386299E 01	-0.18593632E-01
3	0.13848622E-01	-0.88423163E-02
4	0.41820063E-03	0.65311504E-02
5	-0.24893192E-02	-0.10535108E-02
6	0.85542436E-03	-0.77555661E-03
7	0.16199705E-03	0.48186320E-03
8	-0.22916314E-03	-0.16690663E-04

$q_o = 3.25\exp(145^o)$

$t_o = 1.3883571 + i1.7827593$

n	Real A_n	Imag A_n
1	-0.45786219E 01	-0.44850741E 01
2	-0.10363180E 01	-0.18057566E-01
3	0.13010284E-01	-0.79189835E-02
4	0.19157200E-03	0.59359587E-02
5	-0.21674543E-02	-0.10290437E-02
6	0.77774605E-03	-0.63124214E-03
7	0.10976191E-03	0.41650754E-03
8	-0.18787760E-03	-0.28649192E-04

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 3.30\exp(i145^o)$

$t_o = 1.3850945 + i1.7872202$

n	Real A_n	Imag A_n
1	-0.46505665E 01	-0.45595341E 01
2	-0.10341845E 01	-0.17515668E-01
3	0.12223550E-01	-0.71042042E-02
4	0.12450634E-04	0.53991216E-02
5	-0.18893922E-02	-0.99293625E-03
6	0.70401736E-03	-0.51331396E-03
7	0.70755825E-04	0.35919641E-03
8	-0.15376058E-03	-0.35339589E-04

$q_o = 3.35\exp(i145^o)$

$t_o = 1.3819194 + i1.7915103$

n	Real A_n	Imag A_n
1	-0.47224044E 01	-0.46338108E 01
2	-0.10322130E 01	-0.16973265E-01
3	0.11486032E-01	-0.63840523E-02
4	-0.12603816E-03	0.49151176E-02
5	-0.16491113E-02	-0.94931187E-03
6	0.63519484E-03	-0.41698273E-03
7	0.41865315E-04	0.30930389E-03
8	-0.12569413E-03	-0.38433459E-04

$q_o = 3.40\exp(i145^o)$

$t_o = 1.3788290 + i1.7956392$

n	Real A_n	Imag A_n
1	-0.47941465E 01	-0.47079150E 01
2	-0.10303891E 01	-0.16434585E-01
3	0.10796162E-01	-0.57464899E-02
4	-0.23714352E-03	0.44787870E-02
5	-0.14413653E-02	-0.90122464E-03
6	0.57172528E-03	-0.33830739E-03
7	0.20690443E-04	0.26610129E-03
8	-0.10266015E-03	-0.39125869E-04

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 3.45\exp(i145^o)$

$t_o = 1.3758211 + i1.7996162$

n	Real A_n	Imag A_n
1	-0.48658025E 01	-0.47818574E 01
2	-0.10286996E 01	-0.15903011E-01
3	0.10151345E-01	-0.51811139E-02
4	-0.32078498E-03	0.40853695E-02
5	-0.12616122E-02	-0.85091516E-03
6	0.51571504E-03	-0.27405603E-03
7	0.53735118E-05	0.22883779E-03
8	-0.83852606E-04	-0.36263702E-04

$q_o = 3.50\exp(i145^o)$

$t_o = 1.3728925 + i1.8034494$

n	Real A_n	Imag A_n
1	-0.49373803E 01	-0.48556474E 01
2	-0.10271528E 01	-0.15581002E-01
3	0.95490669E-02	-0.46789175E-02
4	-0.38579899E-03	0.37305136E-02
5	-0.11059324E-02	-0.79999913E-03
6	0.46105070E-03	-0.22158436E-03
7	-0.55131951E-05	0.19078722E-03
8	-0.68473867E-04	-0.36440962E-04

$q_o = 3.55\exp(i145^o)$

$t_o = 1.3700408 + i1.8071472$

n	Real A_n	Imag A_n
1	-0.50088878E 01	-0.49292940E 01
2	-0.10256760E 01	-0.14870457E-01
3	0.89867938E-02	-0.42321042E-02
4	-0.43013459E-03	0.34102689E-02
5	-0.97095511E-03	-0.74963500E-03
6	0.41346172E-03	-0.17873184E-03
7	-0.13064817E-04	0.16927404E-03
8	-0.55924583E-04	-0.34069068E-04

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 3.60\exp(i45^o)$

t_o 1.3672636 + i1.8107165

n	Real A_n	Imag A_n
1	-0.50803321E 01	-0.50028058E 01
2	-0.10243256E 01	-0.14372756E-01
3	0.84620083E-02	-0.38339126E-02
4	-0.46301585E-03	0.31210649E-02
5	-0.85578578E-03	-0.70062093E-04
6	0.37067942E-03	-0.14373558E-03
7	-0.18119431E-04	0.14568627E-04
8	-0.45689525E-04	-0.31429048E-04

$q_o = 3.65\exp(i45^o)$

t_o = 1.3645581 + i1.8141641

n	Real A_n	Imag A_n
1	-0.51517186E 01	-0.50761906E 01
2	-0.10230669E 01	-0.13688909E-01
3	0.79722615E-02	-0.34784739E-02
4	-0.48507719E-03	0.28596894E-02
5	-0.75194304E-03	-0.65549316E-04
6	0.33227621E-03	-0.11515796E-03
7	-0.21316995E-04	0.12547878E-04
8	-0.37343342E-04	-0.28708592E-04

$q_o = 3.70\exp(i45^o)$

t_o = 1.3619222 + i1.8174963

n	Real A_n	Imag A_n
1	-0.52230531E 01	-0.51494560E 01
2	-0.10218941E 01	-0.13419077E-01
3	0.75151992E-02	-0.31006881E-02
4	-0.49640887E-03	0.26232601E-02
5	-0.66330115E-03	-0.60859183E-04
6	0.29789285E-03	-0.91826419E-04
7	-0.23144849E-04	0.10817268E-03
8	-0.30536673E-04	-0.26029773E-04

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$$q_o = 3.75\exp(i145^\circ)$$

$$t_o = 1.3593533 + i1.8207189$$

n	Real A_n	Imag A_n
1	-0.52943401E 01	-0.52226063E 01
2	-0.10207999E 01	-0.12965300E-01
3	0.70885750E-02	-0.26761138E-02
4	-0.50494686E-03	0.24071948E-02
5	-0.58605879E-03	-0.56611179E-03
6	0.26715484E-03	-0.72784453E-04
7	-0.23973297E-04	0.93550790E-04
8	-0.24985640E-04	-0.23468540E-04

$$q_o = 3.80\exp(i145^\circ)$$

$$t_o = 1.3568492 + i1.8238375$$

n	Real A_n	Imag A_n
1	-0.53655838E 01	-0.52956545E 01
2	-0.10197762E 01	-0.12526073E-01
3	0.66902680E-02	-0.26208824E-02
4	-0.50594430E-03	0.22151868E-02
5	-0.51059682E-03	-0.52614284E-03
6	0.23970441E-03	-0.57252C96E-04
7	-0.24082687E-04	0.80652606E-04
8	-0.20451079E-04	-0.21069042E-04

$$q_o = 3.85\exp(i145^\circ)$$

$$t_o = 1.3544079 + i1.8268576$$

n	Real A_n	Imag A_n
1	-0.54367881E 01	-0.53086061E 01
2	-0.10188229E 01	-0.12102C67E-01
3	0.63182786E-02	-0.23916164E-02
4	-0.50262973E-03	0.20391754E-02
5	-0.45963939E-03	-0.48869811E-03
6	0.21520581E-03	-0.44592585E-04
7	-0.23684528E-04	0.69767899E-04
8	-0.16749181E-04	-0.18853638E-04

Table C-1 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 3.90\exp(145^\circ)$

$t_o = 1.3520270 + i1.8297833$

n	Real A_n	Imag A_n
1	-0.55079567E 01	-0.54414516E 01
2	-0.10179289E 01	-0.11693042E-01
3	0.59707344E-02	-0.21853626E-02
4	-0.49595626E-03	0.18793207E-02
5	-0.40602091E-03	-0.45373727E-03
6	0.19334895E-03	-0.34265807E-04
7	-0.22937772E-04	0.60430849E-04
8	-0.13723475E-04	-0.16630120E-04

$q_o = 3.95\exp(145^\circ)$

$t_o = 1.3497046 + i1.8326191$

n	Real A_n	Imag A_n
1	-0.55790924E 01	-0.55142124E 01
2	-0.10170912E 01	-0.11298958E-01
3	0.56458866E-02	-0.19995363E-02
4	-0.48669912E-03	0.17339818E-02
5	-0.36275933E-03	-0.42118280E-03
6	0.17385051E-03	-0.25906769E-04
7	-0.21961065E-04	0.52414380E-04
8	-0.11248565E-04	-0.14996757E-04

$q_o = 4.00\exp(145^\circ)$

$t_o = 1.3474386 + i1.8353696$

n	Real A_n	Imag A_n
1	-0.56501981E 01	-0.55868888E 01
2	-0.10163056E 01	-0.10919458E-01
3	0.53421079E-02	-0.18518711E-02
4	-0.47548915E-03	0.16016949E-02
5	-0.32301106E-03	-0.39093304E-03
6	0.15645374E-03	-0.19107663E-04
7	-0.20842511E-04	0.45524789E-04
8	-0.92225168E-05	-0.13345870E-04

Table C-2

VALUES OF q_o, t_o, AND A_n

$q_o = 0.00$

$t_o = 0.50939649 + i\ 0.88230059$

n	Real A_n	Imag A_n
1	0.49077678E-00	-0.85005043E 00
2	-0.52716219E 00	0.
3	0.67202630E-01	0.11639834E-00
4	-0.35801253E-02	0.62010126E-02
5	0.98774254E-02	-0.23255247E-07
6	0.10064451E-02	0.17432125E-02
7	0.67314184E-03	-0.11658314E-02
8	0.56667980E-03	0.82246385E-07

$q_o = 0.05$

$t_o = 0.53512579 + i0.83981265$

n	Real A_n	Imag A_n
1	0.43856310E-00	-0.84917433E 00
2	-0.51712305E 00	0.17552909E-01
3	0.66736226E-01	0.11764263E-00
4	-0.10698256E-02	0.62612829E-02
5	0.10218605E-01	0.40162159E-03
6	0.12807611E-02	0.13532575E-02
7	0.89227157E-03	-0.11828364E-02
8	0.55176835E-03	-0.90049768E-04

$q_o = 0.10$

$t_o = 0.56326947 + i0.79741256$

n	Real A_n	Imag A_n
1	0.38755110E-00	-0.84655358E 00
2	-0.50711574E 00	0.35293913E-01
3	0.66781138E-01	0.11890952E-00
4	0.15369735E-02	0.64214895E-02
5	0.10653329E-01	0.79869235E-03
6	0.16296961E-02	0.91628214E-03
7	0.11004362E-02	-0.12421864E-02
8	0.51116494E-03	-0.21209446E-03

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.15$

$t_o = 0.59387758 + i0.75518902$

n	Real A_n	Imag A_n
1	0.33714148E-00	-0.84210916E 00
2	-0.49706205E-00	0.53227700E-01
3	0.67359254E-01	0.12021575E-00
4	0.42665052E-02	0.66498229E-02
5	0.11203521E-01	0.10048656E-02
6	0.20502741E-02	0.45719365E-03
7	0.14024642E-02	-0.13573345E-02
8	0.50004933E-03	-0.36996638E-04

$q_o = 0.20$

$t_o = 0.62700063 + i0.71323170$

n	Real A_n	Imag A_n
1	0.28794296E-00	-0.83558138E 00
2	-0.46687997E-00	0.71361089E-01
3	0.68496044E-01	0.12157167E-00
4	0.71562634E-02	0.69120374E-02
5	0.11890315E-01	0.10078371E-02
6	0.25407451E-02	-0.48147774E-04
7	0.15008060E-02	-0.15438503E-02
8	0.49126036E-03	-0.57670641E-04

$q_o = 0.25$

$t_o = 0.66268898 + i0.67163127$

n	Real A_n	Imag A_n
1	0.23977263E-00	-0.82783004E 00
2	-0.47648254E-00	0.89701863E-01
3	0.70232038E-01	0.12298039E-00
4	0.10224610E-01	0.71701970E-02
5	0.12734948E-01	0.76787202E-03
6	0.31003407E-02	-0.64410822E-04
7	0.16945567E-02	-0.18209346E-02
8	0.47468224E-03	-0.82429291E-04

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.30$

$t_o = 0.70099524 + i0.63047944$

n	Real A_n	Imag A_n
1	0.19265667E-00	-0.81793389E 00
2	-0.46577841E-00	0.10825760E-00
3	0.72603515E-01	0.12443665E-00
4	0.13532405E-01	0.75800662E-02
5	0.13757250E-01	0.67571150E-03
6	0.37259462E-02	-0.13362344E-02
7	0.18776361E-02	-0.22121786E-02
8	0.43509024E-03	-0.11439832E-02

$q_o = 0.35$

$t_o = 0.74197277 + i0.58986899$

n	Real A_n	Imag A_n
1	0.14663053E-00	-0.80617117E 00
2	-0.45466733E-00	0.12703449E-00
3	0.75663691E-01	0.12592572E-00
4	0.17119835E-01	0.74887159E-02
5	0.14976501E-01	0.15008235E-03
6	0.44118442E-02	-0.21994417E-02
7	0.20366920E-02	-0.27471110E-02
8	0.35001275E-03	-0.15455974E-02

$q_o = 0.40$

$t_o = 0.78567719 + i0.54989380$

n	Real A_n	Imag A_n
1	0.10174035E-00	-0.79251954E 00
2	-0.44304179E-00	0.14603568E-00
3	0.79473514E-01	0.12742114E-00
4	0.21038592E-01	0.74510044E-02
5	0.16409188E-01	-0.66570904E-03
6	0.51459903E-02	-0.32802539E-02
7	0.21472513E-02	-0.34621124E-02
8	0.18598482E-03	-0.20475867E-02

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.45$

$t_o = 0.83216666 + i0.51064889$

n	Real A_n	Imag A_n
1	0.5884328[illegible]E-01	-0.77695664E 00
2	-0.4367[illegible]417E-00	0.16525917E-00
3	0.8410[illegible]37E-01	0.12[illegible]175E-00
4	0.2534[illegible]55E-01	0.71254356E-02
5	0.1[illegible]66501E-01	-0.18469491E-02
6	0.5704[illegible]67E-02	-0.46420848E-02
7	0.21685036E-02	-0.44010455E-02
8	-0.1065[illegible]02E-03	-0.26663819E-02

$q_o = 0.50$

$t_o = 0.88150253 + i0.47223036$

n	Real A_n	Imag A_n
1	0.15608648E-01	-0.75946065E 00
2	-0.[illegible]7765[illegible]E-00	0.18469547E-00
3	0.8964[illegible]354E-01	0.13024806E-00
4	0.3009[illegible]545E-01	0.04690551E-02
5	0.1995[illegible]216E-01	-0.34926626E-02
6	0.6646[illegible]682E-02	-0.63731966E-02
7	0.20355204E-02	-0.56146751E-02
8	-0.6[illegible]3664[illegible]E-03	-0.34224112E-02

$q_o = 0.55$

$t_o = 0.93374994 + i0.43473539$

n	Real A_n	Imag A_n
1	-0.25479670E-01	-0.74001116E 00
2	-0.40384207E-00	0.20432473E-00
3	0.96174251E-01	0.13143735E-00
4	0.35355816E-01	0.53300074E-02
5	0.22045446E-01	-0.57309012E-02
6	0.72999721E-02	-0.86179763E-02
7	0.16477215E-02	-0.71572965E-02
8	-0.14006944E-02	-0.43123186E-02

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.60$

$t_o = 0.98897858 + i0.39826210$

n	Real A_n	Imag A_n
1	-0.65124152E-01	-0.71859045E 00
2	-0.38885761E-00	0.22411231E-00
3	0.10381107E-00	0.13233694E-00
4	0.41127338E-01	0.35407408E-02
5	0.24310289E-01	-0.67240490E-02
6	0.77525707E-02	-0.11447660E-01
7	0.85454329E-03	-0.90793680E-02
8	-0.26562860E-02	-0.54127281E-02

$q_o = 0.65$

$t_o = 1.0472634 + i0.36290942$

n	Real A_n	Imag A_n
1	-0.10320996E-00	-0.69518525E 00
2	-0.37263785E-00	0.24400392E-00
3	0.11266380E-00	0.13279637E-00
4	0.47497964E-01	0.88842097E-03
5	0.26659375E-01	-0.12674160E-01
6	0.78256704E-02	-0.15021605E-01
7	-0.56468973E-03	-0.11411824E-01
8	-0.45704381E-02	-0.63441502E-02

$q_o = 0.70$

$t_o = 1.1086857 + i0.32877678$

n	Real A_n	Imag A_n
1	-0.13960421E-00	-0.66978885E 00
2	-0.35499176E-00	0.26391937E-00
3	0.12284925E-00	0.13261701E-00
4	0.54431606E-01	-0.28962810E-02
5	0.28940794E-01	-0.17625857E-01
6	0.72480750E-02	-0.19482108E-01
7	-0.29245988E-02	-0.14135721E-01
8	-0.74080730E-02	-0.72278067E-02

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.75$

$t_o = 1.1733337 + i0.29596369$

n	Real A_n	Imag A_n
1	-0.17415386E-00	-0.64240444E 00
2	-0.33571065E-00	0.28374455E-00
3	0.13448029E-00	0.13154036E-00
4	0.61942057E-01	-0.81885390E-02
5	0.30903779E-01	-0.24464585E-01
6	0.56192616E-02	-0.24948687E-01
7	-0.66568514E-02	-0.17128546E-01
8	-0.11488824E-01	-0.76155847E-02

$q_o = 0.80$

$t_o = 1.2413039 + i0.26456921$

n	Real A_n	Imag A_n
1	-0.20668434E-00	-0.61304841E 00
2	-0.31457024E-00	0.30332016E-00
3	0.14765367E-00	0.12923298E-00
4	0.69643248E-01	-0.15278535E-01
5	0.32150647E-01	-0.32901528E-01
6	0.23635466E-02	-0.31468274E-01
7	-0.12325263E-01	-0.20077866E-01
8	-0.17140146E-01	-0.65750199E-02

$q_o = 0.85$

$t_o = 1.3127021 + i0.23469114$

n	Real A_n	Imag A_n
1	-0.23679792E-00	-0.56175585E 00
2	-0.29133429E-00	0.32245183E-00
3	0.16242892E-00	0.12527142E-00
4	0.77908545E-01	-0.24774824E-01
5	0.32080334E-01	-0.43440334E-01
6	-0.33133578E-02	-0.35925108E-01
7	-0.20609067E-01	-0.22341965E-01
8	-0.24579514E-01	-0.39073666E-02

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 0.90$

$t_o = 1.3876447 + i0.20642491$

n	Real A_n	Imag A_n
1	-0.26487317E-00	-0.54658702E 00
2	-0.26576160E-00	0.34079287E-00
3	0.17879927E-00	0.11912633E-00
4	0.85702459E-01	-0.37193755E-01
5	0.29817777E-01	-0.56309552E-01
6	-0.12450532E-01	-0.46893570E-01
7	-0.32210658E-01	-0.22749615E-01
8	-0.33650915E-01	0.27294014E-02

$q_o = 0.95$

$t_o = 1.4662595 + i0.17986219$

n	Real A_n	Imag A_n
1	-0.29006452E-00	-0.51363472E 00
2	-0.23762028E-00	0.35802938E-00
3	0.19664627E-00	0.11015970E-00
4	0.92533557E-01	-0.55127269E-01
5	0.24139163E-01	-0.71534265E-01
6	-0.26307254E-01	-0.54409514E-01
7	-0.47618759E-01	-0.19351963E-01
8	-0.43330926E-01	0.15492537E-01

$q_o = 1.00$

$t_o = 1.5486874 + i0.15508894$

n	Real A_n	Imag A_n
1	-0.31230473E-00	-0.47703449E-00
2	-0.20670792E-00	0.37366179E-00
3	0.21567942E-00	0.97605970E-01
4	0.97353712E-01	-0.73127528E-01
5	0.13428456E-01	-0.88732112E-01
6	-0.46207459E-01	-0.59653530E-01
7	-0.66608068E-01	-0.91785008E-02
8	-0.50908878E-01	0.37077615E-01

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.05$

$t_o = 1.6350828 + i0.13218309$

n	Real A_n	Imag A_n
1	-0.33150674E-00	-0.44897590E-00
2	-0.17268365E-00	0.36708962E-00
3	0.23535423E-00	0.80611722E-01
4	0.96664297E-01	-0.97569833E-01
5	-0.42917822E-02	-0.10079898E-00
6	-0.73146594E-01	-0.59573591E-01
7	-0.87348204E-01	0.11813474E-01
8	-0.50659480E-01	0.69866044E-01

$q_o = 1.10$

$t_o = 1.7256153 + i0.11121146$

n	Real A_n	Imag A_n
1	-0.34678288E-00	-0.39971452E-00
2	-0.13611348E-00	0.39757879E-00
3	0.25477710E-00	0.56282649E-01
4	0.94444534E-01	-0.12641752E-00
5	-0.31160440E-01	-0.12349015E-00
6	-0.10701422E-00	-0.49581347E-01
7	-0.10504445E-00	0.48218970E-01
8	-0.33781846E-01	0.11373176E-00

$q_o = 1.15$

$t_o = 1.8204682 + i0.09222614$

n	Real A_n	Imag A_n
1	-0.35843727E-00	-0.35958664E-00
2	-0.96530037E-01	0.40426635E-00
3	0.27259596E-00	0.29800404E-01
4	0.82170694E-01	-0.15883458E-00
5	-0.68952637E-01	-0.13495134E-00
6	-0.14525838E-00	-0.23607950E-01
7	-0.11035906E-00	0.10343695E-00
8	0.13424836E-01	0.16197757E-00

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.20$

$t_o = 1.9198395 + i0.07526016$

n	Real A_n	Imag A_n
1		
2	-0.36600704E-00	-0.31902035E-00
3	-0.54508518E-01	0.40618338E-00
4	0.28691886E-00	-0.53838246E-02
5	0.59015419E-01	-0.19298114E-00
6	-0.11809609E-00	-0.13539547E-00
7	-0.18111543E-00	0.24882849E-01
8	-0.88518978E-01	0.17593172E-00
	0.10370239E-00	0.19563024E-00

$q_o = 1.25$

$t_o = 2.0239391 + i0.06032268$

n	Real A_n	Imag A_n
1	-0.36928052E-00	-0.27854304E-00
2	-0.10750655E-01	0.40231539E-00
3	0.29529958E-00	-0.47260826E-01
4	0.22368342E-01	-0.22504088E-00
5	-0.17613644E-00	-0.11731674E-00
6	-0.20191094E-00	0.10018373E-00
7	-0.21264985E-01	0.25331524E-00
8	0.23899319E-00	0.17913409E-00

$q_o = 1.30$

$t_o = 2.1329851 + i0.04739397$

n	Real A_n	Imag A_n
1	-0.36813535E-00	-0.23678201E-00
2	0.33640496E-01	0.39171547E-00
3	0.29486471E-00	-0.94893631E-01
4	-0.29211505E-01	-0.24943522E-00
5	-0.23588288E-00	-0.72644435E-01
6	-0.18901705E-00	0.19942410E-00
7	0.10518846E-00	0.30624695E-00
8	0.39143325E-00	0.65604385E-01

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.35$

$t_o = 2.2471991 + i0.03642072$

n	Real A_n	Imag A_n
1	-0.36258218E-00	-0.20044874E-00
2	0.77120436E-01	0.47366951E-00
3	0.26266301E-00	-0.14064504E-00
4	-0.94627773E-01	-0.[illegible]14[illegible]E-00
5	-0.28410622E-00	0.31644[illegible]17E-02
6	-0.12198006E-00	0.30631879E-00
7	0.28428029E-00	0.28[illegible]36159E-00
8	0.48534942E-00	-0.17648890E-00

$q_o = 1.40$

$t_o = 2.3667974 + i0.02731259$

n	Real A_n	Imag A_n
1	-0.35280016E-00	-0.16430580E-00
2	0.11773645E-00	0.34791677E-00
3	0.25642068E-00	-0.19689882E-00
4	-0.16877199E-00	-0.24[illegible]7781E-00
5	-0.30259745E-00	0.10[illegible]400940E-00
6	0.10973007E-01	0.38732324E-00
7	0.47026827E-00	0.15202834E-00
8	0.40593039E-00	-0.51228042E-00

$q_o = 1.45$

$t_o = 2.4919823 + i0.01994098$

n	Real A_n	Imag A_n
1	-0.35920253E-00	-0.13110943E-00
2	0.15329533E-00	0.31487478E-00
3	0.21524344E-00	-0.24227031E-00
4	-0.24174544E-00	-0.20[illegible]41[illegible]E-00
5	-0.27201128E-00	0.22642067E-00
6	0.17765565E-00	0.39716548E-00
7	0.[illegible]736907[illegible] 00	-0.11495162E-00
8	0.66566865E-01	-0.79[illegible]086[illegible]E-00

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.50$

$t_o = 2.6229306 + i0.01414122$

n	Real A_n	Imag A_n
1	-0.32236712E-00	-0.10153283E-00
2	0.18163661E-00	0.27580611E-00
3	0.16071396E-00	-0.27627596E-00
4	-0.29984519E-00	-0.13220273E-00
5	-0.18304741E-00	0.35575100E-00
6	0.39108021E-00	0.29879971E-00
7	0.49517125E-00	-0.44792168E-00
8	-0.47707365E-00	-0.80556014E 00

$q_o = 1.55$

$t_o = 2.7597834 + i0.00971885$

n	Real A_n	Imag A_n
1	-0.30317284E-00	-0.76079421E-01
2	0.20105654E-00	0.23282619E-00
3	0.97245142E-01	-0.29368742E-00
4	-0.32903231E-00	-0.39348166E-01
5	-0.44033110E-01	0.39672526E-00
6	0.51831686E 00	0.92667054E-01
7	0.19892638E-00	-0.70283797E 00
8	-0.96562304E 00	-0.40065500E-00

$q_o = 1.60$

$t_o = 2.9026376 + i0.00646011$

n	Real A_n	Imag A_n
1	-0.28250285E-00	-0.55006620E-01
2	0.21070257E-00	0.18868265E-00
3	0.31658978E-01	-0.29156594E-00
4	-0.32017936E-00	0.60113462E-01
5	0.11470491E-00	0.38595626E-00
6	0.51500867E 00	-0.16602948E-00
7	-0.22480080E-00	-0.72881055E 00
8	-0.10676555E 01	0.29171895E-00

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.65$

$t_o = 3.0515411 + 10.00414553$

n	Real A_n	Imag A_n
1	-0.20135071E-00	-0.36283583E-01
2	0.21063341E-00	0.14631177E-00
3	-0.25288849E-01	-0.27046179E-00
4	-0.27301885E-00	0.14737620E-00
5	0.25040152E-00	0.30112602E-00
6	0.10843674E-00	-0.38498398E-00
7	-0.58515148E 00	-0.48197028E-00
8	-0.65784962E 00	0.89123177E 00

$q_o = 1.70$

$t_o = 3.2064956 + 10.00256434$

n	Real A_n	Imag A_n
1	-0.24060726E-00	-0.25998544E-01
2	0.20283372E-00	0.10829071E-00
3	-0.75967736E-01	-0.23450982E-00
4	-0.19994380E-00	0.20653170E-00
5	0.12676045E-00	0.16711468E-00
6	0.13315720E-00	-0.48455469E-00
7	-0.71280074E 00	-0.76573469E-01
8	0.32455586E-01	0.10471603E 01

$q_o = 1.75$

$t_o = 3.3674616 + 10.00152717$

n	Real A_n	Imag A_n
1	-0.22098264E-00	-0.16420528E-01
2	0.18885200E-00	0.76375730E-01
3	-0.10761090E-00	-0.19023661E-00
4	-0.11636575E-00	0.23024898E-00
5	0.33064976E-00	0.24095143E-01
6	-0.99136052E-01	-0.44328199E-00
7	-0.57700423E 00	0.29055967E-00
8	0.59531185E 00	0.72523787E 00

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.80$

$t_o = 3.5343704 + i0.0000087475$

n	Real A_n	Imag A_n
1	-0.20295250E-00	-0.10094626E-01
2	0.17136483E-00	0.51285498E-01
3	-0.12297896E-00	-0.14460298E-00
4	-0.39660082E-01	0.22134382E-00
5	0.27531223E-00	-0.88861914E-01
6	-0.25280646E-00	-0.30294532E-00
7	-0.28909111E-00	0.47559384E-00
8	0.77518813E 00	0.19426705E-00

$q_o = 1.85$

$t_o = 3.7071357 + i0.00048151$

n	Real A_n	Imag A_n
1	-0.18674966E-00	-0.59432433E-02
2	0.15263959E-00	0.32779062E-01
3	-0.12470259E-00	-0.10320780E-00
4	0.18741711E-01	0.18989209E-00
5	0.18957840E-00	-0.15427920E-00
6	-0.30208256E-00	-0.13409275E-00
7	-0.40181538E-02	0.46048797E-00
8	0.60748155E 00	-0.23410610E-00

$q_o = 1.90$

$t_o = 3.8856643 + i0.00025453$

n	Real A_n	Imag A_n
1	-0.17240545E-00	-0.33494539E-02
2	0.13441952E-00	0.19943648E-01
3	-0.11694483E-00	-0.69325527E-01
4	0.54995800E-01	0.14618303E-00
5	0.10251427E-00	-0.17242768E-00
6	-0.26799618E-00	0.49741229E-02
7	0.17773409E-00	0.32196555E-00
8	0.29588679E-00	-0.41601094E-00

Table C-2 (Cont'd)

VALUES OF q_o, t_o, AND A_n

$q_o = 1.95$

$t_o = 4.0698658 + i0.00012912$

n	Real A_n	Imag A_n
1	-0.15981013E-00	-0.18062628E-02
2	0.11780544E-00	0.11554415E-01
3	-0.10400360E-00	-0.43924107E-01
4	0.71440397E-01	0.10648704E-00
5	0.33614268E-01	-0.15098161E-00
6	-0.11988830E-00	0.88272458E-01
7	0.23647793E-00	0.15606169E-00
8	0.24694245E-01	-0.38690419E-00

$q_o = 2.00$

$t_o = 4.2596573 + i0.00006282$

n	Real A_n	Imag A_n
1	-0.14877221E-00	-0.93175689E-03
2	0.10329467E-00	0.63760068E-02
3	-0.69290586E-01	-0.26302908E-01
4	0.43457155E-01	0.71075717E-01
5	-0.12091459E-01	-0.12451768E-00
6	-0.11120382E-00	0.11056965E-00
7	0.21370279E-00	0.25724519E-01
8	-0.12718674E-00	-0.25085050E-00

Appendix D

TABLES OF VALUES OF ROOTS OF $w_1'(t_s) - q\, w_1(t_s) = 0$

In this Appendix we present tables of values for the roots $t_s(q)$ of the equation

$$w_1'(t_s) - q\, w_1(t_s) = 0 \qquad \text{(D-1)}$$

which have been obtained by step-by-step integration of the Riccatti equation

$$\frac{dt}{dq} = \frac{1}{t - q^2} \qquad \text{(D-2)}$$

In each case we have held the phase of the impedance parameter q constant while varying the modulus. In the tables we write

$$t = Q\exp(iP)$$

where P is fixed and Q has been varied. The tables fall into two groups. In the first group, Tables D-1 through D-20, we give values for the roots for s = 1,2,3,4, and 5 for the phase of q equal to 45°, 0°, 90°, and 30°.

In Appendix C we have stressed the importance of the cases in which $P \approx 45^\circ$ and $P \approx 0^\circ$. We presented certain values of the roots and a set of coefficients which could be used in Eq. (C-5) to compute (on a manually-operated desk machine) the numerical value of the roots for values of q which are close to those values of q (for $P = 45^\circ$ and $P = 0^\circ$) which were tabulated. In this Appendix we present further values for the first root $t_1(q)$ for $P = 45^\circ$ and $P = 0^\circ$, as well as values for the next four roots.

For the case of some absorbing surfaces, the phase of q is approximately 90° and therefore we have given tables for the first five roots for this case. In following the literature, the author

has not found any discussion of the case in which the phase of q has been such that $P = 30^\circ$. The fact that no papers have appeared in which this case occurs is most likely due to the fact that only limiting cases have been considered in the theoretical work which has been reported in the literature. However, there are strong arguments for including this case in the present tabulation because it represents a case that lies between the cases $P = 0^\circ$ and $P = 45^\circ$. For example, the case of a perfectly conducting sphere which is coated with a thin (compared to wavelength) dielectric layer leads to a value of the phase of q which is close to 0°. The case of the dielectric interface is described by Eq. (C-3). A study of this case (which corresponds to vertical polarization) reveals that if we express q in terms of its modulus Q and phase P it is found that P lies in the range $45^\circ < P < 90^\circ$. Therefore, if one were to take a metallic sphere surrounded by a dielectric layer and permit the outer radius to remain fixed while permitting the inner core to shrink towards a radius of zero, it should be found that along the way one would observe a continuous transition from the case of $P \approx 0^\circ$ (for the very thin coating) to the range $45^\circ < P < 90^\circ$ for the "thick" coating. The tables for $P = 30^\circ$ which are contained herein will be helpful to anyone who sets out to study this transition.

The second set of tables, Tables D-21 through D-38, describe the loci* of the roots that start from $t_1^\circ \approx 0.5094 + i0.8823$ for $Q = 0$ and then "fan out" as Q increases. If we observe the behavior depicted in Fig. 2-2 we see that for those loci for which $300^\circ < P < 360^\circ$ and $0^\circ < P < 19.2^\circ$ the process of "fanning out" continues to infinity. The reader will observe that Tables D-1, D-6, D-11, and D-16 provide additional tables for this second set of tables for the cases $P = 45^\circ$, 0°, 90°, and 30°.

---

*Tables in this group are for $P = 300^\circ$, 305°, 310°, 315°, 330°, 345°, 350°, 355°, 1°, 5°, 10°, 15°, 20°, 44°, 46°, 60°, 85°, and 120°

Table D-6 deserves some special discussion. This root locus is the one which is associated with the propagation of certain types of surface waves (on thin dielectric-covered surfaces, slightly rough surfaces, etc.). In Section 2 we denoted this root by the subscript "zero." In Eq. (2-25) we gave asymptotic expansions for $t_0(Q)$. Since the imaginary part of this root determines the attenuation of the surface wave, the determination of this imaginary part may be of interest in spite of the fact that it tends rapidly to 0 as Q increases. In the step-by-step integration procedure one finds that one loses accuracy when the root tends toward zero with increasing Q. The degradation is so extreme that in reproducing the computer output we have deleted some of the 8 figures in order to emphasize, in the vicinity of $Q = 2$, that the imaginary part is only known to about four significant figures. The data given in Table D-6b was obtained by means of Eq. (2-25). For $Q > 4$ (or $1/Q < 0.25$), the imaginary part is smaller than 10^{-38} which is the smallest number that the FORTRAN compiler will handle. Because of the extremely rapid variation of this root for large values of Q, we have included in Table D-6b an auxiliary set of numbers which vary suficiently slowly that one can easily interpolate to find the root for intermediate value of $(1/Q)$. The auxiliary functions R(Q) and S(Q) are defined by means of the relation

$$t_1(Q) = t_0(Q) = Q^2 R(Q) + 12Q^2 \exp[-(\tfrac{4}{3}Q^3+1)]S(Q) \qquad (D-3)$$

The functions R(Q) and S(Q) provide good examples of slowly varying functions for which one should have no difficulty in using the theory of trigonometrical interpolation (in particular, the Chebyshev polynomial representation of Eq. (8-42)) to obtain an efficient means of computing $t_0(Q)$ on an electronic computer when Q is large.

The reader will observe that there are two types of computer out-out in the tables. Those numbers which appear in the form

0.50939649E+00

were generated with a single-precision FORTRAN II program which was run on the IBM 7090 electronic computer. Since these operations were carried out with a "word length" of 8 decimal digits, the entries in the tables are generally accurate to no more than 7 significant figures. On the other hand, the entries which appear in the form

.50939649+00

were generated with a double-precision FORTRAN IV program which was run on the UNIVAC 1107.* The results were converted to single precision before output and these entries should be, in most cases, accurate to within a digit in the 8th place.

*The author would like to take this opportunity to point out that he feels that the IBM format is aesthetically superior to the UNIVAC format. The fact that a UNIVAC number requires a field width of 12 characters, whereas the IBM number requires an allowance for 14 characters, permits one to "pack" more output per line of computer output with the UNIVAC 1107.

Table D-1a

THE ROOTS $t_1[Q\exp(i45^\circ)]$

Q	Real t_1	Imag t_1
0.00	.50939648-00	.88230059-00
0.05	.55678568-00	.87077606-00
0.10	.60407776-00	.86158902-00
0.15	.65117583-00	.85471164-00
0.20	.69798310-00	.85011462-00
0.25	.74440256-00	.84776685-00
0.30	.79033688-00	.84763511-00
0.35	.83568820-00	.84968364-00
0.40	.88035798-00	.85387383-00
0.45	.92424691-00	.86016367-00
0.50	.96725480-00	.86850733-00
0.55	.10092806+01	.87885453-00
0.60	.10502224+01	.89114995-00
0.65	.10899776+01	.90533254-00
0.70	.11284431+01	.92133474-00
0.75	.11655156+01	.93908171-00
0.80	.12010925+01	.95849033-00
0.85	.12350718+01	.97946838-00
0.90	.12673541+01	.10019136+01
0.95	.12978431+01	.10257125+01
1.00	.13264475+01	.10507401+01
1.05	.13530829+01	.10768587+01
1.10	.13776745+01	.11039180+01
1.15	.14001590+01	.11317548+01
1.20	.14204884+01	.11601941+01
1.25	.14386330+01	.11890504+01
1.30	.14545842+01	.12181304+01
1.35	.14683577+01	.12472361+01
1.40	.14799952+01	.12761693+01
1.45	.14895654+01	.13047371+01
1.50	.14971640+01	.13327573+01
1.55	.15029108+01	.13600641+01
1.60	.15069468+01	.13865132+01
1.65	.15094288+01	.14119855+01
1.70	.15105235+01	.14363896+01
1.75	.15104014+01	.14596623+01
1.80	.15092312+01	.14817677+01
1.85	.15071744+01	.15026941+01
1.90	.15043815+01	.15224513+01
1.95	.15009896+01	.15410663+01
2.00	.14971207+01	.15585791+01

Table D-1a (Cont'd)

THE ROOTS $t_1[Q\exp(i45^\circ)]$

Q	Real t_1	Imag t_1
2.00	.14971207+01	.15585791+01
2.05	.14928815+01	.15750392+01
2.10	.14883638+01	.15905020+01
2.15	.14836453+01	.16050264+01
2.20	.14787908+01	.16186723+01
2.25	.14738542+01	.16314989+01
2.30	.14688792+01	.16435638+01
2.35	.14639013+01	.16549220+01
2.40	.14589487+01	.16656254+01
2.45	.14540439+01	.16757226+01
2.50	.14492043+01	.16852589+01
2.55	.14444432+01	.16942759+01
2.60	.14397705+01	.17028122+01
2.65	.14351936+01	.17109033+01
2.70	.14307174+01	.17185816+01
2.75	.14263454+01	.17258769+01
2.80	.14220794+01	.17328163+01
2.85	.14179201+01	.17394250+01
2.90	.14138674+01	.17457256+01
2.95	.14099204+01	.17517392+01
3.00	.14060778+01	.17574848+01
3.05	.14023377+01	.17629800+01
3.10	.13986981+01	.17682411+01
3.15	.13951565+01	.17732827+01
3.20	.13917105+01	.17781185+01
3.25	.13883575+01	.17827612+01
3.30	.13850948+01	.17872221+01
3.35	.13819196+01	.17915121+01
3.40	.13788293+01	.17956409+01
3.45	.13758212+01	.17996178+01
3.50	.13728926+01	.18034510+01
3.55	.13700409+01	.18071486+01
3.60	.13672636+01	.18107177+01
3.65	.13645581+01	.18141652+01
3.70	.13619221+01	.18174973+01
3.75	.13593532+01	.18207199+01
3.80	.13568491+01	.18238385+01
3.85	.13544077+01	.18268581+01
3.90	.13520268+01	.18297837+01
3.95	.13497043+01	.18326195+01
4.00	.13474384+01	.18353699+01

Table D-1b

THE ROOTS $t_1[Q\exp(145^\circ)]$

1/Q	Real t_1	Imag t_1
0.00	.11690537+01	.20248604+01
0.01	.11761250+01	.20177886+01
0.02	.11831974+01	.20107123+01
0.03	.11902721+01	.20036268+01
0.04	.11973502+01	.19965279+01
0.05	.12044326+01	.19894107+01
0.06	.12115202+01	.19822706+01
0.07	.12186138+01	.19751030+01
0.08	.12257141+01	.19679031+01
0.09	.12328218+01	.19606660+01
0.10	.12399374+01	.19533867+01
0.11	.12470611+01	.19460603+01
0.12	.12541934+01	.19386815+01
0.13	.12613342+01	.19312450+01
0.14	.12684835+01	.19237455+01
0.15	.12756411+01	.19161772+01
0.16	.12828065+01	.19085346+01
0.17	.12899790+01	.19008116+01
0.18	.12971577+01	.18930022+01
0.19	.13043414+01	.18851002+01
0.20	.13115285+01	.18770991+01
0.21	.13187174+01	.18689923+01
0.22	.13259056+01	.18607731+01
0.23	.13330907+01	.18524344+01
0.24	.13402695+01	.18439690+01
0.25	.13474384+01	.18353698+01
0.26	.13545933+01	.18266292+01
0.27	.13617296+01	.18177397+01
0.28	.13688417+01	.18086935+01
0.29	.13759236+01	.17994830+01
0.30	.13829685+01	.17901004+01
0.31	.13899685+01	.17805380+01
0.32	.13969152+01	.17707884+01
0.33	.14037990+01	.17608440+01
0.34	.14106093+01	.17506978+01
0.35	.14173346+01	.17403433+01
0.36	.14239623+01	.17297743+01
0.37	.14304784+01	.17189854+01
0.38	.14368683+01	.17079720+01
0.39	.14431159+01	.16967306+01
0.40	.14492043+01	.16852587+01

Table D-2a

THE ROOTS $t_2[Q\exp(i145^\circ)]$

Q	Real t_2	Imag t_2
0.00	.16240988+01	.28130215+01
0.05	.16389710+01	.28090750+01
0.10	.16538650+01	.28052077+01
0.15	.16688022+01	.28014266+01
0.20	.16838038+01	.27977397+01
0.25	.16988911+01	.27941562+01
0.30	.17140855+01	.27906864+01
0.35	.17294082+01	.27873422+01
0.40	.17448807+01	.27841372+01
0.45	.17605246+01	.27810871+01
0.50	.17763615+01	.27782098+01
0.55	.17924130+01	.27755260+01
0.60	.18087006+01	.27730592+01
0.65	.18252454+01	.27708366+01
0.70	.18420681+01	.27688890+01
0.75	.18591880+01	.27672519+01
0.80	.18766230+01	.27659655+01
0.85	.18943886+01	.27650756+01
0.90	.19124970+01	.27646335+01
0.95	.19309560+01	.27646975+01
1.00	.19497675+01	.27653318+01
1.05	.19689257+01	.27666074+01
1.10	.19884149+01	.27686017+01
1.15	.20082078+01	.27713972+01
1.20	.20282623+01	.27750804+01
1.25	.20485197+01	.27797394+01
1.30	.20689023+01	.27854607+01
1.35	.20893116+01	.27923246+01
1.40	.21096284+01	.28004010+01
1.45	.21297130+01	.28097428+01
1.50	.21494082+01	.28203803+01
1.55	.21685438+01	.28323163+01
1.60	.21869423+01	.28455209+01
1.65	.22044279+01	.28599295+01
1.70	.22208337+01	.28754424+01
1.75	.22360119+01	.28919274+01
1.80	.22498403+01	.29092241+01
1.85	.22622291+01	.29271516+01
1.90	.22731246+01	.29455166+01
1.95	.22825108+01	.29641222+01
2.00	.22904075+01	.29827774+01

Table D-2a (Cont'd)

THE ROOTS $t_2[Q\exp(i45^\circ)]$

Q	Real t_2	Imag t_2
2.00	.22904075+01	.29827774+01
2.05	.22968672+01	.30013044+01
2.10	.23019697+01	.30195449+01
2.15	.23058152+01	.30373643+01
2.20	.23085181+01	.30546537+01
2.25	.23102000+01	.30713305+01
2.30	.23109842+01	.30873366+01
2.35	.23109907+01	.31026358+01
2.40	.23103331+01	.31172112+01
2.45	.23091161+01	.31310615+01
2.50	.23074342+01	.31441972+01
2.55	.23053713+01	.31566383+01
2.60	.23030005+01	.31684112+01
2.65	.23003852+01	.31795465+01
2.70	.22975796+01	.31900775+01
2.75	.22946294+01	.32000382+01
2.80	.22915733+01	.32094632+01
2.85	.22884433+01	.32183861+01
2.90	.22852661+01	.32268394+01
2.95	.22820639+01	.32348541+01
3.00	.22788545+01	.32424597+01
3.05	.22756530+01	.32496837+01
3.10	.22724708+01	.32565517+01
3.15	.22693178+01	.32630878+01
3.20	.22662015+01	.32693141+01
3.25	.22631279+01	.32752513+01
3.30	.22601014+01	.32809184+01
3.35	.22571258+01	.32863329+01
3.40	.22542034+01	.32915112+01
3.45	.22513363+01	.32964681+01
3.50	.22485254+01	.33012175+01
3.55	.22457714+01	.33057722+01
3.60	.22430748+01	.33101439+01
3.65	.22404353+01	.33143436+01
3.70	.22378527+01	.33183815+01
3.75	.22353263+01	.33222666+01
3.80	.22328554+01	.33260079+01
3.85	.22304392+01	.33296134+01
3.90	.22280765+01	.33330903+01
3.95	.22257665+01	.33364457+01
4.00	.22235079+01	.33396860+01

Table D-2b

THE ROOTS $t_2[Q\exp(i145^\circ)]$

1/Q	Real t_2	Imag t_2
0.00	.20439747+01	.35402680+01
0.01	.20510460+01	.35331956+01
0.02	.20581196+01	.35261153+01
0.03	.20651972+01	.35190192+01
0.04	.20722808+01	.35118992+01
0.05	.20793723+01	.35047472+01
0.06	.20864733+01	.34975548+01
0.07	.20935854+01	.34903137+01
0.08	.21007102+01	.34830151+01
0.09	.21078488+01	.34756500+01
0.10	.21150023+01	.34682090+01
0.11	.21221714+01	.34606826+01
0.12	.21293567+01	.34530606+01
0.13	.21365584+01	.34453326+01
0.14	.21437761+01	.34374876+01
0.15	.21510093+01	.34295138+01
0.16	.21582568+01	.34213993+01
0.17	.21655165+01	.34131312+01
0.18	.21727861+01	.34046961+01
0.19	.21800619+01	.33960796+01
0.20	.21873397+01	.33872671+01
0.21	.21946138+01	.33782428+01
0.22	.22018771+01	.33689900+01
0.23	.22091211+01	.33594918+01
0.24	.22163355+01	.33497299+01
0.25	.22235078+01	.33396859+01
0.26	.22306231+01	.33293405+01
0.27	.22376636+01	.33186742+01
0.28	.22446087+01	.33076674+01
0.29	.22514341+01	.32963005+01
0.30	.22581118+01	.32845550+01
0.31	.22646093+01	.32724132+01
0.32	.22708900+01	.32598596+01
0.33	.22769123+01	.32468816+01
0.34	.22826298+01	.32334702+01
0.35	.22879915+01	.32196216+01
0.36	.22929423+01	.32053382+01
0.37	.22974233+01	.31906301+01
0.38	.23013733+01	.31755161+01
0.39	.23047306+01	.31600253+01
0.40	.23074339+01	.31441972+01

Table D-3a

THE ROOTS $t_3[Q\exp(i145^\circ)]$

Q	Real t_3	Imag t_3
0	.24100496+01	.41743283+01
0.5	.24200711+01	.41716552+01
0.10	.24301028+01	.41690072+01
0.15	.24401549+01	.41663874+01
0.20	.24502376+01	.41637989+01
0.25	.24603611+01	.41612455+01
0.30	.24705358+01	.41587309+01
0.35	.24807721+01	.41562594+01
0.40	.24910806+01	.41538358+01
0.45	.25014721+01	.41514653+01
0.50	.25119575+01	.41491539+01
0.55	.25225481+01	.41469081+01
0.60	.25332553+01	.41447356+01
0.65	.25440909+01	.41426447+01
0.70	.25550667+01	.41406448+01
0.75	.25661951+01	.41387469+01
0.80	.25774883+01	.41369629+01
0.85	.25889591+01	.41353066+01
0.90	.26006201+01	.41337938+01
0.95	.26124838+01	.41324418+01
1.00	.26245628+01	.41312709+01
1.05	.26368689+01	.41303034+01
1.10	.26494135+01	.41295651+01
1.15	.26622069+01	.41290846+01
1.20	.26752578+01	.41288943+01
1.25	.26885727+01	.41290303+01
1.30	.27021552+01	.41295331+01
1.35	.27160052+01	.41304471+01
1.40	.27301175+01	.41318215+01
1.45	.27444809+01	.41337092+01
1.50	.27590764+01	.41361672+01
1.55	.27738757+01	.41392554+01
1.60	.27888399+01	.41430348+01
1.65	.28039176+01	.41475666+01
1.70	.28190434+01	.41529091+01
1.75	.28341380+01	.41591144+01
1.80	.28491071+01	.41662253+01
1.85	.28638435+01	.41742706+01
1.90	.28782289+01	.41832612+01
1.95	.28921374+01	.41931868+01
2.00	.29054411+01	.42040130+01

Table D-3a (Cont'd)

THE ROOTS $t_3[Q\exp(145^\circ)]$

Q	Real t_3	Imag t_3
2.00		
2.05	.29054411+01	.42040130+01
2.10	.29180156+01	.42156801+01
2.15	.29297467+01	.42281035+01
2.20	.29405362+01	.42411766+01
2.25	.29503075+01	.42547756+01
2.30	.29590092+01	.42687640+01
2.35	.29666167+01	.42830008+01
2.40	.29731322+01	.42973460+01
2.45	.29785825+01	.43116672+01
2.50	.29830153+01	.43258446+01
2.55	.29864949+01	.43397739+01
2.60	.29890973+01	.43533688+01
2.65	.29909055+01	.43665612+01
2.70	.29920051+01	.43793006+01
2.75	.29924808+01	.43915524+01
2.80	.29924145+01	.44032964+01
2.85	.29918824+01	.44145237+01
2.90	.29909545+01	.44252352+01
2.95	.29896940+01	.44354391+01
3.00	.29881567+01	.44451491+01
3.05	.29863920+01	.44543829+01
3.10	.29844428+01	.44631606+01
3.15	.29823460+01	.44715042+01
3.20	.29801332+01	.44794361+01
3.25	.29778317+01	.44869789+01
3.30	.29754640+01	.44941549+01
3.35	.29730497+01	.45009860+01
3.40	.29706048+01	.45074927+01
3.45	.29681430+01	.45136950+01
3.50	.29656755+01	.45196117+01
3.55	.29632117+01	.45252604+01
3.60	.29607593+01	.45306577+01
3.65	.29583245+01	.45358189+01
3.70	.29559125+01	.45407587+01
3.75	.29535276+01	.45454903+01
3.80	.29511730+01	.45500264+01
3.85	.29488512+01	.45543785+01
3.90	.29465645+01	.45585576+01
3.95	.29443141+01	.45625737+01
4.00	.29421014+01	.45664360+01
	.29399270+01	.45701533+01

Table D-3b

THE ROOTS $t_3[Q\exp(i145^\circ)]$

1/Q	Real t_3	Imag t_3
0.00	.27602798+01	.47809450+01
0.01	.27673514+01	.47738723+01
0.02	.27744258+01	.47667887+01
0.03	.27815057+01	.47596837+01
0.04	.27885938+01	.47525464+01
0.05	.27956927+01	.47453657+01
0.06	.28028046+01	.47381301+01
0.07	.28099317+01	.47308280+01
0.08	.28170760+01	.47234470+01
0.09	.28242391+01	.47159747+01
0.10	.28314223+01	.47083975+01
0.11	.28386266+01	.47007015+01
0.12	.28458522+01	.46928719+01
0.13	.28530989+01	.46848930+01
0.14	.28603658+01	.46767480+01
0.15	.28676511+01	.46684193+01
0.16	.28749515+01	.46598879+01
0.17	.28822631+01	.46511333+01
0.18	.28895798+01	.46421341+01
0.19	.28968938+01	.46328671+01
0.20	.29041951+01	.46233074+01
0.21	.29114708+01	.46134289+01
0.22	.29187044+01	.46032038+01
0.23	.29258759+01	.45926031+01
0.24	.29329602+01	.45815966+01
0.25	.29399269+01	.45701531+01
0.26	.29467390+01	.45582419+01
0.27	.29533520+01	.45458326+01
0.28	.29597130+01	.45328975+01
0.29	.29657604+01	.45194121+01
0.30	.29714220+01	.45053584+01
0.31	.29766163+01	.44907270+01
0.32	.29812519+01	.44755202+01
0.33	.29852298+01	.44597559+01
0.34	.29884454+01	.44434709+01
0.35	.29907931+01	.44267239+01
0.36	.29921714+01	.44095980+01
0.37	.29924902+01	.43922008+01
0.38	.29916768+01	.43746629+01
0.39	.29896840+01	.43571335+01
0.40	.29864950+01	.43397744+01

Table D-4a

THE ROOTS $t_4[Q\exp(i45^\circ)]$

Q	Real t_4	Imag t_4
0.00	.30816536+01	.53375807+01
0.05	.30894908+01	.53354866+01
0.10	.30973342+01	.53334050+01
0.15	.31051903+01	.53313375+01
0.20	.31130653+01	.53292862+01
0.25	.31209655+01	.53272530+01
0.30	.31288974+01	.53252400+01
0.35	.31368674+01	.53232498+01
0.40	.31448821+01	.53212846+01
0.45	.31529482+01	.53193475+01
0.50	.31610724+01	.53174414+01
0.55	.31692617+01	.53155697+01
0.60	.31775233+01	.53137362+01
0.65	.31858644+01	.53119450+01
0.70	.31942924+01	.53102008+01
0.75	.32028151+01	.53085085+01
0.80	.32114404+01	.53068741+01
0.85	.32201762+01	.53053040+01
0.90	.32290308+01	.53038053+01
0.95	.32380128+01	.53023862+01
1.00	.32471307+01	.53010557+01
1.05	.32563933+01	.52998238+01
1.10	.32658093+01	.52987021+01
1.15	.32753877+01	.52977034+01
1.20	.32851370+01	.52968420+01
1.25	.32950658+01	.52961341+01
1.30	.33051821+01	.52955978+01
1.35	.3315494+01	.52952533+01
1.40	.33260062+01	.52951234+01
1.45	.33367257+01	.52952331+01
1.50	.33476555+01	.52956106+01
1.55	.33587968+01	.52962867+01
1.60	.33701480+01	.52972952+01
1.65	.33817039+01	.52986734+01
1.70	.33934546+01	.53004608+01
1.75	.34053849+01	.53026997+01
1.80	.34174726+01	.53054345+01
1.85	.34296882+01	.53087106+01
1.90	.34419933+01	.53125732+01
1.95	.34543399+01	.53170655+01
2.00	.34666695+01	.53222269+01

Table D-4a (Cont'd)

THE ROOTS $t_4[Q\exp(i145^\circ)]$

Q	Real t_4	Imag t_4
2.00	.34666695+01	.53222269+01
2.05	.34789138+01	.53280903+01
2.10	.34909943+01	.53346793+01
2.15	.35028243+01	.53420057+01
2.20	.35143109+01	.53500666+01
2.25	.35253580+01	.53588418+01
2.30	.35358710+01	.53682936+01
2.35	.35457604+01	.53783657+01
2.40	.35549467+01	.53889853+01
2.45	.35633641+01	.54000645+01
2.50	.35709644+01	.54115051+01
2.55	.35777178+01	.54232025+01
2.60	.35836150+01	.54350505+01
2.65	.35886648+01	.54469458+01
2.70	.35928931+01	.54587923+01
2.75	.35963402+01	.54705039+01
2.80	.35990566+01	.54820062+01
2.85	.36011006+01	.54932383+01
2.90	.36025343+01	.55041519+01
2.95	.36034214+01	.55147113+01
3.00	.36038246+01	.55248919+01
3.05	.36038040+01	.55346790+01
3.10	.36034160+01	.55440660+01
3.15	.36027125+01	.55530529+01
3.20	.36017404+01	.55616453+01
3.25	.36005419+01	.55698524+01
3.30	.35991541+01	.55776863+01
3.35	.35976100+01	.55851613+01
3.40	.35959379+01	.55922930+01
3.45	.35941628+01	.55990975+01
3.50	.35923060+01	.56055910+01
3.55	.35903858+01	.56117901+01
3.60	.35884178+01	.56177107+01
3.65	.35864154+01	.56233682+01
3.70	.35843900+01	.56287776+01
3.75	.35823511+01	.56339531+01
3.80	.35803068+01	.56389079+01
3.85	.35782639+01	.56436550+01
3.90	.35762281+01	.56482062+01
3.95	.35742041+01	.56525727+01
4.00	.35721960+01	.56567650+01

Table D-4b

THE ROOTS $t_4[Q\exp(i145^\circ)]$

1/Q	Real t_4	Imag t_4
0.00	.33933540+01	.58774616+01
0.01	.34004257+01	.58703883+01
0.02	.34075008+01	.58633019+01
0.03	.34145828+01	.58561892+01
0.04	.34216749+01	.58490366+01
0.05	.34287801+01	.58418303+01
0.06	.34359016+01	.58345562+01
0.07	.34430417+01	.58271995+01
0.08	.34502030+01	.58197448+01
0.09	.34573872+01	.58121757+01
0.10	.34645956+01	.58044751+01
0.11	.34718291+01	.57966246+01
0.12	.34790876+01	.57886048+01
0.13	.34863701+01	.57803944+01
0.14	.34936742+01	.57719711+01
0.15	.35009962+01	.57633104+01
0.16	.35083304+01	.57543860+01
0.17	.35156687+01	.57451694+01
0.18	.35230001+01	.57356299+01
0.19	.35303100+01	.57257344+01
0.20	.35375794+01	.57154474+01
0.21	.35447838+01	.57047311+01
0.22	.35518919+01	.56935459+01
0.23	.35588645+01	.56818507+01
0.24	.35656525+01	.56696038+01
0.25	.35721957+01	.56567649+01
0.26	.35784206+01	.56432968+01
0.27	.35842390+01	.56291687+01
0.28	.35895470+01	.56143606+01
0.29	.35942251+01	.55988681+01
0.30	.35981397+01	.55827087+01
0.31	.36011472+01	.55659289+01
0.32	.36031002+01	.55486096+01
0.33	.36038586+01	.55308716+01
0.34	.36033016+01	.55128755+01
0.35	.36013412+01	.54948185+01
0.36	.35979360+01	.54769249+01
0.37	.35930993+01	.54594302+01
0.38	.35869020+01	.54425646+01
0.39	.35794684+01	.54265340+01
0.40	.35709650+01	.54115054+01

Table D-5a

THE ROOTS $t_5[Q\exp(i145^0)]$

Q	Real t_5	Imag t_5
0.00	.36860886+01	.63844927+01
0.05	.36926405+01	.63827407+01
0.10	.36991968+01	.63809961+01
0.15	.37057620+01	.63792601+01
0.20	.37123404+01	.63775342+01
0.25	.37189365+01	.63758196+01
0.30	.37255549+01	.63741177+01
0.35	.37322000+01	.63724302+01
0.40	.37388765+01	.63707587+01
0.45	.37455890+01	.63691048+01
0.50	.37523423+01	.63674708+01
0.55	.37591412+01	.63658585+01
0.60	.37659907+01	.63642703+01
0.65	.37728959+01	.63627087+01
0.70	.37798618+01	.63611765+01
0.75	.37868939+01	.63596769+01
0.80	.37939977+01	.63582131+01
0.85	.38011787+01	.63567889+01
0.90	.38084427+01	.63554086+01
0.95	.38157958+01	.63540766+01
1.00	.38232439+01	.63527982+01
1.05	.38307933+01	.63515790+01
1.10	.38384503+01	.63504255+01
1.15	.38462215+01	.63493447+01
1.20	.38541135+01	.63483445+01
1.25	.38621330+01	.63474337+01
1.30	.38702864+01	.63466223+01
1.35	.38785806+01	.63459210+01
1.40	.38870218+01	.63453422+01
1.45	.38956164+01	.63448997+01
1.50	.39043701+01	.63446084+01
1.55	.39132879+01	.63444854+01
1.60	.39223745+01	.63445492+01
1.65	.39316329+01	.63448206+01
1.70	.39410650+01	.63453225+01
1.75	.39506707+01	.63460796+01
1.80	.39604478+01	.63471194+01
1.85	.39703908+01	.63484710+01
1.90	.39804910+01	.63501662+01
1.95	.39907353+01	.63522380+01
2.00	.40011056+01	.63547213+01

Table D-5a (Cont'd)

THE ROOTS $t_5[Q\exp(i145^\circ)]$

Q	Real t_5	Imag t_5
2.00	.40011056+01	.63547213+01
2.05	.40115779+01	.63576511+01
2.10	.40221217+01	.63610629+01
2.15	.40326992+01	.63649902+01
2.20	.40432654+01	.63694637+01
2.25	.40537669+01	.63745099+01
2.30	.40641435+01	.63801480+01
2.35	.40743282+01	.63863890+01
2.40	.40842489+01	.63932331+01
2.45	.40938309+01	.64006682+01
2.50	.41029990+01	.64086688+01
2.55	.41116813+01	.64171955+01
2.60	.41198119+01	.64261959+01
2.65	.41273343+01	.64356054+01
2.70	.41342041+01	.64453506+01
2.75	.41403906+01	.64553509+01
2.80	.41458779+01	.64655232+01
2.85	.41506647+01	.64757849+01
2.90	.41547638+01	.64860566+01
2.95	.41581993+01	.64962654+01
3.00	.41610056+01	.65063466+01
3.05	.41632242+01	.65162447+01
3.10	.41649015+01	.65259142+01
3.15	.41660864+01	.65353192+01
3.20	.41668285+01	.65444330+01
3.25	.41671766+01	.65532370+01
3.30	.41671775+01	.65617201+01
3.35	.41668751+01	.65698769+01
3.40	.41663099+01	.65777073+01
3.45	.41655190+01	.65852149+01
3.50	.41645354+01	.65924066+01
3.55	.41633891+01	.65992913+01
3.60	.41621061+01	.66058796+01
3.65	.41607098+01	.66121833+01
3.70	.41592205+01	.66182145+01
3.75	.41576556+01	.66239861+01
3.80	.41560305+01	.66295104+01
3.85	.41543584+01	.66348001+01
3.90	.41526508+01	.66398673+01
3.95	.41509172+01	.66447236+01
4.00	.41491662+01	.66493806+01

Table D-5b

THE ROOTS $t_5[Q\exp(145^\circ)]$

1/Q	Real t_5	Imag t_5
0.00	.39720668+01	.68798215+01
0.01	.39791385+01	.68727479+01
0.02	.39862143+01	.68656589+01
0.03	.39932982+01	.68585389+01
0.04	.40003939+01	.68513723+01
0.05	.40075051+01	.68441426+01
0.06	.40146352+01	.68368331+01
0.07	.40217871+01	.68294260+01
0.08	.40289634+01	.68219028+01
0.09	.40361662+01	.68142437+01
0.10	.40433968+01	.68064277+01
0.11	.40506555+01	.67984322+01
0.12	.40579417+01	.67902327+01
0.13	.40652529+01	.67818027+01
0.14	.40725851+01	.67731135+01
0.15	.40799318+01	.67641335+01
0.16	.40872837+01	.67548288+01
0.17	.40946271+01	.67451619+01
0.18	.41019441+01	.67350923+01
0.19	.41092104+01	.67245764+01
0.20	.41163943+01	.67135672+01
0.21	.41234542+01	.67020154+01
0.22	.41303373+01	.66898700+01
0.23	.41369762+01	.66770799+01
0.24	.41432869+01	.66635972+01
0.25	.41491658+01	.66493805+01
0.26	.41544880+01	.66344012+01
0.27	.41591064+01	.66186509+01
0.28	.41628540+01	.66021506+01
0.29	.41655491+01	.65849617+01
0.30	.41670065+01	.65671948+01
0.31	.41670539+01	.65490173+01
0.32	.41655522+01	.65306525+01
0.33	.41624177+01	.65123713+01
0.34	.41576401+01	.64944722+01
0.35	.41512926+01	.64772541+01
0.36	.41435266+01	.64609867+01
0.37	.41345569+01	.64458855+01
0.38	.41246374+01	.64320956+01
0.39	.41140337+01	.64196884+01
0.40	.41029995+01	.64086684+01

Table D-6a

THE ROOTS t_1 [Qexp(10°)]

Q	Real t_1	Imag t_1
0.00	.5093648+00	.8823058-00
0.05	.53512579-00	.83981264-00
0.10	.56326949-00	.79741255-00
0.15	.59387761-00	.75518902-00
0.20	.62700049-00	.71323170-00
0.25	.66268905-00	.67163127-00
0.30	.70099533-00	.63047944-00
0.35	.74197288-00	.58986899-00
0.40	.78567731-00	.54989382-00
0.45	.83216679-00	.51064891-00
0.50	.88150267-00	.47223038-00
0.55	.93375009-00	.43473542-00
0.60	.98897874-00	.39826214-00
0.65	.10472636+01	.36290947-00
0.70	.11086859+01	.32877682-00
0.75	.11733337+01	.29596369-00
0.80	.12413039+01	.26456921-00
0.85	.13127021+01	.2346911[illegible]-00
0.90	.13876447+01	.20642491-00
0.95	.14662595+01	.17986219-00
1.00	.15486874+01	.1550889[illegible]-00
1.05	.16350828+01	.13218309-00
1.10	.17256153+01	.11121146-00
1.15	.18204682+01	.9222614 -01
1.20	.19198395+01	.7526016 -01
1.25	.20239391+01	.6032268 -01
1.30	.21329851+01	.4739397 -01
1.35	.22471991+01	.3642072 -01
1.40	.2366797[illegible]+01	.2731259 -01
1.45	.24919823+01	.1994098 -01
1.50	.26229306+01	.1414122 -01
1.55	.27597834+01	.971885 -02
1.60	.29026376+01	.646011 -02
1.65	.30515411+01	.414353 -02
1.70	.32064956+01	.256434 -02
1.75	.33674616+01	.152717 -02
1.80	.35343704+01	.87475 -03
1.85	.37071357+01	.48151 -03
1.90	.38856643+01	.25453 -03
1.95	.40698658+01	.12912 -03
2.00	.42596573+01	.6282 -04

Table D-6a (Cont'd)

THE ROOTS $t_1[Q\exp(i10^\circ)]$

Q	Real t_1	Imag t_1
2.00	.42596573+01	.6282 -04
2.05	.44549658+01	.2929 -04
2.10	.46557305+01	.1308 -04
2.15	.48618994+01	.5590 -04
2.20	.50734290+01	.22844 -05
2.25	.52902835+01	.89183 -06
2.30	.55124316+01	.33235 -06
2.35	.57398493+01	.11811 -06
2.40	.59725127+01	.400085 -07
2.45	.62104024+01	.128998 -07
2.50	.64535013+01	.395597 -08
2.55	.67017940+01	.115280 -08
2.60	.69552670+01	.318918 -09
2.65	.72139081+01	.836787 -10
2.70	.74777062+01	.208041 -10
2.75	.77466512+01	.4896299 -11
2.80	.80207342+01	.10898139-11
2.85	.82999464+01	.22918461-12
2.90	.85842806+01	.45493291-13
2.95	.88737298+01	.85156374-14
3.00	.91682874+01	.15016693-14

Table D-6b

THE ROOTS $t_1[Q\exp(i0^\circ)]$

1/Q	Real t_1	Imag t_1	R(Q)	S(Q)
0.01	10000.005		1.0000005	0.999999942
0.02	2500.0100		1.0000040	0.99999534
0.03	1111.1261		1.0000134	0.99998426
0.04	625.02000	$< 10^{-38}$	1.0000320	0.99996267
0.05	400.02500		1.0000625	0.99992709
0.06	277.80778		1.0001080	0.99987398
0.07	204.11664		1.0001715	0.99979986
0.08	156.29000		1.0002560	0.99970122
0.09	123.50180		1.0003645	0.99957450
0.10	100.05001		1.0005001	0.99941620
0.11	82.699645		1.0006656	0.99922274
0.12	69.504471		1.0008644	0.99899058
0.13	59.236635		1.0010990	0.99871612
0.14	51.090457		1.0013729	0.99839574
0.15	44.519509		1.0016889	0.99802580
0.16	39.142581		1.0020500	0.99760263
0.17	34.687180		1.0024595	0.997122148
0.18	30.954330		1.0029203	0.99658161
0.19	27.795995		1.0034354	0.99597621
0.20	25.100201		1.0040081	0.99530235
0.21	22.780982		1.0046413	0.99455611
0.22	20.771454		1.0053383	0.993733345
0.23	19.018946		1.0061022	0.99283023
0.24	17.481533		1.0069363	0.99184224
0.25	16.125497	1.0163158×10^{-36}	1.0078435	0.99076514
0.26	14.923483	1.2196076×10^{-32}	1.0088275	0.98959443
0.27	13.853102	3.7987928×10^{-29}	1.0098911	0.98832554
0.28	12.895892	3.8751063×10^{-26}	1.0110379	0.98695366
0.29	12.036519	1.5593392×10^{-23}	1.0122712	0.98547384
0.30	11.262160	2.8760263×10^{-21}	1.0135943	0.98388091
0.31	10.562027	2.7466701×10^{-19}	1.0150108	0.982169145
0.32	9.9269933	1.5007935×10^{-17}	1.0165241	0.98033381
0.33	9.3492912	5.0936593×10^{-16}	1.0181378	0.97836800
0.34	8.8222793	1.1495057×10^{-14}	1.0198554	0.97626571
0.35	8.3402523	1.8256191×10^{-13}	1.0216808	0.97402023
0.36	7.8982858	2.1398222×10^{-12}	1.0236178	0.97162437
0.37	7.4921124	1.9266418×10^{-11}	1.0256701	0.96907046
0.38	7.1180189	1.3784520×10^{-10}	1.0278419	0.96635021
0.39	6.7727626	8.0658878×10^{-10}	1.0301372	0.96345460
0.40	6.4535012	3.9559756×10^{-9}	1.0325602	0.96037380

$$\{t_1[Q\exp(i0^\circ)] = Q^2R(Q) + 12Q^2\exp[-\tfrac{4}{3} - 1]S(Q)\}$$

Table D-7a

THE ROOTS $t_2[Q\exp(i0^\circ)]$

Q	Real t_2	Imag t_2
0.00	•16240988+01	•28130215+01
0.05	•16318500+01	•27996875+01
0.10	•16396229+01	•27863333+01
0.15	•16474067+01	•27729385+01
0.20	•16553502+01	•27594817+01
0.25	•16632023+01	•27459411+01
0.30	•16711912+01	•27322935+01
0.35	•16791245+01	•27185158+01
0.40	•16870426+01	•27045833+01
0.45	•16949492+01	•26904707+01
0.50	•17028063+01	•26761520+01
0.55	•17106187+01	•26616002+01
0.60	•17183521+01	•26467875+01
0.65	•17259415+01	•26316852+01
0.70	•17333450+01	•26162645+01
0.75	•17407587+01	•26004960+01
0.80	•17478774+01	•25843503+01
0.85	•17547540+01	•25677992+01
0.90	•17613570+01	•25508158+01
0.95	•17675692+01	•25333756+01
1.00	•17733890+01	•25154589+01
1.05	•17787097+01	•24970514+01
1.10	•17834593+01	•24781477+01
1.15	•17875463+01	•24587540+01
1.20	•17908516+01	•24388913+01
1.25	•17932512+01	•24186000+01
1.30	•17947232+01	•23979435+01
1.35	•17950087+01	•23770119+01
1.40	•17942070+01	•23559242+01
1.45	•17920085+01	•23346290+01
1.50	•17885916+01	•23139003+01
1.55	•17837388+01	•22930309+01
1.60	•17776510+01	•22733205+01
1.65	•17702621+01	•22540620+01
1.70	•17617956+01	•22357265+01
1.75	•17523491+01	•22184510+01
1.80	•17421203+01	•22025304+01
1.85	•17312959+01	•21874151+01
1.90	•17200526+01	•21737145+01
1.95	•17086091+01	•21612025+01
2.00	•16969774+01	•21498270+01

Table D-7a (Cont'd)

THE ROOTS $t_2[Q\exp(10^{\circ})]$

Q	Real t_2	Imag t_2
2.00	.1696974+01	.2149827 0+01
2.05	.16854026+01	.21395177+01
2.10	.1673941 4+01	.21301945+01
2.15	.16626686+01	.21217728+01
2.20	.16516397+01	.21141686+01
2.25	.1640894 0+01	.2107301 0+01
2.30	.16304577+01	.21010945+01
2.35	.16203465+01	.20954796+01
2.40	.1610568 3+01	.2090393 1+01
2.45	.16011248+01	.20857783+01
2.50	.15920132+01	.20815848+01
2.55	.1583227 5+01	.2077767 3+01
2.60	.1574759 4+01	.20742860+01
2.65	.15665990+01	.20711056+01
2.70	.1558735 5+01	.2068194 9+01
2.75	.15511573+01	.20655261+01
2.80	.15438530+01	.20630749+01
2.85	.1536810 7+01	.2060819 6+01
2.90	.15300190+01	.20587411+01
2.95	.15234666+01	.20568223+01
3.00	.1517142 7+01	.2055048 2+01
3.05	.15110366+01	.20534053+01
3.10	.1505138 3+01	.2051881 6+01
3.15	.1499438 3+01	.2050466 4+01
3.20	.14939274+01	.20491502+01
3.25	.14885968+01	.20479243+01
3.30	.1483438 2+01	.2046781 0+01
3.35	.14784439+01	.20457134+01
3.40	.1473606 4+01	.2044715 4+01
3.45	.1468918 7+01	.2043781 1+01
3.50	.14643740+01	.20429057+01
3.55	.14599661+01	.20420844+01
3.60	.1455689 1+01	.2041313 1+01
3.65	.14515372+01	.20405879+01
3.70	.14475052+01	.20399055+01
3.75	.1443588 0+01	.2039262 7+01
3.80	.14397807+01	.20386565+01
3.85	.14360789+01	.20380845+01
3.90	.1432478 3+01	.2037544 2+01
3.95	.14289748+01	.20370334+01
4.00	.1425664 4+01	.2036550 0+01

Table D-7b

THE ROOTS $t_2[Q\exp(10^\circ)]$

1/Q	Real t_2	Imag t_2
0.00	•20439747+01	•35402680+01
0.01	•20539723+01	•35402692+01
0.02	•20639801+01	•35402775+01
0.03	•20739932+01	•35402999+01
0.04	•20840117+01	•35403439+01
0.05	•20940359+01	•35404164+01
0.06	•21041256+01	•35405252+01
0.07	•21142115+01	•35406778+01
0.08	•21243242+01	•35408821+01
0.09	•21344778+01	•35411461+01
0.10	•21446841+01	•35414784+01
0.11	•21549466+01	•35418878+01
0.12	•21652709+01	•35423835+01
0.13	•21756742+01	•35429751+01
0.14	•21861652+01	•35436727+01
0.15	•21967547+01	•35444871+01
0.16	•22074565+01	•35454294+01
0.17	•22182883+01	•35465116+01
0.18	•22292648+01	•35477462+01
0.19	•22403983+01	•35491463+01
0.20	•22516990+01	•35507260+01
0.21	•22631791+01	•35524998+01
0.22	•22748579+01	•35544830+01
0.23	•22867108+01	•35566917+01
0.24	•22987551+01	•35591425+01
0.25	•23110330+01	•35618525+01
0.26	•23235850+01	•35648393+01
0.27	•23363491+01	•35681206+01
0.28	•23493107+01	•35717142+01
0.29	•23624927+01	•35756373+01
0.30	•23758151+01	•35799067+01
0.31	•23893945+01	•35845379+01
0.32	•24032445+01	•35895443+01
0.33	•24173755+01	•35949374+01
0.34	•24317949+01	•36007255+01
0.35	•24465172+01	•36069129+01
0.36	•24615549+01	•36134996+01
0.37	•24769184+01	•36204804+01
0.38	•24926177+01	•36278445+01
0.39	•25086730+01	•36355749+01
0.40	•25250962+01	•36436481+01

Table D-8a

THE ROOTS $t_3[Q\exp(i10^\circ)]$

Q	Real t_3	Imag t_3
0.00	.24100496+01	.41743283+01
0.05	.24152464+01	.41653433+01
0.10	.24204864+01	.41563491+01
0.15	.24256681+01	.41473360+01
0.20	.24309163+01	.41382948+01
0.25	.24361515+01	.41292158+01
0.30	.24413801+01	.41200889+01
0.35	.24465981+01	.41109043+01
0.40	.24517994+01	.41016517+01
0.45	.24569773+01	.40923210+01
0.50	.24621245+01	.40829017+01
0.55	.24672332+01	.40733832+01
0.60	.24722949+01	.40637550+01
0.65	.24773000+01	.40540062+01
0.70	.24822383+01	.40441262+01
0.75	.24870984+01	.40341043+01
0.80	.24918675+01	.40239301+01
0.85	.24965315+01	.40135931+01
0.90	.25010748+01	.40030836+01
0.95	.25054803+01	.39923923+01
1.00	.25097288+01	.39815106+01
1.05	.25137994+01	.39704312+01
1.10	.25176691+01	.39591479+01
1.15	.25213125+01	.39476566+01
1.20	.25247027+01	.39359550+01
1.25	.25278100+01	.39240438+01
1.30	.25306032+01	.39119271+01
1.35	.25330493+01	.38996127+01
1.40	.25351140+01	.38871125+01
1.45	.25367624+01	.38744444+01
1.50	.25379595+01	.38616313+01
1.55	.25386717+01	.38487026+01
1.60	.25388677+01	.38356940+01
1.65	.25385197+01	.38226476+01
1.70	.25376050+01	.38096116+01
1.75	.25361081+01	.37966395+01
1.80	.25340210+01	.37837891+01
1.85	.25313450+01	.37711204+01
1.90	.25280915+01	.37586945+01
1.95	.25242817+01	.37465703+01
2.00	.25199466+01	.37346039+01

Table D-8a (Cont'd)

THE ROOTS $t_3[Q\exp(i0^\circ)]$

Q	Real t_3	Imag t_3
2.00	.25199466+01	.37348039+01
2.05	.25151267+01	.37234449+01
2.10	.25098630+01	.37125356+01
2.15	.25042193+01	.37021101+01
2.20	.24982412+01	.36921928+01
2.25	.24919868+01	.36827991+01
2.30	.24855126+01	.36739352+01
2.35	.24786775+01	.36655996+01
2.40	.24715287+01	.36577835+01
2.45	.24640998+01	.36504726+01
2.50	.24563444+01	.36436482+01
2.55	.24482591+01	.36372881+01
2.60	.24398536+01	.36313681+01
2.65	.24311045+01	.36258629+01
2.70	.24220761+01	.36207464+01
2.75	.24127416+01	.36159929+01
2.80	.24030826+01	.36115771+01
2.85	.23930861+01	.36074746+01
2.90	.23827434+01	.36036624+01
2.95	.23720478+01	.36001185+01
3.00	.23609873+01	.35968225+01
3.05	.23495512+01	.35937551+01
3.10	.23377286+01	.35908986+01
3.15	.23255095+01	.35882365+01
3.20	.23128846+01	.35857537+01
3.25	.22998450+01	.35834362+01
3.30	.22863827+01	.35812712+01
3.35	.22724903+01	.35792468+01
3.40	.22581611+01	.35773525+01
3.45	.22433888+01	.35755780+01
3.50	.22281676+01	.35739145+01
3.55	.22124923+01	.35723536+01
3.60	.21963579+01	.35708877+01
3.65	.21797600+01	.35695098+01
3.70	.21626944+01	.35682135+01
3.75	.21451574+01	.35669930+01
3.80	.21271457+01	.35658427+01
3.85	.21086562+01	.35647580+01
3.90	.20896862+01	.35637341+01
3.95	.20702335+01	.35627668+01
4.00	.20502961+01	.35618524+01

Table D-8b

THE ROOTS $t_3[Q\exp(i10^\circ)]$

1/Q	Real t_3	Imag t_3
0.00	.27602798+01	.47809450+01
0.01	.27702808+01	.47809466+01
0.02	.27802873+01	.47809578+01
0.03	.27903048+01	.47809882+01
0.04	.28003391+01	.47810475+01
0.05	.28103955+01	.47811458+01
0.06	.28204795+01	.47812934+01
0.07	.28305963+01	.47815008+01
0.08	.28407510+01	.47817788+01
0.09	.28509484+01	.47821390+01
0.10	.28611930+01	.47825933+01
0.11	.28714890+01	.47831545+01
0.12	.28818399+01	.47838358+01
0.13	.28922486+01	.47846512+01
0.14	.29027174+01	.47856158+01
0.15	.29132476+01	.47867452+01
0.16	.29238396+01	.47880560+01
0.17	.29344922+01	.47895660+01
0.18	.29452030+01	.47912937+01
0.19	.29559677+01	.47932585+01
0.20	.29667800+01	.47954807+01
0.21	.29776310+01	.47979812+01
0.22	.29885095+01	.48007817+01
0.23	.29994009+01	.48039038+01
0.24	.30102872+01	.48073689+01
0.25	.30211471+01	.48111980+01
0.26	.30319548+01	.48154106+01
0.27	.30426807+01	.48200244+01
0.28	.30532909+01	.48250545+01
0.29	.30637477+01	.48305114+01
0.30	.30740093+01	.48364016+01
0.31	.30840313+01	.48427255+01
0.32	.30937674+01	.48494770+01
0.33	.31031700+01	.48566422+01
0.34	.31121927+01	.48641996+01
0.35	.31207911+01	.48721197+01
0.36	.31289253+01	.48803654+01
0.37	.31365609+01	.48888928+01
0.38	.31436708+01	.48976528+01
0.39	.31502359+01	.49065921+01
0.40	.31562462+01	.49156556+01

Table D-9a

THE ROOTS $t_4[Q\exp(i10^\circ)]$

Q	Real t_4	Imag t_4
0.00	.30816536+01	.53375807+01
0.05	.30857147+01	.53305541+01
0.10	.30897832+01	.53235219+01
0.15	.30938558+01	.53164781+01
0.20	.30979293+01	.53094171+01
0.25	.31020003+01	.53023331+01
0.30	.31060653+01	.52952201+01
0.35	.31101207+01	.52880721+01
0.40	.31141628+01	.52808832+01
0.45	.31181876+01	.52736473+01
0.50	.31221908+01	.52663583+01
0.55	.31261679+01	.52590100+01
0.60	.31301140+01	.52515963+01
0.65	.31340239+01	.52441109+01
0.70	.31378919+01	.52365478+01
0.75	.31417118+01	.52289008+01
0.80	.31454769+01	.52211639+01
0.85	.31491798+01	.52133310+01
0.90	.31528126+01	.52053963+01
0.95	.31563666+01	.51973543+01
1.00	.31598324+01	.51891996+01
1.05	.31631996+01	.51809274+01
1.10	.31664570+01	.51725331+01
1.15	.31695927+01	.51640130+01
1.20	.31725933+01	.51553639+01
1.25	.31754452+01	.51465837+01
1.30	.31781331+01	.51376711+01
1.35	.31806412+01	.51286265+01
1.40	.31829525+01	.51194513+01
1.45	.31850494+01	.51101491+01
1.50	.31869137+01	.51007251+01
1.55	.31885267+01	.50911868+01
1.60	.31898693+01	.50815443+01
1.65	.31909228+01	.50718102+01
1.70	.31916689+01	.50619999+01
1.75	.31920904+01	.50521316+01
1.80	.31921714+01	.50422268+01
1.85	.31918981+01	.50323094+01
1.90	.31912594+01	.50224061+01
1.95	.31902475+01	.50125460+01
2.00	.31888577+01	.50027600+01

Table D-9a (Cont'd)

THE ROOTS $t_4[Q\exp(i10^\circ)]$

Q	Real t_4	Imag t_4
2.00	.31888577+01	.50027600+01
2.05	.31870901+01	.49930800+01
2.10	.31849487+01	.49835385+01
2.15	.31824422+01	.49741680+01
2.20	.31795837+01	.49649996+01
2.25	.31763905+01	.49560624+01
2.30	.31728840+01	.49473832+01
2.35	.31690884+01	.49389853+01
2.40	.31650309+01	.49308885+01
2.45	.31607402+01	.49231081+01
2.50	.31562463+01	.49156556+01
2.55	.31515795+01	.49085385+01
2.60	.31467695+01	.49017597+01
2.65	.31418452+01	.48953193+01
2.70	.31368340+01	.48892136+01
2.75	.31317614+01	.48834363+01
2.80	.31266510+01	.48779788+01
2.85	.31215240+01	.48728306+01
2.90	.31163994+01	.48679800+01
2.95	.31112940+01	.48634139+01
3.00	.31062222+01	.48591190+01
3.05	.31011967+01	.48550814+01
3.10	.30962279+01	.48512871+01
3.15	.30913246+01	.48477224+01
3.20	.30864942+01	.48443736+01
3.25	.30817425+01	.48412278+01
3.30	.30770738+01	.48382722+01
3.35	.30724920+01	.48354948+01
3.40	.30679993+01	.48328840+01
3.45	.30635975+01	.48304290+01
3.50	.30592877+01	.48281197+01
3.55	.30550703+01	.48259464+01
3.60	.30509451+01	.48238999+01
3.65	.30469117+01	.48219720+01
3.70	.30429693+01	.48201548+01
3.75	.30391166+01	.48184408+01
3.80	.30353524+01	.48168234+01
3.85	.30316751+01	.48152960+01
3.90	.30280830+01	.48138530+01
3.95	.30245743+01	.48124886+01
4.00	.30211472+01	.48111979+01

Table D-9b

THE ROOTS $t_4[Q\exp(i10^\circ)]$

1/Q	Real t_4	Imag t_4
0.00	•33933540+01	•58774616+01
0.01	•34033551+01	•58774635+01
0.02	•34133631+01	•58774773+01
0.03	•34233847+01	•58775146+01
0.04	•34334265+01	•58775878+01
0.05	•34434955+01	•58777090+01
0.06	•34535979+01	•58778910+01
0.07	•34637401+01	•58781473+01
0.08	•34739277+01	•58784914+01
0.09	•34841663+01	•58789381+01
0.10	•34944607+01	•58795028+01
0.11	•35048151+01	•58802017+01
0.12	•35152327+01	•58810521+01
0.13	•35257157+01	•58820723+01
0.14	•35362649+01	•58832820+01
0.15	•35468797+01	•58847018+01
0.16	•35575573+01	•58863534+01
0.17	•35682929+01	•58882601+01
0.18	•35790791+01	•58904456+01
0.19	•35899048+01	•58929350+01
0.20	•36007562+01	•58957536+01
0.21	•36116149+01	•58989267+01
0.22	•36224581+01	•59024795+01
0.23	•36332583+01	•59064355+01
0.24	•36439829+01	•59108163+01
0.25	•36545938+01	•59156404+01
0.26	•36650480+01	•59209213+01
0.27	•36752980+01	•59266670+01
0.28	•36852925+01	•59328786+01
0.29	•36949777+01	•59395482+01
0.30	•37042991+01	•59466589+01
0.31	•37132034+01	•59541842+01
0.32	•37216411+01	•59620877+01
0.33	•37295680+01	•59703237+01
0.34	•37369483+01	•59788392+01
0.35	•37437554+01	•59875751+01
0.36	•37499733+01	•59964689+01
0.37	•37555969+01	•60054573+01
0.38	•37606318+01	•60144786+01
0.39	•37650929+01	•60234742+01
0.40	•37690035+01	•60323913+01

+119

Table D-10a

THE ROOTS $t_5[Q\exp(i10^\circ)]$

Q	Real t_5	Imag t_5
0.00	.36860886+01	.63844927+01
0.05	.3689482 5+01	.63786186+01
0.10	.3692880 3+01	.6372740 3+01
0.15	.3696279 7+01	.6366854 0+01
0.20	.3699678 6+01	.6360955 8+01
0.25	.3703074 4+01	.6355041 4+01
0.30	.3706464 8+01	.6349106 9+01
0.35	.3709847 3+01	.6343148 1+01
0.40	.3713219 2+01	.6337160 9+01
0.45	.3716577 8+01	.6331141 1+01
0.50	.3719920 3+01	.6325084 6+01
0.55	.3723243 7+01	.6318987 1+01
0.60	.3726544 6+01	.6312844 4+01
0.65	.3729819 6+01	.6306652 4+01
0.70	.3733065 1+01	.6300406 9+01
0.75	.3736276 9+01	.6294103 6+01
0.80	.3739450 9+01	.6287738 4+01
0.85	.3742582 4+01	.6281307 2+01
0.90	.3745666 4+01	.6274806 1+01
0.95	.3748697 6+01	.6268231 1+01
1.00	.3751670 1+01	.6261578 4+01
1.05	.3754577 7+01	.6254844 5+01
1.10	.3757413 6+01	.6248026 0+01
1.15	.3760170 6+01	.6241119 7+01
1.20	.3762840 8+01	.6234123 2+01
1.25	.3765415 8+01	.6227033 8+01
1.30	.3767886 9+01	.6219849 9+01
1.35	.3770244 4+01	.6212570 1+01
1.40	.3772478 3+01	.6205193 9+01
1.45	.3774578 0+01	.6197721 6+01
1.50	.3776532 2+01	.6190154 2+01
1.55	.3778329 5+01	.6182493 9+01
1.60	.3779957 8+01	.6174744 0+01
1.65	.3781404 9+01	.6166909 1+01
1.70	.3782658 2+01	.6158995 1+01
1.75	.3783705 5+01	.6151009 6+01
1.80	.3784534 6+01	.6142961 4+01
1.85	.3785133 4+01	.6134861 6+01
1.90	.3785491 1+01	.6126722 3+01
1.95	.3785597 4+01	.6118557 9+01
2.00	.3785443 2+01	.6110384 2+01

Table D-10a (Cont'd)

THE ROOTS $t_5[Q\exp(i10^\circ)]$

Q	Real t_5	Imag t_5
2.00	.3785443+01	.6110384+01
2.05	.3785021+01	.6102218+01
2.10	.3784325+01	.6094079+01
2.15	.3783353+01	.6085987+01
2.20	.3782102+01	.6077961+01
2.25	.3780574+01	.6070025+01
2.30	.3778773+01	.6062198+01
2.35	.3776705+01	.6054501+01
2.40	.3774380+01	.6046955+01
2.45	.3771808+01	.6039579+01
2.50	.3769003+01	.6032391+01
2.55	.3765982+01	.6025406+01
2.60	.3762760+01	.6018637+01
2.65	.3759358+01	.6012096+01
2.70	.3755793+01	.6005791+01
2.75	.3752087+01	.5999729+01
2.80	.3748257+01	.5993915+01
2.85	.3744324+01	.5988349+01
2.90	.3740307+01	.5983032+01
2.95	.3736223+01	.5977962+01
3.00	.3732090+01	.5973134+01
3.05	.3727923+01	.5968543+01
3.10	.3723737+01	.5964183+01
3.15	.3719545+01	.5960047+01
3.20	.3715358+01	.5956126+01
3.25	.3711188+01	.5952412+01
3.30	.3707044+01	.5948897+01
3.35	.3702933+01	.5945570+01
3.40	.3698863+01	.5942423+01
3.45	.3694840+01	.5939448+01
3.50	.3690868+01	.5936634+01
3.55	.3686952+01	.5933973+01
3.60	.3683096+01	.5931458+01
3.65	.3679302+01	.5929079+01
3.70	.3675571+01	.5926828+01
3.75	.3671907+01	.5924699+01
3.80	.3668309+01	.5922685+01
3.85	.3664779+01	.5920778+01
3.90	.3661316+01	.5918972+01
3.95	.3657921+01	.5917261+01
4.00	.3654593+01	.5915640+01

Table D-10b

THE ROOTS $t_5[Q\exp(i10^\circ)]$

1/Q	Real t_5	Imag t_5
0.00	.39720668+01	.68798215+01
0.01	.3982068l+01	.68798237+01
0.02	.39920773+01	.68798398+01
0.03	.4002102$_5$+01	.68798836+01
0.04	.40121514+01	.68799694+01
0.05	.40222318+01	.68801116+01
0.06	.40323509+01	.68803255+01
0.07	.40425158+01	.68806268+01
0.08	.40527331+01	.68810321+01
0.09	.4063008$_4$+01	.68815593+01
0.10	.40733467+01	.68822266+01
0.11	.40837520+01	.68830542+01
0.12	.4094226$_7$+01	.68840633+01
0.13	.4104771$_7$+01	.68852763+01
0.14	.41153859+01	.68867172+01
0.15	.41260657+01	.68884117+01
0.16	.41368047+01	.68903863+01
0.17	.4147592$_6$+01	.68926690+01
0.18	.41584157+01	.68952884+01
0.19	.41692553+01	.68982737+01
0.20	.41800872+01	.69016535+01
0.21	.4190882$_1$+01	.69054553+01
0.22	.42016040+01	.69097043+01
0.23	.42122108+01	.69144218+01
0.24	.42226545+01	.69196239+01
0.25	.42328812+01	.69253198+01
0.26	.4242832$_8$+01	.69315098+01
0.27	.4252448$_5$+01	.69381841+01
0.28	.42616669+01	.69453217+01
0.29	.4270428$_8$+01	.69528897+01
0.30	.42786801+01	.69608439+01
0.31	.42863745+01	.69691294+01
0.32	.4293476$_0$+01	.69776835+01
0.33	.4299960$_7$+01	.69864375+01
0.34	.43058176+01	.69953205+01
0.35	.4311048$_4$+01	.70042621+01
0.36	.43156666+01	.70131952+01
0.37	.43196961+01	.70220589+01
0.38	.4323168$_5$+01	.70307995+01
0.39	.4326121$_7$+01	.70393718+01
0.40	.43285970+01	.70477390+01

Table D-11a

THE ROOTS $t_1[Q\exp(i190^\circ)]$

Q	Real t_1	Imag t_1
0.00	.50939648-00	.88230059-00
0.05	.55073144-00	.90683108-00
0.10	.58978919-00	.93131179-00
0.15	.62665622-00	.95569418-00
0.20	.66141845-00	.97993130-00
0.25	.69416117-00	.10039782+01
0.30	.72496902-00	.10277920+01
0.35	.75392590-00	.10513325+01
0.40	.78111483-00	.10745620+01
0.45	.80661786-00	.10974459+01
0.50	.83051591-00	.11199524+01
0.55	.85288865-00	.11420530+01
0.60	.87381428-00	.11637223+01
0.65	.89336942-00	.11849383+01
0.70	.91162886-00	.12056822+01
0.75	.92866546-00	.12259381+01
0.80	.94454993-00	.12456937+01
0.85	.95935072-00	.12649392+01
0.90	.97313388-00	.12836681+01
0.95	.98596291-00	.13018763+01
1.00	.99789871-00	.13195624+01
1.05	.10089994+01	.13367270+01
1.10	.10193206+01	.13533732+01
1.15	.10289147+01	.13695059+01
1.20	.10378318+01	.13851314+01
1.25	.10461190+01	.14002578+01
1.30	.10538206+01	.14148943+01
1.35	.10609785+01	.14290514+01
1.40	.10676316+01	.14427401+01
1.45	.10738168+01	.14559724+01
1.50	.10795682+01	.14687608+01
1.55	.10849177+01	.14811183+01
1.60	.10898950+01	.14930580+01
1.65	.10945276+01	.15045933+01
1.70	.10988413+01	.15157377+01
1.75	.11028597+01	.15265045+01
1.80	.11066048+01	.15369071+01
1.85	.11100970+01	.15469586+01
1.90	.11133550+01	.15566720+01
1.95	.11163964+01	.15660599+01
2.00	.11192370+01	.15751348+01

Table D-11a (Cont'd)

THE ROOTS $t_1[Q\exp(i90^\circ)]$

Q	Real t_1	Imag t_1
2.00	.11192370+01	.15751348+01
2.05	.11218918+01	.15839086+01
2.10	.11243743+01	.15923932+01
2.15	.11266972+01	.16005997+01
2.20	.11288721+01	.16085393+01
2.25	.11309098+01	.16162225+01
2.30	.11328202+01	.16236594+01
2.35	.11346122+01	.16308599+01
2.40	.11362945+01	.16378335+01
2.45	.11378746+01	.16445892+01
2.50	.11393599+01	.16511357+01
2.55	.11407569+01	.16574814+01
2.60	.11420716+01	.16636341+01
2.65	.11433099+01	.16696017+01
2.70	.11444767+01	.16753913+01
2.75	.11455770+01	.16810100+01
2.80	.11466153+01	.16864645+01
2.85	.11475956+01	.16917612+01
2.90	.11485217+01	.16969062+01
2.95	.11493973+01	.17019053+01
3.00	.11502254+01	.17067642+01
3.05	.11510093+01	.17114882+01
3.10	.11517517+01	.17160823+01
3.15	.11524551+01	.17205515+01
3.20	.11531221+01	.17249004+01
3.25	.11537549+01	.17291335+01
3.30	.11543556+01	.17332550+01
3.35	.11549261+01	.17372690+01
3.40	.11554683+01	.17411794+01
3.45	.11559837+01	.17449899+01
3.50	.11564741+01	.17487041+01
3.55	.11569409+01	.17523253+01
3.60	.11573854+01	.17558569+01
3.65	.11578089+01	.17593019+01
3.70	.11582126+01	.17626633+01
3.75	.11585976+01	.17659440+01
3.80	.11589651+01	.17691468+01
3.85	.11593158+01	.17722742+01
3.90	.11596508+01	.17753287+01
3.95	.11599709+01	.17783128+01
4.00	.11602769+01	.17812288+01

Table D-11b

THE ROOTS $t_1[Q\exp(i190^\circ)]$

1/Q	Real t_1	Imag t_1
0.00	.11690537+01	.20248604+01
0.01	.11690530+01	.20148608+01
0.02	.11690484+01	.20048635+01
0.03	.11690357+01	.19948709+01
0.04	.11690112+01	.19848854+01
0.05	.11689712+01	.19749093+01
0.06	.11689119+01	.19649450+01
0.07	.11688297+01	.19549949+01
0.08	.11687213+01	.19450615+01
0.09	.11685834+01	.19351472+01
0.10	.11684127+01	.19252546+01
0.11	.11682063+01	.19153862+01
0.12	.11679614+01	.19055446+01
0.13	.11676751+01	.18957322+01
0.14	.11673449+01	.18859518+01
0.15	.11669686+01	.18762060+01
0.16	.11665440+01	.18664973+01
0.17	.11660689+01	.18568285+01
0.18	.11655417+01	.18472022+01
0.19	.11649606+01	.18376211+01
0.20	.11643243+01	.18280877+01
0.21	.11636315+01	.18186048+01
0.22	.11628812+01	.18091749+01
0.23	.11620723+01	.17998006+01
0.24	.11612044+01	.17904844+01
0.25	.11602769+01	.17812288+01
0.26	.11592893+01	.17720362+01
0.27	.11582417+01	.17629091+01
0.28	.11571340+01	.17538497+01
0.29	.11559664+01	.17448602+01
0.30	.11547391+01	.17359427+01
0.31	.11534528+01	.17270994+01
0.32	.11521081+01	.17183322+01
0.33	.11507056+01	.17096430+01
0.34	.11492462+01	.17010335+01
0.35	.11477311+01	.16925054+01
0.36	.11461612+01	.16840602+01
0.37	.11445378+01	.16756993+01
0.38	.11428622+01	.16674242+01
0.39	.11411357+01	.16592360+01
0.40	.11393599+01	.16511358+01

Table D-12a

THE ROOTS $t_2[Q\exp(i190^\circ)]$

Q	Real t_2	Imag t_2
0.00	.16240988+01	.28130215+01
0.05	.16373899+01	.28207200+01
0.10	.16505887+01	.28284293+01
0.15	.16636761+01	.28361597+01
0.20	.16766335+01	.28439202+01
0.25	.16894426+01	.28517189+01
0.30	.17020858+01	.28595623+01
0.35	.17145458+01	.28674561+01
0.40	.17268061+01	.28754041+01
0.45	.17388508+01	.28834087+01
0.50	.17506648+01	.28914711+01
0.55	.17622338+01	.28995907+01
0.60	.17735447+01	.29077655+01
0.65	.17845851+01	.29159922+01
0.70	.17953442+01	.29242662+01
0.75	.18058122+01	.29325817+01
0.80	.18159807+01	.29409315+01
0.85	.18258424+01	.29493079+01
0.90	.18353920+01	.29577022+01
0.95	.18446250+01	.29661050+01
1.00	.18535387+01	.29745065+01
1.05	.18621318+01	.29828966+01
1.10	.18704041+01	.29912650+01
1.15	.18783571+01	.29996014+01
1.20	.18859931+01	.30078954+01
1.25	.18933159+01	.30161373+01
1.30	.19003303+01	.30243175+01
1.35	.19070418+01	.30324267+01
1.40	.19134571+01	.30404563+01
1.45	.19195833+01	.30483984+01
1.50	.19254284+01	.30562455+01
1.55	.19310008+01	.30639908+01
1.60	.19363093+01	.30716285+01
1.65	.19413629+01	.30791529+01
1.70	.19461711+01	.30865596+01
1.75	.19507433+01	.30938444+01
1.80	.19550891+01	.31010041+01
1.85	.19592180+01	.31080358+01
1.90	.19631394+01	.31149375+01
1.95	.19668627+01	.31217076+01
2.00	.19703971+01	.31283449+01

Table D-12a (Cont'd)

THE ROOTS $t_2[Q\exp(i190^\circ)]$

Q	Real t_2	Imag t_2
2.00	.19703971+01	.31283449+01
2.05	.19737514+01	.31348488+01
2.10	.19769345+01	.31412192+01
2.15	.19799548+01	.31474563+01
2.20	.19828204+01	.31535608+01
2.25	.19855393+01	.31595334+01
2.30	.19881190+01	.31653754+01
2.35	.19905668+01	.31710882+01
2.40	.19928897+01	.31766734+01
2.45	.19950942+01	.31821331+01
2.50	.19971868+01	.31874689+01
2.55	.19991735+01	.31926833+01
2.60	.20010599+01	.31977783+01
2.65	.20028516+01	.32027563+01
2.70	.20045537+01	.32076198+01
2.75	.20061710+01	.32123711+01
2.80	.20077083+01	.32170129+01
2.85	.20091698+01	.32215475+01
2.90	.20105598+01	.32259775+01
2.95	.20118821+01	.32303054+01
3.00	.20131405+01	.32345339+01
3.05	.20143383+01	.32386653+01
3.10	.20154789+01	.32427022+01
3.15	.20165654+01	.32466470+01
3.20	.20176007+01	.32505021+01
3.25	.20185876+01	.32542700+01
3.30	.20195286+01	.32579529+01
3.35	.20204262+01	.32615531+01
3.40	.20212828+01	.32650730+01
3.45	.20221003+01	.32685146+01
3.50	.20228811+01	.32718801+01
3.55	.20236269+01	.32751718+01
3.60	.20243396+01	.32783915+01
3.65	.20250209+01	.32815412+01
3.70	.20256723+01	.32846230+01
3.75	.20262956+01	.32876386+01
3.80	.20268920+01	.32905900+01
3.85	.20274629+01	.32934788+01
3.90	.20280097+01	.32963068+01
3.95	.20285335+01	.32990758+01
4.00	.20290355+01	.33017873+01

Table D-12b

THE ROOTS $t_2[Q\exp(i90^\circ)]$

1/Q	Real t_2	Imag t_2
0.00	.20439747+01	.35402680+01
0.01	.20439735+01	.35302687+01
0.02	.20439653+01	.35202735+01
0.03	.20439431+01	.35102864+01
0.04	.20439000+01	.35003117+01
0.05	.20438296+01	.34903537+01
0.06	.20437253+01	.34804164+01
0.07	.20435807+01	.34705043+01
0.08	.20433900+01	.34606218+01
0.09	.20431474+01	.34507734+01
0.10	.20428476+01	.34409638+01
0.11	.20424856+01	.34311976+01
0.12	.20420567+01	.34214798+01
0.13	.20415565+01	.34118150+01
0.14	.20409812+01	.34022083+01
0.15	.20403274+01	.33926647+01
0.16	.20395921+01	.33831892+01
0.17	.20387726+01	.33737867+01
0.18	.20378670+01	.33644625+01
0.19	.20368735+01	.33552213+01
0.20	.20357911+01	.33460682+01
0.21	.20346190+01	.33370079+01
0.22	.20333570+01	.33280452+01
0.23	.20320053+01	.33191847+01
0.24	.20305644+01	.33104306+01
0.25	.20290355+01	.33017874+01
0.26	.20274199+01	.32932588+01
0.27	.20257194+01	.32848486+01
0.28	.20239362+01	.32765604+01
0.29	.20220727+01	.32683972+01
0.30	.20201316+01	.32603621+01
0.31	.20181160+01	.32524576+01
0.32	.20160288+01	.32446860+01
0.33	.20138735+01	.32370492+01
0.34	.20116536+01	.32295490+01
0.35	.20093727+01	.32221867+01
0.36	.20070346+01	.32149633+01
0.37	.20046432+01	.32078795+01
0.38	.20022022+01	.32009358+01
0.39	.19997155+01	.31941324+01
0.40	.19971868+01	.31874690+01

Table D-13a

THE ROOTS $t_3[Q\exp(i90^\circ)]$

Q	Real t_3	Imag t_3
0.00	.24100496+01	.41743283+01
0.05	.24190204+01	.41795158+01
0.10	.24279597+01	.41847085+01
0.15	.24368585+01	.41899115+01
0.20	.24457077+01	.41951293+01
0.25	.24544986+01	.42003668+01
0.30	.24632221+01	.42056277+01
0.35	.24718698+01	.42109157+01
0.40	.24804330+01	.42162342+01
0.45	.24889033+01	.42215855+01
0.50	.24972726+01	.42269722+01
0.55	.25055327+01	.42323957+01
0.60	.25136758+01	.42378571+01
0.65	.25216945+01	.42433567+01
0.70	.25295814+01	.42488945+01
0.75	.25373298+01	.42544696+01
0.80	.25449331+01	.42600808+01
0.85	.25523853+01	.42657263+01
0.90	.25596808+01	.42714034+01
0.95	.25668143+01	.42771094+01
1.00	.25737813+01	.42828409+01
1.05	.25805777+01	.42885939+01
1.10	.25872001+01	.42943645+01
1.15	.25936455+01	.43001480+01
1.20	.25999114+01	.43059397+01
1.25	.26059964+01	.43117345+01
1.30	.26118991+01	.43175276+01
1.35	.26176190+01	.43233135+01
1.40	.26231561+01	.43290869+01
1.45	.26285110+01	.43348427+01
1.50	.26336848+01	.43405757+01
1.55	.26386791+01	.43462807+01
1.60	.26434958+01	.43519527+01
1.65	.26481374+01	.43575872+01
1.70	.26526069+01	.43631793+01
1.75	.26569074+01	.43687249+01
1.80	.26610424+01	.43742198+01
1.85	.26650157+01	.43796605+01
1.90	.26688313+01	.43850431+01
1.95	.26724935+01	.43903647+01
2.00	.26760065+01	.43956221+01

Table D-13a (Cont'd)

THE ROOTS $t_3[Q\exp(i90^\circ)]$

Q	Real t_3	Imag t_3
2.00	.26760065+01	.43956221+01
2.05	.26793749+01	.44008129+01
2.10	.26826032+01	.44059348+01
2.15	.26856962+01	.44109857+01
2.20	.26886583+01	.44159638+01
2.25	.26914944+01	.44208678+01
2.30	.26942090+01	.44256963+01
2.35	.26968066+01	.44304484+01
2.40	.26992921+01	.44351234+01
2.45	.27016696+01	.44397208+01
2.50	.27039438+01	.44442401+01
2.55	.27061187+01	.44486815+01
2.60	.27081986+01	.44530447+01
2.65	.27101876+01	.44573302+01
2.70	.27120896+01	.44615381+01
2.75	.27139084+01	.44656691+01
2.80	.27156477+01	.44697236+01
2.85	.27173112+01	.44737026+01
2.90	.27189021+01	.44776066+01
2.95	.27204239+01	.44814365+01
3.00	.27218796+01	.44851935+01
3.05	.27232722+01	.44888784+01
3.10	.27246048+01	.44924925+01
3.15	.27258801+01	.44960366+01
3.20	.27271008+01	.44995121+01
3.25	.27282693+01	.45029200+01
3.30	.27293883+01	.45062618+01
3.35	.27304598+01	.45095385+01
3.40	.27314862+01	.45127513+01
3.45	.27324696+01	.45159016+01
3.50	.27334120+01	.45189906+01
3.55	.27343153+01	.45220196+01
3.60	.27351813+01	.45249898+01
3.65	.27360117+01	.45279025+01
3.70	.27368084+01	.45307588+01
3.75	.27375726+01	.45335601+01
3.80	.27383061+01	.45363075+01
3.85	.27390102+01	.45390023+01
3.90	.27396861+01	.45416456+01
3.95	.27403353+01	.45442387+01
4.00	.27409590+01	.45467824+01

Table D-13b

THE ROOTS $t_3[Q\exp(i90^\circ)]$

1/Q	Real t_3	Imag t_3
0.00	.27602798+01	.47809450+01
0.01	.27602783+01	.47709460+01
0.02	.27602672+01	.47609524+01
0.03	.27602372+01	.47509699+01
0.04	.27601791+01	.47410042+01
0.05	.27600839+01	.47310609+01
0.06	.27599429+01	.47211460+01
0.07	.27597480+01	.47112653+01
0.08	.27594912+01	.47014252+01
0.09	.27591651+01	.46916317+01
0.10	.27587628+01	.46818916+01
0.11	.27582781+01	.46722112+01
0.12	.27577054+01	.46625976+01
0.13	.27570396+01	.46530573+01
0.14	.27562764+01	.46435975+01
0.15	.27554122+01	.46342251+01
0.16	.27544441+01	.46249472+01
0.17	.27533703+01	.46157707+01
0.18	.27521891+01	.46067023+01
0.19	.27509002+01	.45977487+01
0.20	.27495036+01	.45889166+01
0.21	.27480002+01	.45802120+01
0.22	.27463916+01	.45716410+01
0.23	.27446799+01	.45632090+01
0.24	.27428679+01	.45549213+01
0.25	.27409590+01	.45467825+01
0.26	.27389570+01	.45387969+01
0.27	.27368661+01	.45309682+01
0.28	.27346909+01	.45232997+01
0.29	.27324364+01	.45157941+01
0.30	.27301078+01	.45084534+01
0.31	.27277102+01	.45012794+01
0.32	.27252495+01	.44942732+01
0.33	.27227309+01	.44874354+01
0.34	.27201603+01	.44807661+01
0.35	.27175429+01	.44742649+01
0.36	.27148844+01	.44679311+01
0.37	.27121900+01	.44617635+01
0.38	.27094651+01	.44557604+01
0.39	.27067148+01	.44499201+01
0.40	.27039438+01	.44442402+01

Table D-14a

THE ROOTS $t_4[Q\exp(i90^\circ)]$

Q	Real t_4	Imag t_4
0.00	.30816536+01	.53375807+01
0.05	.30886730+01	.53416375+01
0.10	.30956761+01	.53456976+01
0.15	.31026572+01	.53497639+01
0.20	.31096108+01	.53538398+01
0.25	.31165313+01	.53579280+01
0.30	.31234132+01	.53620312+01
0.35	.31302510+01	.53661520+01
0.40	.31370393+01	.53702927+01
0.45	.31437729+01	.53744553+01
0.50	.31504462+01	.53786417+01
0.55	.31570540+01	.53828534+01
0.60	.31635914+01	.53870916+01
0.65	.31700533+01	.53913572+01
0.70	.31764346+01	.53956509+01
0.75	.31827307+01	.53999729+01
0.80	.31889369+01	.54043232+01
0.85	.31950489+01	.54087012+01
0.90	.32010623+01	.54131063+01
0.95	.32069733+01	.54175375+01
1.00	.32127779+01	.54219936+01
1.05	.32184727+01	.54264727+01
1.10	.32240544+01	.54309730+01
1.15	.32295202+01	.54354922+01
1.20	.32348672+01	.54400281+01
1.25	.32400932+01	.54445779+01
1.30	.32451962+01	.54491387+01
1.35	.32501744+01	.54537077+01
1.40	.32550267+01	.54582818+01
1.45	.32597519+01	.54628575+01
1.50	.32643494+01	.54674318+01
1.55	.32688188+01	.54720011+01
1.60	.32731601+01	.54765622+01
1.65	.32773736+01	.54811117+01
1.70	.32814599+01	.54856462+01
1.75	.32854199+01	.54901624+01
1.80	.32892546+01	.54946572+01
1.85	.32929657+01	.54991274+01
1.90	.32965545+01	.55035700+01
1.95	.33000231+01	.55079822+01
2.00	.33033734+01	.55123612+01

Table D-14a (Cont'd)

THE ROOTS $t_4[Q\exp(i190^\circ)]$

Q	Real t_4	Imag t_4
2.00	.33033734+01	.55123612+01
2.05	.33066077+01	.55167043+01
2.10	.33097284+01	.55210094+01
2.15	.33127379+01	.55252739+01
2.20	.33156388+01	.55294959+01
2.25	.33184340+01	.55336734+01
2.30	.33211262+01	.55378047+01
2.35	.33237182+01	.55418881+01
2.40	.33262130+01	.55459224+01
2.45	.33286133+01	.55499062+01
2.50	.33309224+01	.55538385+01
2.55	.33331429+01	.55577182+01
2.60	.33352780+01	.55615447+01
2.65	.33373304+01	.55653173+01
2.70	.33393031+01	.55690355+01
2.75	.33411988+01	.55726989+01
2.80	.33430205+01	.55763072+01
2.85	.33447707+01	.55798603+01
2.90	.33464523+01	.55833582+01
2.95	.33480678+01	.55868008+01
3.00	.33496197+01	.55901884+01
3.05	.33511106+01	.55935213+01
3.10	.33525429+01	.55967996+01
3.15	.33539189+01	.56000238+01
3.20	.33552409+01	.56031942+01
3.25	.33565110+01	.56063114+01
3.30	.33577313+01	.56093760+01
3.35	.33589041+01	.56123884+01
3.40	.33600311+01	.56153493+01
3.45	.33611143+01	.56182595+01
3.50	.33621554+01	.56211194+01
3.55	.33631564+01	.56239299+01
3.60	.33641188+01	.56266917+01
3.65	.33650442+01	.56294055+01
3.70	.33659343+01	.56320722+01
3.75	.33667904+01	.56346925+01
3.80	.33676141+01	.56372671+01
3.85	.33684066+01	.56397968+01
3.90	.33691694+01	.56422825+01
3.95	.33699036+01	.56447249+01
4.00	.33706104+01	.56471249+01

Table D-14b

THE ROOTS $t_4[Q\exp(i90^\circ)]$

Q	Real t_4	Imag t_4
0.00	.33933540+01	.58774616+01
0.01	.33933521+01	.58674627+01
0.02	.33933384+01	.58574706+01
0.03	.33933015+01	.58474922+01
0.04	.33932301+01	.58375344+01
0.05	.33931132+01	.58276044+01
0.06	.33929402+01	.58177093+01
0.07	.33927012+01	.58078567+01
0.08	.33923870+01	.57980543+01
0.09	.33919885+01	.57883102+01
0.10	.33914980+01	.57786323+01
0.11	.33909085+01	.57690293+01
0.12	.33902137+01	.57595095+01
0.13	.33894084+01	.57500817+01
0.14	.33884884+01	.57407545+01
0.15	.33874505+01	.57315366+01
0.16	.33862925+01	.57224368+01
0.17	.33850135+01	.57134634+01
0.18	.33836132+01	.57046246+01
0.19	.33820928+01	.56959285+01
0.20	.33804540+01	.56873823+01
0.21	.33786996+01	.56789931+01
0.22	.33768333+01	.56707674+01
0.23	.33748596+01	.56627108+01
0.24	.33727834+01	.56548285+01
0.25	.33706104+01	.56471250+01
0.26	.33683468+01	.56396038+01
0.27	.33659989+01	.56322680+01
0.28	.33635735+01	.56251196+01
0.29	.33610777+01	.56181600+01
0.30	.33585184+01	.56113901+01
0.31	.33559028+01	.56048098+01
0.32	.33532379+01	.55984186+01
0.33	.33505305+01	.55922150+01
0.34	.33477874+01	.55861974+01
0.35	.33450151+01	.55803634+01
0.36	.33422198+01	.55747103+01
0.37	.33394075+01	.55692349+01
0.38	.33365837+01	.55639337+01
0.39	.33337537+01	.55588029+01
0.40	.33309224+01	.55538385+01

Table D-15a

THE ROOTS $t_5[Q\exp(i190^\circ)]$

Q	Real t_5	Imag t_5
0.00	.36860886+01	.63844927+01
0.05	.36919584+01	.63878842+01
0.10	.36978181+01	.63912781+01
0.15	.37036636+01	.63946763+01
0.20	.37094910+01	.63980813+01
0.25	.37152964+01	.64014950+01
0.30	.37210759+01	.64049195+01
0.35	.37268256+01	.64083566+01
0.40	.37325418+01	.64118081+01
0.45	.37382205+01	.64152755+01
0.50	.37438580+01	.64187605+01
0.55	.37494504+01	.64222641+01
0.60	.37549941+01	.64257877+01
0.65	.37604855+01	.64293320+01
0.70	.37659209+01	.64328977+01
0.75	.37712968+01	.64364855+01
0.80	.37766098+01	.64400955+01
0.85	.37818565+01	.64437281+01
0.90	.37870337+01	.64473829+01
0.95	.37921382+01	.64510596+01
1.00	.37971670+01	.64547577+01
1.05	.38021173+01	.64584764+01
1.10	.38069863+01	.64622149+01
1.15	.38117716+01	.64659718+01
1.20	.38164707+01	.64697460+01
1.25	.38210814+01	.64735358+01
1.30	.38256019+01	.64773397+01
1.35	.38300303+01	.64811558+01
1.40	.38343649+01	.64849821+01
1.45	.38386045+01	.64888166+01
1.50	.38427479+01	.64926571+01
1.55	.38467941+01	.64965013+01
1.60	.38507426+01	.65003470+01
1.65	.38545927+01	.65041917+01
1.70	.38583442+01	.65080331+01
1.75	.38619971+01	.65118687+01
1.80	.38655514+01	.65156960+01
1.85	.38690076+01	.65195129+01
1.90	.38723662+01	.65233168+01
1.95	.38756279+01	.65271054+01
2.00	.38787936+01	.65308765+01

Table D-15a (Cont'd)

THE ROOTS $t_5[Q\exp(i190^\circ)]$

Q	Real t_5	Imag t_5
2.00	.38787936+01	.65308765+01
2.05	.38818643+01	.65346278+01
2.10	.38848413+01	.65383573+01
2.15	.38877258+01	.65420628+01
2.20	.38905195+01	.65457425+01
2.25	.38932239+01	.65493946+01
2.30	.38958406+01	.65530171+01
2.35	.38983713+01	.65566087+01
2.40	.39008180+01	.65601676+01
2.45	.39031826+01	.65636926+01
2.50	.39054671+01	.65671824+01
2.55	.39076734+01	.65706357+01
2.60	.39098036+01	.65740515+01
2.65	.39118596+01	.65774287+01
2.70	.39138439+01	.65807666+01
2.75	.39157581+01	.65840645+01
2.80	.39176047+01	.65873217+01
2.85	.39193855+01	.65905376+01
2.90	.39211027+01	.65937118+01
2.95	.39227584+01	.65968439+01
3.00	.39243545+01	.65999337+01
3.05	.39258929+01	.66029808+01
3.10	.39273758+01	.66059853+01
3.15	.39288050+01	.66089471+01
3.20	.39301823+01	.66118661+01
3.25	.39315097+01	.66147424+01
3.30	.39327889+01	.66175762+01
3.35	.39340217+01	.66203676+01
3.40	.39352096+01	.66231169+01
3.45	.39363545+01	.66258244+01
3.50	.39374579+01	.66284903+01
3.55	.39385214+01	.66311151+01
3.60	.39395465+01	.66336990+01
3.65	.39405345+01	.66362425+01
3.70	.39414871+01	.66387460+01
3.75	.39424054+01	.66412100+01
3.80	.39432908+01	.66436350+01
3.85	.39441445+01	.66460215+01
3.90	.39449679+01	.66483700+01
3.95	.39457621+01	.66506810+01
4.00	.39465282+01	.66529550+01

Table D-15b

THE ROOTS $t_5[Q\exp(i190^\circ)]$

1/Q	Real t_5	Imag t_5
0.00	.39720668+01	.68798215+01
0.01	.39720645+01	.68698228+01
0.02	.39720485+01	.68598321+01
0.03	.39720053+01	.68498574+01
0.04	.39719217+01	.68399069+01
0.05	.39717851+01	.68299889+01
0.06	.39715831+01	.68201120+01
0.07	.39713044+01	.68102854+01
0.08	.39709383+01	.68005180+01
0.09	.39704749+01	.67908193+01
0.10	.39699058+01	.67811994+01
0.11	.39692233+01	.67716679+01
0.12	.39684210+01	.67622352+01
0.13	.39674938+01	.67529117+01
0.14	.39664380+01	.67437074+01
0.15	.39652510+01	.67346327+01
0.16	.39639319+01	.67256975+01
0.17	.39624807+01	.67169115+01
0.18	.39608990+01	.67082839+01
0.19	.39591896+01	.66998232+01
0.20	.39573561+01	.66915375+01
0.21	.39554036+01	.66834339+01
0.22	.39533377+01	.66755190+01
0.23	.39511649+01	.66677980+01
0.24	.39488925+01	.66602755+01
0.25	.39465282+01	.66529551+01
0.26	.39440800+01	.66458394+01
0.27	.39415563+01	.66389300+01
0.28	.39389654+01	.66322275+01
0.29	.39363158+01	.66257318+01
0.30	.39336158+01	.66194420+01
0.31	.39308736+01	.66133560+01
0.32	.39280970+01	.66074716+01
0.33	.39252937+01	.66017856+01
0.34	.39224707+01	.65962943+01
0.35	.39196347+01	.65909936+01
0.36	.39167923+01	.65858792+01
0.37	.39139491+01	.65809460+01
0.38	.39111106+01	.65761890+01
0.39	.39082818+01	.65716030+01
0.40	.39054671+01	.65671824+01

Table D-16a

THE ROOTS $t_1[Q\exp(i130^\circ)]$

Q	Real t_1	Imag t_1
0.00	.50939648-00	.88230058-00
0.05	.55247547-00	.85879382-00
0.10	.59664887-00	.83738442-00
0.15	.64182816-00	.81812159-00
0.20	.68792360-00	.80105397-00
0.25	.73484389-00	.78622971-00
0.30	.78249577-00	.77369668-00
0.35	.83078354-00	.76350248-00
0.40	.87960867-00	.75569461-00
0.45	.92886920-00	.75032056-00
0.50	.97845920-00	.74742780-00
0.55	.10282681+01	.74706384-00
0.60	.10781797+01	.74927623-00
0.65	.11280719+01	.75411252-00
0.70	.11778149+01	.76162010-00
0.75	.12272703+01	.77184601-00
0.80	.12762901+01	.78483663-00
0.85	.13247144+01	.80063715-00
0.90	.13723696+01	.81929082-00
0.95	.14190663+01	.84083785-00
1.00	.14645960+01	.86531381-00
1.05	.15087282+01	.89274730-00
1.10	.15512062+01	.92315654-00
1.15	.15917425+01	.95654450-00
1.20	.16300139+01	.99289173-00
1.25	.16656562+01	.10321461+01
1.30	.16982596+01	.10742079+01
1.35	.17273668+01	.11189089+01
1.40	.17524767+01	.11659839+01
1.45	.17730599+01	.12150335+01
1.50	.17885961+01	.12654821+01
1.55	.17986419+01	.13165425+01
1.60	.18029370+01	.13672092+01
1.65	.18015315+01	.14163190+01
1.70	.17948830+01	.14626985+01
1.75	.17838520+01	.15053746+01
1.80	.17695618+01	.15437548+01
1.85	.17531776+01	.15776854+01
1.90	.17357157+01	.16073740+01
1.95	.17179481+01	.16332454+01
2.00	.17003997+01	.16558069+01

Table D-16a (Cont'd)

THE ROOTS $t_1[Q\exp(i130^\circ)]$

Q	Real t_1	Imag t_1
2.00	.17003997+01	.16558069+01
2.05	.16833934+01	.16755590+01
2.10	.16671073+01	.16929507+01
2.15	.16516230+01	.17083649+01
2.20	.16369626+01	.17221196+01
2.25	.16231125+01	.17344751+01
2.30	.16100395+01	.17456437+01
2.35	.15977006+01	.17557986+01
2.40	.15860488+01	.17650818+01
2.45	.15750364+01	.17736100+01
2.50	.15646173+01	.17814799+01
2.55	.15547478+01	.17887720+01
2.60	.15453874+01	.17955540+01
2.65	.15364984+01	.18018830+01
2.70	.15280464+01	.18078078+01
2.75	.15199997+01	.18133700+01
2.80	.15123295+01	.18186053+01
2.85	.15050096+01	.18235448+01
2.90	.14980159+01	.18282156+01
2.95	.14913265+01	.18326413+01
3.00	.14849214+01	.18368427+01
3.05	.14787821+01	.18408382+01
3.10	.14728920+01	.18446441+01
3.15	.14672356+01	.18482750+01
3.20	.14617988+01	.18517437+01
3.25	.14565684+01	.18550621+01
3.30	.14515327+01	.18582406+01
3.35	.14466804+01	.18612886+01
3.40	.14420013+01	.18642148+01
3.45	.14374860+01	.18670270+01
3.50	.14331257+01	.18697322+01
3.55	.14289121+01	.18723371+01
3.60	.14248379+01	.18748475+01
3.65	.14208958+01	.18772689+01
3.70	.14170794+01	18796063+01
3.75	.14133824+01	.18818644+01
3.80	.14097993+01	.18840474+01
3.85	.14063245+01	.18861593+01
3.90	.14029530+01	.18882037+01
3.95	.13996803+01	.18901842+01
4.00	.13965018+01	.18921037+01

Table D-16b

THE ROOTS $t_1[Q\exp(i30^\circ)]$

1/Q	Real t_1	Imag t_1
0.00	.11690537+01	.20248604+01
0.01	.11777146+01	.20198600+01
0.02	.11863796+01	.20148572+01
0.03	.11950526+01	.20098497+01
0.04	.12037377+01	.20048349+01
0.05	.12124389+01	.19998102+01
0.06	.12211601+01	.19947730+01
0.07	.12299055+01	.19897205+01
0.08	.12386792+01	.19846500+01
0.09	.12474853+01	.19795584+01
0.10	.12563279+01	.19744427+01
0.11	.12652113+01	.19692996+01
0.12	.12741398+01	.19641258+01
0.13	.12831180+01	.19589176+01
0.14	.12921502+01	.19536714+01
0.15	.13012411+01	.19483831+01
0.16	.13103955+01	.19430485+01
0.17	.13196183+01	.19376632+01
0.18	.13289144+01	.19322224+01
0.19	.13382891+01	.19267210+01
0.20	.13477478+01	.19211536+01
0.21	.13572960+01	.19155142+01
0.22	.13669394+01	.19097967+01
0.23	.13766841+01	.19039941+01
0.24	.13865361+01	.18980991+01
0.25	.13965019+01	.18921037+01
0.26	.14065881+01	.18859993+01
0.27	.14168016+01	.18797762+01
0.28	.14271495+01	.18734242+01
0.29	.14376391+01	.18669318+01
0.30	.14482782+01	.18602865+01
0.31	.14590743+01	.18534745+01
0.32	.14700356+01	.18464806+01
0.33	.14811701+01	.18392877+01
0.34	.14924859+01	.18318771+01
0.35	.15039912+01	.18242279+01
0.36	.15156939+01	.18163167+01
0.37	.15276014+01	.18081174+01
0.38	.15397208+01	.17996007+01
0.39	.15520580+01	.17907338+01
0.40	.15646174+01	.17814798+01

Table D-17a

THE ROOTS $t_2[Q\exp(i130^\circ)]$

Q	Real t_2	Imag t_2
0.00	.16240988+01	.28130215+01
0.05	.16374512+01	.28053547+01
0.10	.16508599+01	.27977401+01
0.15	.16643454+01	.27901672+01
0.20	.16779288+01	.27826262+01
0.25	.16916322+01	.27751074+01
0.30	.17054787+01	.27676020+01
0.35	.17194929+01	.27601010+01
0.40	.17337010+01	.27525963+01
0.45	.17481313+01	.27450798+01
0.50	.17628143+01	.27375440+01
0.55	.17777832+01	.27299816+01
0.60	.17930745+01	.27223863+01
0.65	.18087286+01	.27147522+01
0.70	.18247902+01	.27070741+01
0.75	.18413094+01	.26993487+01
0.80	.18583428+01	.26915736+01
0.85	.18759541+01	.26837492+01
0.90	.18942161+01	.26758788+01
0.95	.19132123+01	.26679699+01
1.00	.19330387+01	.26600364+01
1.05	.19538065+01	.26521003+01
1.10	.19756448+01	.26441956+01
1.15	.19987042+01	.26363732+01
1.20	.20231599+01	.26287080+01
1.25	.20492156+01	.26213090+01
1.30	.20771051+01	.26143342+01
1.35	.21070924+01	.26080109+01
1.40	.21394638+01	.26026639+01
1.45	.21745084+01	.25987510+01
1.50	.22124757+01	.25969035+01
1.55	.22535001+01	.25979593+01
1.60	.22974873+01	.26029634+01
1.65	.23439753+01	.26131012+01
1.70	.23920241+01	.26295376+01
1.75	.24402045+01	.26531870+01
1.80	.24867208+01	.26845004+01
1.85	.25296171+01	.27233545+01
1.90	.25669681+01	.27690477+01
1.95	.25970158+01	.28203254+01
2.00	.26183032+01	.28753721+01

Table D-17a (Cont'd)

THE ROOTS $t_2[Q\exp(i30^\circ)]$

Q	Real t_2	Imag t_2
2.00	.26183032+01	.28753721+01
2.05	.26299094+01	.29318044+01
2.10	.26318134+01	.29868337+01
2.15	.26251809+01	.30377754+01
2.20	.26121923+01	.30827685+01
2.25	.25953516+01	.31211828+01
2.30	.25767729+01	.31534132+01
2.35	.25578839+01	.31803638+01
2.40	.25394989+01	.32030272+01
2.45	.25220152+01	.32222806+01
2.50	.25055866+01	.32388306+01
2.55	.24902362+01	.32532249+01
2.60	.24759234+01	.32658826+01
2.65	.24625793+01	.32771245+01
2.70	.24501255+01	.32871978+01
2.75	.24384837+01	.32962952+01
2.80	.24275799+01	.33045684+01
2.85	.24173462+01	.33121382+01
2.90	.24077214+01	.33191023+01
2.95	.23986506+01	.33255401+01
3.00	.23900852+01	.33315171+01
3.05	.23819812+01	.33370879+01
3.10	.23743003+01	.33422980+01
3.15	.23670077+01	.33471862+01
3.20	.23600727+01	.33517855+01
3.25	.23534675+01	.33561242+01
3.30	.23471674+01	.33602268+01
3.35	.23411503+01	.33641146+01
3.40	.23353959+01	.33678062+01
3.45	.23298860+01	.33713181+01
3.50	.23246043+01	.33746646+01
3.55	.23195358+01	.33778585+01
3.60	.23146667+01	.33809114+01
3.65	.23099848+01	.33838336+01
3.70	.23054786+01	.33866342+01
3.75	.23011377+01	.33893215+01
3.80	.22969525+01	.33919030+01
3.85	.22929141+01	.33943855+01
3.90	.22890145+01	.33967753+01
3.95	.22852461+01	.33990779+01
4.00	.22816020+01	.34012985+01

Table D-17b

THE ROOTS $t_2[Q\exp(i30^\circ)]$

1/Q	Real t_2	Imag t_2
0.00	.20439747+01	.35402680+01
0.01	.20526361+01	.35352673+01
0.02	.20613046+01	.35302626+01
0.03	.20699873+01	.35252494+01
0.04	.20786912+01	.35202237+01
0.05	.20874236+01	.35151809+01
0.06	.20961917+01	.35101167+01
0.07	.21050031+01	.35050261+01
0.08	.21138652+01	.34999043+01
0.09	.21227861+01	.34947462+01
0.10	.21317738+01	.34895464+01
0.11	.21408367+01	.34842990+01
0.12	.21499836+01	.34789980+01
0.13	.21592237+01	.34736368+01
0.14	.21685667+01	.34682085+01
0.15	.21780229+01	.34627055+01
0.16	.21876028+01	.34571197+01
0.17	.21973182+01	.34514421+01
0.18	.22071813+01	.34456631+01
0.19	.22172052+01	.34397718+01
0.20	.22274040+01	.34337564+01
0.21	.22377930+01	.34276038+01
0.22	.22483885+01	.34212994+01
0.23	.22592087+01	.34148265+01
0.24	.22702727+01	.34081665+01
0.25	.22816020+01	.34012984+01
0.26	.22932199+01	.33941978+01
0.27	.23051517+01	.33868369+01
0.28	.23174255+01	.33791834+01
0.29	.23300723+01	.33711996+01
0.30	.23431260+01	.33628412+01
0.31	.23566242+01	.33540556+01
0.32	.23706077+01	.33447800+01
0.33	.23851214+01	.33349385+01
0.34	.24002136+01	.33244389+01
0.35	.24159356+01	.33131678+01
0.36	.24323398+01	.33009843+01
0.37	.24494766+01	.32877123+01
0.38	.24673882+01	.32731295+01
0.39	.24860974+01	.32569542+01
0.40	.25055870+01	.32388290+01

Table D-18a

THE ROOTS $t_3[Q\exp(i130^\circ)]$

Q	Real t_3	Imag t_3
0.00	.24100496+01	.41743283+01
0.05	.24190401+01	.41691505+01
0.10	.24280512+01	.41639868+01
0.15	.24370920+01	.41588320+01
0.20	.24461722+01	.41536812+01
0.25	.24553016+01	.41485292+01
0.30	.24644904+01	.41433711+01
0.35	.24737491+01	.41382020+01
0.40	.24830888+01	.41330168+01
0.45	.24925210+01	.41278105+01
0.50	.25020580+01	.41225780+01
0.55	.25117127+01	.41173140+01
0.60	.25214990+01	.41120132+01
0.65	.25314319+01	.41066701+01
0.70	.25415273+01	.41012792+01
0.75	.25518027+01	.40958346+01
0.80	.25622770+01	.40903302+01
0.85	.25729711+01	.40847600+01
0.90	.25839080+01	.40791173+01
0.95	.25951130+01	.40733957+01
1.00	.26066144+01	.40675880+01
1.05	.26184440+01	.40616874+01
1.10	.26306373+01	.40556867+01
1.15	.26432350+01	.40495785+01
1.20	.26562833+01	.40433559+01
1.25	.26698352+01	.40370119+01
1.30	.26839520+01	.40305404+01
1.35	.26987051+01	.40239369+01
1.40	.27141783+01	.40171984+01
1.45	.27304707+01	.40103255+01
1.50	.27477003+01	.40033245+01
1.55	.27660090+01	.39962097+01
1.60	.27855688+01	.39890088+01
1.65	.28065889+01	.39817704+01
1.70	.28293265+01	.39745761+01
1.75	.28540967+01	.39675600+01
1.80	.28812850+01	.39609402+01
1.85	.29113527+01	.39550694+01
1.90	.29448268+01	.39505135+01
1.95	.29822454+01	.39481645+01
2.00	.30240083+01	.39493725+01

Table D-18a (Cont'd)

THE ROOTS $t_3[Q\exp(i130^\circ)]$

Q	Real t_3	Imag t_3
2.00	.30240083+01	.39493725+01
2.05	.30700730+01	.39560249+01
2.10	.31195027+01	.39703957+01
2.15	.31700949+01	.39945961+01
2.20	.32184828+01	.40297535+01
2.25	.32607992+01	.40754271+01
2.30	.32934771+01	.41295420+01
2.35	.33139005+01	.41885970+01
2.40	.33210863+01	.42480398+01
2.45	.33163414+01	.43032195+01
2.50	.33030265+01	.43509350+01
2.55	.32850573+01	.43903140+01
2.60	.32654504+01	.44222204+01
2.65	.32459583+01	.44481330+01
2.70	.32274046+01	.44694521+01
2.75	.32100924+01	.44872850+01
2.80	.31940723+01	.45024518+01
2.85	.31792845+01	.45155488+01
2.90	.31656281+01	.45270112+01
2.95	.31529924+01	.45371602+01
3.00	.31412712+01	.45462369+01
3.05	.31303676+01	.45544253+01
3.10	.31201953+01	.45618679+01
3.15	.31106785+01	.45686768+01
3.20	.31017511+01	.45749419+01
3.25	.30933554+01	.45807356+01
3.30	.30854409+01	.45861172+01
3.35	.30779635+01	.45911361+01
3.40	.30708844+01	.45958332+01
3.45	.30641694+01	.46002432+01
3.50	.30577881+01	.46043957+01
3.55	.30517139+01	.46083159+01
3.60	.30459229+01	.46120257+01
3.65	.30403937+01	.46155441+01
3.70	.30351070+01	.46188875+01
3.75	.30300458+01	.46220706+01
3.80	.30251945+01	.46251060+01
3.85	.30205390+01	.46280056+01
3.90	.30160666+01	.46307792+01
3.95	.30117656+01	.46334360+01
4.00	.30076254+01	.46359841+01

Table D-18b

THE ROOTS $t_3[Q\exp(i30^\circ)]$

1/Q	Real t_3	Imag t_3
0.00	.27602798+01	.47809450+01
0.01	.27689417+01	.47759441+01
0.02	.27776131+01	.47709376+01
0.03	.27863037+01	.47659199+01
0.04	.27950231+01	.47608853+01
0.05	.28037813+01	.47558276+01
0.06	.28125881+01	.47507410+01
0.07	.28214542+01	.47456188+01
0.08	.28303902+01	.47404543+01
0.09	.28394073+01	.47352404+01
0.10	.28485174+01	.47299696+01
0.11	.28577327+01	.47246335+01
0.12	.28670666+01	.47192237+01
0.13	.28765329+01	.47137304+01
0.14	.28861468+01	.47081433+01
0.15	.28959244+01	.47024512+01
0.16	.29058836+01	.46966412+01
0.17	.29160433+01	.46906995+01
0.18	.29264248+01	.46846104+01
0.19	.29370511+01	.46783560+01
0.20	.29479480+01	.46719162+01
0.21	.29591440+01	.46652680+01
0.22	.29706713+01	.46583846+01
0.23	.29825658+01	.46512350+01
0.24	.29948684+01	.46437825+01
0.25	.30076256+01	.46359839+01
0.26	.30208907+01	.46277869+01
0.27	.30347248+01	.46191283+01
0.28	.30491990+01	.46099301+01
0.29	.30643955+01	.46000952+01
0.30	.30804104+01	.45895003+01
0.31	.30973553+01	.45779864+01
0.32	.31153598+01	.45653447+01
0.33	.31345718+01	.45512947+01
0.34	.31551537+01	.45354516+01
0.35	.31772678+01	.45172755+01
0.36	.32010361+01	.44959942+01
0.37	.32264362+01	.44704911+01
0.38	.32530577+01	.44391806+01
0.39	.32795877+01	.43999964+01
0.40	.33030050+01	.43509232+01

Table D-19a

THE ROOTS $t_4[Q\exp(i30^\circ)]$

Q	Real t_4	Imag t_4
0.00	.30816536+01	.53375807+01
0.05	.30886829+01	.53335285+01
0.10	.30957232+01	.53294823+01
0.15	.31027803+01	.53254390+01
0.20	.31098599+01	.53213952+01
0.25	.31169681+01	.53173479+01
0.30	.31241109+01	.53132937+01
0.35	.31312947+01	.53092296+01
0.40	.31385258+01	.53051523+01
0.45	.31458110+01	.53010586+01
0.50	.31531574+01	.52969449+01
0.55	.31605723+01	.52928079+01
0.60	.31680634+01	.52886441+01
0.65	.31756388+01	.52844498+01
0.70	.31833076+01	.52802213+01
0.75	.31910786+01	.52759547+01
0.80	.31989620+01	.52716456+01
0.85	.32069683+01	.52672900+01
0.90	.32151092+01	.52628834+01
0.95	.32233971+01	.52584210+01
1.00	.32318456+01	.52538977+01
1.05	.32404697+01	.52493083+01
1.10	.32492856+01	.52446472+01
1.15	.32583114+01	.52399084+01
1.20	.32675672+01	.52350855+01
1.25	.32770752+01	.52301718+01
1.30	.32868605+01	.52251599+01
1.35	.32969513+01	.52200425+01
1.40	.33073794+01	.52148110+01
1.45	.33181813+01	.52094571+01
1.50	.33293987+01	.52039717+01
1.55	.33410798+01	.51983453+01
1.60	.33532806+01	.51925684+01
1.65	.33660670+01	.51866313+01
1.70	.33795167+01	.51805252+01
1.75	.33937223+01	.51742423+01
1.80	.34087958+01	.51677776+01
1.85	.34248731+01	.51611308+01
1.90	.34421217+01	.51543099+01
1.95	.34607498+01	.51473370+01
2.00	.34810195+01	.51402584+01

Table D-19a (Cont'd)

THE ROOTS $t_4[Q\exp(i130^\circ)]$

Q	Real t_4	Imag t_4
2.00	.34810195+01	.51402584+01
2.05	.35032633+01	.51331623+01
2.10	.35279060+01	.51262084+01
2.15	.35554887+01	.51196823+01
2.20	.35866860+01	.51140899+01
2.25	.36222872+01	.51103179+01
2.30	.36630611+01	.51098797+01
2.35	.37093508+01	.51151808+01
2.40	.37602566+01	.51294973+01
2.45	.38127143+01	.51561013+01
2.50	.38615120+01	.51965401+01
2.55	.39007554+01	.52493513+01
2.60	.39257294+01	.53100950+01
2.65	.39344409+01	.53722598+01
2.70	.39287383+01	.54291839+01
2.75	.39135679+01	.54768170+01
2.80	.38941148+01	.55147400+01
2.85	.38737795+01	.55446123+01
2.90	.38541975+01	.55684316+01
2.95	.38359728+01	.55878211+01
3.00	.38192324+01	.56039493+01
3.05	.38039112+01	.56176317+01
3.10	.37898793+01	.56294383+01
3.15	.37769945+01	.56397747+01
3.20	.37651225+01	.56489351+01
3.25	.37541439+01	.56571379+01
3.30	.37439546+01	.56645481+01
3.35	.37344652+01	.56712933+01
3.40	.37255989+01	.56774732+01
3.45	.37172897+01	.56831678+01
3.50	.37094810+01	.56884414+01
3.55	.37021235+01	.56933466+01
3.60	.36951748+01	.56979271+01
3.65	.36885975+01	.57022194+01
3.70	.36823591+01	.57062541+01
3.75	.36764310+01	.57100576+01
3.80	.36707878+01	.57136523+01
3.85	.36654072+01	.57170577+01
3.90	.36602689+01	.57202904+01
3.95	.36553551+01	.57233655+01
4.00	.36506499+01	.57262959+01

Table D-19b

THE ROOTS $t_4[Q\exp(i30^\circ)]$

1/Q	Real t_4	Imag t_4
0.00	.33933540+01	.58774616+01
0.01	.34020162+01	.58724605+01
0.02	.34106902+01	.58674525+01
0.03	.34193878+01	.58624308+01
0.04	.34281209+01	.58573882+01
0.05	.34369019+01	.58523174+01
0.06	.34457434+01	.58472107+01
0.07	.34546584+01	.58420602+01
0.08	.34636606+01	.58368574+01
0.09	.34727646+01	.58315931+01
0.10	.34819856+01	.58262579+01
0.11	.34913401+01	.58208410+01
0.12	.35008456+01	.58153311+01
0.13	.35105213+01	.58097155+01
0.14	.35203879+01	.58039802+01
0.15	.35304687+01	.57981095+01
0.16	.35407888+01	.57920860+01
0.17	.35513766+01	.57858895+01
0.18	.35622638+01	.57794970+01
0.19	.35734862+01	.57728823+01
0.20	.35850848+01	.57660140+01
0.21	.35971060+01	.57588558+01
0.22	.36096044+01	.57513639+01
0.23	.36226428+01	.57434857+01
0.24	.36362953+01	.57351564+01
0.25	.36506502+01	.57262956+01
0.26	.36658125+01	.57168016+01
0.27	.36819101+01	.57065429+01
0.28	.36990982+01	.56953464+01
0.29	.37175683+01	.56829779+01
0.30	.37375564+01	.56691108+01
0.31	.37593526+01	.56532750+01
0.32	.37833036+01	.56347667+01
0.33	.38097867+01	.56124876+01
0.34	.38390822+01	.55846553+01
0.35	.38709112+01	.55483254+01
0.36	.39030024+01	.54989717+01
0.37	.39280373+01	.54320033+01
0.38	.39330282+01	.53497346+01
0.39	.39094511+01	.52660362+01
0.40	.38616068+01	.51965476+01

Table D-20a

THE ROOTS $t_5[Q\exp(i30^\circ)]$

Q	Real t_5	Imag t_5
0.00	.36860886+01	.63844927+01
0.05	.36919644+01	.63811040+01
0.10	.36978474+01	.63777183+01
0.15	.37037414+01	.63743335+01
0.20	.37096506+01	.63709473+01
0.25	.37155791+01	.63675574+01
0.30	.37215311+01	.63641616+01
0.35	.37275109+01	.63607576+01
0.40	.37335230+01	.63573430+01
0.45	.37395718+01	.63539157+01
0.50	.37456621+01	.63504730+01
0.55	.37517989+01	.63470126+01
0.60	.37579873+01	.63435320+01
0.65	.37642326+01	.63400284+01
0.70	.37705406+01	.63364994+01
0.75	.37769171+01	.63329420+01
0.80	.37833685+01	.63293534+01
0.85	.37899014+01	.63257305+01
0.90	.37965231+01	.63220702+01
0.95	.38032412+01	.63183691+01
1.00	.38100639+01	.63146237+01
1.05	.38170002+01	.63108301+01
1.10	.38240594+01	.63069846+01
1.15	.38312522+01	.63030829+01
1.20	.38385899+01	.62991205+01
1.25	.38460849+01	.62950926+01
1.30	.38537509+01	.62909941+01
1.35	.38616029+01	.62868194+01
1.40	.38696577+01	.62825628+01
1.45	.38779338+01	.62782176+01
1.50	.38864522+01	.62737771+01
1.55	.38952363+01	.62692339+01
1.60	.39043125+01	.62645797+01
1.65	.39137109+01	.62598060+01
1.70	.39234661+01	.62549033+01
1.75	.39336176+01	.62498615+01
1.80	.39442117+01	.62446697+01
1.85	.39553019+01	.62393163+01
1.90	.39669520+01	.62337892+01
1.95	.39792371+01	.62280761+01
2.00	.39922481+01	.62221648+01

Table D-20a (Cont'd)

THE ROOTS $t_5[Q\exp(i30^\circ)]$

Q	Real t_5	Imag t_5
2.00	.39922481+01	.62221648+01
2.05	.40060949+01	.62160441+01
2.10	.40209129+01	.62097059+01
2.15	.40368702+01	.62031475+01
2.20	.40541793+01	.61963770+01
2.25	.40731118+01	.61894222+01
2.30	.40940199+01	.61823474+01
2.35	.41173665+01	.61752838+01
2.40	.41437627+01	.61684883+01
2.45	.41740085+01	.61624543+01
2.50	.42091029+01	.61581213+01
2.55	.42501146+01	.61572386+01
2.60	.42976388+01	.61628367+01
2.65	.43504927+01	.61792988+01
2.70	.44041432+01	.62108760+01
2.75	.44510215+01	.62586676+01
2.80	.44835041+01	.63188283+01
2.85	.44972403+01	.63834389+01
2.90	.44934171+01	.64434163+01
2.95	.44781222+01	.64929494+01
3.00	.44580112+01	.65314188+01
3.05	.44371939+01	.65610217+01
3.10	.44174349+01	.65842213+01
3.15	.43992761+01	.66028871+01
3.20	.43827611+01	.66182957+01
3.25	.43677612+01	.66313031+01
3.30	.43541038+01	.66424917+01
3.35	.43416197+01	.66522669+01
3.40	.43301578+01	.66609187+01
3.45	.43195882+01	.66686599+01
3.50	.43098007+01	.66756499+01
3.55	.43007020+01	.66820111+01
3.60	.42922134+01	.66878391+01
3.65	.42842676+01	.66932096+01
3.70	.42768078+01	.66981839+01
3.75	.42697846+01	.67028116+01
3.80	.42631559+01	.67071340+01
3.85	.42568850+01	.67111856+01
3.90	.42509397+01	.67149952+01
3.95	.42452920+01	.67185877+01
4.00	.42399172+01	.67219841+01

Table D-20b

THE ROOTS $t_5[Q\exp(i130^\circ)]$

1/Q	Real t_5	Imag t_5
0.00	.39720668+01	.68798215+01
0.01	.39807294+01	.68748201+01
0.02	.39894056+01	.68698109+01
0.03	.39981096+01	.68647854+01
0.04	.40068553+01	.68597355+01
0.05	.40156574+01	.68546526+01
0.06	.40245306+01	.68495275+01
0.07	.40334909+01	.68443509+01
0.08	.40425546+01	.68391125+01
0.09	.40517394+01	.68338014+01
0.10	.40610643+01	.68284059+01
0.11	.40705496+01	.68229130+01
0.12	.40802177+01	.68173083+01
0.13	.40900930+01	.68115760+01
0.14	.41002028+01	.68056981+01
0.15	.41105775+01	.67996541+01
0.16	.41212513+01	.67934211+01
0.17	.41322633+01	.67869715+01
0.18	.41436579+01	.67802741+01
0.19	.41554870+01	.67732913+01
0.20	.41678108+01	.67659780+01
0.21	.41807006+01	.67582797+01
0.22	.41942415+01	.67501286+01
0.23	.42085360+01	.67414402+01
0.24	.42237096+01	.67321059+01
0.25	.42399178+01	.67219835+01
0.26	.42573560+01	.67108820+01
0.27	.42762739+01	.66985368+01
0.28	.42969945+01	.66845676+01
0.29	.43199405+01	.66684037+01
0.30	.43456624+01	.66491415+01
0.31	.43748431+01	.66252597+01
0.32	.44081480+01	.65940139+01
0.33	.44453201+01	.65502166+01
0.34	.44812629+01	.64849944+01
0.35	.44974878+01	.63925967+01
0.36	.44713742+01	.62912076+01
0.37	.44071042+01	.62130000+01
0.38	.43306223+01	.61715592+01
0.39	.42628334+01	.61579540+01
0.40	.42090850+01	.61581145+01

Table D-21

THE ROOTS $t_1[Q\exp(i300^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.48425817E-00	0.83875976E 00
0.10	0.45788710E-00	0.79308371E 00
0.15	0.43023229E-00	0.74518418E 00
0.20	0.40124264E-00	0.69497262E 00
0.25	0.37086708E-00	0.64236061E 00
0.30	0.33905483E-00	0.58726015E 00
0.35	0.30575553E-00	0.52958407E 00
0.40	0.27091953E-00	0.46924635E-00
0.45	0.23449802E-00	0.40616243E-00
0.50	0.19644319E-00	0.34024952E-00
0.55	0.15670846E-00	0.27142695E-00
0.60	0.11524856E-00	0.19961629E-00
0.65	0.72019693E-01	0.12474169E-00
0.70	0.26979624E-01	0.46730001E-01
0.75	-0.19912204E-01	-0.34489032E-01
0.80	-0.68694678E-01	-0.11898274E-00
0.85	-0.11940488E-00	-0.20681539E-00
0.90	-0.17207811E-00	-0.29804809E-00
0.95	-0.22674790E-00	-0.39273890E-00
1.00	-0.28344598E-00	-0.49094280E-00
1.05	-0.34220229E-00	-0.59271164E 00
1.10	-0.40304510E-00	-0.69809435E 00
1.15	-0.46600099E-00	-0.80713695E 00
1.20	-0.53109486E 00	-0.91988255E 00
1.25	-0.59835024E 00	-0.10363717E 01
1.30	-0.66778903E 00	-0.11566425E 01
1.35	-0.73943189E 00	-0.12807303E 01
1.40	-0.81329826E 00	-0.14086686E 01
1.45	-0.88940658E 00	-0.15404887E 01
1.50	-0.96777467E 00	-0.16762204E 01
1.55	-0.10484203E 01	-0.18158925E 01
1.60	-0.11313625E 01	-0.19595339E 01
1.65	-0.12166236E 01	-0.21071758E 01
1.70	-0.13042349E 01	-0.22588560E 01
1.75	-0.13942472E 01	-0.24146291E 01
1.80	-0.14867578E 01	-0.25745891E 01
1.85	-0.15819736E 01	-0.27389237E 01
1.90	-0.16803730E 01	-0.29080495E 01
1.95	-0.17831670E 01	-0.30829714E 01
2.00	-0.18938587E 01	-0.32663264E 01

Table D-22

THE ROOTS $t_1[Q\exp(i305^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.48824671E-00	0.83669402E 00
0.10	0.46626214E-00	0.78876795E 00
0.15	0.44341645E-00	0.73842390E 00
0.20	0.41968318E-00	0.68556309E 00
0.25	0.39503582E-00	0.63008691E 00
0.30	0.36944775E-00	0.57189726E 00
0.35	0.34289240E-00	0.51089697E 00
0.40	0.31534317E-00	0.44699021E-00
0.45	0.28677358E-00	0.38008288E-00
0.50	0.25715727E-00	0.31008293E-00
0.55	0.22646802E-00	0.23690071E-00
0.60	0.19467987E-00	0.16044921E-00
0.65	0.16176715E-00	0.80644389E-01
0.70	0.12770452E-00	-0.25946805E-02
0.75	0.92467077E-01	-0.89345606E-01
0.80	0.56030356E-01	-0.17968255E-00
0.85	0.18370441E-01	-0.27367620E-00
0.90	-0.20536020E-01	-0.37139367E-00
0.95	-0.60711714E-01	-0.47289844E-00
1.00	-0.10217869E-00	-0.57825056E 00
1.05	-0.14495825E-00	-0.68750644E 00
1.10	-0.18907097E-00	-0.80071922E 00
1.15	-0.23453663E-00	-0.91793869E 00
1.20	-0.28137419E-00	-0.10392114E 01
1.25	-0.32960181E-00	-0.11645809E 01
1.30	-0.37923677E-00	-0.12940878E 01
1.35	-0.43029550E-00	-0.14277703E 01
1.40	-0.48279350E-00	-0.15656637E 01
1.45	-0.53674508E 00	-0.17078011E 01
1.50	-0.59216332E 00	-0.18542141E 01
1.55	-0.64905923E 00	-0.20049318E 01
1.60	-0.70744068E 00	-0.21599827E 01
1.65	-0.76730975E 00	-0.23193938E 01
1.70	-0.82865708E 00	-0.24831901E 01
1.75	-0.89145035E 00	-0.26513918E 01
1.80	-0.95560941E 00	-0.28240015E 01
1.85	-0.10209555E 01	-0.30009662E 01
1.90	-0.10871132E 01	-0.31820671E 01
1.95	-0.11533467E 01	-0.33666115E 01
2.00	-0.12184072E 01	-0.35526696E 01

Table D-23

THE ROOTS $t_1[Q\exp(i310^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.49240567E-00	0.83500167E 00
0.10	0.47500439E-00	0.78527314E 00
0.15	0.45719276E-00	0.73301307E 00
0.20	0.43897082E-00	0.67811918E 00
0.25	0.42033835E-00	0.62048927E 00
0.30	0.40129482E-00	0.56002155E 00
0.35	0.38183913E-00	0.49661510E-00
0.40	0.36196961E-00	0.43017018E-00
0.45	0.34168380E-00	0.36058863E-00
0.50	0.32097848E-00	0.28777424E-00
0.55	0.29984948E-00	0.21163306E-00
0.60	0.27829172E-00	0.13207366E-00
0.65	0.25629914E-00	0.49007501E-01
0.70	0.23386472E-00	-0.37650917E-01
0.75	0.21098050E-00	-0.12798378E-00
0.80	0.18763766E-00	-0.22206990E-00
0.85	0.16382650E-00	-0.31998455E-00
0.90	0.13953663E-00	-0.42179927E-00
0.95	0.11475698E-00	-0.52758197E 00
1.00	0.89475925E-01	-0.63739695E 00
1.05	0.63681416E-01	-0.75130469E 00
1.10	0.37361048E-01	-0.86936221E 00
1.15	0.10502201E-01	-0.99162298E 00
1.20	-0.16907834E-01	-0.11181369E 01
1.25	-0.44881810E-01	-0.12489508E 01
1.30	-0.73432365E-01	-0.13841081E 01
1.35	-0.10257192E-00	-0.15236495E 01
1.40	-0.13231253E-00	-0.16676124E 01
1.45	-0.16266576E-00	-0.18160316E 01
1.50	-0.19364257E-00	-0.19689389E 01
1.55	-0.22525304E-00	-0.21263628E 01
1.60	-0.25750633E-00	-0.22883283E 01
1.65	-0.29041118E-00	-0.24548545E 01
1.70	-0.32397731E-00	-0.26259528E 01
1.75	-0.35822076E-00	-0.28016209E 01
1.80	-0.39317731E-00	-0.29818370E 01
1.85	-0.42893478E-00	-0.31665549E 01
1.90	-0.46570464E-00	-0.33557170E 01
1.95	-0.50396582E 00	-0.35493349E 01
2.00	-0.54473260E 00	-0.37477726E 01

Table D-24

THE ROOTS t_1 [Qexp(i315°)]

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.49669862E-00	0.83369682E 00
0.10	0.48402708E-00	0.78263102E 00
0.15	0.47140844E-00	0.72900476E 00
0.20	0.45886923E-00	0.67271921E 00
0.25	0.44643567E-00	0.61367548E 00
0.30	0.43413338E-00	0.55177484E 00
0.35	0.42198709E-00	0.48691906E-00
0.40	0.41002037E-00	0.41901073E-00
0.45	0.39825537E-00	0.34795349E-00
0.50	0.38671260E-00	0.27365240E-00
0.55	0.37541074E-00	0.19601418E-00
0.60	0.36436646E-00	0.11494756E-00
0.65	0.35359431E-00	0.30363480E-01
0.70	0.34310667E-00	-0.57824623E-01
0.75	0.33291370E-00	-0.14970052E-00
0.80	0.32302330E-00	-0.24534516E-00
0.85	0.31344124E-00	-0.34483628E-00
0.90	0.30417116E-00	-0.44824830E-00
0.95	0.29521473E-00	-0.55565240E 00
1.00	0.28657177E-00	-0.66711618E 00
1.05	0.27824040E-00	-0.78270371E 00
1.10	0.27021722E-00	-0.90247561E 00
1.15	0.26249748E-00	-0.10264889E 01
1.20	0.25507531E-00	-0.11547971E 01
1.25	0.24794378E-00	-0.12874503E 01
1.30	0.24109524E-00	-0.14244954E 01
1.35	0.23452132E-00	-0.15659760E 01
1.40	0.22821315E-00	-0.17119323E 01
1.45	0.22216141E-00	-0.18624017E 01
1.50	0.21635631E-00	-0.20174186E 01
1.55	0.21078748E-00	-0.21770147E 01
1.60	0.20544371E-00	-0.23412193E 01
1.65	0.20031247E-00	-0.25100598E 01
1.70	0.19537951E-00	-0.26835637E 01
1.75	0.19062909E-00	-0.28617609E 01
1.80	0.18604647E-00	-0.30446893E 01
1.85	0.18162673E-00	-0.32324005E 01
1.90	0.17739627E-00	-0.34249634E 01
1.95	0.17345548E-00	-0.36224477E 01
2.00	0.17003874E-00	-0.38248476E 01

Table D-25

THE ROOTS $t_1[Q\exp(i330^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.51000212E 00	0.83219077E 00
0.10	0.51187769E 00	0.77998306E 00
0.15	0.51511170E 00	0.72562680E 00
0.20	0.51979306E 00	0.66907095E 00
0.25	0.52601074E 00	0.61026385E 00
0.30	0.53385338E 00	0.54915298E 00
0.35	0.54340898E 00	0.48568481E-00
0.40	0.55476443E 00	0.41980458E-00
0.45	0.56800520E 00	0.35145611E-00
0.50	0.58321481E 00	0.28058167E-00
0.55	0.60047454E 00	0.20712193E-00
0.60	0.61986291E 00	0.13101584E-00
0.65	0.64145532E 00	0.52200682E-01
0.70	0.66532360E 00	-0.29387904E-01
0.75	0.69153564E 00	-0.11381571E-00
0.80	0.72015504E 00	-0.20114993E-00
0.85	0.75124077E 00	-0.29145877E-00
0.90	0.78484690E 00	-0.38481111E-00
0.95	0.82102241E 00	-0.48127632E-00
1.00	0.85981107E 00	-0.58092370E 00
1.05	0.90125130E 00	-0.68382210E 00
1.10	0.94537632E 00	-0.79003951E 00
1.15	0.99221415E 00	-0.89964259E 00
1.20	0.10417877E 01	-0.10126961E 01
1.25	0.10941155E 01	-0.11292630E 01
1.30	0.11492112E 01	-0.12494035E 01
1.35	0.12070847E 01	-0.13731752E 01
1.40	0.12677421E 01	-0.15006327E 01
1.45	0.13311864E 01	-0.16318273E 01
1.50	0.13974176E 01	-0.17668073E 01
1.55	0.14664335E 01	-0.19056176E 01
1.60	0.15382298E 01	-0.20482999E 01
1.65	0.16128006E 01	-0.21948925E 01
1.70	0.16901387E 01	-0.23454308E 01
1.75	0.17702358E 01	-0.24999468E 01
1.80	0.18530831E 01	-0.26584703E 01
1.85	0.19386708E 01	-0.28210282E 01
1.90	0.20269890E 01	-0.29876448E 01
1.95	0.21180280E 01	-0.31583431E 01
2.00	0.22117778E 01	-0.33331431E 01

Table D-26

THE ROOTS $t_1[Q\exp(i345^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.52313869E 00	0.83430852E 00
0.10	0.53902606E 00	0.78516090E 00
0.15	0.55715689E 00	0.73488475E 00
0.20	0.57763024E 00	0.68350731E 00
0.25	0.60054610E 00	0.63105556E 00
0.30	0.62600543E 00	0.57755590E 00
0.35	0.65411032E 00	0.52303369E 00
0.40	0.68496415E 00	0.46751276E-00
0.45	0.71867163E 00	0.41101496E-00
0.50	0.75533883E 00	0.35355965E-00
0.55	0.79507322E 00	0.29516312E-00
0.60	0.83798364E 00	0.23583809E-00
0.65	0.88418011E 00	0.17559306E-00
0.70	0.93377370E 00	0.11443170E-00
0.75	0.98687612E 00	0.52352326E-01
0.80	0.10435994E 01	-0.10652851E-01
0.85	0.11040553E 01	-0.74598142E-01
0.90	0.11683545E 01	-0.13950493E-00
0.95	0.12366059E 01	-0.20540212E-00
1.00	0.13089158E 01	-0.27232634E-00
1.05	0.13853861E 01	-0.34032205E-00
1.10	0.14661144E 01	-0.40944146E-00
1.15	0.15511917E 01	-0.47974406E-00
1.20	0.16407014E 01	-0.55129609E 00
1.25	0.17347192E 01	-0.62416948E 00
1.30	0.18333108E 01	-0.69844089E 00
1.35	0.19365325E 01	-0.77419027E 00
1.40	0.20444301E 01	-0.85149921E 00
1.45	0.21570391E 01	-0.93044962E 00
1.50	0.22743854E 01	-0.10111220E 01
1.55	0.23964852E 01	-0.10935944E 01
1.60	0.25233468E 01	-0.11779407E 01
1.65	0.26549710E 01	-0.12642302E 01
1.70	0.27913523E 01	-0.13525265E 01
1.75	0.29324803E 01	-0.14428876E 01
1.80	0.30783417E 01	-0.15353658E 01
1.85	0.32289194E 01	-0.16300070E 01
1.90	0.33841958E 01	-0.17268525E 01
1.95	0.35441513E 01	-0.18259376E 01
2.00	0.37087664E 01	-0.19272934E 01

Table D-27

THE ROOTS $t_1[Q\exp(i350^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.52730741E 00	0.83578679E 00
0.10	0.54752740E 00	0.78851534E 00
0.15	0.57014437E 00	0.74053787E 00
0.20	0.59524691E 00	0.69190650E 00
0.25	0.62292461E 00	0.64267351E 00
0.30	0.65326834E 00	0.59289110E 00
0.35	0.68637057E 00	0.54261092E 00
0.40	0.72232564E 00	0.49188385E-00
0.45	0.76123014E 00	0.44075949E-00
0.50	0.80318311E 00	0.38928573E-00
0.55	0.84828637E 00	0.33750825E-00
0.60	0.89664481E 00	0.28546997E-00
0.65	0.94836644E 00	0.23321036E-00
0.70	0.10035627E 01	0.18076472E-00
0.75	0.10623483E 01	0.12816345E-00
0.80	0.11248414E 01	0.75431076E-01
0.85	0.11911632E 01	0.22585367E-01
0.90	0.12614375E 01	-0.30363645E-01
0.95	0.13357901E 01	-0.83414733E-01
1.00	0.14143480E 01	-0.13657641E-00
1.05	0.14972385E 01	-0.18986784E-00
1.10	0.15845870E 01	-0.24331959E-00
1.15	0.16765160E 01	-0.29697409E-00
1.20	0.17731429E 01	-0.35088566E-00
1.25	0.18745777E 01	-0.40512038E-00
1.30	0.19809213E 01	-0.45975504E-00
1.35	0.20922633E 01	-0.51487599E 00
1.40	0.22086799E 01	-0.57057687E 00
1.45	0.23302338E 01	-0.62695663E 00
1.50	0.24569722E 01	-0.68411651E 00
1.55	0.25889280E 01	-0.74215757E 00
1.60	0.27261198E 01	-0.80117789E 00
1.65	0.28685538E 01	-0.86127046E 00
1.70	0.30162253E 01	-0.92252108E 00
1.75	0.31691208E 01	-0.98500760E 00
1.80	0.33272206E 01	-0.10487990E 01
1.85	0.34905001E 01	-0.11139555E 01
1.90	0.36589321E 01	-0.11805287E 01
1.95	0.38324884E 01	-0.12485625E 01
2.00	0.40111401E 01	-0.13180937E 01

Table D-27 (Cont'd)

THE ROOTS $t_1[Q\exp(i350^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.41948597E 01	-0.13891532E 01
2.10	0.43836205E 01	-0.14617665E 01
2.15	0.45773977E 01	-0.15359551E 01
2.20	0.47761680E 01	-0.16117364E 01
2.25	0.49799103E 01	-0.16891254E 01
2.30	0.51886044E 01	-0.17681344E 01
2.35	0.54022331E 01	-0.18487737E 01
2.40	0.56207791E 01	-0.19310521E 01
2.45	0.58442281E 01	-0.20149770E 01
2.50	0.60725667E 01	-0.21005550E 01
2.55	0.63057818E 01	-0.21877914E 01
2.60	0.65438624E 01	-0.22766910E 01
2.65	0.67867981E 01	-0.23672579E 01
2.70	0.70345793E 01	-0.24594960E 01
2.75	0.72871969E 01	-0.25534081E 01
2.80	0.75446432E 01	-0.26489973E 01
2.85	0.78069106E 01	-0.27462660E 01
2.90	0.80739928E 01	-0.28452166E 01
2.95	0.83458821E 01	-0.29458509E 01
3.00	0.86225738E 01	-0.30481708E 01
3.05	0.89040617E 01	-0.31521779E 01
3.10	0.91903421E 01	-0.32578740E 01
3.15	0.94814083E 01	-0.33652599E 01
3.20	0.97772570E 01	-0.34743372E 01
3.25	0.10077885E 02	-0.35851072E 01
3.30	0.10383286E 02	-0.36975702E 01
3.35	0.10693458E 02	-0.38117277E 01
3.40	0.11008398E 02	-0.39275806E 01
3.45	0.11328103E 02	-0.40451295E 01
3.50	0.11652568E 02	-0.41643751E 01
3.55	0.11981792E 02	-0.42853184E 01
3.60	0.12315772E 02	-0.44079597E 01
3.65	0.12654506E 02	-0.45322997E 01
3.70	0.12997991E 02	-0.46583391E 01
3.75	0.13346225E 02	-0.47860783E 01
3.80	0.13699206E 02	-0.49155176E 01
3.85	0.14056931E 02	-0.50466578E 01
3.90	0.14419401E 02	-0.51794992E 01
3.95	0.14786611E 02	-0.53140417E 01
4.00	0.15158561E 02	-0.54502864E 01

Table D-28

THE ROOTS $t_1[Q\exp(i355^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.53131361E 00	0.83762775E 00
0.10	0.55563027E 00	0.79261623E 00
0.15	0.58241788E 00	0.74733871E 00
0.20	0.61174843E 00	0.70186862E 00
0.25	0.64369476E 00	0.65628002E 00
0.30	0.67833108E 00	0.61064740E 00
0.35	0.71573333E 00	0.56504557E 00
0.40	0.75597963E 00	0.51954938E 00
0.45	0.79915078E 00	0.47423360E-00
0.50	0.84533074E 00	0.42917255E-00
0.55	0.89460714E 00	0.38443977E-00
0.60	0.94707184E 00	0.34010768E-00
0.65	0.10028214E 01	0.29624700E-00
0.70	0.10619577E 01	0.25292611E-00
0.75	0.11245885E 01	0.21021041E-00
0.80	0.11908277E 01	0.16816122E-00
0.85	0.12607960E 01	0.12683485E-00
0.90	0.13346209E 01	0.86281236E-01
0.95	0.14124369E 01	0.46542361E-01
1.00	0.14943855E 01	0.76506638E-02
1.05	0.15806143E 01	-0.30372903E-01
1.10	0.16712762E 01	-0.67520928E-01
1.15	0.17665277E 01	-0.10380143E-00
1.20	0.18665268E 01	-0.13923988E-00
1.25	0.19714298E 01	-0.17388061E-00
1.30	0.20813880E 01	-0.20778771E-00
1.35	0.21965434E 01	-0.24104501E-00
1.40	0.23170245E 01	-0.27375481E-00
1.45	0.24429415E 01	-0.30603542E-00
1.50	0.25743841E 01	-0.33801726E-00
1.55	0.27114173E 01	-0.36983784E-00
1.60	0.28540824E 01	-0.40163629E-00
1.65	0.30023966E 01	-0.43354771E-00
1.70	0.31563564E 01	-0.46569832E-00
1.75	0.33159410E 01	-0.49820174E-00
1.80	0.34811168E 01	-0.53115678E 00
1.85	0.36518419E 01	-0.56464661E 00
1.90	0.38280699E 01	-0.59873942E 00
1.95	0.40097535E 01	-0.63348979E 00
2.00	0.41968466E 01	-0.66894055E 00

Table D-28 (Cont'd)

THE ROOTS $t_1[Q\exp(1355^{\circ})]$

Q	Real t_1	Imag t_1
2.05	0.43893055E 01	-0.70512490E 00
2.10	0.45870909E 01	-0.74206850E 00
2.15	0.47901667E 01	-0.77979091E 00
2.20	0.49985008E 01	-0.81830736E 00
2.25	0.52120642E 01	-0.85762940E 00
2.30	0.54308319E 01	-0.89776643E 00
2.35	0.56547813E 01	-0.93872569E 00
2.40	0.58838923E 01	-0.98051303E 00
2.45	0.61181471E 01	-0.10231332E 01
2.50	0.63575301E 01	-0.10665905E 01
2.55	0.66020263E 01	-0.11108878E 01
2.60	0.68516232E 01	-0.11560282E 01
2.65	0.71063093E 01	-0.12020140E 01
2.70	0.73660733E 01	-0.12488473E 01
2.75	0.76309060E 01	-0.12965298E 01
2.80	0.79007985E 01	-0.13450634E 01
2.85	0.81757426E 01	-0.13944490E 01
2.90	0.84557310E 01	-0.14446882E 01
2.95	0.87407574E 01	-0.14957820E 01
3.00	0.90308147E 01	-0.15477312E 01
3.05	0.93258969E 01	-0.16005369E 01
3.10	0.96260001E 01	-0.16541998E 01
3.15	0.99311178E 01	-0.17087206E 01
3.20	0.10241245E 02	-0.17641000E 01
3.25	0.10556380E 02	-0.18203385E 01
3.30	0.10876516E 02	-0.18774368E 01
3.35	0.11201651E 02	-0.19353952E 01
3.40	0.11531781E 02	-0.19942144E 01
3.45	0.11866902E 02	-0.20538945E 01
3.50	0.12207011E 02	-0.21144361E 01
3.55	0.12552106E 02	-0.21758395E 01
3.60	0.12902185E 02	-0.22381050E 01
3.65	0.13257243E 02	-0.23012330E 01
3.70	0.13617279E 02	-0.23652238E 01
3.75	0.13982291E 02	-0.24300775E 01
3.80	0.14352276E 02	-0.24957945E 01
3.85	0.14727233E 02	-0.25623750E 01
3.90	0.15107160E 02	-0.26298191E 01
3.95	0.15492054E 02	-0.26981271E 01
4.00	0.15881915E 02	-0.27672993E 01

Table D-29

THE ROOTS $t_1[Q\exp(i11^6)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.53586233E 00	0.84028906E 00
0.10	0.56473643E 00	0.79845038E 00
0.15	0.59606411E 00	0.75687576E 00
0.20	0.62989088E 00	0.71565741E 00
0.25	0.66626280E 00	0.67488857E 00
0.30	0.70522692E 00	0.63466366E 00
0.35	0.74683178E 00	0.59507831E 00
0.40	0.79112784E 00	0.55622942E 00
0.45	0.83816805E 00	0.51821530E 00
0.50	0.88800842E 00	0.48113564E-00
0.55	0.94070870E 00	0.44509157E-00
0.60	0.99633317E 00	0.41018563E-00
0.65	0.10549512E 01	0.37652168E-00
0.70	0.11166386E 01	0.34420469E-00
0.75	0.11814782E 01	0.31334051E-00
0.80	0.12495613E 01	0.28403536E-00
0.85	0.13209886E 01	0.25639518E-00
0.90	0.13958719E 01	0.23052464E-00
0.95	0.14743352E 01	0.20652590E-00
1.00	0.15565167E 01	0.18449675E-00
1.05	0.16425694E 01	0.16452830E-00
1.10	0.17326630E 01	0.14670191E-00
1.15	0.18269844E 01	0.13108536E-00
1.20	0.19257376E 01	0.11772807E-00
1.25	0.20291422E 01	0.10665561E-00
1.30	0.21374312E 01	0.97863609E-01
1.35	0.22508450E 01	0.91311550E-01
1.40	0.23696236E 01	0.86917552E-01
1.45	0.24939957E 01	0.84555227E-01
1.50	0.26241653E 01	0.84054349E-01
1.55	0.27602994E 01	0.85206583E-01
1.60	0.29025146E 01	0.87776831E-01
1.65	0.30508714E 01	0.91519276E-01
1.70	0.32053736E 01	0.96195456E-01
1.75	0.33659758E 01	0.10159086E-00
1.80	0.35325966E 01	0.10752660E-00
1.85	0.37051331E 01	0.11386447E-00
1.90	0.38834761E 01	0.12050562E-00
1.95	0.40675204E 01	0.12738470E-00
2.00	0.42571715E 01	0.13446207E-00

Table D-29 (Cont'd)

THE ROOTS t_1 Qexp(i1°)

Q	Real t_1	Imag t_1
2.05	0.44523486E 01	0.14171591E-00
2.10	0.46529850E 01	0.14913570E-00
2.15	0.48590256E 01	0.15671743E-00
2.20	0.50704252E 01	0.16446042E-00
2.25	0.52871472E 01	0.17236555E-00
2.30	0.55091601E 01	0.18043413E-00
2.35	0.57364381E 01	0.18866756E-00
2.40	0.59689590E 01	0.19706707E-00
2.45	0.62067029E 01	0.20563369E-00
2.50	0.64496528E 01	0.21436831E-00
2.55	0.66977932E 01	0.22327162E-00
2.60	0.69511110E 01	0.23234427E-00
2.65	0.72095934E 01	0.24158671E-00
2.70	0.74732296E 01	0.25099942E-00
2.75	0.77420100E 01	0.26058277E-00
2.80	0.80159245E 01	0.27033707E-00
2.85	0.82949657E 01	0.28026263E-00
2.90	0.85791261E 01	0.29035969E-00
2.95	0.88683976E 01	0.30062846E-00
3.00	0.91627748E 01	0.31106917E-00
3.05	0.94622514E 01	0.32168198E-00
3.10	0.97668222E 01	0.33246706E-00
3.15	0.10076481E 02	0.34342454E-00
3.20	0.10391225E 02	0.35455458E-00
3.25	0.10711049E 02	0.36585729E-00
3.30	0.11035948E 02	0.37733276E-00
3.35	0.11365918E 02	0.38898110E-00
3.40	0.11700958E 02	0.40080243E-00
3.45	0.12041062E 02	0.41279680E-00
3.50	0.12386228E 02	0.42496430E-00
3.55	0.12736453E 02	0.43730500E-00
3.60	0.13091735E 02	0.44981898E-00
3.65	0.13452070E 02	0.46250629E-00
3.70	0.13817456E 02	0.47536698E-00
3.75	0.14187891E 02	0.48840111E-00
3.80	0.14563373E 02	0.50160875E 00
3.85	0.14943900E 02	0.51498991E 00
3.90	0.15329470E 02	0.52854465E 00
3.95	0.15720080E 02	0.54227304E 00
4.00	0.16115730E 02	0.55617508E 00

Table D-29 (Cont'd)

THE ROOTS $t_1[Q\exp(i1^\circ)]$

Q	Real t_1	Imag t_1
2.0	0.42570173E 01	0.13450941E-00
2.1	0.46529218E 01	0.14915717E-00
2.2	0.50703993E 01	0.16446946E-00
2.3	0.55091490E 01	0.18043794E-00
2.4	0.59689538E 01	0.19706874E-00
2.5	0.64496504E 01	0.21436910E-00
2.6	0.69511098E 01	0.23234465E-00
2.7	0.74732292E 01	0.25099963E-00
2.8	0.80159244E 01	0.27033719E-00
2.9	0.85791258E 01	0.29035974E-00
3.0	0.91627748E 01	0.31106920E-00
3.1	0.97668220E 01	0.33246708E-00
3.2	0.10391225E 02	0.35455460E-00
3.3	0.11035948E 02	0.37733278E-00
3.4	0.11700958E 02	0.40080244E-00
3.5	0.12386228E 02	0.42496431E-00
3.6	0.13091735E 02	0.44981899E-00
3.7	0.13817456E 02	0.47536699E-00
3.8	0.14563373E 02	0.50160876E 00
3.9	0.15329470E 02	0.52854468E 00
4.0	0.16115730E 02	0.55617509E 00
4.1	0.16922140E 02	0.58450028E 00
4.2	0.17748690E 02	0.61352053E 00
4.3	0.18595367E 02	0.64323607E 00
4.4	0.19462162E 02	0.67364711E 00
4.5	0.20349065E 02	0.70475381E 00
4.6	0.21256070E 02	0.73655640E 00
4.7	0.22183166E 02	0.76905494E 00
4.8	0.23130351E 02	0.80224965E 00
4.9	0.24097616E 02	0.83614061E 00
5.0	0.25084955E 02	0.87072793E 00

Table D-30

THE ROOTS $t_1[Q\exp(i15^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.55871473E 00	0.84232002E 00
0.10	0.57038693E 00	0.80284781E 00
0.15	0.60443838E 00	0.76398256E 00
0.20	0.64089420E 00	0.72582394E 00
0.25	0.67977976E 00	0.68847287E 00
0.30	0.72112097E 00	0.65203176E 00
0.35	0.76494474E 00	0.61660475E 00
0.40	0.81127940E 00	0.58229791E 00
0.45	0.86015518E 00	0.54921965E 00
0.50	0.91160472E 00	0.51748081E 00
0.55	0.96566369E 00	0.48719514E-00
0.60	0.10223715E 01	0.45847946E-00
0.65	0.10817723E 01	0.43145397E-00
0.70	0.11439154E 01	0.40624256E-00
0.75	0.12088571E 01	0.38297296E-00
0.80	0.12766615E 01	0.36177697E-00
0.85	0.13474023E 01	0.34279038E-00
0.90	0.14211646E 01	0.32615291E-00
0.95	0.14980470E 01	0.31200776E-00
1.00	0.15781645E 01	0.30050079E-00
1.05	0.16616514E 01	0.29177903E-00
1.10	0.17486645E 01	0.28598859E-00
1.15	0.18393870E 01	0.28327103E-00
1.20	0.19340325E 01	0.28375841E-00
1.25	0.20328485E 01	0.28756609E-00
1.30	0.21361182E 01	0.29478282E-00
1.35	0.22441611E 01	0.30545802E-00
1.40	0.23573292E 01	0.31958605E-00
1.45	0.24759898E 01	0.33708924E-00
1.50	0.26005156E 01	0.35780281E-00
1.55	0.27312435E 01	0.38146701E-00
1.60	0.28684420E 01	0.40773373E-00
1.65	0.30122749E 01	0.43619268E-00
1.70	0.31627791E 01	0.46641569E-00
1.75	0.33198686E 01	0.49800924E-00
1.80	0.34833638E 01	0.53065798E 00
1.85	0.36530364E 01	0.56414604E 00
1.90	0.38286541E 01	0.59835336E 00
1.95	0.40100119E 01	0.63323421E 00
2.00	0.41969465E 01	0.66878983E 00

Table D-30 (Cont'd)

THE ROOTS $t_1[Q\exp(i15^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.43893366E 01	0.70504472E 00
2.10	0.45870964E 01	0.74203002E 00
2.15	0.47901648E 01	0.77977456E 00
2.20	0.49984981E 01	0.81830158E 00
2.25	0.52120622E 01	0.85762823E 00
2.30	0.54308308E 01	0.89776705E 00
2.35	0.56547807E 01	0.93872695E 00
2.40	0.58838919E 01	0.98051453E 00
2.45	0.61181469E 01	0.10231348E 01
2.50	0.63575300E 01	0.10665921E 01
2.55	0.66020259E 01	0.11108895E 01
2.60	0.68516230E 01	0.11560300E 01
2.65	0.71063089E 01	0.12020159E 01
2.70	0.73660729E 01	0.12488492E 01
2.75	0.76309057E 01	0.12965319E 01
2.80	0.79007982E 01	0.13450654E 01
2.85	0.81757422E 01	0.13944511E 01
2.90	0.84557308E 01	0.14446905E 01
2.95	0.87407570E 01	0.14957842E 01
3.00	0.90308142E 01	0.15477335E 01
3.05	0.93258966E 01	0.16005393E 01
3.10	0.96259999E 01	0.16542023E 01
3.15	0.99311174E 01	0.17087232E 01
3.20	0.10241245E 02	0.17641027E 01
3.25	0.10556380E 02	0.18203414E 01
3.30	0.10876516E 02	0.18774396E 01
3.35	0.11201651E 02	0.19353982E 01
3.40	0.11531780E 02	0.19942174E 01
3.45	0.11866901E 02	0.20538977E 01
3.50	0.12207010E 02	0.21144393E 01
3.55	0.12552106E 02	0.21758428E 01
3.60	0.12902184E 02	0.22381085E 01
3.65	0.13257242E 02	0.23012366E 01
3.70	0.13617278E 02	0.23652274E 01
3.75	0.13982290E 02	0.24300813E 01
3.80	0.14352275E 02	0.24957983E 01
3.85	0.14727233E 02	0.25623789E 01
3.90	0.15107159E 02	0.26298232E 01
3.95	0.15492053E 02	0.26981312E 01
4.00	0.15881914E 02	0.27673035E 01

Table D-30 (Cont'd)

THE ROOTS $t_1[Q\exp(i15^\circ)]$

Q	Real t_1	Imag t_1
2.0	0.41969408E 01	0.66898195E 00
2.1	0.45871055E 01	0.74210170E 00
2.2	0.49985020E 01	0.81832830E 00
2.3	0.54308318E 01	0.89777808E 00
2.4	0.58838920E 01	0.98051957E 00
2.5	0.63575298E 01	0.10665945E 01
2.6	0.68516230E 01	0.11560312E 01
2.7	0.73660731E 01	0.12488499E 01
2.8	0.79007983E 01	0.13450658E 01
2.9	0.84557310E 01	0.14446906E 01
3.0	0.90308142E 01	0.15477336E 01
3.1	0.96259997E 01	0.16542023E 01
3.2	0.10241245E 02	0.17641027E 01
3.3	0.10876516E 02	0.18774397E 01
3.4	0.11531781E 02	0.19942174E 01
3.5	0.12207011E 02	0.21144393E 01
3.6	0.12902185E 02	0.22381084E 01
3.7	0.13617279E 02	0.23652274E 01
3.8	0.14352276E 02	0.24957984E 01
3.9	0.15107160E 02	0.26298232E 01
4.0	0.15881914E 02	0.27673034E 01
4.1	0.16676527E 02	0.29082408E 01
4.2	0.17490985E 02	0.30526365E 01
4.3	0.18325279E 02	0.32004917E 01
4.4	0.19179398E 02	0.33518074E 01
4.5	0.20053334E 02	0.35065846E 01
4.6	0.20947078E 02	0.36648241E 01
4.7	0.21860623E 02	0.38265266E 01
4.8	0.22793964E 02	0.39916929E 01
4.9	0.23747091E 02	0.41603234E 01
5.0	0.24720001E 02	0.43324189E 01

Table D-31

THE ROOTS $t_1[Q\exp(i110^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.54205377E 00	0.84512601E 00
0.10	0.57693213E 00	0.80886113E 00
0.15	0.61403035E 00	0.77360766E 00
0.20	0.65334666E 00	0.73946831E 00
0.25	0.69487895E 00	0.70654710E 00
0.30	0.73862506E 00	0.67494974E 00
0.35	0.78458298E 00	0.64478402E 00
0.40	0.83275110E 00	0.61616022E 00
0.45	0.88312855E 00	0.58919162E 00
0.50	0.93571553E 00	0.56399497E 00
0.55	0.99051370E 00	0.54069111E 00
0.60	0.10475266E 01	0.51940558E 00
0.65	0.11067604E 01	0.50026932E 00
0.70	0.11682244E 01	0.48341949E-00
0.75	0.12319318E 01	0.46900029E-00
0.80	0.12979010E 01	0.45716406E-00
0.85	0.13661571E 01	0.44807230E-00
0.90	0.14367330E 01	0.44189689E-00
0.95	0.15096724E 01	0.43882143E-00
1.00	0.15850321E 01	0.43904270E-00
1.05	0.16628863E 01	0.44277193E-00
1.10	0.17433311E 01	0.45023623E-00
1.15	0.18264917E 01	0.46167929E-00
1.20	0.19125301E 01	0.47736143E-00
1.25	0.20016582E 01	0.49755768E-00
1.30	0.20941512E 01	0.52255260E 00
1.35	0.21903668E 01	0.55262919E 00
1.40	0.22907673E 01	0.58804732E 00
1.45	0.23959403E 01	0.62900607E 00
1.50	0.25066108E 01	0.67558241E 00
1.55	0.26236220E 01	0.72764245E 00
1.60	0.27478562E 01	0.78473941E 00
1.65	0.28800628E 01	0.84604859E 00
1.70	0.30206237E 01	0.91042999E 00
1.75	0.31693781E 01	0.97667549E 00
1.80	0.33256678E 01	0.10438470E 01
1.85	0.34886098E 01	0.11114877E 01
1.90	0.36574047E 01	0.11795870E 01
1.95	0.38315036E 01	0.12483915E 01
2.00	0.40106160E 01	0.13182141E 01

Table D-31 (Cont'd)

THE ROOTS $t_1[Q\exp(i110^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.41946356E 01	0.13893206E 01
2.10	0.43835534E 01	0.14618903E 01
2.15	0.45773943E 01	0.15360238E 01
2.20	0.47761803E 01	0.16117664E 01
2.25	0.49799208E 01	0.16891357E 01
2.30	0.51886096E 01	0.17681371E 01
2.35	0.54022343E 01	0.18487749E 01
2.40	0.56207787E 01	0.19310535E 01
2.45	0.58442272E 01	0.20149790E 01
2.50	0.60725658E 01	0.21005574E 01
2.55	0.63057806E 01	0.21877938E 01
2.60	0.65438615E 01	0.22766937E 01
2.65	0.67867971E 01	0.23672608E 01
2.70	0.70345782E 01	0.24594988E 01
2.75	0.72871958E 01	0.25534111E 01
2.80	0.75446422E 01	0.26490004E 01
2.85	0.78069095E 01	0.27462692E 01
2.90	0.80739911E 01	0.28452199E 01
2.95	0.83458810E 01	0.29458544E 01
3.00	0.86225725E 01	0.30481744E 01
3.05	0.89040606E 01	0.31521817E 01
3.10	0.91903404E 01	0.32578778E 01
3.15	0.94814069E 01	0.33652639E 01
3.20	0.97772560E 01	0.34743415E 01
3.25	0.10077883E 02	0.35851111E 01
3.30	0.10383284E 02	0.36975745E 01
3.35	0.10693456E 02	0.38117322E 01
3.40	0.11008397E 02	0.39275852E 01
3.45	0.11328101E 02	0.40451343E 01
3.50	0.11652566E 02	0.41643800E 01
3.55	0.11981791E 02	0.42853235E 01
3.60	0.12315771E 02	0.44079649E 01
3.65	0.12654504E 02	0.45323050E 01
3.70	0.12997989E 02	0.46583445E 01
3.75	0.13346224E 02	0.47860838E 01
3.80	0.13699204E 02	0.49155234E 01
3.85	0.14056930E 02	0.50466637E 01
3.90	0.14419398E 02	0.51795051E 01
3.95	0.14786609E 02	0.53140481E 01
4.00	0.15158559E 02	0.54502928E 01

Table D-31 (Cont'd)

THE ROOTS $t_1[Q\exp(i110^\circ)]$

Q	Real t_1	Imag t_1
2.0	0.40109725E 01	0.13181297E 01
2.1	0.43836296E 01	0.14618231E 01
2.2	0.47761938E 01	0.16117514E 01
2.3	0.51886161E 01	0.17681370E 01
2.4	0.56207831E 01	0.19310543E 01
2.5	0.60725679E 01	0.21005576E 01
2.6	0.65438627E 01	0.22766938E 01
2.7	0.70345789E 01	0.24594989E 01
2.8	0.75446427E 01	0.26490004E 01
2.9	0.80739915E 01	0.28452198E 01
3.0	0.86225725E 01	0.30481744E 01
3.1	0.91903404E 01	0.32578778E 01
3.2	0.97772562E 01	0.34743414E 01
3.3	0.10383284E 02	0.36975745E 01
3.4	0.11008397E 02	0.39275853E 01
3.5	0.11652566E 02	0.41643800E 01
3.6	0.12315771E 02	0.44079649E 01
3.7	0.12997989E 02	0.46583445E 01
3.8	0.13699204E 02	0.49155234E 01
3.9	0.14419399E 02	0.51795051E 01
4.0	0.15158559E 02	0.54502928E 01
4.1	0.15916672E 02	0.57278895E 01
4.2	0.16693725E 02	0.60122977E 01
4.3	0.17489709E 02	0.63035199E 01
4.4	0.18304615E 02	0.66015578E 01
4.5	0.19138432E 02	0.69064133E 01
4.6	0.19991153E 02	0.72180882E 01
4.7	0.20862772E 02	0.75365836E 01
4.8	0.21753281E 02	0.78619015E 01
4.9	0.22662675E 02	0.81940422E 01
5.0	0.23590948E 02	0.85330077E 01

Table D-32

THE ROOTS t_1 [Qexp(i15°)]

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.54511897E 00	0.84820469E 00
0.10	0.58286220E 00	0.81538835E 00
0.15	0.62259859E 00	0.78394940E 00
0.20	0.66429963E 00	0.75398639E 00
0.25	0.70793594E 00	0.72559905E 00
0.30	0.75347733E 00	0.69888870E 00
0.35	0.80089285E 00	0.67395864E 00
0.40	0.85015082E 00	0.65091478E 00
0.45	0.90121883E 00	0.62986609E 00
0.50	0.95406388E 00	0.61092525E 00
0.55	0.10086522E 01	0.59420940E 00
0.60	0.10649499E 01	0.57984087E 00
0.65	0.11229222E 01	0.56794818E 00
0.70	0.11825542E 01	0.55866706E 00
0.75	0.12437510E 01	0.55214176E 00
0.80	0.13065375E 01	0.54852656E 00
0.85	0.13708597E 01	0.54798758E 00
0.90	0.14366841E 01	0.55070504E 00
0.95	0.15039796E 01	0.55687594E 00
1.00	0.15727177E 01	0.56671755E 00
1.05	0.16428749E 01	0.58047155E 00
1.10	0.17144350E 01	0.59840963E 00
1.15	0.17873929E 01	0.62084026E 00
1.20	0.18617604E 01	0.64811797E 00
1.25	0.19375756E 01	0.68065530E 00
1.30	0.20149175E 01	0.71893819E 00
1.35	0.20939311E 01	0.76354700E 00
1.40	0.21748681E 01	0.81519222E 00
1.45	0.22581613E 01	0.87469430E 00
1.50	0.23445567E 01	0.94310667E 00
1.55	0.24353559E 01	0.10215885E 01
1.60	0.25328308E 01	0.11112340E 01
1.65	0.26407389E 01	0.12122399E 01
1.70	0.27639673E 01	0.13218964E 01
1.75	0.29048711E 01	0.14330178E 01
1.80	0.30589309E 01	0.15385686E 01
1.85	0.32189775E 01	0.16378651E 01
1.90	0.33811571E 01	0.17342103E 01
1.95	0.35448062E 01	0.18307966E 01
2.00	0.37106340E 01	0.19295880E 01

Table D-32 (Cont'd)

THE ROOTS $t_1[Q\exp(i115^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.38796823E 01	0.20314784E 01
2.10	0.40528535E 01	0.21366268E 01
2.15	0.42307068E 01	0.22447891E 01
2.20	0.44134344E 01	0.23556141E 01
2.25	0.46009796E 01	0.24688467E 01
2.30	0.47932009E 01	0.25843885E 01
2.35	0.49899842E 01	0.27022515E 01
2.40	0.51912750E 01	0.28224852E 01
2.45	0.53970608E 01	0.29451295E 01
2.50	0.56073425E 01	0.30702033E 01
2.55	0.58221170E 01	0.31977122E 01
2.60	0.60413779E 01	0.33276594E 01
2.65	0.62651153E 01	0.34600480E 01
2.70	0.64933200E 01	0.35948820E 01
2.75	0.67259841E 01	0.37321661E 01
2.80	0.69631007E 01	0.38719041E 01
2.85	0.72046623E 01	0.40140989E 01
2.90	0.74506642E 01	0.41587542E 01
2.95	0.77010994E 01	0.43058719E 01
3.00	0.79559629E 01	0.44554551E 01
3.05	0.82152497E 01	0.46075056E 01
3.10	0.84789559E 01	0.47620259E 01
3.15	0.87470762E 01	0.49190171E 01
3.20	0.90196065E 01	0.50784815E 01
3.25	0.92965437E 01	0.52404206E 01
3.30	0.95778836E 01	0.54048353E 01
3.35	0.98636232E 01	0.55717278E 01
3.40	0.10153759E 02	0.57410986E 01
3.45	0.10448288E 02	0.59129489E 01
3.50	0.10747208E 02	0.60872801E 01
3.55	0.11050516E 02	0.62640929E 01
3.60	0.11358210E 02	0.64433884E 01
3.65	0.11670288E 02	0.66251673E 01
3.70	0.11986746E 02	0.68094303E 01
3.75	0.12307583E 02	0.69961782E 01
3.80	0.12632797E 02	0.71854119E 01
3.85	0.12962386E 02	0.73771320E 01
3.90	0.13296349E 02	0.75713386E 01
3.95	0.13634682E 02	0.77680326E 01
4.00	0.13977388E 02	0.79672150E 01

Table D-32 (Cont'd)

THE ROOTS $t_1[Q\exp(115^\circ)]$

Q	Real t_1	Imag t_1
2.0	0.37096836E 01	0.19283783E 01
2.1	0.40523782E 01	0.21367379E 01
2.2	0.44134422E 01	0.23557464E 01
2.3	0.47932413E 01	0.25843836E 01
2.4	0.51912767E 01	0.28224722E 01
2.5	0.56073410E 01	0.30702011E 01
2.6	0.60413782E 01	0.33276586E 01
2.7	0.64933203E 01	0.35948814E 01
2.8	0.69631007E 01	0.38719036E 01
2.9	0.74506642E 01	0.41587537E 01
3.0	0.79559632E 01	0.44554549E 01
3.1	0.84789559E 01	0.47620255E 01
3.2	0.90196066E 01	0.50784814E 01
3.3	0.95778836E 01	0.54048353E 01
3.4	0.10153759E 02	0.57410984E 01
3.5	0.10747208E 02	0.60872801E 01
3.6	0.11358210E 02	0.64433882E 01
3.7	0.11986746E 02	0.68094302E 01
3.8	0.12632797E 02	0.71854119E 01
3.9	0.13296349E 02	0.75713386E 01
4.0	0.13977388E 02	0.79672150E 01
4.1	0.14675900E 02	0.83730450E 01
4.2	0.15391877E 02	0.87888329E 01
4.3	0.16125306E 02	0.92145818E 01
4.4	0.16876180E 02	0.96502946E 01
4.5	0.17644489E 02	0.10095973E 02
4.6	0.18430226E 02	0.10551621E 02
4.7	0.19233384E 02	0.11017240E 02
4.8	0.20053958E 02	0.11492832E 02
4.9	0.20891940E 02	0.11978399E 02
5.0	0.21747325E 02	0.12473942E 02

Table D-33

THE ROOTS $t_1[Q\exp(i120^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.54788925E 00	0.85152842E 00
0.10	0.58814208E 00	0.82236311E 00
0.15	0.63010296E 00	0.79489185E 00
0.20	0.67371870E 00	0.76920222E 00
0.25	0.71893478E 00	0.74538255E 00
0.30	0.76569525E 00	0.72352232E 00
0.35	0.81394254E 00	0.70371262E 00
0.40	0.86361726E 00	0.68604656E 00
0.45	0.91465793E 00	0.67061978E 00
0.50	0.96700083E 00	0.65753107E 00
0.55	0.10205795E 01	0.64688294E 00
0.60	0.10753246E 01	0.63878234E 00
0.65	0.11311634E 01	0.63334156E 00
0.70	0.11880191E 01	0.63067911E 00
0.75	0.12458103E 01	0.63092095E 00
0.80	0.13044506E 01	0.63420185E 00
0.85	0.13638472E 01	0.64066709E 00
0.90	0.14239002E 01	0.65047454E 00
0.95	0.14845009E 01	0.66379725E 00
1.00	0.15455304E 01	0.68082676E 00
1.05	0.16068573E 01	0.70177733E 00
1.10	0.16683343E 01	0.72689169E 00
1.15	0.17297951E 01	0.75644836E 00
1.20	0.17910484E 01	0.79077213E 00
1.25	0.18518706E 01	0.83024863E 00
1.30	0.19119938E 01	0.87534557E 00
1.35	0.19710898E 01	0.92664495E 00
1.40	0.20287409E 01	0.98489403E 00
1.45	0.20843913E 01	0.10510906E 01
1.50	0.21372553E 01	0.11266363E 01
1.55	0.21861168E 01	0.12136395E 01
1.60	0.22288193E 01	0.13155930E 01
1.65	0.22605315E 01	0.14391842E 01
1.70	0.22638530E 01	0.15999239E 01
1.75	0.21735803E 01	0.17132550E 01
1.80	0.20310266E 01	0.17349278E 01
1.85	0.19598942E 01	0.17546788E 01
1.90	0.19061477E 01	0.17673491E 01
1.95	0.18629371E 01	0.17769817E 01
2.00	0.18268178E 01	0.17849336E 01

+ 147

Table D-33 (Cont'd)

THE ROOTS $t_1[Q\exp(i120^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.17958205E 01	0.17918021E 01
2.10	0.17697099E 01	0.17978998E 01
2.15	0.17446566E 01	0.18034106E 01
2.20	0.17230751E 01	0.18084536E 01
2.25	0.17035351E 01	0.18131100E 01
2.30	0.16857107E 01	0.18174394E 01
2.35	0.16693490E 01	0.18214867E 01
2.40	0.16542488E 01	0.18252864E 01
2.45	0.16402481E 01	0.18288672E 01
2.50	0.16272143E 01	0.18322517E 01
2.55	0.16150365E 01	0.18354592E 01
2.60	0.16036223E 01	0.18385062E 01
2.65	0.15928932E 01	0.18414064E 01
2.70	0.15827818E 01	0.18441720E 01
2.75	0.15732303E 01	0.18468135E 01
2.80	0.15641880E 01	0.18493403E 01
2.85	0.15556112E 01	0.18517604E 01
2.90	0.15474609E 01	0.18540817E 01
2.95	0.15397031E 01	0.18563104E 01
3.00	0.15323074E 01	0.18584526E 01
3.05	0.15252466E 01	0.18605138E 01
3.10	0.15184965E 01	0.18624987E 01
3.15	0.15120351E 01	0.18644119E 01
3.20	0.15058434E 01	0.18662579E 01
3.25	0.14999027E 01	0.18680399E 01
3.30	0.14941972E 01	0.18697616E 01
3.35	0.14887121E 01	0.18714263E 01
3.40	0.14834340E 01	0.18730369E 01
3.45	0.14783506E 01	0.18745963E 01
3.50	0.14734502E 01	0.18761065E 01
3.55	0.14687229E 01	0.18775706E 01
3.60	0.14641589E 01	0.18789904E 01
3.65	0.14597493E 01	0.18803681E 01
3.70	0.14554863E 01	0.18817056E 01
3.75	0.14513617E 01	0.18830047E 01
3.80	0.14473687E 01	0.18842670E 01
3.85	0.14435009E 01	0.18854943E 01
3.90	0.14397521E 01	0.18866882E 01
3.95	0.14361165E 01	0.18878496E 01
4.00	0.14325889E 01	0.18889805E 01

Table D-33 (Cont'd)

THE ROOTS $t_1[Q\exp(i120^\circ)]$

Q	Real t_1	Imag t_1
2.0	0.33104712E 01	C.24694440E 01
2.1	0.35990214E 01	C.27407619E 01
2.2	0.39190886E 01	0.30292844E 01
2.3	0.42577693E 01	0.33221941E 01
2.4	0.46087171E 01	C.36269369E 01
2.5	0.49759050E 01	C.39457256E 01
2.6	0.53594987E 01	C.42766944E 01
2.7	0.57587586E 01	0.46201694E 01
2.8	0.61738424E 01	0.49763100E 01
2.9	0.66046794E 01	C.53450790E 01
3.0	0.70512225E 01	0.57265215E 01
3.1	0.75134376E C1	0.61206563E 01
3.2	0.79912909E 01	0.65275021E 01
3.3	0.84847539E 01	C.69470746E 01
3.4	0.89938012E 01	0.73793877E C1
3.5	0.95184101E 01	0.78244522E 01
3.6	0.10058561E 02	0.82822781E 01
3.7	0.10614236E 02	C.87528755E 01
3.8	0.11185418E 02	0.92362508E 01
3.9	0.11772093E 02	0.97324110E 01
4.0	0.12374249E 02	0.10241361E 02
4.1	0.12991873E 02	C.10763108E 02
4.2	0.13624956E 02	0.11297656E 02
4.3	0.14273486E 02	0.11845008E C2
4.4	0.14937457E 02	0.12405170E 02
4.5	0.15616859E 02	C.12978143E 02
4.6	0.16311585E 02	0.13563931E 02
4.7	0.17021929E 02	0.14162537E 02
4.8	0.17747585E 02	C.14773965E 02
4.9	0.18488649E 02	C.15398215E 02
5.0	0.19245112E 02	0.16035289E C2

Table D-34

THE ROOTS $t_1[Q\exp(144^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.55659781E 00	0.86995077E 00
0.10	0.60378324E 00	0.85994212E 00
0.15	0.65085455E 00	0.85225192E 00
0.20	0.69771345E 00	0.84685599E 00
0.25	0.74426132E 00	0.84372834E 00
0.30	0.79039906E 00	0.84284094E 00
0.35	0.83602680E 00	0.84416331E 00
0.40	0.88104378E 00	0.84766213E 00
0.45	0.92534816E 00	0.85330080E 00
0.50	0.96883687E 00	0.86103895E 00
0.55	0.10114055E 01	0.87083185E 00
0.60	0.10529487E 01	0.88262985E 00
0.65	0.10933592E 01	0.89637755E 00
0.70	0.11325293E 01	0.91201306E 00
0.75	0.11703501E 01	0.92946712E 00
0.80	0.12067122E 01	0.94866216E 00
0.85	0.12415066E 01	0.96951113E 00
0.90	0.12746254E 01	0.99191654E 00
0.95	0.13059630E 01	0.10157692E 01
1.00	0.13354173E 01	0.10409474E 01
1.05	0.13628925E 01	0.10673155E 01
1.10	0.13883007E 01	0.10947239E 01
1.15	0.14115651E 01	0.11230082E 01
1.20	0.14326237E 01	0.11519897E 01
1.25	0.14514323E 01	0.11814768E 01
1.30	0.14679686E 01	0.12112668E 01
1.35	0.14822359E 01	0.12411496E 01
1.40	0.14942658E 01	0.12709122E 01
1.45	0.15041198E 01	0.13003443E 01
1.50	0.15118899E 01	0.13292450E 01
1.55	0.15176971E 01	0.13574295E 01
1.60	0.15216870E 01	0.13847347E 01
1.65	0.15240251E 01	0.14110252E 01
1.70	0.15248899E 01	0.14361960E 01
1.75	0.15244661E 01	0.14601739E 01
1.80	0.15229364E 01	0.14829162E 01
1.85	0.15204774E 01	0.15044086E 01
1.90	0.15172531E 01	0.15246608E 01
1.95	0.15134124E 01	0.15437026E 01
2.00	0.15090879E 01	0.15615778E 01

Table D-34 (Cont'd)

THE ROOTS $t_1[Q\exp(i144^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.15043942E 01	0.15783416E 01
2.10	0.14994296E 01	0.15940554E 01
2.15	0.14942764E 01	0.16087842E 01
2.20	0.14890031E 01	0.16225935E 01
2.25	0.14836653E 01	0.16355483E 01
2.30	0.14783081E 01	0.16477112E 01
2.35	0.14729675E 01	0.16591417E 01
2.40	0.14676718E 01	0.16698954E 01
2.45	0.14624430E 01	0.16800247E 01
2.50	0.14572979E 01	0.16895778E 01
2.55	0.14522489E 01	0.16985987E 01
2.60	0.14473052E 01	0.17071286E 01
2.65	0.14424732E 01	0.17152043E 01
2.70	0.14377566E 01	0.17228604E 01
2.75	0.14331582E 01	0.17301275E 01
2.80	0.14286789E 01	0.17370340E 01
2.85	0.14243182E 01	0.17436059E 01
2.90	0.14200755E 01	0.17498668E 01
2.95	0.14159489E 01	0.17558384E 01
3.00	0.14119365E 01	0.17615400E 01
3.05	0.14080359E 01	0.17669903E 01
3.10	0.14042440E 01	0.17722052E 01
3.15	0.14005581E 01	0.17772002E 01
3.20	0.13969750E 01	0.17819890E 01
3.25	0.13934916E 01	0.17865846E 01
2.30	0.13901050E 01	0.17909983E 01
3.35	0.13868119E 01	0.17952412E 01
3.40	0.13836090E 01	0.17993234E 01
3.45	0.13804937E 01	0.18032541E 01
3.50	0.13774627E 01	0.18070416E 01
3.55	0.13745131E 01	0.18106941E 01
3.60	0.13716422E 01	0.18142187E 01
3.65	0.13688470E 01	0.18176223E 01
3.70	0.13661250E 01	0.18209114E 01
3.75	0.13634738E 01	0.18240912E 01
3.80	0.13608906E 01	0.18271681E 01
3.85	0.13583730E 01	0.18301468E 01
3.90	0.13559192E 01	0.18330323E 01
3.95	0.13535266E 01	0.18358283E 01
4.00	0.13511929E 01	0.18385402E 01

Table D-35

THE ROOTS $t_1[Q\exp(i146^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.55695913E 00	0.87160375E 00
0.10	0.60434368E 00	0.86323792E 00
0.15	0.65145490E 00	0.85717026E 00
0.20	0.69819770E 00	0.85336641E 00
0.25	0.74447697E 00	0.85179024E 00
0.30	0.79019745E 00	0.85240344E 00
0.35	0.83526354E 00	0.85516517E 00
0.40	0.87957921E 00	0.86003161E 00
0.45	0.92304796E 00	0.86695559E 00
0.50	0.96557277E 00	0.87588596E 00
0.55	0.10070561E 01	0.88676716E 00
0.60	0.10474002E 01	0.89953852E 00
0.65	0.10865069E 01	0.91413372E 00
0.70	0.11242783E 01	0.93047989E 00
0.75	0.11606171E 01	0.94849707E 00
0.80	0.11954270E 01	0.96809721E 00
0.85	0.12286140E 01	0.98918354E 00
0.90	0.12600867E 01	0.10116496E 01
0.95	0.12897585E 01	0.10353787E 01
1.00	0.13175484E 01	0.10602434E 01
1.05	0.13433833E 01	0.10861047E 01
1.10	0.13672003E 01	0.11128125E 01
1.15	0.13889492E 01	0.11402062E 01
1.20	0.14085951E 01	0.11681148E 01
1.25	0.14261208E 01	0.11963599E 01
1.30	0.14415300E 01	0.12247574E 01
1.35	0.14548487E 01	0.12531213E 01
1.40	0.14661273E 01	0.12812677E 01
1.45	0.14754399E 01	0.13090196E 01
1.50	0.14828841E 01	0.13362120E 01
1.55	0.14885785E 01	0.13626959E 01
1.60	0.14926589E 01	0.13883434E 01
1.65	0.14952738E 01	0.14130496E 01
1.70	0.14965792E 01	0.14367350E 01
1.75	0.14967334E 01	0.14593449E 01
1.80	0.14958920E 01	0.14808489E 01
1.85	0.14942037E 01	0.15012382E 01
1.90	0.14918070E 01	0.15205223E 01
1.95	0.14888285E 01	0.15387264E 01
2.00	0.14853809E 01	0.15558864E 01

Table D-35 (Cont'd)

THE ROOTS $t_1[Q\exp(146^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.14815637E 01	0.15720475E 01
2.10	0.14774625E 01	0.15872605E 01
2.15	0.14731507E 01	0.16015783E 01
2.20	0.14686901E 01	0.16150559E 01
2.25	0.14641321E 01	0.16277479E 01
2.30	0.14595193E 01	0.16397069E 01
2.35	0.14548863E 01	0.16509841E 01
2.40	0.14502615E 01	0.16616277E 01
2.45	0.14456673E 01	0.16716832E 01
2.50	0.14411214E 01	0.16811930E 01
2.55	0.14366380E 01	0.16901962E 01
2.60	0.14322276E 01	0.16987297E 01
3.65	0.14278982E 01	0.17068268E 01
2.70	0.14236559E 01	0.17145185E 01
2.75	0.14195047E 01	0.17218333E 01
2.80	0.14154472E 01	0.17287977E 01
2.85	0.14114852E 01	0.17354348E 01
2.90	0.14076189E 01	0.17417680E 01
2.95	0.14038485E 01	0.17478166E 01
3.00	0.14001732E 01	0.17535996E 01
3.05	0.13965916E 01	0.17591333E 01
3.10	0.13931024E 01	0.17644347E 01
3.15	0.13897038E 01	0.17695175E 01
3.20	0.13863936E 01	0.17743950E 01
3.25	0.13831699E 01	0.17790797E 01
3.30	0.13800301E 01	0.17835831E 01
3.35	0.13769723E 01	0.17879153E 01
3.40	0.13739940E 01	0.17920867E 01
3.45	0.13710928E 01	0.17961055E 01
3.50	0.13682663E 01	0.17999807E 01
3.55	0.13655124E 01	0.18037197E 01
3.60	0.13628287E 01	0.18073298E 01
3.65	0.13602129E 01	0.18108178E 01
3.70	0.13576629E 01	0.18141899E 01
3.75	0.13551766E 01	0.18174518E 01
3.80	0.13527516E 01	0.18206090E 01
3.85	0.13503865E 01	0.18236671E 01
3.90	0.13480789E 01	0.18266304E 01
3.95	0.13458268E 01	0.18295027E 01
4.00	0.13436288E 01	0.18322894E 01

Table D-36

THE ROOTS $t_1[Q\exp(i60^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.55787475E 00	0.88330974E 00
0.10	0.60512120E 00	0.88627898E 00
0.15	0.65108723E 00	0.89112059E 00
0.20	0.69572591E 00	0.89774631E 00
0.25	0.73899173E 00	0.90606698E 00
0.30	0.78084134E 00	0.91599235E 00
0.35	0.82123359E 00	0.92743080E 00
0.40	0.86012989E 00	0.94028924E 00
0.45	0.89749453E 00	0.95447290E 00
0.50	0.93329515E 00	0.96988524E 00
0.55	0.96750307E 00	0.98642788E 00
0.60	0.10000937E 01	0.10040007E 01
0.65	0.10310474E 01	0.10225017E 01
0.70	0.10603493E 01	0.10418278E 01
0.75	0.10879903E 01	0.10618740E 01
0.80	0.11139675E 01	0.10825350E 01
0.85	0.11382845E 01	0.11037046F 01
0.90	0.11609519E 01	0.11252772E 01
0.95	0.11819877E 01	0.11471478E 01
1.00	0.12014173E 01	0.11692132E 01
1.05	0.12192739E 01	0.11913729E 01
1.10	0.12355982E 01	0.12135297E 01
1.15	0.12504381E 01	0.12355911E 01
1.20	0.12638487E 01	0.12574701E 01
1.25	0.12758910E 01	0.12790859E 01
1.30	0.12866317E 01	0.13003649L 01
1.35	0.12961422E 01	0.13212414E 01
1.40	0.13044969E 01	0.13416579E 01
1.45	0.13117730E 01	0.13615653E 01
1.50	0.13180488E 01	0.13809235E 01
1.55	0.13234024E 01	0.13997006E 01
1.60	0.13279115E 01	0.14178729E 01
1.65	0.13316514E 01	0.14354243E 01
1.70	0.13346948E 01	0.14523458E 01
1.75	0.13371111E 01	0.14686345E 01
1.80	0.13389657E 01	0.14842934E 01
1.85	0.13403198E 01	0.14993300E 01
1.90	0.13412301E 01	0.15137556E 01
1.95	0.13417490E 01	0.15275852E 01
2.00	0.13419241E 01	0.15408358F 01

Table D-36 (Cont'd)

THE ROOTS $t_1[Q\exp(i160^\circ)]$

Q	Real t_1	Imag t_1
2.05	0.13417988E 01	0.15535264E 01
2.10	0.13414124E 01	0.15656778E 01
2.15	0.13407996E 01	0.15773114E 01
2.20	0.13399923E 01	0.15884490E 01
2.25	0.13390183E 01	0.15991122E 01
2.30	0.13379026E 01	0.16093231E 01
2.35	0.13366670E 01	0.16191033E 01
2.40	0.13353314E 01	0.16284736E 01
2.45	0.13339123E 01	0.16374546E 01
2.50	0.13324250E 01	0.16460656E 01
2.55	0.13308825E 01	0.16543256E 01
2.60	0.13292961E 01	0.16622528E 01
2.65	0.13276760E 01	0.16698641E 01
2.70	0.13260309E 01	0.16771757E 01
2.75	0.13243681E 01	0.16842032E 01
2.80	0.13226943E 01	0.16909612E 01
2.85	0.13210151E 01	0.16974635E 01
2.90	0.13193352E 01	0.17037234E 01
2.95	0.13176589E 01	0.17097531E 01
3.00	0.13159896E 01	0.17155641E 01
3.05	0.13143306E 01	0.17211673E 01
3.10	0.13126843E 01	0.17265733E 01
3.15	0.13110528E 01	0.17317916E 01
3.20	0.13094380E 01	0.17368315E 01
3.25	0.13078413E 01	0.17417016E 01
3.30	0.13062641E 01	0.17464100E 01
3.35	0.13047074E 01	0.17509641E 01
3.40	0.13031720E 01	0.17553715E 01
3.45	0.13016584E 01	0.17596387E 01
3.50	0.13001673E 01	0.17637721E 01
3.55	0.12986987E 01	0.17677780E 01
3.60	0.12972533E 01	0.17716620E 01
3.65	0.12958307E 01	0.17754293E 01
3.70	0.12944315E 01	0.17790854E 01
3.75	0.12930553E 01	0.17826346E 01
3.80	0.12917022E 01	0.17860818E 01
3.85	0.12903719E 01	0.17894312E 01
3.90	0.12890643E 01	0.17926870E 01
3.95	0.12877794E 01	0.17958525E 01
4.00	0.12865165E 01	0.17989323E 01

Table D-37

THE ROOTS $t_1[Q\exp(i185^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.55272371E 00	0.90323513E 00
0.10	0.59379385E 00	0.92450963E 00
0.15	0.63267811E 00	0.94605481E 00
0.20	0.66944774E 00	0.96780308E 00
0.25	0.70417416E 00	0.98968890E 00
0.30	0.73692904E 00	0.10116489E 01
0.35	0.76778425E 00	0.10336226E 01
0.40	0.79681196E 00	0.10555521E 01
0.45	0.82408451E 00	0.10773826E 01
0.50	0.84967438E 00	0.10990626E 01
0.55	0.87365402E 00	0.11205443E 01
0.60	0.89609571E 00	0.11417833E 01
0.65	0.91707145E 00	0.11627391E 01
0.70	0.93665271E 00	0.11833750E 01
0.75	0.95491019E 00	0.12036581E 01
0.80	0.97191364E 00	0.12235598E 01
0.85	0.98773161E 00	0.12430547E 01
0.90	0.10024311E 01	0.12621218E 01
0.95	0.10160779E 01	0.12807435E 01
1.00	0.10287353E 01	0.12989057E 01
1.05	0.10404652E 01	0.13165977E 01
1.10	0.10513267E 01	0.13338120E 01
1.15	0.10613769E 01	0.13505439E 01
1.20	0.10706703E 01	0.13667914E 01
1.25	0.10792591E 01	0.13825552E 01
1.30	0.10871927E 01	0.13978378E 01
1.35	0.10945178E 01	0.14126440E 01
1.40	0.11012786E 01	0.14269800E 01
1.45	0.11075169E 01	0.14408536E 01
1.50	0.11132713E 01	0.14542738E 01
1.55	0.11185788E 01	0.14672507E 01
1.60	0.11234731E 01	0.14797952E 01
1.65	0.11279862E 01	0.14919189E 01
1.70	0.11321477E 01	0.15036336E 01
1.75	0.11359849E 01	0.15149520E 01
1.80	0.11395232E 01	0.15258866E 01
1.85	0.11427864E 01	0.15364501E 01
1.90	0.11457959E 01	0.15466552E 01
1.95	0.11485717E 01	0.15565148E 01
2.00	0.11511327E 01	0.15660413E 01

Table D-37 (Cont'd)

THE ROOTS t_1 [Qexp(185°)]

Q	Real t_1	Imag t_1
2.05	0.11534956E 01	0.15752470E 01
2.10	0.11556762E 01	0.15841441E 01
2.15	0.11576888E 01	0.15927444E 01
2.20	0.11595468E 01	0.16010594E 01
2.25	0.11612622E 01	0.16091004E 01
2.30	0.11628464E 01	0.16168781E 01
2.35	0.11643097E 01	0.16244030E 01
2.40	0.11656614E 01	0.16316851E 01
2.45	0.11669100E 01	0.16387344E 01
2.50	0.11680640E 01	0.16455600E 01
2.55	0.11691304E 01	0.16521710E 01
2.60	0.11701158E 01	0.16585760E 01
2.65	0.11710268E 01	0.16647833E 01
2.70	0.11718684E 01	0.16708008E 01
2.75	0.11726464E 01	0.16766361E 01
2.80	0.11733655E 01	0.16822965E 01
2.85	0.11740302E 01	0.16877888E 01
2.90	0.11746443E 01	0.16931199E 01
2.95	0.11752114E 01	0.16982958E 01
3.00	0.11757355E 01	0.17033230E 01
3.05	0.11762194E 01	0.17082068E 01
3.10	0.11766662E 01	0.17129532E 01
3.15	0.11770783E 01	0.17175671E 01
3.20	0.11774585E 01	0.17220537E 01
3.25	0.11778086E 01	0.17264180E 01
3.30	0.11781315E 01	0.17306643E 01
3.35	0.11784288E 01	0.17347971E 01
3.40	0.11787021E 01	0.17388207E 01
3.45	0.11789533E 01	0.17427392E 01
3.50	0.11791839E 01	0.17465561E 01
3.55	0.11793955E 01	0.17502753E 01
3.60	0.11795888E 01	0.17539003E 01
3.65	0.11797659E 01	0.17574345E 01
3.70	0.11799274E 01	0.17608808E 01
3.75	0.11800747E 01	0.17642426E 01
3.80	0.11802084E 01	0.17675228E 01
3.85	0.11803295E 01	0.17707242E 01
3.90	0.11804390E 01	0.17738494E 01
3.95	0.11805380E 01	0.17769009E 01
4.00	0.11806262E 01	0.17798814E 01

Table D-37 (Cont'd)

THE ROOTS $t_1[Q\exp(i185^{\circ})]$

Q	Real t_1	Imag t_1
4.05	0.11807057E 01	0.17827930E 01
4.10	0.11807755E 01	0.17856382E 01
4.15	0.11808374E 01	0.17884190E 01
4.20	0.11808920E 01	0.17911376E 01
4.25	0.11809392E 01	0.17937960E 01
4.30	0.11809799E 01	0.17963959E 01
4.35	0.11810139E 01	0.17989394E 01
4.40	0.11810424E 01	0.18014282E 01
4.45	0.11810656E 01	0.18038640E 01
4.50	0.11810828E 01	0.18062485E 01
4.55	0.11810960E 01	0.18085830E 01
4.60	0.11811046E 01	0.18108693E 01
4.65	0.11811090E 01	0.18131086E 01
4.70	0.11811099E 01	0.18153025E 01
4.75	0.11811069E 01	0.18174522E 01
4.80	0.11811005E 01	0.18195591E 01
4.85	0.11810905E 01	0.18216244E 01
4.90	0.11810781E 01	0.18236492E 01
4.95	0.11810635E 01	0.18256346E 01
5.00	0.11810452E 01	0.18275822E 01

Table D-38

THE ROOTS $t_1[Q\exp(i120^\circ)]$

Q	Real t_1	Imag t_1
0	0.50939649E 00	0.88230059E 00
0.5	0.53335264E 00	0.92379388E 00
0.10	0.55617675E 00	0.96332640E 00
0.15	0.57791819E 00	0.10009836E 01
0.20	0.59862544E 00	0.10368496E 01
0.25	0.61834597E 00	0.10710067E 01
0.30	0.63712609E 00	0.11035348E 01
0.35	0.65501083E 00	0.11345121E 01
0.40	0.67204385E 00	0.11640141E 01
0.45	0.68826737E 00	0.11921141E 01
0.50	0.70372204E 00	0.12188824E 01
0.55	0.71844704E 00	0.12443868E 01
0.60	0.73247990E 00	0.12686925E 01
0.65	0.74585659E 00	0.12918615E 01
0.70	0.75861146E 00	0.13139536E 01
0.75	0.77077727E 00	0.13350254E 01
0.80	0.78238522E 00	0.13551310E 01
0.85	0.79346499E 00	0.13743217E 01
0.90	0.80404471E 00	0.13926463E 01
0.95	0.81415112E 00	0.14101511E 01
1.00	0.82380947E 00	0.14268798E 01
1.05	0.83304375E 00	0.14428741E 01
1.10	0.84187656E 00	0.14581729E 01
1.15	0.85032927E 00	0.14728134E 01
1.20	0.85842206E 00	0.14868306E 01
1.25	0.86617402E 00	0.15002573E 01
1.30	0.87360308E 00	0.15131249E 01
1.35	0.88072622E 00	0.15254624E 01
1.40	0.88755932E 00	0.15372977E 01
1.45	0.89411753E 00	0.15486568E 01
1.50	0.90041494E 00	0.15595642E 01
1.55	0.90646499E 00	0.15700433E 01
1.60	0.91228019E 00	0.15801154E 01
1.65	0.91787246E 00	0.15898015E 01
1.70	0.92325296E 00	0.15991207E 01
1.75	0.92843213E 00	0.16080914E 01
1.80	0.93342005E 00	0.16167306E 01
1.85	0.93822598E 00	0.16250547E 01
1.90	0.94285875E 00	0.16330789E 01
1.95	0.94732666E 00	0.16408174E 01
2.00	0.95163757E 00	0.16482842E 01

Table D-38 (Cont'd)

THE ROOTS t_1 [Qexp(i120)]

Q	Real t_1	Imag t_1
2.05	0.95579901E 00	0.16554920E 01
2.10	0.95981773E 00	0.16624525E 01
2.15	0.96370045E 00	0.16691776E 01
2.20	0.96745347E 00	0.16756780E 01
2.25	0.97108258E 00	0.16819637E 01
2.30	0.97459336E 00	0.16880445E 01
2.35	0.97799111E 00	0.16939297E 01
2.40	0.98128080E 00	0.16996276E 01
2.45	0.98446722E 00	0.17051465E 01
2.50	0.98755471E 00	0.17104941E 01
2.55	0.99054762E 00	0.17156778E 01
2.60	0.99344977E 00	0.17207047E 01
2.65	0.99626514E 00	0.17255810E 01
2.70	0.99899732E 00	0.17303131E 01
2.75	0.10016494E 01	0.17349068E 01
2.80	0.10042252E 01	0.17393681E 01
2.85	0.10067274E 01	0.17437021E 01
2.90	0.10091592E 01	0.17479140E 01
2.95	0.10115231E 01	0.17520082E 01
3.00	0.10138218E 01	0.17559898E 01
3.05	0.10160580E 01	0.17598630E 01
3.10	0.10182339E 01	0.17636319E 01
3.15	0.10203519E 01	0.17673001E 01
3.20	0.10224140E 01	0.17708720E 01
3.25	0.10244226E 01	0.17743508E 01
3.30	0.10263794E 01	0.17777400E 01
3.35	0.10282863E 01	0.17810431E 01
3.40	0.10301454E 01	0.17842627E 01
3.45	0.10319579E 01	0.17874021E 01
3.50	0.10337257E 01	0.17904643E 01
3.55	0.10354506E 01	0.17934518E 01
3.60	0.10371338E 01	0.17963672E 01
3.65	0.10387769E 01	0.17992129E 01
3.70	0.10403810E 01	0.18019915E 01
3.75	0.10419478E 01	0.18047052E 01
3.80	0.10434782E 01	0.18073560E 01
3.85	0.10449735E 01	0.18099459E 01
3.90	0.10464349E 01	0.18124771E 01
3.95	0.10478636E 01	0.18149519E 01
4.00	0.10492603E 01	0.18173711E 01

Appendix E

THE HEIGHT GAIN FUNCTIONS FOR $q = 0$ AND $q = \infty$

In this appendix we will discuss a set of FORTRAN programs which can be employed to obtain values for the height gain functions for the two limiting cases of $q = 0$ and $q = \infty$. The case $q = 0$ is important because it occurs in the theory of the propagation of vertically polarized waves around a convex surface which is perfectly conducting. The case $q = \infty$ is encountered in the theory of the propagation of horizontally polarized waves around a perfectly conducting surface. However, these limiting cases are also important in certain other problems in which the convex surface is the interface between a dielectric with the propagation constant

$$k_1^2 = \omega^2 \varepsilon_1 \mu_1 + i\omega\mu_1\sigma_1 \quad , \quad r < a \tag{E-1}$$

and "free space" with the propagation constant

$$k^2 = \omega^2 \varepsilon_o \mu_o \quad , \quad r > a \tag{E-2}$$

Let q_v and q_h denote the impedance parameters for vertical and horizontal polarization, respectively. For $ka \to \infty$, the impedance parameters are given by

$$q_v = i(ka/2)^{\frac{1}{3}}(k/k_1)\sqrt{1 - (k/k_1)^2} \tag{E-3}$$

$$q_h = i(ka/2)^{\frac{1}{3}}(k_1/k)\sqrt{1 - (k/k_1)^2} = (k_1/k)^2 q_v \tag{E-4}$$

These are actually very complex relationships since the parameters ε_1, μ_1 and σ_1 are functions of frequency ω. However, if $(k_1/k) \to \infty$, we observe that $q_v \to 0$ and $q_h \to \infty$. Therefore, if the frequency ω is held fixed and the conductivity $\sigma_1 \to \infty$, we have $q_v \to 0$ and $q_h \to \infty$. However, if the conductivity σ_1 is held "fixed" and we

permit the frequency ω to vary, it is possible for ω to become so large that $q_v \to \infty$ even though σ_c may be large numerically. Therefore, the case $q = \infty$ plays a very important role in problems involving the convex interface between a dielectric body and free space. This case is also an interesting one because of the fact that the expansion given in Eq. (2-22) for the roots $t_s(q)$, namely

$$t_s(q) = t_s^\infty + \frac{1}{q} + \frac{t_s^\infty}{3q^3} + \cdots \qquad \text{(E-5)}$$

can be, for very large values of q, adequately represented by*

$$t_s(q) \approx t_s^\infty + \frac{1}{q} \qquad \text{(E-6)}$$

---

*We must be careful to observe that Eq. (E-6) holds for only a finite number of the roots $t_s(q)$ because eventually the modulus of the root t_s^∞ will become so large that the neglected terms will have to be considered. If one examines the highest powers of t_s^∞ in the coefficients $B_n(t_s^\infty)$ in Eq. (2-22) one will recognize that these are the coefficients in the well known expansion

$$\frac{\tanh^{-1} x}{x} = 1 + \frac{x^2}{3} + \frac{x^4}{5} + \frac{x^6}{7} + \cdots, \quad x < 1$$

and therefore we will have to replace Eq. (E-6) by the result

$$t_s(q) \approx t_s^\infty + \frac{1}{\sqrt{t_s^\infty}} \tanh^{-1}(\sqrt{t_s^\infty}/q) \quad , |\sqrt{t_s^\infty}/q| < 1$$

in which the modulus of q is very large, and $30^\circ < \arg(q) < 210^\circ$. One will always find that as s is increased that one will reach the situation in which Eq. (2-22) is no longer a convergent series. However, if ξ in the diffraction functions $V_0(\xi,q)$, $V_1(\xi,q)$ and $V_{11}(\xi,q)$ is positive, we can generally observe that the number of terms in the residue series which must be used is finite and hence we can determine how to treat the computation of the root $t_s(q)$ for large values of q.

The approximation in Eq. (E-6) reveals that we can sometimes use the approximation

$$\exp(i\xi t_s) \approx \exp(i\xi t_s^\infty)\exp(i\xi/q) \qquad \text{(E-7)}$$

which permits us to extend the usefulness of the functions $V_o(\xi,\infty)$ $V_1(\xi,\infty)$ and $V_{11}(\xi,\infty)$ by using the approximations

$$V_o(\xi,q) \approx \exp(i\xi/q)V_o(\xi,\infty) \qquad \text{(E-8a)}$$

$$V_1(\xi,q) \approx \exp(i\xi/q)V_1(\xi,\infty) \qquad \text{(E-8b)}$$

$$V_{11}(\xi,q) \approx \exp(i\xi/q)V_{11}(\xi,\infty) \qquad \text{(E-8c)}$$

In each of these expressions we must have $\xi > 0$ and q must have its argument in the range $30^o < \arg(q) < 210^o$ and the modulus of q must be "very large". More precise descriptions of the applicability of the approximations in Eq. (E-8) have not yet been established. This is really a very complex problem since Eq. (E-8) can be used only if $\xi > 0$, but it is clear from considering the form

$$\exp[i\xi(t_s - t_s^\infty - \frac{1}{q})] = \exp[i\xi(\frac{t_s^\infty}{3q^3} + \frac{1}{4q^4} + \frac{t_s^{\infty 2}}{5q^5} + \cdots)] \qquad \text{(E-9)}$$

that one must be certain that the expression in parenthesis must have a positive real part if Eq. (E-6) is to be used when $\xi \to \infty$.

The situation is somewhat different when we consider the case in which $q \to 0$. From Eq. (2-21) we find that in this case we have the approximation

$$t_s(q) \approx t_s^o + \frac{1}{t_s^o}\, q \qquad \text{(E-10)}$$

and therefore the "correction term" is dependent upon s and hence we do not find that the functions $V_o(\xi,0)$, $V_1(\xi,0)$ and $V_{11}(\xi,0)$ suffice to describe the behavior when q is very small. We find that we have to define three additional functions $V_o^{(-1)}(\xi,0)$,

$$V_o^{(-1)}(\xi,q) = -i\sqrt{i\pi\xi} \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{t_s(t_s - q^2)} \qquad \text{(E-11a)}$$

$$V_1^{(-1)}(\xi,q) = -i2\sqrt{i\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{t_s(t_s - q^2)w_1(t_s)} \qquad \text{(E-11b)}$$

$$V_{11}^{(-1)}(\xi,q) = i2\sqrt{-i\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{t_s(t_s - q^2)[w_1(t_s)]^2} \qquad \text{(E-11c)}$$

We can employ these functions when q is in the vicinity of 0 so as to obtain the approximations

$$V_o(\xi,q) \approx V_o(\xi,0) - \xi q V_o^{(-1)}(\xi,0) \qquad \text{(E-12a)}$$

$$V_1(\xi,q) \approx V_1(\xi,0) - \xi q V_1^{(-1)}(\xi,0) \qquad \text{(E-12b)}$$

$$V_{11}(\xi,q) \approx V_{11}(\xi,0) - \xi q V_{11}^{(-1)}(\xi,0) \qquad \text{(E-12c)}$$

As in the case of the approximations given in Eq. (E-6) and Eq. (E-8), the exact conditions under which the approximations* in Eq. (E-10) and Eq. (E-12) can be used cannot be stated explicitly and we leave undefined what we mean by "small values of q." Since

---

*For $s \to \infty$, the approximation in Eq. (E-10) will apparently be valid provided the modulus of q is such that the modulus of the combination $(q/\sqrt{t_s^o})$ is less than unity. This seems to be the case since the dominant terms as $s \to \infty$ are

$$t_s(q) \approx t_s^o + \frac{1}{t_s^o} q \left[1 + \frac{1}{3} z + \frac{1}{5} z^2 + \frac{1}{7} z^3 + \cdots\right], \quad z = \frac{q^2}{t_s^o}$$

and therefore we see that the series converges like $\tanh^{-1} z$ and that $t_s(q) \approx t_s^o + (1/\sqrt{t_s^o})\tanh^{-1}(q/\sqrt{t_s^o})$ and we must require that $(q/\sqrt{t_s^o})$ be less than unity in absolute value.

the form of the coefficients $B_n(t_s^o)$ in Eq. (2-21) reveals that t_s^o occurs only in inverse powers, we can be assured that as s increases that the approximations given in Eq. (E-10) and Eq. (E-12) become increasingly accurate representations provided the modulus of q is "small" and $30^o < \arg(q) < 210^o$,

The above discussion has emphasized the importance of the diffraction functions $V_o(\xi,0)$, $V_1(\xi,0)$ and $V_{11}(\xi,0)$ for $q = 0$ and $V_o(\xi,\infty)$, $V_1(\xi,\infty)$ and $V_{11}(\xi,\infty)$ for $q = \infty$. These functions are dependent only upon the distance parameter ξ and the impedance parameter q. We want now to bring in the dependance of the diffraction phenomenon upon height. In Section 1 we have defined the functions

$$V(x,y,y_o,q) = 2\sqrt{i\pi x}\sum_{s=1}^{\infty}\frac{\exp(ixt_s)w_1(t_s - y)w_1(t_s - y_o)}{t_s[w_1(t_s)]^2 - [w_1'(t_s)]^2} \qquad \text{(E-13)}$$

$$E(x,y,q) = i2\sqrt{\pi}\sum_{s=1}^{\infty}\frac{\exp(ixt_s)w_1(t_s - y)}{t_s[w_1(t_s)]^2 - [w_1'(t_s)]^2} \qquad \text{(E-14)}$$

We can make use of the equation defining t_s to write

$$w_1'(t_s) = qw_1(t_s) \qquad \text{(E-15)}$$

to express the functions $V(x,y,y_o,q)$ and $E(x,y,q)$ in the forms

$$V(x,y,y_o,q) = 2\sqrt{i\pi x}\sum_{s=1}^{\infty}\frac{\exp(ixt_s)}{t_s - q^2} f_s(y,q)f_s(y_o,q) \qquad \text{(E-16)}$$

$$E(x,y,q) = i2\sqrt{\pi}\sum_{s=1}^{\infty}\frac{\exp(ixt_s)}{(t_s - q^2)w_1(t_s)} f_s(y,q) \qquad \text{(E-17)}$$

where $f_s(y,q)$ is the height gain function

$$f_s(y,q) = \frac{w_1(t_s - y)}{w_1(t_s)} = \frac{Ai[-a_s + y\exp(-i\frac{2}{3}\pi)]}{Ai(-a_s)} \qquad \text{(E-18)}$$

and

$$V(x,y,y_o,q) = -2\sqrt{i\pi x}\; q^{-2} \sum_{s=1}^{\infty} \frac{\exp(ixt_s)}{1 - t_s q^{-2}}\, g_s(y,q) g_s(y_o,q) \qquad \text{((E-19)}$$

$$E(x,y,q) = -i2\sqrt{\pi}\; q^{-2} \sum_{s=1}^{\infty} \frac{\exp(ixt_s)}{(1 - t_s q^{-2}) w_1'(t_s)}\, g_s(y,q) \qquad \text{(E-20)}$$

where $g_s(y,q)$ is the height gain function

$$g_s(y,q) = \frac{w_1(t_s - y)}{w_1'(t_s)} = \frac{Ai[-a_s + y\exp(-i\frac{2}{3}\pi)]}{\exp(i\frac{1}{3}\pi) Ai(-a_s)} \qquad \text{(E-21)}$$

The relationship between the t_s and the a_s is

$$t_s = a_s \exp(i\tfrac{1}{3}\pi)$$

and the form taken by Eq. (E-15) when expressed in terms of the functions $Ai(z)$ and $Ai'(z)$ is

$$Ai'(-a_s) + q\exp(i\tfrac{1}{3}\pi) Ai(-a_s) = 0$$

Since the Airy functions are solution of the differential equation $d^2w(z)/dz^2 = zw(z)$, we observe that the height gain functions $f_s(y,q)$ and $g_s(y,q)$ are solutions of the differential equation

$$\frac{d^2h(y)}{dy^2} = (t_s - y)h(y) \qquad \text{(E-22)}$$

which satisfy the boundary conditions

$$f_s(0,q) = 1 \quad , \quad \frac{df_s(0,q)}{dy} = -q \qquad \text{(E-23)}$$

$$g_s(0,q) = \frac{1}{q} \quad , \quad \frac{dg_s(0,q)}{dy} = -1 \tag{E-24}$$

The functions $V(x,y,y_o,q)$ and $E(x,y,q)$ can be referred to as the diffraction functions for vertical polarization because they often occur in situations in which the electric vector the wave is perpendicular to the surface over which the wave is being propagated. In the description of the propagation of horizontally polarized waves in which the electric vector is parallel to the surface, we usually encounter the diffraction functions

$$W(x,y,y_o,q) = 4\sqrt{-i\pi}\, x^{3/2} \sum_{s=1}^{\infty} \frac{\exp(ixt_s)}{1 - t_s q^{-2}}\, g_s'(y,q) g_s'(y_o,q) \tag{E-25}$$

$$F(x,y,q) = -i2\sqrt{\pi} \sum_{s=1}^{\infty} \frac{\exp(ixt_s)}{(1 - t_s q^{-2}) w_1'(t_s)}\, g_s'(y,q) \tag{E-26}$$

where

$$g_s'(y,q) = \frac{\partial g_s(y,q)}{\partial y}$$

If we compare Eqs. (E-25) and (E-26) with Eqs. (E-19) and (E-20), respectively, we see that

$$W(x,y,y_o,q) = i2xq^2 \frac{\partial^2 V(x,y,y_o,q)}{\partial y\, \partial y_o} \tag{E-27}$$

$$F(x,y,q) = q^2 \frac{\partial E(x,y,q)}{\partial y} \tag{E-28}$$

The functions $V_o(\xi,q)$ and $V_1(\xi,q)$ have been defined so that

$$V(x,0,0,q) = 2V_o(x,q) \tag{E-29}$$

$$E(x,0,q) = V_1(x,q) \tag{E-30}$$

By analogy with these results, we define the functions $U_o(\xi,q)$ and $U_1(\xi,q)$ by means of the relations

$$W(x,0,0,q) = 2U_o(x,q) \tag{E-31}$$

$$F(x,0,q) = U_1(x,q) \tag{E-32}$$

where

$$U_o(\xi,q) = 2\sqrt{-i\pi}\, x^{3/2} \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{1 - t_s q^{-2}} \tag{E-34}$$

$$U_1(\xi,q) = i2\sqrt{\pi} \sum_{s=1}^{\infty} \frac{\exp(i\xi t_s)}{(1 - t_s q^{-2}) w_1'(t_s)} \tag{E-35}$$

We also observe that

$$[-2i\xi q^2 V_o(\xi,q)] \xrightarrow[q \to \infty]{} U_o(\xi,q) \tag{E-36}$$

$$[-qV_1(\xi,q)] \xrightarrow[q \to \infty]{} U_1(\xi,q) \tag{E-37}$$

Let us now return to a consideration of the height gain functions and designate by $f_s(y)$ and $g_s(y)$ the special cases which arise when $q = 0$ and $q = \infty$, respectively. We then have

$$f_s(y) = \frac{w_1(t_s^o - y)}{w_1(t_s^o)} = \frac{Ai[-\beta_s + y\exp(-i\frac{2}{3}\pi)]}{Ai(-\beta_s)} \tag{E-38}$$

where

$$w_1'(t_s^o) = Ai(-\beta_s) = 0$$

and

$$g_s(y) = -\frac{w_1(t_s^\infty - y)}{w_1'(t_s^\infty)} = \frac{Ai[-\alpha_s + y\exp(-i\frac{2}{3}\pi)]}{\exp(-i\frac{2}{3}\pi) Ai'(-\alpha_s)} \tag{E-39}$$

where

$$w_1(t_s^\infty) = Ai(-\alpha_s) = 0$$

We observe that these functions satisfy the differential equations

$$\frac{d^2 f_s(y)}{dy^2} = (t_s^o - y) f_s(y) \qquad \text{(E-40)}$$

$$\frac{d^2 g_s(y)}{dy^2} = (t_s^\infty - y) g_s(y) \qquad \text{(E-41)}$$

with the boundary conditions

$$f_s(0) = 1 \quad , \quad \frac{df_s(0)}{dy} = 0 \qquad \text{(E-42)}$$

$$g_s(0) = 0 \quad , \quad \frac{dg_s(0)}{dy} = 1 \qquad \text{(E-43)}$$

We have chosen to define $f_s(y,0) = f_s(y)$ and $g_s(y,\infty) = -g_s(y)$ in order that the functions $f_s(y)$ and $g_s(y)$ agree with the definitions in the tables published by Azriliant and Belkina (Ref. 1).

For moderately small values of we can evaluate the height gain functions by means of a Taylor series. Let us develop the theory of the Taylor series by considering solutions of the differential equation

$$\frac{d^2 Y(z)}{dz^2} = zY(z) \qquad \text{(E-44)}$$

which are expressed in the form

$$Y(t + x) = Y(t)u(x,t) + Y'(t)v(x,t) \qquad \text{(E-45a)}$$

$$Y'(t + x) = Y(t)u'(x,t) + Y'(t)v'(x,t) \qquad \text{(E-45b)}$$

The functions $u(x,t)$ and $v(x,t)$ are solutions of the differential equation

$$\frac{d^2y(x)}{dx^2} = (x + t)y(x) \qquad \text{(E-46)}$$

We will let a prime denote differentiation with respect to x, i.e.,

$$u'(x,t) = \frac{\partial u(x,t)}{\partial x} \quad , \quad v'(x,t) = \frac{\partial v(x,t)}{\partial x}$$

The boundary conditions on $u(x,t)$ and $v(x,t)$ are

$$u(0,t) = 1 \quad , \; u'(0,t) = 0 \quad , \quad v(0,t) = 0 \quad , \; v'(0,t) = 1$$

We can express the solution for these functions in the form of Taylor series which involve a set of coefficients $a_n(t)$ and $b_n(t)$ in the following manner:

$$u(x,t) = 1 + \sum_{n=2}^{\infty} a_n(t)\frac{x^n}{n!} \qquad \text{(E-47a)}$$

$$v(x,t) = x + \sum_{n=2}^{\infty} b_n(t)\frac{x^n}{n!} \qquad \text{(E-47b)}$$

$$u'(x,t) = \sum_{n=2}^{\infty} n a_n(t)\frac{x^{n-1}}{n!} \qquad \text{(E-47c)}$$

$$v'(x,t) = 1 + \sum_{n=2}^{\infty} n b_n(t)\frac{x^{n-1}}{n!} \qquad \text{(E-47d)}$$

When the expansions in Eq. (E-47) are inserted in Eq. (E-46), the $a_n(t)$ and $b_n(t)$ are found to satisfy the recursion relations

$$c_2(t) - tc_o(t) = 0$$

$$c_3(t) - 1c_o(t) - tc_1(t) = 0$$

$$c_4(t) - 2c_1(t) - tc_2(t) = 0$$

$$c_5(t) - 3c_2(t) - tc_3(t) = 0$$

and, in general,

$$c_n(t) = (n - 2)c_{n-3}(t) + tc_{n-2}(t) \quad , n > 3 \qquad \text{(E-48)}$$

where $c_n(t)$ denotes either $a_n(t)$ or $b_n(t)$. The first few terms are

$$a_0(t) = 1 \quad , \quad b_0(t) = 0$$
$$a_1(t) = 0 \quad , \quad b_1(t) = 1$$
$$a_2(t) = t \quad , \quad b_2(t) = 0$$
$$a_3(t) = 1 \quad , \quad b_3(t) = t$$
$$a_4(t) = t^2 \quad , \quad b_4(t) = 2$$
$$a_5(t) = 4t \quad , \quad b_5(t) = t^2$$

In Program E-1 we have presented an algorithm for determining the numerical values of the $a_n(x,t)$ and $b_n(x,t)$. The subroutine is identified as SUBRØUTINE STOKES(N,A,B,C,D,U,V). The integer N is the largest value of n for which a coefficient is desired. We have used the FORTRAN variables in the following manner:

$$a_I(x,t) = A(I) + iC(I)$$
$$b_I(x,t) = B(I) + iD(I)$$
$$t = U + iV$$

In Program E-2 we employ the coefficients obtained from this subroutine to compute Y(t+x). We have restricted this program to real values of t and x, which we denote in FORTRAN by T and X, respectively. The reader will observe that we let Y(t) = WT, Y'(t) = WTP, and Y(t+x) = WTPX. The integer N is the number of terms which we ask SUBRØUTINE TAYSUM(T,X,WT,WPT,WTPX,N) to employ. We wrote a main program in which N = 25 and

```
X    = 1.0D0
T    = 0.0D0
WT   = 0.35502805388781 7D0
WPT  =-0.25881940379280 7D0
```

Program E-1

```
      SUBRØUTINE STOKES(N,A,B,C,D,U,V)
C-----THIS SUBRØUTINE EVALUATES THE CØEFFICIENTS IN THE TAYLOR
C-----SERIES OF THE AIRY FUNCTION
      DØUBLE PRECISIØN A(N),B(N),C(N),D(N),U,V,ZERØ,ØNE,TWØ,DBIM2
      DATA ZERØ,ØNE,TWØ/ 0.0D0,1.0D0,2.0D0/
      DBIM2(K) = DBLE(FLØAT(K-2))
      IF(N.LT.5) GØ TØ 20
      A(1) = ZERØ
      C(1) = ZERØ
      B(1) = ØNE
      D(1) = ZERØ
      A(2) = U
      C(2) = ZERØ
      B(2) = ZERØ
      D(2) = ZERØ
      A(3) = ØNE
      C(3) = ZERØ
      B(3) = U
      D(3) = ZERØ
      A(4) = U*U
      C(4) = ZERØ
      B(4) = TWØ
      D(4) = ZERØ
      IF(DABS(V).GT.1.0D-37) GØ TØ  10
C-----TRANSFER TØ 10 IF T = CMPLX(U,V) IS NØT REAL
      DØ 2 I = 5,N
      A(I) = DBIM2(I)*A(I-3) + U*A(I-2)
      C(I) = ZERØ
      B(I) = DBIM2(I)*B(I-3) + U*B(I-2)
      D(I) = ZERØ
    2 CØNTINUE
      RETURN
   10 C(2) = V
      D(3) = V
      A(4) = A(4) - V*V
      C(4) = TWØ*U*V
      DØ 1 I = 5,N
      A(I) = DBIM2(I)*A(I-3) + U*A(I-2) - V*C(I-2)
      C(I) = DBIM2(I)*C(I-3) + U*C(I-2) + V*A(I-2)
      B(I) = DBIM2(I)*B(I-3) + U*B(I-2) - V*D(I-2)
      D(I) = DBIM2(I)*D(I-3) + U*D(I-2) + V*B(I-2)
    1 CØNTINUE
      RETURN
   20 WRITE(6,110) N
  110 FØRMAT(43H1SUBRØUTINE STØKES IS BEING ASKED TO USE N =,I5,
     A       34HPRØGRAM EXITS UNDER THIS CØNDITIØN)
      CALL EXIT
      END
```

Program E-2

```
      SUBRØUTINE TAYSUM(T,X,WT,WPT,WTPX,N)
C-----THIS SUBRØUTINE COMPUTES THE AIRY FUNCTION W(T+X) = WT*
C-----U(X,T) + WTP*V(X,T)
      DØUBLE PRECISIØN T,X,WT,WPT,WTPX,WPTPX,SUM,XTNØNF(100)
      DØUBLE PRECISIØN A(100),B(100),C(100),D(100), DFLØAT
      DFLØAT(K) = DBLE(FLØAT(K))
      IF(N.GT.100) GØ TØ 20
      CALL STØKES(N,A,B,C,D,T,0.0D0)
      XTNØNF(1) = X
      DØ 1 I = 2,N
      XTNØNF(I) = X*XTNØNF(I-1)/DFLØAT(I)
    1 CØNTINUE
      SUM = 1.0D0
      DØ 2 I = 1,N
      SUM    = SUM + XTNØNF(I)*A(I)
    2 CØNTINUE
      WTPX   = WT*SUM
      SUM    = X
      DØ 3 I = 1,N
      SUM    = SUM + XTNØNF(I)*B(I)
    3 CØNTINUE
      WTPX   = WTPX + WPT*SUM
      RETURN
   20 WRITE(6,120) N
  120 FØRMAT(43H1SUBRØUTINE TAYSUM IS BEING ASKED TO USE N =,I5,
     A       34HPRØGRAM EXITS UNDER THIS CØNDITIØN)
      CALL EXIT
      END
```

Since this means that we have taken WT = Ai(0) and WTP = Ai'(0), the result obtained, namely

WTPX = .13529241631288O8+00

represents the value of the Airy function

Ai(1) = 0.135292416312881415 5

In Table E-1 we give the values of the coefficients which were generated during the course of this calculation.

The method employed in Program E-2 to compute Y(t+x) does not represent an application of good programming technique if the desired end product is a set of numerical values for the functions

Table E-1

TABLE OF COEFFICIENTS $a_n(0)$ AND $b_n(0)$ FOR $u(x,0)$ AND $v(x,0)$

n	$a_n(0)$	$b_n(0)$
3	.1000000000000000+01	.0000000000000000
4	.0000000000000000	.1999999999999999+01
5	.0000000000000000	.0000000000000000
6	.3999999999999999+01	.0000000000000000
7	.0000000000000000	.1000000000000000+02
8	.0000000000000000	.0000000000000000
9	.2799999999999999+02	.0000000000000000
10	.0000000000000000	.7999999999999999+02
11	.0000000000000000	.0000000000000000
12	.2799999999999999+03	.0000000000000000
13	.0000000000000000	.8799999999999999+03
14	.0000000000000000	.0000000000000000
15	.3639999999999999+04	.0000000000000000
16	.0000000000000000	.1231999999999999+05
17	.0000000000000000	.0000000000000000
18	.5823999999999999+05	.0000000000000000
19	.0000000000000000	.2094399999999999+06
20	.0000000000000000	.0000000000000000
21	.1106559999999999+07	.0000000000000000
22	.0000000000000000	.4188799999999999+07
23	.0000000000000000	.0000000000000000
24	.2434431999999999+08	.0000000000000000
25	.0000000000000000	.9634239999999998+08

$u(x,t)$ and $v(x,t)$. It is more convenient to compute the terms in the series directly rather than to evaluate the $a_n(t)$ and $b_n(t)$ and then sum the polynomial which results when the series is truncated. This is also desirable because the $a_n(t)$ and $b_n(t)$ may "overflow" on the computer (i.e., become so large that they exceed the maximum magnitude of 10^{38} which most FORTRAN compilers permit) and yet the product of the coefficients $a_n(t)$ and $x^n/n!$ may still be significant.*

Let us introduce the notation

$$A_n(x,t) = a_n(t)x^n/n! \qquad \text{(E-49a)}$$

$$B_n(x,t) = b_n(t)x^n/n! \qquad \text{(E-49b)}$$

The $A_n(x,t)$ and $B_n(x,t)$ can be shown to satisfy the recursion formula

$$n(n-1)C_n(x,t) = x^3C_{n-3}(x,t) + tx^2C_{n-2}(x,t) \qquad \text{(E-50)}$$

with the initial values

$$A_0(x,t) = 1 \quad , \quad B_0(x,t) = 0$$

$$A_1(x,t) = 0 \quad , \quad B_1(x,t) = x$$

$$A_2(x,t) = \frac{tx^2}{2} \quad , \quad B_2(x,t) = 0$$

$$A_3(x,t) = \frac{x^3}{6} \quad , \quad B_3(x,t) = \frac{tx^3}{6}$$

$$A_4(x,t) = \frac{t^2x^4}{24} \quad , \quad B_4(x,t) = \frac{x^4}{12}$$

---

*The factor $x^n/n!$ may also "underflow." For example, since $80! = 0.71569457E+119$, we would have $10^{80}/80! \approx 1.4 \times 10^{-39}$ and this is outside the range 10^{-38} to 10^{38} permitted by most compilers.

The coefficients $A_n(x,t)$ and $B_n(x,t)$ can be evaluated with the SUBRØUTINE TERMS(N,AR,AI,BR,BI,XR,XI,TR,TI) which is given in Program E-3. The reader will observe that we have used the following FORTRAN representations for the values of x, t, A_n and B_n:

$$x = XR + iXI \quad , \quad A_I(x,t) = AR(I) + iAI(I)$$
$$t = TR + iTI \quad , \quad B_I(x,t) = BR(I) + IBI(I)$$

Program E-3

```
      SUBROUTINE TERMS(N,AR,AI,BR,BI,XR,XI,TR,TI)
C-----COMPUTE TERMS IN TAYLOR SERIES EXPANSION OF AIRY FUNCTION
      DOUBLE PRECISION AR(N),AI(N),BR(N),BI(N),XR,XI,TR,TI,DFLOAT,
     A                 ARI,AII,BRI,BII,XSQR,XSQI
      DFLOAT(K)= DBLE(FLOAT(K))
      XSQR = XR*XR - XI*XI
      XSQI = 2.0D0*XR*XI
      AR(1)= 0.0D0
      AI(1)= 0.0D0
      AR(2)= (TR*XSQR - TI*XSQI)/(2.0D0)
      AI(2)= (TR*XSQI + TI*XSQR)/(2.0D0)
      AR(3)= (XR*XSQR - XI*XSQI)/(6.0D0)
      AI(3)= (XR*XSQI + XI*XSQR)/(6.0D0)
      AR(4)= (AR(2)*AR(2) - AI(2)*AI(2))/(6.0D0)
      AI(4)= (AR(2)*AI(2))/(3.0D0)
      BR(1)= XR
      BI(1)= XI
      BR(2)= 0.0D0
      BI(2)= 0.0D0
      BR(3)= TR*AR(3) - TI*AI(3)
      BI(3)= TR*AI(3) + TI*AR(3)
      BR(4)  = (XSQR*XSQR - XSQI*XSQI)/12.0D0
      BI(4)  = (XSQR*XSQI)/6.0D0
      DO 1 I=5,N
C-----THE NEXT STEP GENERATES AN EXPRESSION FRØM WHICH AR(I),ETC.
C-----CAN THEN BE CØNSTRUCTED
      ARI  = XR*AR(I-3) - XI*AI(I-3) + TR*AR(I-2) - TI*AI(I-2)
      AII  = XR*AI(I-3) + XI*AR(I-3) + TR*AI(I-2) + TI*AR(I-2)
      BRI  = XR*BR(I-3) - XI*BI(I-3) + TR*BR(I-2) - TI*BI(I-2)
      BII  = XR*BI(I-3) + XI*BR(I-3) + TR*BI(I-2) + TI*BR(I-2)
      AR(I)= (XSQR*ARI - XSQI*AII)/DFLOAT(I*(I-1))
      AI(I)= (XSQR*AII + XSQI*ARI)/DFLOAT(I*(I-1))
      BR(I)= (XSQR*BRI - XSQI*BII)/DFLOAT(I*(I-1))
      BI(I)= (XSQR*BII + XSQI*BRI)/DFLOAT(I*(I-1))
    1 CONTINUE
      RETURN
      END
```

In Program E-4 we present a main program which we used to sum the series of terms obtained from Program E-3. The series is of the form

$$u(x,t) = 1 + A_2(x,t) + A_3(x,t) + A_4(x,t) + \ldots \qquad \text{(E-51a)}$$

$$v(x,t) = x + B_2(x,t) + B_3(x,t) + B_4(x,t) + \ldots \qquad \text{(E-51b)}$$

Program E-4

```
C-----THIS IS A TEST OF SUBRØUTINE TERMS(N,AR,AI,BR,BI,XR,XI,TR,TI)
      DØUBLE PRECISIØN AR(100),AI(100),BR(100),BI(100),XR,XI,TR,TI,
     A        DFLØAT,WTR,WTI,WPTR,WPTI,UR,UI,VR,VI,WTPXR,WTPXI
      WTR     =-0.7258727400
      WTI     = 3.4419277000
      WPTR    =-3.7668668500
      WPTI    =+4.8506065800
      CALL TERMS(40,AR,AI,BR,BI, 1.0D0, 1.0D0, 2.5D0, 2.5D0)
      WRITE(6,110) (I,AR(I),AI(I),BR(I),BI(I),I=1,40)
  110 FORMAT(1H0,I4,4D23.16)
      XR = 1.0D0
      XI = 1.0D0
      UR      = 1.0D0
      UI      = 0.0D0
      DO 10 I= 2,40
      UR      = UR+ AR(I)
      UI      = UI+ AI(I)
   10 CONTINUE
      WRITE(6,500) UR,UI
  500 FORMAT (1H0, 5HUR,UI, 2D25.16)
      VR      = XR
      VI      = XI
      DO 20 I= 2,40
      VR      = VR + BR(I)
      VI      = VI + BI(I)
   20 CONTINUE
      WRITE(6,400) VR,VI
  400 FORMAT (1H0, 5HVR,VI, 2D25.16)
      WTPXR   = WTR*UR-WTI*UI + WPTR*VR-WPTI*VI
      WTPXI   = WTR*UI+WTI*UR + WPTR*VI+WPTI*VR
      WRITE(6,111) WTPXR, WTPXI
  111 FORMAT(1H1,2X,2D25.16)
      CALL EXIT
      END
```

The test that we have conducted in Prog ram E-4 was based upon entries in Harvard Computation Laboratory (Ref. 2) tables of the functions $h_1(z)$ and $h_2(z)$ which satisfy the differential equation

$$\frac{d^2 h_{1,2}(z)}{dz^2} + z h_{1,2}(z) = 0 \qquad \text{(E-52)}$$

If we compare Eq. (E-52) with Eq. (E-44), we see that we can take

$$Y(t+x) = h_1(-t-x)$$
$$Y(t) = h_1(-t)$$
$$Y'(t) = -h_1'(-t)$$

In the test we took

$$t = TR + iTI = 2.5 + i2.5$$
$$x = XR + iXI = 1.0 + i1.0$$

with the initial conditions

$$Y(t) = h_1(-t) = WTR + iWTI = -0.72587274 + i3.44192770$$
$$Y'(t) = -h_1'(-t) = WPTR+iWPTI = 3.76686685 - i4.85060658$$

The computer output

-.215728908898266 3+01 -.9312479583220205+01

is in agreement with the entry from the tables, namely

$$Y(t+x) = h_1(-t-x) = WTPXR + iWTPXI = -2.15728909 - i9.31247959$$

The SUBRØUTINE TERMS(N,AR,AI,BR,BI,XR,XI,TR,TI) can be used as a building block for the development of a step-by-step method of integrating the differential equation in Eq. (E-22) to obtain the height gain functions $f_s(y,q)$ and $g_s(y,q)$.

Since we have shown that the limiting cases $q = 0$ and $q = \infty$ play such important roles, we want now to present programs which permit one to evaluate the Taylor series for the height gain functions $f_s(y)$ and $g_s(y)$. These functions are readily expressed in terms of the functions $u(x,t)$ and $v(x,t)$.

Let us express the four height gain functions in the form

$$f_s(y) = 1 + A_1(t_s^o,-y) + A_2(t_s^o,-y) + A_3(t_s^o,-y) + \cdots \quad \text{(E-53)}$$

$$f_s'(y) = \quad B_1(t_s^o,-y) + B_2(t_s^o,-y) + B_3(t_s^o,-y) + \cdots \quad \text{(E-54)}$$

$$g_s(y) = y + C_1(t_s^\infty,-y) + C_2(t_s^\infty,-y) + C_3(t_s^\infty,-y) + \cdots \quad \text{(E-55)}$$

$$g_s'(y) = 1 + D_1(t_s^\infty,-y) + D_2(t_s^\infty,-y) + D_3(t_s^\infty,-y) + \cdots \quad \text{(E-56)}$$

where the $A_n(t,-y)$, $B_n(t,-y)$, $C_n(t,-y)$ and $D_n(t,-y)$ satisfy the recursion relation

$$n(n-1)K_n(t,-y) = y^2[-yK_{n-3}(t,-y) + tK_{n-2}(t,-y)] \quad \text{(E-57)}$$

with the initial terms

$$A_1 = 0 \quad , A_2 = \frac{ty^2}{2} \quad , A_3 = -\frac{y^3}{6} \quad , A_4 = \frac{t^2y^4}{24}$$

$$B_1 = ty \quad , B_2 = -\frac{y^2}{2} \quad , B_3 = \frac{t^2y^3}{6} \quad , B_4 = -\frac{ty^4}{6}$$

$$C_1 = y \quad , C_2 = 0 \quad , C_3 = \frac{ty^3}{6} \quad , C_4 = -\frac{y^4}{12}$$

$$D_1 = 0 \quad , D_2 = \frac{ty^2}{2} \quad , D_3 = -\frac{y^3}{3} \quad , D_4 = \frac{t^2y^4}{24}$$

Program E-5 contains SUBRØUTINE HGFY(N,Y,FYR,FYI) which will compute

$$f_I(Y) = FYR(I) + iFYI(I) \quad , I = 1,2,3,\ldots,N \quad \text{(E-58)}$$

The SUBRØUTINE HGFPY(N,Y,FPYR,FPYI) in Program E-6 will generate values for

$$f_I'(Y) = -[FPYR(I) + iFPYI(I)] \quad , I = 1,2,\ldots,N \quad \text{(E-59)}$$

Each of these program require the roots β_s of the Airy function

which are tabulated in Table 2-1. These values are transferred to the subroutine by means of a labeled CØMMØN/DABETA/BETA(50), AIRYF(50). Output from these programs is presented in Tables E-4 and E-5. The results are consistent with the tables prepared by Azriliant and Belkina (Ref. 1). A sample of the tables of Ref. 1 is presented in Tables E-2 and E-3.

Table E-2

SAMPLES OF TABLES OF $f_s(y)$ and $f_s'(y)$ PUBLISHED BY SOVIET AUTHORS

y	Re $f_s(y)$	Im $f_s(y)$	Re $f_s(y)$	Im $f_s(y)$	Re $f_s'(y)$	Im $f_s'(y)$	Re $f_s'(y)$	Im $f_s'(y)$
0,0	1	0	1	0	0	0	0	0
0,1	1,0024	0,0044	1,0079	0,0141	0,0458	0,0894	0,1565	0,2828
0,2	1,0098	0,0177	1,0308	0,0568	0,0811	0,1774	0,2973	0,5740
0,3	1,0182	0,0399	1,0666	0,1294	0,1148	0,2675	0,4157	0,8811
0,4	1,0294	0,0712	1,1129	0,2338	0,1163	0,3587	0,5041	1,2102
0,5	1,0410	0,1117	1,1661	0,3724	0,1144	0,4585	0,5534	1,5658
0,6	1,0518	0,1613	1,2219	0,5480	0,0982	0,5422	0,5530	1,9506
0,7	1,0612	0,2201	1,2747	0,7636	0,0666	0,6327	0,4911	2,3653
0,8	1,0646	0,2878	1,3177	1,0220	0,0187	0,7203	0,3514	2,8078
0,9	1,0633	0,3640	1,3426	1,3259	−0,0463	0,8031	0,1283	3,2728
1,0	1,0547	0,4482	1,3398	1,6770	−0,1287	0,8787	−0,2024	3,7515
1,1	1,0370	0,5394	1,2981	2,0762	−0,2288	0,9443	−0,6532	4,2308
1,2	1,0084	0,6366	1,2017	2,5226	−0,3412	0,9969	−1,2394	4,6926
1,3	0,9672	0,7382	1,0453	3,0134	−0,4810	1,0331	−1,9746	5,1138
1,4	0,9119	0,8426	0,8044	3,5431	−0,6248	1,0496	−2,8703	5,4653
1,5	0,8410	0,9474	0,4656	4,1030	−0,7903	1,0428	−3,9337	5,7124
1,6	0,7535	1,0502	0,0120	4,6807	−0,9615	1,0092	−5,1667	5,8144
1,7	0,6485	1,1482	− 0,5732	5,2595	−1,1385	0,9456	−6,5639	5,7253
1,8	0,5257	1,2382	− 1,3058	5,8177	−1,3168	0,8494	−8,1105	5,3943
1,9	0,3853	1,3179	− 2,1995	6,3285	−1,4907	0,7182	−9,7803	4,7674
2,0	0,2279	1,3907	− 3,2647	6,7595	−1,6541	0,5508	−11,534	3,789
2,1	0,0551	1,4259	− 4,5073	7,0728	−1,7999	0,3469	−13,317	2,405
2,2	−0,1312	1,4489	− 5,9267	7,2252	−1,9206	0,1075	−15,057	0,564
2,3	−0,3280	1,4463	− 7,5144	7,1190	−2,0086	−0,1651	−16,666	−1,774
2,4	−0,5416	1,4149	− 9,2520	6,8528	−2,0558	−0,4667	−18,036	−4,640
2,5	−0,7375	1,3521	−11,109	6,223	−2,0548	−0,7916	−19,041	−8,043
2,6	−0,9407	1,2560	−13,043	5,227	−1,9986	−1,1323	−19,541	−11,971
2,7	−1,1353	1,1255	−14,996	3,813	−1,8813	−1,4793	−19,383	−16,382
2,8	−1,3148	0,9603	−16,892	1,937	−1,6986	−1,8217	−18,405	−21,196
2,9	−1,4727	0,7617	−18,644	− 0,436	−1,4480	−2,1469	−16,446	−26,294
3,0	−1,6021	0,5320	−20,144	− 3,326	−1,1293	−2,4410	−13,353	−31,510
3,1	−1,6963	0,2750	−21,272	− 6,735	−0,7452	−2,6896	−8,990	−36,631
3,2	−1,7491	−0,0039	−21,896	−10,640	−0,3014	−2,8779	−3,256	−41,393
3,3	−1,7549	−0,2981	−21,876	−14,991	0,1931	−2,9915	3,908	−45,485
3,4	−1,7092	−0,5993	−21,067	−19,703	0,7259	−3,0170	12,498	−48,751
3,5	−1,6080	−0,8982	−19,331	−24,654	1,2811	−2,9432	22,435	−50,205
3,6	−1,4529	−1,1843	−16,541	−29,783	1,8499	−2,7614	33,548	−50,040
3,7	−1,2416	−1,4467	−12,591	−34,598	2,3805	−2,4665	45,561	−47,646
3,8	−0,9782	−1,6738	− 7,411	−38,126	2,8793	−2,0579	58,084	−42,940
3,9	−0,6680	−1,8546	− 0,974	−43,017	3,3112	−1,5399	70,607	−34,689
4,0	−0,3190	−1,9785	6,689	−45,966	3,6512	−0,9224	82,198	−23,542

The subroutines in Programs E-7 and E-8 evaluate the functions

$$g_I(Y) = - [GYR(I) + iGYI(I)] \quad , I = 1,2,3,\ldots,N \qquad (E\text{-}60)$$

$$g'_I(Y) = GPYR(I) + iGPYI(I) \quad , I = 1,2,3,\ldots,N \qquad (E\text{-}61)$$

These programs have been compared with the Soviet tables by means of the sample output given in Tables E-6 and E-7.

Table E-3

SAMPLES OF TABLES OF $g_s(y)$ AND $g'_s(y)$ PUBLISHED BY SOVIET AUTHORS

y	Re $g_s(y)$	Im $g_s(y)$	Re $g_s(y)$	Im $g_s(y)$	Re $g_s(y)$	Re $g_s'(y)$	Im $g_s'(y)$	Re $g_s'(y)$	Im $g_s'(y)$
0,0	0	0	0	0	0	1	0	1	0
0,1	0,10019	0,00084	0,10033	0,00069	0,1004	1,00550	0,01014	1,00995	0,01776
0,2	0,20141	0,00271	0,20257	0,00476	0,2035	1,02451	0,04078	1,03762	0,07171
0,3	0,30452	0,00920	0,30834	0,01620	0,3114	1,04254	0,09247	1,07985	0,16376
0,4	0,41006	0,02193	0,41887	0,03887	0,4259	1,05867	0,16599	1,13193	0,29680
0,5	0,51828	0,04315	0,53486	0,07705	0,5477	1,09452	0,26219	1,18777	0,47443
0,6	0,62906	0,07516	0,65630	0,13538	0,6768	1,11915	0,38159	1,23965	0,70666
0,7	0,74186	0,12033	0,78234	0,21893	0,8114	1,13508	0,52553	1,27788	0,97951
0,8	0,85565	0,18109	0,91103	0,33315	0,9479	1,13822	0,69357	1,29057	1,31448
0,9	0,96889	0,25964	1,03913	0,48378	1,0804	1,12242	0,88523	1,26847	1,70902
1,0	1,07942	0,35888	1,16184	0,67674	1,1999	1,08299	1,09933	1,17974	2,16078
1,1	1,18444	0,48037	1,27254	0,91785	1,2939	1,01181	1,33353	1,02101	2,67080
1,2	1,28051	0,62614	1,36258	1,21262	1,3457	0,90248	1,58420	0,7 214	3,23257
1,3	1,36344	0,79760	1,42098	1,56576	1,3337	0,74800	1,84025	0,38212	3,83602
1,4	1,42837	0,99555	1,43430	1,98070	1,2310	0,54 56	2,11290	— 0,14343	4,46575
1,5	1,46990	1,22005	1,39651	2,45895	1,0046	0,27688	2,37552	— 0,84428	5,09787
1,6	1,48163	1,47017	1,29901	2,99837	0,6160	—0,05140	2,62352	— 1,74187	5,70293
1,7	1,46727	1,74365	1,03074	3,59731	0,0212	—0,44724	2,84430	— 2,86233	6,24682
1,8	1,38986	2,03764	0,67852	4,24356	— 0,4324	—0,91206	3,02331	— 4,22329	6,66901
1,9	1,27243	2,34658	0,17766	4,92384	— 1,0956	—1,44717	3,14452	— 5,83135	6,90864
2,0	1,09823	2,66398	—0,49722	5,61677	— 3,3255	—2,04720	3,18946	— 7,70296	6,90453
2,1	0,86104	2,98132	— 1,3710	6,2939	— 5,476	—2,70541	3,13997	— 9,8110	6,5759
2,2	0,55559	3,28818	— 2,4665	6,9181	— 7,896	—3,41024	2,97070	—12,1199	5,8628
2,3	0,17798	3,57224	— 3,8026	7,4435	—10,523	—4,14530	2,68092	—14,6115	4,5818
2,4	—0,27377	3,81937	— 5,3917	7,8144	—14,277	—4,88910	2,28537	—17,1765	2,7277
2,5	—0,79924	4,01383	— 7,2372	7,9858	—18,250	—5,61480	1,62545	—19,7202	0,1783
2,6	—1,39506	4,13881	— 9,3306	7,8241	—22,699	—6,29628	0,84052	—22,1052	— 3,1481
2,7	—2,05440	4,17501	—11,6973	7,3082	—27,531	—6,87844	— 0,12462	—24,1996	— 7,3157
2,8	—2,76647	4,10773	—14,1445	6,3317	—32,965	—7,33797	— 1,26817	—25,6772	—12,36 4
2,9	—3,51622	3,91686	—16,7869	4,8069	—37,965	—7,62433	— 2,5799	—26,4193	—18,2797
3,0	—4,28403	3,58680	—19,3937	2,6493	—42,435	—7,6913	— 4,0395	—26,1188	—25,0208
3,1	—5,04564	3,10484	—21,8363	— 0,2209	—46,499	—7,4931	— 5,6158	—24,4880	—32,4675
3,2	—5,77236	2,46117	—24,2370	— 3,8825	—49,251	—6,9865	— 7,2654	—21,2304	—40,4280
3,3	— 6,43130	1,65115	—26,1184	— 8,3147	—50,251	—6,1338	— 8,9325	—16,0658	—48,6381
3,4	—6,99861	0,67844	—27,3754	—13,6837	—48,879	—4,9863	—10,5490	— 8,7010	—56,6884
3,5	—7,39867	—0,45433	—27,7787	—19,6334	—44,045	—3,2872	—12,0764	1,0460	—64,1537
3,6	—7,63092	—1,72346	—27,082	—26,375	—35,62	—1,2753	—13,3087	13,3176	—70,4387
3,7	—7,64210	—3,10436	—25,030	—33,880	—20,92	1,1114	—14,2540	28,1343	—74,8880
3,8	—7,39743	—4,58065	—21,374	—41,267	— 0,05	3,8330	—14,7930	45,3716	—76,7861
3,9	—6,86644	—6,04592	—15,885	—48,099	25,88	6,8251	—14,8192	64,7857	—75,2879
4,0	—6,02837	—7,50449	— 8,375	—56,179	59,85	9,9973	—14,9432	88,982	—69,586

The roots α_s which are required in Programs E-7 and E-8 have been transmitted to the program by means of a labeled block of COMMON which is obtained by means of the non-executable statement which reads CØMMØN/DALPHA/ALPHA(50),AIRYD(50).

For $q \neq 0$ and $q \neq \infty$, one may find it convenient to use Program E-9 to compute the Airy functions. This program is similar to Program B-1; the difference ebing that this program returns the complex numbers which represent the Airy functions $v(z)$ and $w_1(z)$ which are employed in the papers by Fock and other Soviet writers. The program is called SUBRØUTINE VTWTZ(Z,VT,WT,VTP,WTP) and the relation of the FORTRAN variables to the mathematical quantities are

$$z = Z \ , \ v(z) = VT \ , \ w_1(z) = WT \ , \ v'(z) = VTP \ , \ w_1'(z) = WTP$$

The program should only be used when the modulus of z is less than 6.1.

The author has done extensive work on the case of obtaining the height gain functions when the roots have a large modulus. However, this study is incomplete and no attempt will be made to discuss the methods and the results which have been obtained. This work will be carried to completion and reported in a later publication. The author would like to make one observation regarding this work. In the book by Azriliant and Belkina (Ref. 1) use is made of an approximation which is due to Fock, namely

$$f_s(y,q) \approx \cosh(y\sqrt{t_s}) - \frac{q}{\sqrt{t_s}} \sinh(y\sqrt{t_s}) \ , \quad \left| \frac{y^2}{2\sqrt{t_s}} \right| << 1 \quad \text{(E-62)}$$

When one attempts to find an asymptotic series which has these terms as the leading terms, the second term (which vanishes when $q = 0$) is apparently cancelled out by the higher order terms. Therefore, until further work has been done and comparisons made with other methods of approximation, the second term in Eq. (E-62) is of questionable value.

```
      SUBROUTINE HGFY(N,Y,FYR,FYI)
      COMMON/DABETA/BETA(50),AIRYF(50)
      DOUBLE PRECISION BETA,AIRYF,TR,TI,A1R,A1I,A2R,A2I,A3R,A3I,AR,AI,
     1        DFLOAT,SR,SI,X,X2,STR,STI,A4R,A4I
      DIMENSION FYR(N),FYI(N)
  199 FORMAT(11H1  FOR Y = ,E16.8,27H  PROGRAM EXIT BECAUSE N = ,I4,
     A        51H  EXCEEDS 50 WHICH IS THE CAPACITY OF HGFY(N,Y,...))
  198 FORMAT(11H1  FOR Y = ,E16.8,62H  PROGRAM EXIT BECAUSE OF LOSS OF A
     ACCURACY DUE TO CANCELLATION//10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     B   10X,54HHGFY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
  197 FORMAT(11H1  FOR Y = ,E16.8,88H  PROGRAM EXIT BECAUSE COMPLETION O
     AF 50 ITERATIONS FAILED TO YIELD CONVERGENCE OF SERIES  //
     B                                    10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     C   10X,54HHGFY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
      DFLOAT(J) = DBLE(FLOAT(J))
      IF(N.GT.50) GO TO 99
C-----PROGRAM WILL CAUSE CALL EXIT WHEN N EXCEEDS 50
      X       = DBLE(Y)
      X2      = X*X
      DO 10 K=1,N
      KDEX    = K
      TR      = 0.5D0*BETA(K)
      TI      = 1.73205080756887D0*TR
      A2R     = TR*(X**2)/2.0D0
      A2I     = 1.73205080756887D0*A2R
      A3R     =-(X**3)/6.0D0
      A3I     = 0.0D0
      A4R     = (A2R*A2R - A2I*A2I)/6.0D0
      A4I     = (A2R*A2I)/3.0D0
      SR      = 1.0D0 + A2R + A3R + A4R
      SI      =         A2I + A3I + A4I
      STR     = 1.0D0 + DABS(A2R) + DABS(A3R) + DABS(A4R)
      STI     =         DABS(A2I) + DABS(A3I) + DABS(A4I)
C-----STR AND STI ARE TEST QUANTITIES TO TEST FOR EXCESSIVE CANCELLATION
```

```
      DO 20 L=5,50
      LDEX   = L
      AR     = A2*(-X*A2R  + TR*A3R - TI*A3I)/DFLOAT(L*(L-1))
      AI     = A2*(-X*A2I  + TR*A3I + TI*A3R)/DFLOAT(L*(L-1))
      SR     = SR + AR
      SI     = SI + AI
      STR    = STR + DABS(AR)
      STI    = STI + DABS(AI)
      IF((STR/DABS(SR) + STI/DABS(SI)).GT.1.0D12)  GO TO 98
C-----TRANSFER OUT IF CANCELLATION SUGGESTS EXCESSIVE LOSS OF ACCURACY
      IF(DABS(AR + A4R + A3R).LT.1.0D-8) GO TO 89
C-----TRANSFER OUT IF SUM OF LAST THREE TERMS OF REAL PART IS LESS THAN
C-----1.0D-8 WHERE AS INITIAL TERM WAS UNITY.
      A2R    = A3R
      A2I    = A3I
      A3R    = A4R
      A3I    = A4I
      A4R    = AR
      A4I    = AI
   20 CONTINUE
      GO TO 97
C-----WRITE A MESSAGE THAT PROGRAM FAILED TO LEAVE THE LOOP AFTER FIFTY
C-----ITERATIONS AND THEN CALL EXIT
   89 FYR(K) = SNGL(SR)
      FYI(K) = SNGL(SI)
   10 CONTINUE
C-----THIS IS THE NORMAL RETURN WHEN PROGRAM SUCCESSFULLY ESTABLISHES AN
C-----ARRAY FYR(N) AND FYI(N) WHICH CONTAINS THE N HEIGHT GAIN FUNCTIONS
      RETURN
   99 WRITE(6,199) Y,I
      CALL EXIT
   98 WRITE(6,198) Y,KDEX,LDEX
      CALL EXIT
   97 WRITE(6,197) Y,KDEX,LDEX
      CALL EXIT
      END
```

```
      SUBROUTINE HGFPY(N,Y,FPYR,FPYI)
      DIMENSION FPYR(N),FPYI(N)
      COMMON/DABETA/BETA(50),AIRYF(50)
      DOUBLE PRECISION BETA,AIRYF,DFLOAT,TR,TI,X,X2,A1R,A1I,A2R,A2I,A3R,
     A       A3I,SR,SI,STR,STI,TEST,AR,AI
  199 FORMAT(11H1  FOR Y = ,E16.8,27H  PROGRAM EXIT BECAUSE N = ,I4,
     A       52H  EXCEEDS 50 WHICH IS THE CAPACITY OF HGFPY(N,Y,...))
  198 FORMAT(11H1  FOR Y = ,E16.8,62H  PROGRAM EXIT BECAUSE OF LOSS OF A
     ACCURACY DUE TO CANCELLATION//10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     B   10X,55HHGFPY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
  197 FORMAT(11H1  FOR Y = ,E16.8,88H  PROGRAM EXIT BECAUSE COMPLETION O
     AF 50 ITERATIONS FAILED TO YIELD CONVERGENCE OF SERIES//
     B                                   10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     C   10X,55HHGFPY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
      DFLOAT(J) = DBLE(FLOAT(J))
      IF(N.GT.50) GO TO 99
C-----PROGRAM WILL CAUSE EXIT WHEN N EXCEEDS 50
      X       = DBLE(Y)
      X2      = X*X
      DO 10 K=1,N
      KDEX    = K
      TEST    = X*BETA(K)*1.0D-8
      TR      = 0.5D0*BETA(K)
      TI      = 1.732050807568877D0*TR
      A1R     =-TR*X
      A1I     =-TI*X
      A2R     = 0.5D0*(X*X)
      A2I     = 0.0D0
      A3R     = (-X/6.0D0)*(A1R*A1R - A1I*A1I)
      A3I     = (-X/3.0D0)*(A1R*A1I)
      SR      = A1R + A2R + A3R
      SI      = A1I + A2I + A3I
      STR     = DABS(A1R) + DABS(A2R) + DABS(A3R)
      STI     = DABS(A1I) + DABS(A2I) + DABS(A3I)
C-----STR AND STI ARE TEST QUANTITIES TO TEST FOR EXCESSIVE CANCELLATION
```

```
      DO 20 L=4,50
      LDEX    = L
      AR      =-X2*(X*DFLOAT(L-1)*A1R - DFLOAT(L-2)*(TR*A2R - TI*A2I))/
     A          DFLOAT(L*(L-1)*(L-2))
      AI      =-X2*(X*DFLOAT(L-1)*A1I - DFLOAT(L-2)*(TR*A2I + TI*A2R))/
     A          DFLOAT(L*(L-1)*(L-2))
      SR      = SR + AR
      SI      = SI + AI
      STR     = STR + DABS(AR)
      STI     = STI + DABS(AI)
      IF((STR/DABS(SR) + STI/DABS(SI)).GT.1.0D12) GO TO 98
C-----TRANSFER OUT IF CANCELLATION SUGGESTS EXCESSIVE LOSS OF ACCURACY
      IF(DABS(AR + A3R + A2R).LT.TEST) GO TO 89
C-----TRANSFER OUT IF SUM OF LAST THREE TERMS OF REAL PART IS LESS THAN
C-----TEST = 1.0D-8*(MAGNITUDE OF FIRST TERM)
      A1R     = A2R
      A1I     = A2I
      A2R     = A3R
      A2I     = A3I
      A3R     = AR
      A3I     = AI
   20 CONTINUE
      GO TO 97
C-----WRITE A MESSAGE THAT PROGRAM FAILED TO LEAVE THE LOOP AFTER FIFTY
C-----ITERATIONS AND THEN CALL EXIT
   89 FPYR(K)= SNGL(SR)
      FPYI(K)= SNGL(SI)
   10 CONTINUE
      RETURN
C-----THIS IS THE NORMAL RETURN WHEN PROGRAM SUCCESSFULLY ESTABLISHES AN
C-----ARRAY FPYR(N),FPYI(N) WHICH CONTAINS THE N HEIGHT GAIN FUNCTIONS
   99 WRITE(6,199) Y,N
      CALL EXIT
   98 WRITE(6,198) Y,KDEX,LDEX
      CALL EXIT
   97 WRITE(6,197) Y,KDEX,LDEX
      CALL EXIT
      END
```

```
      SUBROUTINE HGGY(N,Y,GYR,GYI)
      DIMENSION GYR(N),GYI(N)
      COMMON/DALPHA/ALPHA(50),AIRYD(50)
      DOUBLE PRECISION ALPHA,AIRYD,DFLOAT,TR,TI,X,X2,A2R,A2I,A3R,A3I,A4R
     A                 ,A4I,SR,SI,STR,STI,TEST,AR,AI
  199 FORMAT(11H1  FOR Y = ,E16.8,27H  PROGRAM EXIT BECAUSE N = ,I4,
     A       51H  EXCEEDS 50 WHICH IS THE CAPACITY OF HGGY(N,Y,...))
  198 FORMAT(11H1  FOR Y = ,E16.8,62H  PROGRAM EXIT BECAUSE OF LOSS OF A
     ACCURACY DUE TO CANCELLATION//10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     B   10X,54HHGGY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
  197 FORMAT(11H1  FOR Y = ,E16.8,88H  PROGRAM EXIT BECAUSE COMPLETION O
     AF 50 ITERATIONS FAILED TO YIELD CONVERGENCE OF SERIES//
     B                                 10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     C   10X,54HHGGY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
      DFLOAT(J) = DBLE(FLOAT(J))
      IF(N.GT.50) GO TO 99
C-----PROGRAM WILL CAUSE EXIT WHEN N EXCEEDS 50
      X       = DBLE(Y)
      TEST    = (1.0D-9)*DABS(X)
      DO 10 K=1,N
      KDEX    = K
      TR      = 0.5*ALPHA(K)
      TI      = 1.732050807568877D0*TR
      X2      = X*X
      A2R     = 0.0D0
      A2I     = 0.0D0
      A3R     =-X*X2*TR/6.0D0
      A3I     = 1.732050807568877D0*A3R
      A4R     = X2*X2/12.0D0
      A4I     = 0.0D0
      SR      =-X + A2R + A3R + A4R
      SI      =     A2I + A3I + A4I
      STR     = DABS(X) + DABS(A2R) + DABS(A3R) + DABS(A4R)
      STI     =           DABS(A2I) + DABS(A3I) + DABS(A4I)
C-----STR AND STI ARE TEST QUANTITIES TO TEST FOR EXCESSIVE CANCELLATION
```

```
      DO 20 L=5,50
      LDEX   = L
      AR     = X2*(-X*A2R + TR*A3R - TI*A3I)/DFLOAT(L*(L-1))
      AI     = X2*(-X*A2I + TR*A3I + TI*A3R)/DFLOAT(L*(L-1))
      SR     = SR + AR
      SI     = SI + AI
      STR    = STR + DABS(AR)
      STI    = STI + DABS(AI)
      IF((STR/DABS(SR) + STI/DABS(SI)).GT.1.0D12) GO TO 98
C-----TRANSFER OUT IF CANCELLATION SUGGESTS EXCESSIVE LOSS OF ACCURACY
      IF(DABS(AR + A4R + A3R).LT.TEST) GO TO 89
C-----TRANSFER OUT IF SUM OF LAST THREE TERMS OF REAL PART IS LESS THAN
C-----TEST = 1.0D-8*(MAGNITUDE OF FIRST TERM)
      A2R    = A3R
      A2I    = A3I
      A3R    = A4R
      A3I    = A4I
      A4R    = AR
      A4I    = AI
   20 CONTINUE
      GO TO 97
C-----WRITE A MESSAGE THAT PROGRAM FAILED TO LEAVE THE LOOP AFTER FIFTY
C-----ITERATIONS AND THEN CALL EXIT
   89 GYR(K) = SNGL(SR)
      GYI(K) = SNGL(SI)
   10 CONTINUE
C-----THIS IS THE NORMAL RETURN WHEN PROGRAM SUCCESSFULLY ESTABLISHES AN
C-----ARRAY GYR(N) AND GYI(N) WHICH CONTAINS THE N HEIGHT GAIN FUNCTIONS
      RETURN
   99 WRITE(6,199) Y,N
      CALL EXIT
   98 WRITE(6,198) Y,KDEX,LDEX
      CALL EXIT
   97 WRITE(6,197) Y,KDEX,LDEX
      CALL EXIT
      END
```

```
      SUBROUTINE HGGPY(N,Y,GPYR,GPYI)
      DIMENSION GPYR(N),GPYI(N)
      COMMON/DALPHA/ALPHA(50),AIRYD(50)
      DOUBLE PRECISION ALPHA,AIRYD,DFLOAT,TR,TI,X,X2,A1R,A1I,A2R,A2I,
     A       A3R,A3I,SR,SI,STR,STI,TEST,AR,AI
  199 FORMAT(11H1  FOR Y = ,E16.8,27H  PROGRAM EXIT BECAUSE N = ,I4,
     A       52H  EXCEEDS 50 WHICH IS THE CAPACITY OF HGGPY(N,Y,...))
  198 FORMAT(11H1  FOR Y = ,E16.8,62H  PROGRAM EXIT BECAUSE OF LOSS OF A
     ACCURACY DUE TO CANCELLATION//10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     B   10X,55HHGGPY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
  197 FORMAT(11H1  FOR Y = ,E16.8,88H  PROGRAM EXIT BECAUSE COMPLETION O
     AF 50 ITERATIONS FAILED TO YIELD CONVERGENCE OF SERIES//
     B                                10X,7HKDEX = ,I4,9H  LDEX = ,I4//
     C   10X,55HHGGPY(N,Y,...) IS NOT CAPABLE OF HANDLING THIS ARGUMENT)
      DFLOAT(J) = DBLE(FLOAT(J))
      TEST   = 1.0D-8
      IF(N.GT.50) GO TO 99
C-----PROGRAM WILL CAUSE EXIT WHEN N EXCEEDS 50
      X      = DBLE(Y)
      X2     = X*X
      DO 10 K=1,N
      KDEX   = K
      TR     = 0.5D0*ALPHA(K)
      TI     = 1.732050807568877D0*TR
      A1R    = 0.0D0
      A1I    = 0.0D0
      A2R    = 0.5*X2*TR
      A2I    = 1.732050807568877D0*A2R
      A3R    =-X*X2/3.0D0
      A3I    = 0.0D0
      SR     = 1.0D0 + A1R + A2R + A3R
      SI     =         A1I + A2I + A3I
      STR    = 1.0D0 + DABS(A1R) + DABS(A2R) + DABS(A3R)
      STI    =         DABS(A1I) + DABS(A2I) + DABS(A3I)
C-----STR AND STI ARE TEST QUANTITIES TO TEST FOR EXCESSIVE CANCELLATION
```


LOCKHEED MISSILES & SPACE COMPANY

```
      DO 20 L=4,50
      LDEX   = L
      AR     = X2*(-X*DFLOAT(L-1)*A1R + DFLOAT(L-2)*(TR*A2R - TI*A2I))/
     A           DFLOAT(L*(L-1)*(L-2))
      AI     = X2*(-X*DFLOAT(L-1)*A1I + DFLOAT(L-2)*(TR*A2I + TI*A2R))/
     A           DFLOAT(L*(L-1)*(L-2))
      SR     = SR + AR
      SI     = SI + AI
      STR    = STR + DABS(AR)
      STI    = STI + DABS(AI)
      IF((STR/DABS(SR) + STI/DABS(SI)).GT.1.0D12) GO TO 98
C-----TRANSFER OUT IF CANCELLATION SUGGESTS LOSS OF ACCURACY
      IF(DABS(AR + A3R + A2R).LT.TEST) GO TO 89
C-----TRANSFER OUT IF SUM OF LAST THREE TERMS OF REAL PART IS LESS THAN
C-----TEST = 1.0D-8*(MAGNITUDE OF FIRST TERM)
      A1R    = A2R
      A1I    = A2I
      A2R    = A3R
      A2I    = A3I
      A3R    = AR
      A3I    = AI
   20 CONTINUE
      GO TO 97
C-----WRITE A MESSAGE THAT PROGRAM FAILED TO LEAVE THE LOOP AFTER FIFTY
C-----ITERATIONS AND THEN CALL EXIT
   89 GPYR(K)= SNGL(SR)
      GPYI(K)= SNGL(SI)
   10 CONTINUE
C-----THIS IS THE NORMAL RETURN WHEN PROGRAM SUCCESSFULLY ESTABLISHES AN
C-----ARRAY GPYR(N) AND GPYI(N) WHICH CONTAINS N HEIGHT GAIN FUNCTIONS
      RETURN
   99 WRITE(6,199) Y,N
      CALL EXIT
   98 WRITE(6,198) Y,KDEX,LDEX
      CALL EXIT
   97 WRITE(6,197) Y,KDEX,LDEX
      CALL EXIT
      END
```

Table E-4

THE HEIGHT GAIN FUNCTION $f_s(y)$

$$f_s(y) = \frac{w_1(t_s^o - y)}{w_1(t_s^o)} = \frac{Ai[-\beta_s + y\exp(-i\frac{2}{3}\pi)]}{Ai(-\beta_s)} = FYR(J) + iFYI(J)$$

$$f_s(0) = 1 \ , \ s = J = 1,2,3,\ldots$$

```
TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.EQ.5 AND Y = 0.2
J =  1  FYR(J) =     .10088148+01   FYI(J) =     .17696507-01
J =  2  FYR(J) =     .10307771+01   FYI(J) =     .56839366-01
J =  3  FYR(J) =     .10460582+01   FYI(J) =     .84782937-01
J =  4  FYR(J) =     .10589806+01   FYI(J) =     .10888700+00
J =  5  FYR(J) =     .10705028+01   FYI(J) =     .13075837-00

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.EQ.5 AND Y = 0.4
J =  1  FYR(J) =     .10293761+01   FYI(J) =     .71239454-01
J =  2  FYR(J) =     .11129232+01   FYI(J) =     .23380095-00
J =  3  FYR(J) =     .11683412+01   FYI(J) =     .35391892-00
J =  4  FYR(J) =     .12132932+01   FYI(J) =     .46016063-00
J =  5  FYR(J) =     .12517816+01   FYI(J) =     .55860890-00

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.EQ.5 AND Y = 0.6
J =  1  FYR(J) =     .10517905+01   FYI(J) =     .16134623-00
J =  2  FYR(J) =     .12219528+01   FYI(J) =     .54793072-00
J =  3  FYR(J) =     .13223805+01   FYI(J) =     .84805959-00
J =  4  FYR(J) =     .13940288+01   FYI(J) =     .11224667+01
J =  5  FYR(J) =     .14464756+01   FYI(J) =     .13834634+01

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.EQ.5 AND Y = 0.8
J =  1  FYR(J) =     .10645694+01   FYI(J) =     .28780445-00
J =  2  FYR(J) =     .13177004+01   FYI(J) =     .10219096+01
J =  3  FYR(J) =     .14261585+01   FYI(J) =     .16246052+01
J =  4  FYR(J) =     .14651560+01   FYI(J) =     .21942076+01
J =  5  FYR(J) =     .14528968+01   FYI(J) =     .27487827+01

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.EQ.5 AND Y = 1.0
J =  1  FYR(J) =     .10547342+01   FYI(J) =     .44819684-00
J =  2  FYR(J) =     .13399089+01   FYI(J) =     .16769136+01
J =  3  FYR(J) =     .13447475+01   FYI(J) =     .27419598+01
J =  4  FYR(J) =     .11899459+01   FYI(J) =     .37751160+01
J =  5  FYR(J) =     .90555649-00   FYI(J) =     .47952715+01
```

Table E-4 (Cont'd)

```
TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 1.2
J =  1  FYR(J) =    .10083951+01  FYI(J) =    .63663021-00
J =  2  FYR(J) =    .12048182+01  FYI(J) =    .25224828+01
J =  3  FYR(J) =    .87463003-00  FYI(J) =    .42331848+01
J =  4  FYR(J) =    .19362763-00  FYI(J) =    .59131758+01
J =  5  FYR(J) =   -.79359463-00  FYI(J) =    .75674323+01

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 1.4
J =  1  FYR(J) =    .91188672-00  FYI(J) =    .84261201-00
J =  2  FYR(J) =    .80464242-00  FYI(J) =    .35429361+01
J =  3  FYR(J) =   -.26646827-00  FYI(J) =    .60652344+01
J =  4  FYR(J) =   -.20648704+01  FYI(J) =    .85151403+01
J =  5  FYR(J) =   -.45288368+01  FYI(J) =    .10846764+02

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 1.6
J =  1  FYR(J) =    .75350671-00  FYI(J) =    .10502922+01
J =  2  FYR(J) =    .12381600-01  FYI(J) =    .46806542+01
J =  3  FYR(J) =   -.24350606+01  FYI(J) =    .80816305+01
J =  4  FYR(J) =   -.62691234+01  FYI(J) =    .11211196+02
J =  5  FYR(J) =   -.11461491+02  FYI(J) =    .13887667+02

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 1.8
J =  1  FYR(J) =    .52575394-00  FYI(J) =    .12383454+01
J =  2  FYR(J) =   -.13052221+01  FYI(J) =    .58178070+01
J =  3  FYR(J) =   -.60298929+01  FYI(J) =    .99303788+01
J =  4  FYR(J) =   -.13273552+02  FYI(J) =    .13173055+02
J =  5  FYR(J) =   -.22877772+02  FYI(J) =    .15048999+02

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 2.0
J =  1  FYR(J) =    .22794384-00  FYI(J) =    .13807892+01
J =  2  FYR(J) =   -.32639299+01  FYI(J) =    .67598190+01
J =  3  FYR(J) =   -.11416890+02  FYI(J) =    .10985952+02
J =  4  FYR(J) =   -.23715350+02  FYI(J) =    .12903678+02
J =  5  FYR(J) =   -.39777827+02  FYI(J) =    .11356582+02

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 2.2
J =  1  FYR(J) =   -.13122203-00  FYI(J) =    .14489918+01
J =  2  FYR(J) =   -.59257936+01  FYI(J) =    .72258984+01
J =  3  FYR(J) =   -.18795560+02  FYI(J) =    .10284675+02
J =  4  FYR(J) =   -.37854578+02  FYI(J) =    .80492896+01
J =  5  FYR(J) =   -.62064302+02  FYI(J) =   -.18881157+01
```

Table E-4 (Cont'd)

```
TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 2.4
J =  1  FYR(J) =    -.53160498-00  FYI(J) =     .14150044+01
J =  2  FYR(J) =    -.92510889+01  FYI(J) =     .68539393+01
J =  3  FYR(J) =    -.27996995+02  FYI(J) =     .65062997+01
J =  4  FYR(J) =    -.54895929+02  FYI(J) =    -.46683320+01
J =  5  FYR(J) =    -.87200189+02  FYI(J) =    -.31212131+02

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 2.6
J =  1  FYR(J) =    -.94075157-00  FYI(J) =     .12561362+01
J =  2  FYR(J) =    -.13042705+02  FYI(J) =     .52283652+01
J =  3  FYR(J) =    -.38217935+02  FYI(J) =    -.19512682+01
J =  4  FYR(J) =    -.72204016+02  FYI(J) =    -.29243561+02
J =  5  FYR(J) =    -.10829514+03  FYI(J) =    -.84380431+02

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 2.8
J =  1  FYR(J) =    -.13146679+01  FYI(J) =     .96038737-00
J =  2  FYR(J) =    -.16892447+02  FYI(J) =     .19391098+01
J =  3  FYR(J) =    -.47721217+02  FYI(J) =    -.16746924+02
J =  4  FYR(J) =    -.94347471+02  FYI(J) =    -.69692163+02
J =  5  FYR(J) =    -.11160126+03  FYI(J) =    -.16852822+03

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 3.0
J =  1  FYR(J) =    -.16022247+01  FYI(J) =     .53199985-00
J =  2  FYR(J) =    -.20144670+02  FYI(J) =    -.33231721+01
J =  3  FYR(J) =    -.53571033+02  FYI(J) =    -.39164963+02
J =  4  FYR(J) =    -.82208068+02  FYI(J) =    -.12863547+03
J =  5  FYR(J) =    -.75405886+02  FYI(J) =    -.28632849+03

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 3.2
J =  1  FYR(J) =    -.17492878+01  FYI(J) =    -.39692703-02
J =  2  FYR(J) =    -.21898223+02  FYI(J) =    -.10637400+02
J =  3  FYR(J) =    -.51518709+02  FYI(J) =    -.69443727+02
J =  4  FYR(J) =    -.52569112+02  FYI(J) =    -.20497868+03
J =  5  FYR(J) =     .32677336+02  FYI(J) =    -.42961048+03

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y = 3.4
J =  1  FYR(J) =    -.17093749+01  FYI(J) =    -.59938923-00
J =  2  FYR(J) =    -.21070597+02  FYI(J) =    -.19700594+02
J =  3  FYR(J) =    -.36201143+02  FYI(J) =    -.10588472+03
J =  4  FYR(J) =     .21174397+02  FYI(J) =    -.29067785+03
J =  5  FYR(J) =     .25105467+03  FYI(J) =    -.57035594+03
```

Table E-4 (Cont'd)

```
TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y =  3.6
J =  1  FYR(J) =    -.14529858+01  FYI(J) =    -.11844860+01
J =  2  FYR(J) =    -.16545933+02  FYI(J) =    -.29681987+02
J =  3  FYR(J) =    -.18415579+01  FYI(J) =    -.14385580+03
J =  4  FYR(J) =     .15635155+03  FYI(J) =    -.36695578+03
J =  5  FYR(J) =     .61646407+03  FYI(J) =    -.65015223+03

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y =  3.8
J =  1  FYR(J) =    -.97820794-00  FYI(J) =    -.16740365+01
J =  2  FYR(J) =    -.74179436+01  FYI(J) =    -.39125926+02
J =  3  FYR(J) =     .56376314+02  FYI(J) =    -.17493213+03
J =  4  FYR(J) =     .36553682+03  FYI(J) =    -.40090809+03
J =  5  FYR(J) =     .11461494+04  FYI(J) =    -.57114778+03

TEST OF SUBROUTINE HGFY(N,Y,FYR,FYI) FOR N.LE.5 AND Y =  4.0
J =  1  FYR(J) =    -.31894834-00  FYI(J) =    -.19787108+01
J =  2  FYR(J) =     .66820224+01  FYI(J) =    -.45959718+02
J =  3  FYR(J) =     .14018605+03  FYI(J) =    -.18656795+03
J =  4  FYR(J) =     .64770570+03  FYI(J) =    -.34419971+03
J =  5  FYR(J) =     .18093090+04  FYI(J) =    -.19430102+03
```

Table E-5

THE HEIGHT GAIN FUNCTION $f_s'(y)$

$$f_s'(y) = -\frac{w_1'(t_s^o - y)}{w_1(t_s^o)} = \frac{Ai'[-\beta_s + y\exp(-i\frac{2}{3}\pi)]}{\exp(i\frac{2}{3}\pi)Ai(-\beta_s)} = -[FPYR(J)+iFPYI(J)]$$

$$f_s'(0) = 0 \ , \ s = J = 1,2,3,\ldots$$

```
TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 0.2
J =  1 FPYR(J) =    -.81059795-01 FPYI(J) =    -.17742269-00
J =  2 FPYR(J) =    -.29727631-00 FPYI(J) =    -.57402983-00
J =  3 FPYR(J) =    -.44559926-00 FPYI(J) =    -.86056310-00
J =  4 FPYR(J) =    -.56958488-00 FPYI(J) =    -.11099266+01
J =  5 FPYR(J) =    -.67896298-00 FPYI(J) =    -.13379091+01

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 0.4
J =  1 FPYR(J) =    -.11633140+00 FPYI(J) =    -.35870559-00
J =  2 FPYR(J) =    -.50413105-00 FPYI(J) =    -.12101973+01
J =  3 FPYR(J) =    -.74116873-00 FPYI(J) =    -.18653974+01
J =  4 FPYR(J) =    -.91811085-00 FPYI(J) =    -.24611172+01
J =  5 FPYR(J) =    -.10560531+01 FPYI(J) =    -.30254193+01

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 0.6
J =  1 FPYR(J) =    -.98231439-01 FPYI(J) =    -.54229230-00
J =  2 FPYR(J) =    -.55307698-00 FPYI(J) =    -.19505852+01
J =  3 FPYR(J) =    -.73219433-00 FPYI(J) =    -.31251052+01
J =  4 FPYR(J) =    -.77875100-00 FPYI(J) =    -.42461856+01
J =  5 FPYR(J) =    -.72787657-00 FPYI(J) =    -.53462727+01

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 0.8
J =  1 FPYR(J) =    -.18694966-01 FPYI(J) =    -.72038312-00
J =  2 FPYR(J) =    -.35449129-00 FPYI(J) =    -.28076883+01
J =  3 FPYR(J) =    -.19452000-00 FPYI(J) =    -.46911680+01
J =  4 FPYR(J) =     .26149514-00 FPYI(J) =    -.65586106+01
J =  5 FPYR(J) =     .96223660-00 FPYI(J) =    -.84316372+01

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 1.0
J =  1 FPYR(J) =     .12871639-00 FPYI(J) =    -.87872599-00
J =  2 FPYR(J) =     .20213357-00 FPYI(J) =    -.37514367+01
J =  3 FPYR(J) =     .11772721+01 FPYI(J) =    -.65132960+01
J =  4 FPYR(J) =     .28002930+01 FPYI(J) =    -.92977919+01
J =  5 FPYR(J) =     .50045191+01 FPYI(J) =    -.12083924+02
```

Table E-5 (Cont'd)

```
TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 1.2
J =  1 FPYR(J) =     .34620561-00 FPYI(J) =    -.99693047-00
J =  2 FPYR(J) =     .12389611+01 FPYI(J) =    -.46925993+01
J =  3 FPYR(J) =     .37593541+01 FPYI(J) =    -.83762085+01
J =  4 FPYR(J) =     .76136740+01 FPYI(J) =    -.12016350+02
J =  5 FPYR(J) =     .12723719+02 FPYI(J) =    -.15477475+02

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 1.4
J =  1 FPYR(J) =     .62880537-00 FPYI(J) =    -.10496925+01
J =  2 FPYR(J) =     .28696762+01 FPYI(J) =    -.54654790+01
J =  3 FPYR(J) =     .79511998+01 FPYI(J) =    -.98199940+01
J =  4 FPYR(J) =     .15563002+02 FPYI(J) =    -.13710465+02
J =  5 FPYR(J) =     .25607763+02 FPYI(J) =    -.16725555+02

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 1.6
J =  1 FPYR(J) =     .96148112-00 FPYI(J) =    -.10092729+01
J =  2 FPYR(J) =     .51659364+01 FPYI(J) =    -.58149509+01
J =  3 FPYR(J) =     .14073071+02 FPYI(J) =    -.10057363+02
J =  4 FPYR(J) =     .27356528+02 FPYI(J) =    -.12577946+02
J =  5 FPYR(J) =     .44823103+02 FPYI(J) =    -.12353588+02

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 1.8
J =  1 FPYR(J) =     .13166399+01 FPYI(J) =    -.84944202-00
J =  2 FPYR(J) =     .81095923+01 FPYI(J) =    -.53953089+01
J =  3 FPYR(J) =     .22191188+02 FPYI(J) =    -.79146220+01
J =  4 FPYR(J) =     .43098140+02 FPYI(J) =    -.58099227+01
J =  5 FPYR(J) =     .70239089+02 FPYI(J) =     .31833582+01

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 2.0
J =  1 FPYR(J) =     .16541683+01 FPYI(J) =    -.55085786-00
J =  2 FPYR(J) =     .11533250+02 FPYI(J) =    -.37905148+01
J =  3 FPYR(J) =     .31861902+02 FPYI(J) =    -.18401523+01
J =  4 FPYR(J) =     .61567689+02 FPYI(J) =     .10457679+02
J =  5 FPYR(J) =     .98791426+02 FPYI(J) =     .37642753+02

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y = 2.2
J =  1 FPYR(J) =     .19207697+01 FPYI(J) =    -.10749185+00
J =  2 FPYR(J) =     .15057028+02 FPYI(J) =    -.56630688-00
J =  3 FPYR(J) =     .41806538+02 FPYI(J) =     .99586570+01
J =  4 FPYR(J) =     .79229468+02 FPYI(J) =     .40832852+02
J =  5 FPYR(J) =     .12212040+03 FPYI(J) =     .10028559+03
```

Table E-5 (Cont'd)

```
TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  2.4
J =  1 FPYR(J) =     .20559415+01 FPYI(J) =     .46673623-00
J =  2 FPYR(J) =     .18036326+02 FPYI(J) =     .46370593+01
J =  3 FPYR(J) =     .49564185+02 FPYI(J) =     .29178223+02
J =  4 FPYR(J) =     .89070005+02 FPYI(J) =     .89726979+02
J =  5 FPYR(J) =     .12369509+03 FPYI(J) =     .19957245+03

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  2.6
J =  1 FPYR(J) =     .19987497+01 FPYI(J) =     .11324218+01
J =  2 FPYR(J) =     .19542961+02 FPYI(J) =     .11968350+02
J =  3 FPYR(J) =     .51221023+02 FPYI(J) =     .56815356+02
J =  4 FPYR(J) =     .79563832+02 FPYI(J) =     .15948731+03
J =  5 FPYR(J) =     .76154087+02 FPYI(J) =     .33836818+03

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  2.8
J =  1 FPYR(J) =     .16987926+01 FPYI(J) =     .18219288+01
J =  2 FPYR(J) =     .18408219+02 FPYI(J) =     .21192943+02
J =  3 FPYR(J) =     .41376768+02 FPYI(J) =     .92267553+02
J =  4 FPYR(J) =     .34314187+02 FPYI(J) =     .24737615+03
J =  5 FPYR(J) =    -.59604295+02 FPYI(J) =     .50586680+03

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  3.0
J =  1 FPYR(J) =     .11294327+01 FPYI(J) =     .24412707+01
J =  2 FPYR(J) =     .13357986+02 FPYI(J) =     .31508587+02
J =  3 FPYR(J) =     .13566644+02 FPYI(J) =     .13218696+03
J =  4 FPYR(J) =    -.66771097+02 FPYI(J) =     .34141783+03
J =  5 FPYR(J) =    -.33102658+03 FPYI(J) =     .66608839+03

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  3.2
J =  1 FPYR(J) =     .30141609-00 FPYI(J) =     .28782056+01
J =  2 FPYR(J) =     .32632682+01 FPYI(J) =     .41393879+02
J =  3 FPYR(J) =    -.38615334+02 FPYI(J) =     .16927116+03
J =  4 FPYR(J) =    -.24366706+03 FPYI(J) =     .41563716+03
J =  5 FPYR(J) =    -.78266474+03 FPYI(J) =     .74429638+03

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  3.4
J =  1 FPYR(J) =    -.72602319-00 FPYI(J) =     .30173384+01
J =  2 FPYR(J) =    -.12489749+02 FPYI(J) =     .48554537+02
J =  3 FPYR(J) =    -.11948773+03 FPYI(J) =     .19135496+03
J =  4 FPYR(J) =    -.50865792+03 FPYI(J) =     .42600988+03
J =  5 FPYR(J) =    -.14331648+04 FPYI(J) =     .61530291+03
```

Table E-5 (Cont'd)

```
TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  3.6
J =  1 FPYR(J) =    -.18401798+01 FPYI(J) =     .27616575+01
J =  2 FPYR(J) =    -.33539942+02 FPYI(J) =     .50046442+02
J =  3 FPYR(J) =    -.22829863+03 FPYI(J) =     .18136887+03
J =  4 FPYR(J) =    -.85433155+03 FPYI(J) =     .30948775+03
J =  5 FPYR(J) =    -.22376838+04 FPYI(J) =     .10136946+03

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  3.8
J =  1 FPYR(J) =    -.28796604+01 FPYI(J) =     .20580650+01
J =  2 FPYR(J) =    -.58079117+02 FPYI(J) =     .42651922+02
J =  3 FPYR(J) =    -.35555709+03 FPYI(J) =     .11888645+03
J =  4 FPYR(J) =    -.12376604+04 FPYI(J) =    -.10479266+02
J =  5 FPYR(J) =    -.30367860+04 FPYI(J) =    -.10083432+04

TEST OF SUBROUTINE HGFPY(N,Y,FPYR,FPYI) FOR N.LE.5 AND Y =  4.0
J =  1 FPYR(J) =    -.36516627+01 FPYI(J) =     .92241727-00
J =  2 FPYR(J) =    -.82499113+02 FPYI(J) =     .23556542+02
J =  3 FPYR(J) =    -.47938444+03 FPYI(J) =    -.16003852+02
J =  4 FPYR(J) =    -.15630375+04 FPYI(J) =    -.60788676+03
J =  5 FPYR(J) =    -.35006523+04 FPYI(J) =    -.29016662+04
```

Table E-6

THE HEIGHT GAIN FUNCTION $g_s(y)$

$$g_s(y) = -\frac{w_1(t_s^\infty - y)}{w_1'(t_s^\infty)} = \frac{Ai[-a_s + y\exp(-i\frac{2}{3}\pi)]}{\exp(-i\frac{2}{3}\pi)Ai'(-a_s)} = -[GYR(J) + iGYI(J)]$$

$$g_s(0) = 0\ ,\ s = J = 1,2,3,\ldots$$

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 0.2

J =	1	GYR(J) =	-.20141748-00	GYI(J) =	-.27113553-02
J =	2	GYR(J) =	-.20256845-00	GYI(J) =	-.47570512-02
J =	3	GYR(J) =	-.20350454-00	GYI(J) =	-.64424053-02
J =	4	GYR(J) =	-.20432716-00	GYI(J) =	-.79398178-02
J =	5	GYR(J) =	-.20507526-00	GYI(J) =	-.93151245-02

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 0.4

J =	1	GYR(J) =	-.41006308-00	GYI(J) =	-.21932527-01
J =	2	GYR(J) =	-.41886896-00	GYI(J) =	-.38874135-01
J =	3	GYR(J) =	-.42586648-00	GYI(J) =	-.53080253-01
J =	4	GYR(J) =	-.43188708-00	GYI(J) =	-.65887243-01
J =	5	GYR(J) =	-.43725257-00	GYI(J) =	-.77800848-01

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 0.6

J =	1	GYR(J) =	-.62905992-00	GYI(J) =	-.75154160-01
J =	2	GYR(J) =	-.65629295-00	GYI(J) =	-.13537752-00
J =	3	GYR(J) =	-.67681631-00	GYI(J) =	-.18721087-00
J =	4	GYR(J) =	-.69353877-00	GYI(J) =	-.23490903-00
J =	5	GYR(J) =	-.70759956-00	GYI(J) =	-.28005432-00

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 0.8

J =	1	GYR(J) =	-.85565010-00	GYI(J) =	-.18108702-00
J =	2	GYR(J) =	-.91101921-00	GYI(J) =	-.33314802-00
J =	3	GYR(J) =	-.94792631-00	GYI(J) =	-.46808085-00
J =	4	GYR(J) =	-.97355965-00	GYI(J) =	-.59506267-00
J =	5	GYR(J) =	-.99077038-00	GYI(J) =	-.71739352-00

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 1.0

J =	1	GYR(J) =	-.10794139+01	GYI(J) =	-.35888197-00
J =	2	GYR(J) =	-.11618280+01	GYI(J) =	-.67672353-00
J =	3	GYR(J) =	-.11999053+01	GYI(J) =	-.96758191-00
J =	4	GYR(J) =	-.12082572+01	GYI(J) =	-.12468703+01
J =	5	GYR(J) =	-.11927380+01	GYI(J) =	-.15197225+01

Table E-6 (Cont'd)

```
TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 1.2
J =  1  GYR(J) =   -.12805049+01  GYI(J) =   -.62613682-00
J =  2  GYR(J) =   -.13625718+01  GYI(J) =   -.12126042+01
J =  3  GYR(J) =   -.13457168+01  GYI(J) =   -.17639314+01
J =  4  GYR(J) =   -.12559362+01  GYI(J) =   -.23006450+01
J =  5  GYR(J) =   -.11037300+01  GYI(J) =   -.28281220+01

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 1.4
J =  1  GYR(J) =   -.14283726+01  GYI(J) =   -.99554102-00
J =  2  GYR(J) =   -.14342927+01  GYI(J) =   -.19806679+01
J =  3  GYR(J) =   -.12309604+01  GYI(J) =   -.29242437+01
J =  4  GYR(J) =   -.86024064-00  GYI(J) =   -.38448675+01
J =  5  GYR(J) =   -.33914699-00  GYI(J) =   -.47421474+01

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 1.6
J =  1  GYR(J) =   -.14816288+01  GYI(J) =   -.14701586+01
J =  2  GYR(J) =   -.12590161+01  GYI(J) =   -.29993241+01
J =  3  GYR(J) =   -.01592690-00  GYI(J) =   -.44711455+01
J =  4  GYR(J) =    .38519678-00  GYI(J) =   -.58824038+01
J =  5  GYR(J) =    .17178123+01  GYI(J) =   -.72100006+01

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 1.8
J =  1  GYR(J) =   -.13898627+01  GYI(J) =   -.20376276+01
J =  2  GYR(J) =   -.67854464-00  GYI(J) =   -.42435969+01
J =  3  GYR(J) =    .83254799-00  GYI(J) =   -.63290778+01
J =  4  GYR(J) =    .30534214+01  GYI(J) =   -.82225474+01
J =  5  GYR(J) =    .59392569+01  GYI(J) =   -.98378531+01

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 2.0
J =  1  GYR(J) =   -.10982454+01  GYI(J) =   -.26639637+01
J =  2  GYR(J) =    .49716704-00  GYI(J) =   -.56166993+01
J =  3  GYR(J) =    .35258340+01  GYI(J) =   -.82487614+01
J =  4  GYR(J) =    .78516312+01  GYI(J) =   -.10322998+02
J =  5  GYR(J) =    .13381091+02  GYI(J) =   -.11603099+02

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 2.2
J =  1  GYR(J) =   -.55561009-00  GYI(J) =   -.32881657+01
J =  2  GYR(J) =    .24664162+01  GYI(J) =   -.69180658+01
J =  3  GYR(J) =    .78964868+01  GYI(J) =   -.97147733+01
J =  4  GYR(J) =    .15494031+02  GYI(J) =   -.11088800+02
J =  5  GYR(J) =    .25022206+02  GYI(J) =   -.10484983+02
```

Table E-6 (Cont'd)

```
TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 2.4
J =  1  GYR(J) =     .27374032-00  GYI(J) =    -.38193570+01
J =  2  GYR(J) =     .53915074+01  GYI(J) =    -.78143335+01
J =  3  GYR(J) =     .14277759+02  GYI(J) =    -.98524704+01
J =  4  GYR(J) =     .26422244+02  GYI(J) =    -.86644955+01
J =  5  GYR(J) =     .41188855+02  GYI(J) =    -.30870224+01

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 2.6
J =  1  GYR(J) =     .13950253+01  GYI(J) =    -.41386146+01
J =  2  GYR(J) =     .93302394+01  GYI(J) =    -.78241112+01
J =  3  GYR(J) =     .22700072+02  GYI(J) =    -.73669500+01
J =  4  GYR(J) =     .40315197+02  GYI(J) =    -.30035844-00
J =  5  GYR(J) =     .60531960+02  GYI(J) =     .15570195+02

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 2.8
J =  1  GYR(J) =     .27664256+01  GYI(J) =    -.41077460+01
J =  2  GYR(J) =     .14144237+02  GYI(J) =    -.63317790+01
J =  3  GYR(J) =     .32596669+02  GYI(J) =    -.56693160-00
J =  4  GYR(J) =     .55356056+02  GYI(J) =     .17565259+02
J =  5  GYR(J) =     .78479660+02  GYI(J) =     .51758324+02

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 3.0
J =  1  GYR(J) =     .42839799+01  GYI(J) =    -.35868418+01
J =  2  GYR(J) =     .19393325+02  GYI(J) =    -.26485033+01
J =  3  GYR(J) =     .42437174+02  GYI(J) =     .12454100+02
J =  4  GYR(J) =     .67293973+02  GYI(J) =     .48798756+02
J =  5  GYR(J) =     .85253765+02  GYI(J) =     .11175233+03

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 3.2
J =  1  GYR(J) =     .57723219+01  GYI(J) =    -.24612336+01
J =  2  GYR(J) =     .24236689+02  GYI(J) =     .38620754+01
J =  3  GYR(J) =     .49352720+02  GYI(J) =     .33397172+02
J =  4  GYR(J) =     .68471184+02  GYI(J) =     .96383370+02
J =  5  GYR(J) =     .63874614+02  GYI(J) =     .19890821+03

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y = 3.4
J =  1  GYR(J) =     .69865948+01  GYI(J) =    -.67642115-00
J =  2  GYR(J) =     .27375179+02  GYI(J) =     .13583117+02
J =  3  GYR(J) =     .48879109+02  GYI(J) =     .63027858+02
J =  4  GYR(J) =     .47189535+02  GYI(J) =     .16031080+03
J =  5  GYR(J) =    -.10878451+02  GYI(J) =     .30849614+03
```

Table E-6 (Cont'd)

```
TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y =  3.6
J =  1  GYR(J) =     .76309413+01   GYI(J) =     .17233598+01
J =  2  GYR(J) =     .27081870+02   GYI(J) =     .26374540+02
J =  3  GYR(J) =     .35017028+02   GYI(J) =     .10011961+03
J =  4  GYR(J) =    -.11951250+02   GYI(J) =     .23445623+03
J =  5  GYR(J) =    -.17060078+03   GYI(J) =     .42005070+03

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y =  3.8
J =  1  GYR(J) =     .73975016+01   GYI(J) =     .45605559+01
J =  2  GYR(J) =     .21374433+02   GYI(J) =     .41265697+02
J =  3  GYR(J) =     .86936059-00   GYI(J) =     .14016021+03
J =  4  GYR(J) =    -.12586035+03   GYI(J) =     .30268367+03
J =  5  GYR(J) =    -.44767795+03   GYI(J) =     .48795804+03

TEST OF SUBROUTINE HGGY(N,Y,GYR,GYI) FOR N.LE.5 AND Y =  4.0
J =  1  GYR(J) =     .60264868+01   GYI(J) =     .75043356+01
J =  2  GYR(J) =     .83760788+01   GYI(J) =     .56178005+02
J =  3  GYR(J) =    -.59869993+02   GYI(J) =     .17408120+03
J =  4  GYR(J) =    -.30790649+03   GYI(J) =     .33502712+03
J =  5  GYR(J) =    -.85910077+03   GYI(J) =     .43275864+03
```

Table E-7

THE HEIGHT GAIN FUNCTION $g_s'(y)$

$$g_s'(y) = \frac{w_1'(t_s^\infty - y)}{w_1'(t_s^\infty)} = \frac{Ai'[-\alpha_s + y\exp(-i\frac{2}{3}\pi)]}{Ai'(-\alpha_s)} = GPYR(J) + iGPYI(J)$$

$$g_s'(0) = 1 \ , \ s = J = 1,2,3,\ldots$$

```
TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N = 5 AND Y = 0.2
J =  1 GPYR(J) =     .10205133+01 GPYI(J) =     .40780293-01
J =  2 GPYR(J) =     .10376182+01 GPYI(J) =     .71713101-01
J =  3 GPYR(J) =     .10514853+01 GPYI(J) =     .97301131-01
J =  4 GPYR(J) =     .10635846+01 GPYI(J) =     .12011312+00
J =  5 GPYR(J) =     .10745647+01 GPYI(J) =     .14112810-00

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N = 5 AND Y = 0.4
J =  1 GPYR(J) =     .10686725+01 GPYI(J) =     .16598491-00
J =  2 GPYR(J) =     .11319266+01 GPYI(J) =     .29678874-00
J =  3 GPYR(J) =     .11809780+01 GPYI(J) =     .40806884-00
J =  4 GPYR(J) =     .12221912+01 GPYI(J) =     .50955490-00
J =  5 GPYR(J) =     .12580464+01 GPYI(J) =     .60489720-00

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N = 5 AND Y = 0.6
J =  1 GPYR(J) =     .11191494+01 GPYI(J) =     .38188630-00
J =  2 GPYR(J) =     .12396467+01 GPYI(J) =     .70064884-00
J =  3 GPYR(J) =     .13225516+01 GPYI(J) =     .98251755-00
J =  4 GPYR(J) =     .13829884+01 GPYI(J) =     .12471880+01
J =  5 GPYR(J) =     .14269534+01 GPYI(J) =     .15017994+01

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N = 5 AND Y = 0.8
J =  1 GPYR(J) =     .11382220+01 GPYI(J) =     .69356551-00
J =  2 GPYR(J) =     .12905714+01 GPYI(J) =     .13144618+01
J =  3 GPYR(J) =     .13572264+01 GPYI(J) =     .18865691+01
J =  4 GPYR(J) =     .13666646+01 GPYI(J) =     .24389009+01
J =  5 GPYR(J) =     .13303431+01 GPYI(J) =     .29811059+01

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N = 5 AND Y = 1.0
J =  1 GPYR(J) =     .10829915+01 GPYI(J) =     .10993282+01
J =  2 GPYR(J) =     .11797408+01 GPYI(J) =     .21607415+01
J =  3 GPYR(J) =     .10917066+01 GPYI(J) =     .31750151+01
J =  4 GPYR(J) =     .86537374-00 GPYI(J) =     .41735974+01
J =  5 GPYR(J) =     .51959375-00 GPYI(J) =     .51638765+01
```

Table E-7 (Cont'd)

```
TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 1.2
J =  1 GPYR(J) =     .90248087-00 GPYI(J) =     .15841909+01
J =  2 GPYR(J) =     .76245495-00 GPYI(J) =     .32325230+01
J =  3 GPYR(J) =     .23929378-00 GPYI(J) =     .48471341+01
J =  4 GPYR(J) =    -.59906833-00 GPYI(J) =     .64430758+01
J =  5 GPYR(J) =    -.17250233+01 GPYI(J) =     .80119323+01

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 1.4
J =  1 GPYR(J) =     .54156506-00 GPYI(J) =     .21128957+01
J =  2 GPYR(J) =    -.14339948-00 GPYI(J) =     .44652860+01
J =  3 GPYR(J) =    -.15857376+01 GPYI(J) =     .67792939+01
J =  4 GPYR(J) =    -.36947826+01 GPYI(J) =     .90105704+01
J =  5 GPYR(J) =    -.64321790+01 GPYI(J) =     .11100826+02

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 1.6
J =  1 GPYR(J) =    -.51387051-01 GPYI(J) =     .26235091+01
J =  2 GPYR(J) =    -.17416135+01 GPYI(J) =     .57028639+01
J =  3 GPYR(J) =    -.48432740+01 GPYI(J) =     .86331017+01
J =  4 GPYR(J) =    -.92393076+01 GPYI(J) =     .11218970+02
J =  5 GPYR(J) =    -.14869005+02 GPYI(J) =     .13273506+02

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 1.8
J =  1 GPYR(J) =    -.91263425-00 GPYI(J) =     .30233315+01
J =  2 GPYR(J) =    -.42231832+01 GPYI(J) =     .66619482+01
J =  3 GPYR(J) =    -.99887680+01 GPYI(J) =     .97463735+01
J =  4 GPYR(J) =    -.18042143+02 GPYI(J) =     .11747671+02
J =  5 GPYR(J) =    -.28245572+02 GPYI(J) =     .12184427+02

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 2.0
J =  1 GPYR(J) =    -.20471672+01 GPYI(J) =     .31894595+01
J =  2 GPYR(J) =    -.77028220+01 GPYI(J) =     .69044852+01
J =  3 GPYR(J) =    -.17314343+02 GPYI(J) =     .90290804+01
J =  4 GPYR(J) =    -.30557677+02 GPYI(J) =     .83626230+01
J =  5 GPYR(J) =    -.47034778+02 GPYI(J) =     .38175676+01

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 2.2
J =  1 GPYR(J) =    -.34102030+01 GPYI(J) =     .29767052+01
J =  2 GPYR(J) =    -.12130638+02 GPYI(J) =     .58328312+01
J =  3 GPYR(J) =    -.26686734+02 GPYI(J) =     .49078859+01
J =  4 GPYR(J) =    -.46274705+02 GPYI(J) =    -.22380004+01
J =  5 GPYR(J) =    -.69732343+02 GPYI(J) =    -.17808027+02
```

Table E-7 (Cont'd)

```
TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 2.4
J =  1 GPYR(J) =    -.48890671+01 GPYI(J) =     .22353970+01
J =  2 GPYR(J) =    -.17176183+02 GPYI(J) =     .27277597+01
J =  3 GPYR(J) =    -.37176061+02 GPYI(J) =    -.46111128+01
J =  4 GPYR(J) =    -.62801281+02 GPYI(J) =    -.24300828+02
J =  5 GPYR(J) =    -.90939591+02 GPYI(J) =    -.60339312+02

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 2.6
J =  1 GPYR(J) =    -.62902455+01 GPYI(J) =     .84056528-00
J =  2 GPYR(J) =    -.22104870+02 GPYI(J) =    -.31478631+01
J =  3 GPYR(J) =    -.46606030+02 GPYI(J) =    -.21667563+02
J =  4 GPYR(J) =    -.74694300+02 GPYI(J) =    -.62347681+02
J =  5 GPYR(J) =    -.98851490+02 GPYI(J) =    -.13155881+03

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 2.8
J =  1 GPYR(J) =    -.75379594+01 GPYI(J) =    -.12681035+01
J =  2 GPYR(J) =    -.25676934+02 GPYI(J) =    -.12359819+02
J =  3 GPYR(J) =    -.51117337+02 GPYI(J) =    -.47955532+02
J =  4 GPYR(J) =    -.72257309+02 GPYI(J) =    -.11962638+03
J =  5 GPYR(J) =    -.72654373+02 GPYI(J) =    -.23578718+03

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 3.0
J =  1 GPYR(J) =    -.76913535+01 GPYI(J) =    -.40394570+01
J =  2 GPYR(J) =    -.26116650+02 GPYI(J) =    -.25020012+02
J =  3 GPYR(J) =    -.44920700+02 GPYI(J) =    -.83732820+02
J =  4 GPYR(J) =    -.40800705+02 GPYI(J) =    -.19536877+03
J =  5 GPYR(J) =     .13956369+02 GPYI(J) =    -.36737613+03

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 3.2
J =  1 GPYR(J) =    -.69865280+01 GPYI(J) =    -.72653465+01
J =  2 GPYR(J) =    -.21230481+02 GPYI(J) =    -.40427937+02
J =  3 GPYR(J) =    -.20512066+02 GPYI(J) =    -.12642486+03
J =  4 GPYR(J) =     .30794494+02 GPYI(J) =    -.28073326+03
J =  5 GPYR(J) =     .21595064+03 GPYI(J) =    -.50088553+03

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 3.4
J =  1 GPYR(J) =    -.49064146+01 GPYI(J) =    -.10548927+02
J =  2 GPYR(J) =    -.87015509+01 GPYI(J) =    -.56692151+02
J =  3 GPYR(J) =     .30297857+02 GPYI(J) =    -.16897067+03
J =  4 GPYR(J) =     .18702579+03 GPYI(J) =    -.35377117+03
J =  5 GPYR(J) =     .55864176+03 GPYI(J) =    -.57875136+03
```

Table E-7 (Cont'd)

```
TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR  N=5 AND Y = 3.6
J =  1 GPYR(J) =    -.12754405+01 GPYI(J) =    -.13302737+02
J =  2 GPYR(J) =     .13316375+02 GPYI(J) =    -.70437594+02
J =  3 GPYR(J) =     .11419202+03 GPYI(J) =    -.19829828+03
J =  4 GPYR(J) =     .41888331+03 GPYI(J) =    -.37457464+03
J =  5 GPYR(J) =     .10686715+04 GPYI(J) =    -.49956673+03

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR  N=5 AND Y = 3.8
J =  1 GPYR(J) =     .38328255+01 GPYI(J) =    -.14793096+02
J =  2 GPYR(J) =     .45369728+02 GPYI(J) =    -.76755306+02
J =  3 GPYR(J) =     .23275085+03 GPYI(J) =    -.19462313+03
J =  4 GPYR(J) =     .73207103+03 GPYI(J) =    -.28293075+03
J =  5 GPYR(J) =     .17176650+04 GPYI(J) =    -.11363028+03

TEST OF SUBROUTINE HGGPY(N,Y,GPYR,GPYI) FOR N=5 AND Y = 4.0
J =  1 GPYR(J) =     .99971268+01 GPYI(J) =    -.14243368+02
J =  2 GPYR(J) =     .85679182+02 GPYI(J) =    -.69585072+02
J =  3 GPYR(J) =     .37759458+03 GPYI(J) =    -.13256712+03
J =  4 GPYR(J) =     .10903817+04 GPYI(J) =    -.21427482+01
J =  5 GPYR(J) =     .23790268+04 GPYI(J) =     .76325448+03
```

```
      SUBROUTINE VTWTZ(Z,VT,WT,VTP,WTP)
      COMPLEX          Z,VT,WT,VTP,WTP
      DOUBLE PRECISION AZERO,A(22),BZERO,B(22),CZERO,C(22),DZERO,D(22)
      DOUBLE PRECISION DRZ,DIZ,D2RZ,D2IZ,D3RZ,D3IZ,FR,FI,FPR,FPI,GR,GI,
     A        GPR,GPI,AREAL(23),AIMAG(23),ZERO,TWO,CA,CB,
     B        DVTR,DVTI,DVTPR,DVTPI,DWTR,DWTI,DWTPR,DWTPI
      DATA AZERO,BZERO,(A(M),B(M),M=1,11) /
     A             0.93043671693   D  0 ,        0.67829872514   D  0 ,
     B            31.01455723097   D  0 ,       11.30497875240   D  0 ,
     C           206.76371487316   D  0 ,       53.83323215431   D  0 ,
     D           574.34365242545   D  0 ,      119.62940478735   D  0 ,
     E           870.21765519008   D  0 ,      153.37103177865   D  0 ,
     F           828.77871922864   D  0 ,      127.80919314888   D  0 ,
     G           541.68543740434   D  0 ,       74.74221821572   D  0 ,
     H           257.94544638302   D  0 ,       32.35593862152   D  0 ,
     I            93.45849506631   D  0 ,       10.78531287384   D  0 ,
     J            26.62635187074   D  0 ,        2.85325737403   D  0 ,
     K             6.121000430056  D  0 ,        0.613603736351  D  0 ,
     L             1.159280384480  D  0 ,        0.109376780098  D  0 /
      DATA (A(M),B(M),M=12,22)                                  /
     A             0.184012759441  D  0 ,        0.16422939955   D- 1 ,
     B             0.24833030964   D- 1 ,        0.2105505122    D- 2 ,
     C             0.2884208010    D- 2 ,        0.233167788     D- 3 ,
     D             0.291334142     D- 3 ,        0.22528289      D- 4 ,
     E             0.25827495      D- 4 ,        0.1915671       D- 5 ,
     F             0.2025686       D- 5 ,        0.144470        D- 6 ,
     G             0.141557        D- 6 ,        0.9729          D- 8 ,
     H             0.8870          D- 8 ,        0.589           D- 9 ,
     I             0.501           D- 9 ,        0.32            D-10 ,
     J             0.26            D-10 ,        0.2             D-11 ,
     K             0.1             D-11 ,        0.0             D  0 /
      DATA CZERO,DZERO,(C(M),D(M),M=1,11) /
     A             0.46521835846   D  0 ,        0.67829872514   D  0 ,
     B             6.20291144619   D  0 ,       45.21991500962   D  0 ,
     C            25.84546435915   D  0 ,      376.83262508015   D  0 ,
     D            52.21305931140   D  0 ,     1196.29404787350   D  0 ,
     E            62.15840394215   D  0 ,     1993.82341312250   D  0 ,
     F            48.75168936639   D  0 ,     2044.94709038206   D  0 ,
     G            27.08427187022   D  0 ,     1420.10214609865   D  0 ,
```

```
      H            11.21501940796    D  0 ,      711.83064967351    D  0 ,
      I             3.59455750255    D  0 ,      269.63282184603    D  0 ,
      J             0.91815006451    D  0 ,       79.89120647290    D  0 ,
      K             0.191281263439   D  0 ,       19.021715826880   D  0 ,
      L             0.33122296699    D- 1 ,        3.718810523339   D  0 /
       DATA (C(M),D(M),M=12,22)                 /
      A             0.4842441038     D- 2 ,        0.607648778323   D  0 ,
      B             0.605683682      D- 3 ,        0.84220204896    D- 1 ,
      C             0.65550182       D- 4 ,        0.10026214869    D- 1 ,
      D             0.6198599        D- 5 ,        0.1036301278     D- 2 ,
      E             0.516550         D- 6 ,        0.93867869       D- 4 ,
      F             0.38220          D- 7 ,        0.7512435        D- 5 ,
      G             0.2528           D- 8 ,        0.535074         D- 6 ,
      H             0.150            D- 9 ,        0.34135          D- 7 ,
      I             0.8              D-11 ,        0.1962           D- 8 ,
      J             0.0              D  0 ,        0.102            D- 9 ,
      K             0.0              D  0 ,        0.5              D-11 /
       DATA ZERO,TWO,CA,CB/0.0D0,2.0D0,1.171416657372D0,0.676317722467D0/
       IF(CABS(Z).GT.(6.1)) GO TO 99
       M=3+INT(20.*CABS(Z)/6.1)
       DRZ=DBLE(REAL(-Z))
       DIZ=DBLE(REAL((0.,1.)*Z))
       D2RZ=DRZ*DRZ-DIZ*DIZ
       D2IZ=TWO*DRZ*DIZ
       D3RZ=DRZ*D2RZ-DIZ*D2IZ
       D3IZ=DRZ*D2IZ+DIZ*D2RZ
       D3RZ = D3RZ/(-200.0D0)
       D3IZ = D3IZ/(-200.0D0)
       DO 39  I=1,23
       AREAL(I)= ZERO
       AIMAG(I)= ZERO
    39 CONTINUE
       IF(M.GT.22)  M=22
       DO 1 N=M,1,-1
       AREAL(N)=D3RZ*AREAL(N+1) - D3IZ*AIMAG(N+1) + A(N)
       AIMAG(N)=D3RZ*AIMAG(N+1) + D3IZ*AREAL(N+1)
     1 CONTINUE
       FR = D3RZ*AREAL(1)-D3IZ*AIMAG(1)+AZERO
       FI = D3RZ*AIMAG(1)+D3IZ*AREAL(1)
```

```
      DO 2 N=M,1,-1
      AREAL(N)=D3RZ*AREAL(N+1) - D3IZ*AIMAG(N+1) + B(N)
      AIMAG(N)=D3RZ*AIMAG(N+1) + D3IZ*AREAL(N+1)
    2 CONTINUE
      GPR= D3RZ*AREAL(1)-D3IZ*AIMAG(1)+BZERO
      GPI= D3RZ*AIMAG(1)+D3IZ*AREAL(1)
      GR = DRZ*GPR-DIZ*GPI
      GI = DRZ*GPI+DIZ*GPR
      DO 3 N=M,1,-1
      AREAL(N)=D3RZ*AREAL(N+1)-D3IZ*AIMAG(N+1)+C(N)
      AIMAG(N)=D3RZ*AIMAG(N+1)+D3IZ*AREAL(N+1)
    3 CONTINUE
      GPR = D3RZ*AREAL(1)-D3IZ*AIMAG(1)+CZERO
      GPI = D3RZ*AIMAG(1)+D3IZ*AREAL(1)
      FPR =-D2RZ*GPR + D2IZ*GPI
      FPI =-D2RZ*GPI - D2IZ*GPR
      DO 4 N=M,1,-1
      AREAL(N)=D3RZ*AREAL(N+1)-D3IZ*AIMAG(N+1)+D(N)
      AIMAG(N)=D3RZ*AIMAG(N+1)+D3IZ*AREAL(N+1)
    4 CONTINUE
      GPR = D3RZ*AREAL(1)-D3IZ*AIMAG(1)+DZERO
      GPI = D3RZ*AIMAG(1)+D3IZ*AREAL(1)
C-----NOW CONSTRUCT FOCK AIRY FUNCTIONS
      DVTR   = CB*(FR + GR)
      DVTI   = CB*(FI + GI)
      DVTPR  =-CB*(FPR + GPR)
      DVTPI  =-CB*(FPI + GPI)
      DWTR   = CA*(FR - GR) - DVTI
      DWTI   = CA*(FI - GI) + DVTR
      DWTPR  =-CA*(FPR - GPR) - DVTPI
      DWTPI  =-CA*(FPI - GPI) + DVTPR
      VT     = CMPLX(SNGL(DVTR),SNGL(DVTI))
      VTP    = CMPLX(SNGL(DVTPR),SNGL(DVTPI))
      WT     = CMPLX(SNGL(DWTR),SNGL(DWTI))
      WTP    = CMPLX(SNGL(DWTPR),SNGL(DWTPI))
      RETURN
   99 WRITE(6,110) Z
  110 FØRMAT(25H1VTWTZ CANNØT HANDLE Z = ,(E15.8,E15.8))
      CALL EXIT
      END
```

UNITED STATES AIR FORCE
AFGL Research Library

REFERENCES FOR APPENDIX E

1. P. A. Azriliant and M. G. Belkina, Numerical Results of the Theory of Diffraction of Radio Waves Around the Earth's Surface, Moscow, Soviet Radio Press, 1957. (The author is endebted to his colleague Mr. M. D. Friedman for preparing an English translation of this important publication. The would like to see this report made available to people interested in diffraction theory but the cost of printing the text, tables, and graphs is prohibitive and the demand for this material so limited that no attempts have been made to seek a publisher.)

2. Harvard University Computation Laboratory, Tables of the Modified Hankel Functions of Order One-Third and of Their Derivatives, Annals of the Computation Laboratory of Harvard University, Vol. II, Cambridge, Mass., Harvard University Press, 1945

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY *(Corporate author)*
Lockheed Aircraft Corp., Lockheed Missiles & Space Co., 1111 Lockheed Way, Sunnyvale, Calif.

2a. REPORT SECURITY CLASSIFICATION
Unclassified

2b. GROUP

3. REPORT TITLE
Numerical Investigation of Electromagnetic Scattering and Diffraction by Convex Objects

4. DESCRIPTIVE NOTES *(Type of report and inclusive dates)* Scientific Report. Final. Period covered: 16 Nov 64 through 15 Nov 65. Accepted: 14 Mar 66.

5. AUTHOR(S) *(Last name, first name, initial)*
Logan, Nelson A.

6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
Dec 65	707	240

8a. CONTRACT OR GRANT NO. AF19(628)-4393	9a. ORIGINATOR'S REPORT NUMBER(S)
b. PROJECT AND TASK NO. 460004	M-76-66-1
c. DOD ELEMENT 62405304	9b. OTHER REPORT NO(S) *(Any other numbers that may be assigned this report)*
d. DOD SUBELEMENT 674600	AFCRL-66-153

10. AVAILABILITY/LIMITATION NOTICES
Distribution of this document is unlimited.

11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY
	Hq AFCRL, OAR (CRD) United States Air Force L.G. Hanscom Field, Bedford, Mass.

13. ABSTRACT

The solutions $D(x,y,y_o,q)$ of the partial differential equation $D_{yy} + iD_x + yD = 0$ are employed as a mathematical model for the fields reflected from and diffracted around an opaque convex surface. This solution involves the Airy function $w_1(t)$ which is a solution of $w''_1(t) - tw_1(t) = 0$. FORTRAN computer programs and tables are given for the appropriate Airy functions. Representations for the roots of the equation $w'_1(t_s) - qw_1(t_s) = 0$ which play a fundamental role in the theory are discussed. As a function of q, these roots satisfy the Riccati equation $(t - q^2)(dt/dq) = 1$. The Riccati equation is used to develop several representations for these roots. FORTRAN programs are presented for the evaluation of these constants. Tables of the roots are given for certain cases which have been shown to be of practical significance. Asymptotic expansions are developed for the height gain function $F(y)$ which is a solution of $F''(y) + (k^2f(y) - \lambda_s)F(y) = 0$. The theory of representing functions by a series of Chebyshev polynomials is reviewed and a discussion is given of the significance of these expansions for future work in diffraction theory.

DD FORM 1 JAN 64 1473

Unclassified
Security Classification

Unclassified
Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Diffraction Functions	8	3				
Airy Functions	8	3				
Electromagnetic Theory	8	2				
Height Gain Function	8	3				
Ricatti Equation	8	1				
Asymptotic Expansions	8	3				
FORTRAN programs	8	2				
Chebyshev Polynomials	8	2				
Propagation Constants	8	3				
WKB Approximation	8	2				
Convex Surface	5	3				
Inhomogeneous Medium	5	2				
Theory	10	3				
Airy	0	3				
Fock	0	3				
Bremmer	0	2				
Keller	0	2				

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization *(corporate author)* issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers *(either by the originator or by the sponsor)*, also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

(1) "Qualified requesters may obtain copies of this report from DDC."

(2) "Foreign announcement and dissemination of this report by DDC is not authorized."

(3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."

(4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."

(5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring *(paying for)* the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as *(TS), (S), (C), or (U).*

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

Unclassified
Security Classification

UNITED STATES AIR FORCE
AFGL Research Library