

AD 663845

AD

Report Number 3

The Role of Adrenal Cortex in the Process of Acclimatization to
High Altitudes

FINAL REPORT



JAN 11 1968

By: Federico Moncloa

October 1967

DA-ARO-49-092-65-G89

Army Project Number: 2N014501B71D 00 025 LA

Instituto de Investigaciones de la Altura, Universidad Peruana Cayetano Heredia
Lima, Peru

U. S. Army Element

Defense Research Office, Latin America

Rio de Janeiro, Brazil

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

15

Distribution of this document is unlimited.

Destroy this report when it is no longer needed. Do not return it to the originator.

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

ACCESSION for	
CPSTI	WHITE SECTION ☒
DDC	BWP SECTION ☐
UNAUTHORISED	☐
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
DIST.	AVAIL. and/or SPECIAL

REPORT NUMBER 3

AD

THE ROLE OF ADRENAL CORTEX IN THE PROCESS OF ACCLIMATIZATION TO
HIGH ALTITUDES

FINAL REPORT

BY

Federico Moncloa

OCTOBER 1967

U. S. Army Element
Defense Research Office for Latin America
Rio de Janeiro, Brazil
DA-ARO-49-092-61-00

U. S. Department of the Army Project Number 2N014501B71D 00 025 LA

Instituto de Investigaciones de la Altura
Universidad Peruana Cayetano Heredia
Apartado 6083, Lima, Perú.

Distribution of this Document is Unlimited

ABSTRACT

The first part refers to some indexes of adrenal function and cortisol metabolism in acute exposure of sea level natives to 4,300 meters of altitude. The results show the following: (a) Cortisol secretion rate is temporarily increased correlating with increases in its plasma concentration and in the urinary excretion of its metabolites; (b) Hypoxia does not result in a maximal stimulus for the adrenal cortex since it can be further stimulated by exogenous corticotrophin; (c) The adrenocortical response to high altitude can be blocked by the administration of dexamethasone; (d) The second compartment in which cortisol is distributed is diminished. This may be related with the contraction of the radiosulfate space, observed under the same experimental conditions; and (e) The metabolic clearance rate of cortisol is not significantly altered.

The second part refers to the study of corticotrophin half-life ($t_{1/2}$). Our results indicate that a 24 aminoacids ACTH has a $t_{1/2}$ slightly longer than a 39 aminoacids ACTH. Using the same type of experiments we have studied the $t_{1/2}$ of the 39 aminoacids ACTH in 11 high altitude natives, the results were similar to the ones obtained in the sea level controls.

ADRENAL FUNCTION AND CORTISOL METABOLISM

This part comprises four papers; two have been published and the other two have been accepted for publication. A summary of each one follows:

1.- Endocrine Studies at High Altitude. II Adrenal Cortical Function in Sea Level Natives Exposed to High Altitude (4300 meters) for Two Weeks

Moncloa, F., Donayre, J., Sobrevilla, L.A., & Guerra-García, R.

J. Clin. Endocrinol. 12:1640-1642, 1965.

Ten young men, sailors, sea level natives were exposed for two weeks to 4300 meters of altitude. The ascent to high altitude was associated with a transitory but definite increase in the cortisol secretion rate and in the urinary excretion of 17,21-dihydroxy-20-ketocorticosteroids and in 17 ketogenic steroids. The increase in cortisol secretion rate suggests that the greater urinary excretion of the urinary steroids was due to that increase. Fig. 1.

2.- ACTH Stimulation and Dexamethasone Inhibition in Newcomers to High Altitude

Moncloa, F., Beteta, L., Velasco, I., & Goñez, C.
Proc. Soc. Exp. Biol. & Med. 122:1029-1031, 1966.

We studied 3 groups of young adult males, natives and residents of sea level. The first group was formed by 12 subjects; this group was stimulated with ACTH twice, one at sea level and the other during a trip to high altitude. The increase in urinary steroids was of the same magnitude in both circumstances and significantly higher than the one obtained by simple exposure as reported above (i.e. in another group). The second and third group were formed by 7 subjects each. Both groups were taken to 4300 meters under dexamethasone administration; in the second group the dose was 0.5 mg/6 hours and in the third group it was 2.0 mg/6 hours. The results indicate that when dexamethasone is given the hypoxic stimuli are easily blocked. Fig. 2.

3.- Radiosulfate Space in Humans at High Altitude

Malpartida, M. & Moncloa, F.
Proc. Soc. Exp. Biol. & Med.

We studied 14 sea level natives, 9 of which were observed again after 36 hours of exposure to high altitude (4,300 m). Seven high altitude natives were also studied. The results indicated that extracellular space was significantly decreased (1.4 liters). Body weight was also decreased (1.2 kg.). However, there was no statistical cor-

relation between these variables. This was interpreted as⁴ meaning that the decrease in body weight is a result of several changes, the decrease in extracellular fluid being only one of them. Table 1.

4.- Plasma Cortisol Concentration and Plasma Dissappearance
Rate of 4-¹⁴C-cortisol in Newcomers to High Altitude

Moncloa, F., Velasco, I., & Beteta, L.
J. Clin. Endocrinol.

Plasma cortisol was measured in 10 sea level natives before and during the first day of their stay at 4,300 meters. There was a significant increase in the mean concentration and the normal circadian rhythm was altered. In the control observations the lower value was obtained at 12 p.m. but under hypoxia at that time there was a 3.5 fold increase. The plasma disappearance curve of 4-¹⁴C-cortisol was studied in another group of nine sea level subjects. The experiment was done twice; a control observation at sea level and the second one after 36 hours of exposure to high altitude. The curves obtained were double exponentials; therefore, a two-compartment model describes the data. In brief, the results show that the second compartment was smaller in acute exposure to high altitude; this finding may be partially explained by the contraction in the extracellular space. The other indexes, namely the transfer constants and the metabolic clearance rate were not statistically different. Tables 2 and 3.

HALF-LIFE OF CORTICOTROPHIN (ACTH)

This part comprises one published paper and a summary of recent observations.

1.- Biological Half-Life of a Synthetic Corticotrophin in Men Moncloa, F., Velasco, I., & Beteta, L. Acta Endocrinologica (Kbh) 52:337-340, 1966.

We studied nine male students of our School of Medicine. Two experiments were performed in each subject. The first time they were intravenously injected with 0.05 IU of ACTH/kg of body weight and plasma samples were analyzed for cortisol after 60, 75, 90, 105, 180 and 240 minutes. The second time, at least three days later, the injection was of 0.25 IU of ACTH/kg; and the plasma samples obtained at 120, 135, 150, 165, 240 and 300 minutes afterwards. The $t_{1/2}$ was calculated using the following equation

$$t_{1/2} = \frac{.693 (h-h')}{\log a - \log a'}$$

where a and a' were the doses injected; h the time when cortisol starts to fall with dose a ; and h' with dose a' . The $t_{1/2}$ ranged from 19.4 to 38.7 minutes, the mode being 38.7 minutes, the mode being 32.3 minutes.

2.- Comparison of $t_{1/2}$ of Two Synthetic ACTH Preparations and Further Observations in High Altitude Natives

The method used is described in the previous paper (vide supra). In 10 subjects we have studied the $t_{1/2}$ of two different ACTH preparations; the 24 aminoacids ACTH (Ciba 30920) and the 39 aminoacids ACTH (Ferring A B). The results show that the 39 a.a. preparation has the same $t_{1/2}$ in four subjects and shorter in the other six. With the 39 aminoacids ACTH we have compared the $t_{1/2}$ in 14 natives of sea level versus 11 high altitude natives. The results (Fig. 3) indicate that in high altitude natives the preparation tested has the same $t_{1/2}$.

Table 1.- Extracellular space measured with S³⁵

	Body weight kg	Radiosulfate space liters	% body weight
High Altitude Natives (7)	55.8 $\pm$ 1.5	9.2 $\pm$ 0.5	16.4 $\pm$ 0.6
Sea Level Natives Contents (14)	65.6 $\pm$ 1.2	9.8 $\pm$ 0.3	15.0 $\pm$ 0.4

Nine Sea Level Natives

	Body weight kg	Radiosulfate space liters
Before exposure to high altitude	66.3 $\pm$ 1.4	10.2 $\pm$ 0.4
2nd day of exposure to 4,300 meters	65.1 $\pm$ 1.4	8.8 $\pm$ 0.2
Decrease	1.2 $\pm$ 0.2 ^a	1.4 $\pm$ 0.2 ^a

^a P < 0.001

Table 2.- Plasma cortisol concentration in acute exposure to high altitude ug/100 ml

	12 p.m.	8 a.m.	12 m.	4 p.m.	8 p.m.	Mean concentrations
Sea level controls	4.3 ± 0.9	17.7 ± 1.3	10.5 ± 1.5	8.4 ± 0.9	7.6 ± 1.3	9.9 ± 0.6
High altitude exposure	15.1 ± 4.0	20.0 ± 2.2	18.0 ± 4.6	12.5 ± 1.5	8.7 ± 2.1	15.5 ± 2.1
p Values	< 0.05	> 0.20	< 0.20 > 0.10	< 0.05	> 0.20	< 0.05

The figures are the means ± S.E.

Table 3.- Results obtained from the plasma disappearance curve of radio active cortisol using a two-compartment model in 9 sea level volunteers before and during exposure to 4,300 m. Figures are the means $\pm$ SE.

	Control at sea level	2nd day of exposure at high altitude
V_m (liters)	5.45 ± 0.59	6.55 ± 1.01
V_n (liters)	11.71 ± 1.47	7.08 ± 0.97^a
k_{13} (min^{-1})	0.1032 ± 0.0146	0.0749 ± 0.0163
k_{31} (min^{-1})	0.0476 ± 0.0063	0.0557 ± 0.0088
k_{12} (min^{-1})	0.0321 ± 0.0011	0.0349 ± 0.0062
α (min^{-1})	0.1742 ± 0.0194	0.1534 ± 0.0251
β (min^{-1})	0.00885 ± 0.00116	0.01220 ± 0.00131
MCR liters/ 24 h	239.5 ± 22.7	259.7 ± 15.7

^a $p < 0.02$. In all the other instances the differences were not statistically different.

URINARY STEROIDS AND CORTISOL SECRETION RATE




Fig. 1

URINARY STEROIDS IN SEA LEVEL NATIVES TAKEN TO
AN ALTITUDE OF 4,300 m. UNDER ACTH & DEXAME-
THASONE MEDICATION






Fig. 2

Fig. 3 BIOLOGICAL HALF-LIFE OF ACNH 11



DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Instituto de Investigaciones de la Altura Universidad Peruana "Cayetano Heredia" Lima, Peru		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP NA
3. REPORT TITLE THE ROLE OF ADRENAL CORTEX IN THE PROCESS OF ACCLIMATIZATION TO HIGH ALTITUDES		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Final Report		
5. AUTHOR(S) (Last name, first name, initial) Moncloa, Federico		
6. REPORT DATE October 1967	7a. TOTAL NO. OF PAGES 12	7b. NO. OF REFS
8a. CONTRACT OR GRANT NO. DA-ARO-49-092-65-089	8a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO. 2N014501B71D 00 025 1A		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.	Report Number 3	
10. AVAILABILITY/LIMITATION NOTICES Unlimited		
11. SUPPLEMENTARY NOTES None	12. SPONSORING MILITARY ACTIVITY U. S. Army Element Defense Research Office, Latin America U. S. Embassy, APO New York 09676	
13. ABSTRACT Indexes of adrenal function and cortisol metabolism in acute exposure of sea level natives to 4,300 meters of altitude prepared. a) Cortisone secretion rate is temporarily increased correlating with increases in its plasma concentration and in the urinary excretion of its metabolites; b) Hypoxia does not result in a maximal corticotrophin; c) The adrenocortical response to high altitude can be blocked by administration of dexamethasone; d) Second compartment in which cortisol is distributed is diminished. May be related with the contraction of radiosulfate space observed under same experimental conditions; e) Metabolic clearance rate of cortisol is not significantly altered. Corticotrophin half-life ($t_{1/2}$). Synthetic 24 aminoacid ACTH has $t_{1/2}$ slightly longer than synthetic 39 aminoacids ACTH. Using same type of experiments we studied $t_{1/2}$ of the 39 aminoacids ACTH in 11 high altitude natives; results were similar to those obtained in sea level controls.		

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
	Endocrinology Adrenal Cortex Adrenal Hormones Acclimatization Steroids Tolerance to Hypoxia Metabolic Clearance					

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.
- 2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.
- 8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, &c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).
10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:
- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."
- If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.
11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.
12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.
13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.
- It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).
- There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.
14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.