

AD 682634

TRANSLATION NO. 2387

DATE: Jan 1969

~~DDC AVAILABLE FOR NOTICE~~

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of commanding Officer, Fort Detrick, ATMC SMUPD-AET, Frederick, Md. 21701

DDC
FEB 27 1969
E

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

This document is approved
for public release and sale; its
distribution is unlimited

UDC 616.935-078.4
+ 576.851.49.094.1

BIOLOGICAL PECULIARITIES OF BOYD-10
SHIGELLA, ISOLATED IN GROUP INCIDENCE OF
DYSENTERY

Following is a translation of an author's abstract by A.Ya. Borovaya, Ye.Ye. Verkholomov and V.A. Arbuzova (Institute of Epidemiology and Microbiology imeni Pasteur) from the Russian-language periodical Zhurnal Mikrobiologii Epidemiologii i Immunobiologii (Journal of Microbiology, Epidemiology and Immunobiology), No 9, Moscow, September 1967, page 147.7

During the bacteriological examination of feces of 23 patients obtained from a focus of severe dysentery, dysentery bacteria were isolated from ten specimens which were identified as a Boyd subspecies, serological type 10. As dysentery Boyd-10 cultures are isolated and described extremely rarely (we did not find any reports of their isolation in the Soviet Union), and also because the cultivating peculiarities of these strains render their serological identification quite difficult, we subjected to a detailed study 25 cultures stemming from ten patients. Study was made of the morphological, biochemical properties of the cultures, of their biological characteristics, and a serological analysis was also performed.

It was established as a result of this work that the enzymatic and serological characteristics of the cultures were identical to the standard strain Boyd 10. At the same time, their cultural and morphological peculiarities were identified (the articulation of colonies on agar media and the ability of cultures to non-specifically clot in certain immune sera). Also established were their abilities to produce keratoconjunctivitis in guinea pigs, to reproduce in thin white muscle with intranasal infection and to cause

cell degeneration when multiplying in tissue cultures.

In specially conducted experimental research, prolonged (up to 200 days) viability and virulence of the Boyd-10 cultures in water kept at 4° were established.

Received by Editors
1 December 1966

- END -

1471
CSO: 1870-N