


Distribution of this document is unlimited

AD 709477

NAMRU-2

FASCILOPSIASIS IN SOUTHEAST ASIA
AND THE FAR EAST : A REVIEW



DDC
RECEIVED
AUG 5 1970
B

NAMRU-2-TR-395
(MRC05.20.01-0098.80)
1969

United States Naval
Medical Research Unit No. Two
Taipei, Taiwan

**Best
Available
Copy**

MEDICAL ECOLOGY DEPARTMENT

John H. Cross, Jr., Ph.D., Head

ADMINISTRATIVE INFORMATION

This study was accomplished under the Bureau of Medicine and Surgery Work Unit MR005.20-0098. The study was supported through funds provided by the Bureau of Medicine and Surgery, and by funding under Public Law 480, Section 104(c).

Distribution of this document is unlimited

R. H. WATTEN
CAPT MC USN
Commanding Officer

ADL	SECTION	☒
OFST	DIFF SECTION	☐
BBC		☐
UNA		☐
JA		☐
		☐
		☐
D. P. B.	AVAILABILITY CODES	
DISC	AVAIL. OF SPECIAL	
1	20	

**REPRINTS FROM PROCEEDINGS
OF
THE FOURTH SOUTHEAST ASIAN SEMINAR
ON
PARASITOLOGY AND TROPICAL MEDICINE,
SCHISTOSOMIASIS AND OTHER
SNAIL-TRANSMITTED HELMINTHIASIS**

FASCIOLOPSIASIS IN SOUTHEAST ASIA AND THE FAR EAST: A REVIEW†

JOHN H. CROSS

U. S. Naval Medical Research Unit No. 2
Taipei, Taiwan.

In his classic paper, "This Wormy World", Stoll (1947) estimated 10 million human infections with the giant intestinal fluke *Fasciolopsis buski* in Asia. Nearly half the cases, however, were ascribed to Chekiang Province of mainland China. The remaining, according to Stoll, were elsewhere in China, Taiwan, Southeast Asia, and Assam. At the present time it is almost impossible to establish the present distribution of this parasite on mainland China. It is equally difficult to estimate the infection rate and distribution of the parasite in other areas of the Far East since there exist only a few scattered reports of the parasite and the disease in the current literature. An attempt has been made to assemble the available information and to present the distribution of this parasitosis by various countries. The main sources of reference were the Tropical Disease Bulletin and the Index Catalog of Medical and Veterinary Zoology. Although a number of references, mostly old, were found, the availability of published papers/reports has been less than one would desire in preparing a review article. The reference section contains a great number of articles which have not been cited in the text. They are listed simply to assist future investigators who may wish to have a bibliography of some of the past works. This bibliography should not be considered complete, however.

MAINLAND CHINA

Fasciolopsis buski has been reported from ten provinces of mainland China. In certain areas of Chekiang and Kiangsi the prevalence rate has been reported as high as 85 percent and in other areas it may vary from less than 1 to 5 percent (Hsü and Li, 1953). Man and pigs, and possibly dogs (Chen, 1934), are reported to be the main definitive hosts for the parasite, and the aquatic snails, *Segmentina hemisphaerula* and *Hippeutis cantori*, the intermediate hosts. Metacercariae are found encysted on aquatic plants such as water caltrop (*Trapa natans*), water chestnut (*Eliocharis tuberosa*), water hyacinth (*Eichornia crassipes*), and water bamboo (*Zizania aquatica*, *Salvinia natans*, *Lemna polyrhiza*). These plants are considered the main sources of human infections. Most water-bodies in which these plants are cultivated are fertilized with human excrement. In addition, there are other plants upon which the metacercariae may encyst and these plants may serve as vehicles of transmission to pigs and other animals. Infections in mainland China, as elsewhere, are acquired most commonly by people who eat water plants raw and peel off the outer layers of the plants with their teeth. Recent reports from mainland China indicate that *Areca* nut may be of therapeutic value in the treatment of fasciolopsiasis (Fang, 1955; Teng, *et al.* 1959; Liao, 1959).

† This study was supported in part through funds provided by the Bureau of Medicine and Surgery, Navy Department, for Work Unit MR 005.20-0098.

The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Navy Department or the Naval Service at large.

REPUBLIC OF CHINA - TAIWAN

Human infection with *F. buski* on Taiwan was first recognized by Maxwell in 1911. This report was followed by the first description of the life cycle of the parasite in Taiwan pigs by Nakagawa (1921) and additional reports of human infections by Oi (1924), Suzuki (1925), Huang (1947), and Hsü *et al.* (1953). It was not until the report by Hsieh (1959), however, that fasciolopsiasis was considered a significant parasite of man in certain areas of the island.

Pigs from many areas of Taiwan appear to be infected (Lee, 1957) but the prevalence of human infections, except for a few areas, is minimal. The most highly endemic area is in Taiwan Hsieh in Southern Taiwan where stool surveys have shown infection rates of 28 percent (Hsieh, 1959), 48 percent (Hsieh, 1960), and 24 percent (Kuntz *et al.* 1961).

Hsieh (1960) carried out extensive studies in this area of Taiwan and found the highest infection rate to occur in children in the age group of 10 to 14 who resided in villages situated near ponds where water caltrop and other edible water plants are cultivated. Families in villages closer to ponds demonstrated even higher infections than those living some distance from them. Although seven species of snails were recovered from ponds in the area, only *S. haemisphaerula* was found positive (1.9%). Two edible water plants, *Trapa bispinosa* (caltrop) and *Ipomoea reptans*, (Kung-shin tsai), were also grown in the water and encysted metacercariae were found on these plants in varying numbers. The latter is commonly fed to pigs and is considered the main source of pig infections. Pigs in the area were also studied and the infection rate by stool examination was determined to be 29 percent and at the time of slaughtering, 52 percent. Of interest in the reports by Hsieh (1960) and Kuntz *et al.* (1961) was the finding of great numbers of cercariae-producing snails in ponds in the vicinity of slaughter houses and the observation that the abattoir drains emptied into nearby ponds. This, of course, leads to a continuous "seeding" of the ponds.

A few studies on the treatment of *F. buski* have been carried out in Taiwan. Brown *et al.* (1959) found tetrachlorethylene to be effective in the treatment of the disease, and Hsieh *et al.* (1953) reported a cure-rate of 62 percent with stilbazium iodide (Monopar).

JAPAN

Kobayashi (1915) was one of the first to report the parasite in Japan and later there were a few additional reports of *F. buski* eggs in stools of Japanese (Morishita *et al.* 1964). Whether or not these infections were acquired in Japan is not known.

KOREA

Although Korea has not been considered endemic for *F. buski*, there are reports in the literature which suggest the possible presence of the parasite in this country. Kobayashi (1920) reported abnormal-appearing eggs of the parasite in human stools. Many years later Brooke *et al.* (1956) reported eggs of either *F. buski* or *Fasciola hepatica* in six North Korean

prisoners of war. In another report in the same year, Maschke (1956) reported *F. buski* in a child who had recently immigrated from Korea to Germany. The most interesting report, however, was by Kuntz *et al.* (1958) who found *F. buski* eggs in the feces of a retired Turkish soldier who had served with the United Nation Forces in Korea during the Korean War. In the light of these reports it seems appropriate to suggest that further studies be undertaken to determine whether *F. buski* is truly endemic in Korea. It is entirely possible that the parasite might be present in the human and pig population in isolated areas and has gone unnoticed. On the other hand, it may also be possible that the eggs found in the feces in the above reports were of *F. hepatica* rather than *F. buski* since it is often very difficult to differentiate between the eggs of these two species of trematodes.

RYUKYU ISLANDS

Reports of *F. buski* could not be found in any scientific journal; however, a statement appears in a textbook (Simmons *et al.* 1944) on the occurrence of the parasite in some of the inhabitants of these islands.

PHILIPPINE ISLANDS

While there have been reports of fasciolopsiasis in the Philippines there is some question whether the infections were acquired there or imported. Garrison (1910) reported *F. buski* collections from the Philippines, and Schwartz (1924) found eggs in the stools of several Chinese in Sulu (Tubangui, 1933). It was not known whether these latter infections were acquired locally or in China. Africa *et al.* (1940) did not believe that the infection was endemic to their country and reported no authentic records of infection among native Filipinos. In recent studies carried out jointly by NAMRU-2 and the Philippine Health Department on nearly 4000 residents in Northern Luzon, no infections of *F. buski* were found. It is probable that any infections reported in the past in the Philippines were from immigrants from China, and the Philippines, therefore, should probably be considered a nonendemic area.

VIETNAM

There are a few reports on *F. buski* in Vietnam, the earliest of which was by Brau and Bruyant (1913). In North Vietnam, Galliard (1948) reported infections in pigs, and Bauge (1954) found nearly 15 percent and 3 percent, respectively, of Asians and Europeans infected with the parasite. Landmann *et al.* (1961) using *F. buski* skin-test antigen reported a very low prevalence of infection, however, Fournier (1954) reported the parasite in Asians as well as Europeans in Saigon. In studies by NAMRU-2 in DaNang, South Vietnam, only one of several hundred people examined was found to be passing eggs of the parasite in his feces. Snails of the genera *Planorbis* as well as *Segmentina* have been reported as the intermediate host in North Vietnam, and more than likely they are also the intermediate host in South Vietnam. Buffle and canines have also been listed as possible definitive host in this country (Segal *et al.* 1968).

LAOS

No information was found in a search of the literature regarding the prevalence of *F. buski* in Laos. It has been stated, however, in a health data bulletin distributed to U. S. military personnel and by Segal *et al.* (1968), that the parasite is present in humans of this country.

CAMBODIA

Cambodia is another country where very little information is available on fasciolopsiasis. The only paper seen was that of Brumpt and Kong-kim-chuon (1965) in which the parasite was reported in 0.04 percent of the human and 5 percent of the pig populations in the region of Phnom-Penh.

THAILAND

It has been only in recent times that human fasciolopsiasis was found to be endemic in Thailand. Daensvang and Mangalasmaya (1941) reported the first infections in five people from Dhonburi Province, in Central Thailand. In 1951 an epidemiologic investigation of the disease was initiated by Sadun and Maiphoom (1953) following the death of a 15-year-old girl infected with over 500 adult worms. The study was carried out in three provinces of Central Thailand in areas of water caltrop (*Trapa bicornis*) cultivation. Over 1500 people were examined and 13 percent were found to be infected. The prevalence of infection was highest in children 5 to 14 years old with the intensity of infection, determined by egg counts, to be greatest among children between 10 and 14 years of age. The authors found that children often ate raw water caltrops during their walk to and from school. The people in these areas for the greater part of the year live more or less in floating villages where the dwellings are built on stilts. Lacking toilet facilities, people commonly defecate into the water surrounding the houses. Although the people did not defecate directly into the water caltrop beds, the same waters surrounding the dwellings also flowed into the cultivated areas. Snails in the area (*Lymnaea pignis (sulvaria)*, *Lymnaea plicatula*, and *Caridia tenuicostata*), however, were found to be negative for cercariae of *F. buski*.

In another aspect of this study, Sadun and Maiphoom (1953) examined people in Bangkok and Ayuthya who commonly eat water caltrop purchased in local markets. *Fasciolopsis buski* eggs were not found in any of those examined. This information was presented by the authors to further show that the greatest numbers of infections take place in localities where water caltrops are cultivated and that the consumption of the plant some distances from the caltrop plantations is of little epidemiological importance. In the studies of Saovakontha *et al.* (1965) and Manning (1969) other endemic areas of fasciolopsiasis in Thailand were described.

INDONESIA

Although statements are made regarding the occurrence of *F. buski* in Indonesia there are very few actual reports on the parasite in this country. Hegner *et al.* (1923), Hsü and Li

(1953), and Faust and Russell (1964) refer to the parasite as being found in man and pigs in Borneo and Sumatra. The only reference in the more recent literature, however, is that by Ressang *et al.* (1959) who mentioned the parasite in pigs.

MALAYSIA

Sandosham (1955) reported *F. buski* to be rare in man and pigs in Malaysia but in a later publication he (Sandosham and bin Keing, 1967) suggested that reports of the eggs of the parasite in human feces in Malaysia were probably due to misidentification. Lie (1964), on the other hand, reported the recovery of adult worms from two Chinese. In the first case worms were recovered from an adult in Kuala Lumpur following treatment with carbon tetrachloride. The second case was a 13-year-old Singapore girl who died of other causes and worms were recovered at autopsy. It was believed by Lie that the adult acquired the infection in China. In the second case, however, the 13-year-old girl was born in Singapore and had never left the country. Lie suggested that the infection of the latter may have resulted from eating water chestnuts imported from China. This seems unlikely since this would have required the water chestnuts to be maintained in a wet or moist condition for an extended period of time in order to enable the encysted metacercariae to remain viable. Sadun and Maiphoom (*vide supra*) showed that even at distances of from 10 to 20 miles from endemic areas in Thailand no infections could be found in people who commonly ate water caltrop purchased from street peddlers or markets. Lie (1964) also examined pigs in Kuala Lumpur and failed to find the parasite.

BURMA

The only reports of *F. buski* that could be found in a literature search regarding Burma referred only to the presence of the trematode in pigs (Bhalerao, 1924; Bhattacharjee, 1937; Chatterjee, 1938; Griffiths, 1957). It seems likely that human infections would also be present but references to such infections could not be uncovered.

EAST PAKISTAN


Two reports show fasciolopsiasis to be endemic in East Pakistan. Kuntz (1960) reported one human case and Muazzan and Ali (1961) five additional cases. No other reports of the disease or the parasite were found.

INDIA

The prevalence of *F. buski* in pigs and man in India seems to be variable. While Buckley (1939) found nearly 60 percent of the people examined in Assam infected with the parasite, Sataya Prakash and Varmani (1967) failed to find a single human infection in the Ghaziabad area of Uttar Pradesh. These latter authors, however, reported 1.4 percent of the pigs and 24

percent of *Planorbis* sp. infected with the parasite. In other studies in Bihar, Kant and Rama (1954) found 7 percent of the population infected and, following treatment with tetrachlorethylene, recovered from 1-57 adult worms from the patients. Other reports indicate the parasite to be in pigs and man in Calcutta (DeRivas and Lucke, 1915; Thapar 1956). According to a recent publication, Shah *et al.* (1966) found 29 percent of a population in Bombay infected with the parasite and they suggest that infections may also be present in parts of Bihar, Orissa, and Madras.

Additional studies by Buckley (1939) in Assam have shown *Segmentina trochoideus* to be an intermediate host for the parasite. Although Buckley was unable to determine the principal sources of infection, he listed the following water plants as possibilities: *Ottelia alismoides*, *Nymphaea lotus*, *Trapa natans*, *Euryale ferox*, and *Nehumbo nucifera*. He noted that pigs were rare in the area because of religious prejudices against eating pork, but examination of 30 pigs in a slaughter house failed to reveal the parasite. Similarly, Shah *et al.* (1966) reported the absence of the parasite from animals in Bombay. This is not unusual, however, as Cameron (1927) stated that in certain areas where *F. buski* is common in man the parasite may be uncommon in pigs and vice versa.



Distribution of *Fasciolopsis buski* in the Far East.

REPORTS OF FASCILOPSIASIS OUTSIDE THE FAR EAST

Unusual reports of human infection with *F. buski* have been made from several nonendemic areas of the world. For example, there are references to fasciolopsiasis in the United States (Moore and Terrill, 1905), South Africa (Nicol, 1910), Venezuela (Risquez, 1911), Australia (Nicoll, 1914), Russia (Seminov and Garmatti, 1927), Melanesia (Russell and Scott, 1945), and Guatemala (Penalver *et al.* 1955). It is probable that reports of *F. buski* outside the Far East are due either to misidentification of the egg or to instances where the parasite was imported with emigrants from the Far East.

COMMENT

In searching the literature and in preparing this review the author was impressed with the paucity of information available regarding *F. buski*. There appears to be little interest in this trematode today. Except in drug studies or an occasional report of finding the eggs in faces, reports of experimental studies are meager. Only two papers of recent vintage concerning experimental investigations (Kuntz and Lo, 1967; Lo, 1967) were found. This is incongruous as the parasite should be of great value to the experimentalist. The worm is large, is easily obtainable from pigs in endemic areas, and would offer those interested in biochemistry, physiology, and immunology of helminths a ready source of material.

SUMMARY

In 1947 it was estimated that over 10 million people in the Far East were infected with *Fasciolopsis buski*. Most of these infections were reported from mainland China. Although fasciolopsiasis is endemic in Taiwan, Thailand, Vietnam, Laos, East Pakistan and India, the distribution of infections in these countries is limited and the overall prevalence rates are considerably lower than those of mainland China.

While human fasciolopsiasis has not been reported from Indonesia and Burma, pigs in these countries have been found infected with the parasite. Further studies should be carried out in these countries to determine whether or not the parasite is present in man. Studies are recommended in Cambodia where little information is available and in Korea in order to determine whether or not the parasite is endemic.

Human infections have been reported from Japan, the Philippines, Malaysia, and a number of western countries, but it is probable these were from people who immigrated from endemic areas or were due to misidentification of the egg.

Fasciolopsiasis seems to be restricted to areas where populations raise water plants, such as water caltrop, water chestnuts, water hyacinth and water bamboo; and to populations that commonly ingest the uncooked metacercariae-laden plants. In addition, in most but not all the endemic countries, pigs appear to be an important reservoir of infection and species of snails of the genera *Segmentina*, *Hippeutis*, and *Planorbis*, the intermediate hosts.

Investigations show infections are found most commonly in individuals and families that live in the immediate vicinity of water culture plantations and that prevalence rates are highest in children.

REFERENCES

- Abbas, C. M., de Leon, W. and Garcia, E. Y : Visceral complications in intestinal fasciolopsiasis of man. *Acta Med. Philipp. Monogr. Ser.*, 1: 132, 1948.
- Abbas, S. A. : A survey of the helminth parasites of the domestic pigs in Western U.P. with a note on *Parasitocryptus ovalis* Baylis. *Indian J. Vet. Sci.*, 28: 180, 1939.
- Barlow, C. H. : Difference between specimens of *Fasciolopsis buski* collected from a 9-year-old child and those from a man. *J. Parasit.*, 7: 199, 1921.
- Barlow, C. H. : Experimental ingestion of the ova of *Fasciolopsis buski*; also the ingestion of adult *Fasciolopsis buski* for the purpose of artificial infestation. *J. Parasit.*, 8: 40, 1921.
- Barlow, C. H. : Life cycle of *Fasciolopsis buski* (Lankester) in China. *Chin. Med. J.*, 37: 453, 1923.
- Barlow, C. H. : The life cycle of the human intestinal fluke, *Fasciolopsis buski* (Lankester). *Amer. J. Hyg., Monogr. Ser.*, 4, 96, 1925.
- Barlow, C. H. : *Fasciolopsis buski* infestation in the region of Shaohsing. *J. Parasit.*, 15: 290, 1929.
- Barlow, C. H. : The prevalence of fasciolopsiasis in the Hangchow region. *J. Parasit.*, 15: 285, 1929.
- Barrois, T. C., Née, F. : Sur la fréquence de *Fasciolopsis buski* (Lank, 1957) en Cochinchine. *Bull. Soc. Path. Exot.*, 1: 216, 1968.
- Buop, R. : Incidence of intestinal parasites in Hongay region, North Vietnam in February-May 1953. *Bull. Soc. Path. Exot.*, 47: 720, 1954.
- Bhalerao, G. : A contribution to the knowledge of the trematode parasites of the food animals of Rangoon. *Ann. Trop. Med. Parasit.*, 18: 139, 1924.
- Bhalerao, G. : Trematode parasites of pigs in Bengal. *Rec. Indian Mus.*, 33: 475, 1931.
- Bhattacharjee, J. : A checklist of the trematode and cestode parasites of the domesticated animals of Burma. *Indian Vet. J.*, 14: 1, 1937.
- Brau, P. and Bruyant, L. A. : Que notes sur les helminthes du porc en Cochinchine. *Bull. Soc. Path. Exot.*, 15: 834, 1913.
- Brooke, M. M., Swartzweider, C., Payne, F. J., Weinstein, P. and Frye, W. W. : Intestinal parasite survey of Korean prisoner-of-war camp. Extract from *U. S. Armed Forces Medical J.*, 7: 708, 1956.

- Brown, H. W., Cheng, C. S., Chen, C. Y., Lin, K. H., Chang, C. R., Lee, L. T., Hsü, C. C., Chow, L. P. and Hsieh, H. C. : The treatment of *Fasciolopsis buski* infections with dichlorzinc iodide, 1-bromonaphthol (2), ascaridol, piperazine, and tetrachlorethylene. *J. Form. Med. Ass.*, 38 : 792, 1959.
- Brown, N. W. : The Fasciolopsinae of China. A study of two species from Chekiang Province. *John's Hopkins Hosp. Bull.*, 28 : 322, 1917.
- Braun, V. and Kong-Kim-Choon : Les helminthoses intestinales chez L'homme au cambodge. *Bull. Soc. Path. Exot.*, 58 : 501, 1965.
- Bocley, J. J. C. : Observations on *Gastrodiscoides hominis* and *Fasciolopsis buski* in Assam. *J. Helminth.*, 17 : 1, 1939.
- Cadbury, W. W. : The research report from the South China Branch. *China Med. J.*, 26 : 12, 1912.
- Cameron, T. W. M. : The pig and human disease. *Proc. Roy. Soc. Med.*, 17 : 31, 1924.
- Cameron, T. W. M. : The helminth parasites of animals and human disease. *Proc. Roy. Soc. Med.*, 20 : 15, 1927.
- Chandler, A. C. : A new schistosome infection of man, with notes on other human fluke infections in India. *Indian J. Med. Res.*, 14 : 179, 1926.
- Chandler, A. C. : The prevalence and epidemiology of hookworm and other helminthic infections in India. Part XII. General summary and conclusions. *Indian J. Med. Res.*, 15 : 695, 1928.
- Chang, K. : Researches in parasitology in Free China. A brief review. *Acta Brevia Sinensia*, 4 : 1, 1943.
- Chang, H. P., Yeh, K. T., Sun-f'u, T. C., Mo, W. C. and Huang, S. H. : The endemicity of Fasciolopsiasis (*F. buski*) and a clinical report on 65 cases in children. *Chinese J. Pediatrics*, 7 : 355, 1956.
- Chatterjee, R. C. : Annotated list of the helminths recorded from domesticated animals of Burma. Part I. Trematoda. *Proc. Nat. Acad. Sc. India*, 8 : 93, 1938.
- Chen, H. T. : Helminths of dogs in Canton with a list of those occurring in China. *Lingnan Sc. J.*, 13 : 75, 1934.
- Chen, T. : Parasites in slaughter houses in Canton Part I. Helminths of Kwangtung hogs. *Lingnan Sc. J.*, 15 : 31, 1936.
- Chen, W. L. and Poo, L. C. : Preliminary report on the prevention and treatment of *Fasciolopsis buski* in Chekiang Province. *Nat. Med. J. China*, 23 : 1105, 1937.

- Chen, H. H. and Wu, P. H. : Incidence of intestinal helminthiasis in the Wu Chia-ling area and vicinity, Changsha, Hunan, with special reference to *Fasciolopsis buski*. *Chinese Med. J.*, 68 : 173, 1950.
- Chien, C. : Untersuchungen über *Fasciolopsis buski*. *Jgaku Kenkyu*, 14 : 97, 1940.
- Chiu, J. K. and Kau, C. T. : The incidence of intestinal parasites among the elementary school children in Hualien Prefecture, Taiwan., *J. Form. Med. Ass.*, 58 : 101, 1959.
- Cau, S. H. : Mass treatment of *Fasciolopsis buski* infestation among school children in Canton. *Chinese Med. J.*, 78 : 273, 1959.
- Chu, S. H., Chou, C. C., and Liu, T. C. : The treatment of severe fasciolopsiasis. *Chinese J. Pediatrics*, 9 : 335, 1958.
- Chung, H. L. : The efficacy of hexachloroparoxylol and hexachlorophene in the treatment of two human cases of fasciolopsiasis *buski*. *China Med. J. (Peking)*, 84 : 533, 1956.
- Cobbold, T. S. : On some new forms of entozoa. *Tr. Linn Soc. London*, 22 : 363, 1859.
- Cobbold, T. S. : Synopsis of the distomidae. *J. Proc. Linn. Soc. London Zool.*, 5 : 1, 1860.
- Cole, A. : Five cases of *Fasciolopsis* infection, with remarks. *China Med. J.*, 36 : 391, 1921.
- Daengsvang, S. and Mangalasmaya, M. : Record of some cases of human infection with *Fasciolopsis buski* occurring in Thailand. *Ann. Trop. Med. Parasit.*, 35 : 43, 1941.
- De Rivas, D. and Lucke, B. : Eggs of hookworm and *Fasciolopsis buski*. *Med. Rec. N.Y.*, 87 : 93, 1915.
- Fan, P. C. : Prevalence of parasitic disease in Taiwan. *J. Chinese Nursing*, 12 : 50, 1965.
- Fa, N. T. : *Fasciolopsis buski* in Shantung. *Chinese Int. Med.*, 3 : 651, 1955.
- Faust, E. C. : The distribution and differentiation of human helminths in China and adjacent territory. *China Med. J.*, 37 : 1, 1923.
- Faust, E. C. : Notes on larval flukes from China. II Studies on some larval flukes from the central and south coast provinces of China. *Amer. J. Hyg.*, 4 : 1, 1924.
- Faust, E. C. : Some facts regarding the relationship between nightsoil disposal in China and the propagation of helminthic diseases. *Amer. J. Trop. Med.*, 4 : 487, 1924.
- Faust, E. C. : Parasitic infections and human disease in China. *Arch. Path. Lab. Med.* 2 : 223, 1926.

- Faust, E. C. and Kellogg, C. R. : Parasitic infections in the Foochow area, Fukien Province, China., *J. Parasit.*, 14 : 196, 1928.
- Faust, E. C. and Wassell, C. A. : Preliminary survey of the intestinal parasites of man in the central Yangtze Valley. *China Med. J.*, 35 : 532, 1921.
- Fournier, J. : Services partiques. *Rap. Inst. Pasteur Saigon 1953*. pp. 109, 1954.
- Fukuda, K. : *Fasciolopsis buski* (Lank). *Distoma crassum* (Cobb.). *Mitt. Med. Gesellsch. Tokyo*, 21 : 53, 1907.
- Gaehlinger H. : Un cas de pluriparasitisme avec distomatose intestinale (*Fasciolopsis buski*). *Marseille Med.*, 61 : 525, 1924.
- Galliard, H. : Human infection by *Fasciolopsis buski* in Tong King, Vietnam. *Bull. Soc. Path. Exot.*, 41 : 214, 1948.
- Galliard, H. and Dang-van-Ngu : Quelques trematodes parasites des animaux domestiques au Tonkin. *Rev. Med. Franc. Extreme-Orient*, 18 : 131, 1940.
- Galliard, H. and Dang-van-Ngu : L'infestation de l'homme et du chien par *Fasciolopsis buski* au Tonkin. *Rev. Med. Franc. Extreme-Orient*. 19 : 31, 1941.
- Galliard H. and Dang-van-Ngu : Recherches sur la specificite parasitaire de *Fasciolopsis buski*. *Ann. Parasit.*, 22 : 16, 1947.
- Garrison, P. E. and Lago, F. P. : Specimens added to the helminthological collection U. S. Naval Medical School, June and August. *U. S. Naval Med. Bull.*, 4 : 511, 1910.
- Germain, L. and Neveu-Lemaire, M. Essai de malacologie medicale. *Ann. Parasit.*, 4 : 286, 1926.
- Goddard, F. W. : Two rare Fasciolidae. *China Med. J.*, 21 : 195, 1907.
- Goddard, F. W. : Further notes on the Shaohsing fluke. *China Med. J.*, 23 : 387, 1909.
- Goddard, F. W. : *Fasciolopsis buski* a parasite of man as seen in Shaohsing, China. *J. Parasit.*, 5 : 141, 1919.
- Griffiths, R. B. : Report to the government of Burma on parasitic diseases of livestock. *Expand. Techn. Assist. Program*, 1, 1957.
- Heanley, C. M. : A large fluke of man probably not hitherto described. *Fasciolopsis buski* as a parasite of man in Hong Kong ; its usual host the pig. *J. Trop. Med.*, 11 : 122, 1908.
- Heanley, C. M. : Report of the research work completed, in progress, and in contemplation at the Bacteriological Institute and Public Mortuary, Hong Kong, October 1907 to February 1908. *Rep. Advis. Com. Trop. Dis. Research Fund (1908)*, 65, 1090.

- Hedblom, C. A. : On disease incidence in China. *China Med. J.*, 31 : 271, 1917.
- Hegner, R. W., Cort, W. W. and Root, F. M. : Outline of Medical Zoology with special reference to laboratory and field diagnosis, N. Y., N. Y., 175, 1923.
- Hoeppli, R. J. C. and Hou, H. C. : Reaction of tissue against parasites and their eggs as observed in the transparent chamber of the rabbit's ear. *Chinese J. Physiol.*, 5 : 19, 1931.
- Hou, T. C., Li, W. Y., Chung, H. L. and Li, Y. F. : Differential significance of intradermal tests with serial high dilutions of trematode antigens on various trematode diseases : Observations on 80 cases of paragonimiasis. *Jen Min Pao Chien*, 1 : 505, 1959.
- Houdemer, F. E. and Tran-tho-Huy : Notes sur quelques parasites observes chez les animaux de boucherie à Hanoi, *Bull. Med. Vet. Indochin.* 1 : 145, 1925.
- Hsieh, H. C. : Outline of parasitic zoonoses in Taiwan. *Form. Sci.*, 13 : 99, 1959.
- Hsieh, H. C. : Fasciolopsiasis in Taiwan. *J. Form. Med. Ass.*, 58 : 623, 1959.
- Hsieh, H. C. : Studies on the epidemiology of *Fasciolopsis buski* in South Taiwan. *Form. Sci.*, 14 : 95, 1960.
- Hsieh, H. C., Brown, H. W., Chen E. R., Chen, C. Y. and Shih, C. C. : Treatment of *Fasciolopsis buski*, *Ancylostoma duodenale*, *Ascaris lumbricoides*, *Trichuris trichiura*, and *Enterobius vermicularis* with stilbazium iodide. *J. Parasit.*, 49 : 425, 1963.
- Hsü, H. F. and Chow, C. Y. : Studies on human intestinal helminths in 809 autopsy cases. *Bull. Fan Memorial Inst. Biol.*, 8 : 245, 1938.
- Hsü, H. F. and Li, S. Y. : Studies on the food and the digestive system of certain parasites. VI. On the food of certain helminths living in the digestive tract of vertebrates. *Chinese Med. J.*, 57 : 559, 1940.
- Hsü, H. F. and Li, S. Y. : Helminths of dogs in Peiping. *Peking Nat. Hist. Bull.*, 15 : 201, 1941.
- Hsü, H. F. and Li, S. Y. : Notes on *Fasciolopsis buski* in China. Tharpar Commemoration Volume, 1953. Department of Zoology, National Taiwan University, Taipei, Taiwan, China, 1953.
- Hsü, P. J. : A survey of fasciolopsiasis of swine in Kwangtung Province. *Chu Mu Shou I Hsueh Pao.*, 7 : 143, 1964.
- Huang, T. Y. : Investigation on the human parasites among the inhabitants in Wantan District, Kaohsiung Prefecture, South Formosa. *J. Form. Med. Ass.*, 46 : 119, 1947.
- Huang, W. H. and Chiu, J. K. : Resurvey of the incidence of intestinal helminths among school children in a rural area in Northern Taiwan. *J. Form. Med. Ass.*, 60 : 216, 1961.

- Hung, S. L. : Über die morphologie von *Fasciolopsis buski* beim menschen in Hsiao-Shan. *Arch. Schiffs- u. Tropen-Hyg.*, 37 : 315, 1933.
- Ishii, S. : Human and dog cases of *Fasciolopsis buski* in Canton, China. *J. Form. Med. Ass.*, 288 : 242, 1929.
- Ishii, Y. : Studies on the development of *Fasciolopsis buski*. Part 1. The development of normal eggs outside the host. *J. Form. Med. Ass.*, 33 : 349, 1934.
- Ishii, Y. : Studies on the development of *Fasciolopsis buski*. Part 2. Morphology and biology of the miracidium. *J. Form. Med. Ass.*, 33 : 379, 1934.
- Ishii, Y. : Studies on the development of *Fasciolopsis buski*. Part 3. Development in the intermediate host of *Fasciolopsis buski*. *J. Form. Med. Ass.*, 33 : 391, 1934.
- Ishii, Y. : Experimental infections of *Fasciolopsis buski* in man and its egg production. *J. Form. Med. Ass.*, 33 : 137, 1934.
- Isshiki, O. : Investigations on internal parasites of domestic animals from the Mongolia steppe countries and North China (Preliminary report II). *J. Chosen Nat. Hist. Soc.*, 26 : 69, 1939.
- Joint WHO/FAO Expert Committee on Zoonoses Wld. Hlth. Org. Techn. Rep. Ser. No. 169 : 5, 1959.
- Joyeux, C. E. and Houdemer, F. E. : Recherches sur la faune helminthologique de l'Indochine (cestodes et trematodes). *Ann. Parasit.*, 5 : 289, 1927.
- Joyeux, C. E. and Houdemer, F. E. : Researches sur la faune helminthologique de l'Indochine (cestodes et trematodes). *Ann. Parasit.*, 6 : 27, 1928.
- Juzuki, S. : On an inquiry into the conditions of the spread of *Fasciolopsis buski*. *Taiwan. J. Gakkwai Zasshi, Taihoku*, 274. May 1920.
- Kamisaka, T. : How to differentiate between the eggs of *Fasciola hepatica* and *Fasciolopsis buski*. *J. Form. Med. Ass.*, 38 : 1250, 1930.
- Kant, L. and Rama, K. : A field survey of *Fasciolopsis buski* in Chandrain area of Saharsa District (Bihar). *Indian Med. Gaz.*, 89 : 89, 1954.
- Kau, L. S. and Wu, K. : Pathological findings among pigs experimentally infected with *Fasciolopsis buski*. *Ann. Trop. Med. Parasit.*, 32 : 133, 1938.
- Khaw, O. K. : Historical notes on *Fasciolopsis buski* (Lankester, 1857) Odhner, 1902. *Nat. Med. J. China*, 17 : 475, 1931.
- Kobayashi, Harujiro : On some digenetic trematodes in Japan. *Dobuts Zasshi, Tokyo*, 27 : 109, 1915.

- Kobayashi, Harujiro : On intestinal parasites occurring in Choeans, on the abnormal forms of the eggs. *Nippon No Ikai, Tokyo*, 10 : 1015, 1920.
- Kobayashi, Harujiro : On some digenetic trematodes in Japan. *Parasit.*, 12 : 380, 1921.
- Kobayashi, Hidekazu : Studies on the structure of the reproductive organs of the trematoda (report 1) on the male genital organs of *Fasciolopsis buski* and *Fasciola hepatica*. *J. Form. Med. Ass.*, 308, 1229, 1930.
- Krijgsman, B. J. : De in zoogdieren en vogels parasiteerende wormen van Nederlandsch-Indie. *Nederl.-Ind. Blad. Diergeneesk.*, 45 : 366, 1933.
- Kuntz, R. E. : Intestinal protozoa and helminths in school children of Dacca, East Pakistan (East Bengal). *Amer. J. Trop. Med. Hyg.*, 9 : 168, 1960.
- Kuntz, R. E., Burke, J. C., Lin, S. and Watten, R. H. : Protozoa and helminth parasites in peoples of Taiwan (Formosan). *J. Form. Med. Ass.*, 60 : 809, 1961.
- Kuntz, R. E., Lawless, D. K. and Langbehn H. R. : Intestinal protozoa and helminths in peoples of Western (Anatolia) Turkey. *Amer. J. Med. Hyg.*, 7 : 298, 1958.
- Kuntz, R. E. and Li, C. T. : Preliminary studies on *Fasciolopsis buski* (Lankester, 1857) (Giant Asian Intestinal fluke) in the United States. *Trans. Am. Microsc. Soc.*, 86 : 163, 1967.
- Lagrange, E. : Observations sur les trematodes l'Indochine. *Bull. Soc. Path. Exot.*, 16 : 173.
- Lal, S. M. : A case of *Fasciolopsis buski* infection. *Indian Med. Gaz.*, 58 : 586, 1923.
- Landmann, H., Thai, D. D. and Than, D. T. : Diagnosis of paragonimiasis by the intradermal reaction with the antigens of *Fasciolopsis buski* and *Fasciola hepatica*. *Zent. f. Bakt. I Abt. Orig.*, 182 : 417, 1961.
- Lane, C. : Intestinal animal parasites in Monghyr. *Indian Med. Gaz.*, 44 : 131, 1909.
- Lane, C. : Discussion of Cole - five cases of *Fasciolopsis* infection. *Lancet*, 200 : 540, 1921.
- Lane, C. : Some little recognized features in common helminths. *Fasciolopsis buski*. *Tr. Roy. Soc. Trop. Med. Hyg.*, 16 : 21, 1922.
- Lane, C. and Low, G. C. : Trematode infestations other than haemai (In Byam. W., and Archibald, R. G. The practice of medicine in the tropics. London), 3 : 1788, 1923.
- Lankester, E. : On the occurrence of species of Distoma in the human body (Appendix B to Kuchenmeister, Gottlob Friedrich Heinrich. 433,). 1857.
- Lee, Y. C. : Investigation on the internal parasites of domestic animals in Taiwan. *J. Agric. Ass. China*, 18 : 56, 1957.

- Leiper, R. T. : On Kwan's fluke and the presence of spines in *Fasciolopsis*. *J. Trop. Hyg.*, 14: 119, 1911.
- Liang, P. H. : Two cases of *Fasciolopsis* in Tali, Yunnan. *Chinese J. Int. Med.*, 7 : 33, 1959.
- Liao, Y. C. : Fasciolopsiasis treated with the traditional *Areca catechu* preparation : clinical observations on 124 cases. *J. Chinese Traditional Med.*, 4 : 52, 1959.
- Lie, K. J. : Prevalence of intestinal helminths among patients of the General Hospital in Kuala Lumpur, Malaysia. *Trop. Geogr. Med.*, 16 : 225, 1964.
- Lo, C. T. : Life history of the snail, *Segmentina hemisphaerula* (Benson) and its experimental infection with *Fasciolopsis buski* (Lankester). *J. Parasitol.*, 53 : 735, 1967.
- McCoy, O. R. and Chu, T. C. : *Fasciolopsis buski* infection among school children in Shao-hsiung and treatment with hexylresocinol. *Chinese Med. J.*, 51 : 937, 1937.
- Manning, G. S. : (Personal communication) 1959.
- Maplestone, P. A. : Report of the hookworm research department. *Ann. Rep. Calcutta School Trop. Med.* (1932), 105, 1933.
- Maschke, R. : Darm- und leberegelbefall des menschen. *Problems Parasitol.* (Borchert) 218, 1956.
- Mathis, C. and Leger, M. : Parasitisme intestinal et hepaticque chez le indigenes de l'Indochine du nord (nematodes et trematodes). *Ann. Hyg. et Med. Colon.*, 13 : 459, 1910.
- Mathis C. and Leger, M. : *Fasciolopsis buski* au Tonkin. Son extreme rarete chez l'homme; sa frequence chez le porc. *Bull. Soc. Path. Exot.*, 4 : 200, 1911.
- Maxwell, J. L. : *Fasciolopsis buski* and related species. A criticism of professor Ward's paper in the China Medical Journal of January, 1910. *China Med. J.* 24 : 270, 1910.
- Maxwell, J. L. : A few notes from Taiwan Formosa. *Chinese Med. Sci.*, 25 : 172, 1911.
- Maxwell, J. L. : The disease of China. *J. Trop. Med. Hyg.*, 19 : 237, 1916.
- Moore, J. T. and Terrill, J. J. : *Fasciolopsis buski* (*Distomum buski*) The first reported case in the United States. Found in a patient dying of typhoid fever. *J. Am. Med. Ass.*, 45 : 1002, 1905.
- Morishita, K., Komiya, Y. and Matsubayashi, H. : *Progress of Medical Parasitology in Japan*, 1 : 41, 1964.
- Morisita, T. and Kimura, Z. : Studies on *Fasciolopsis buski* (giant fluke) in the region of Hsiao-shan. *Dozinkai Igaku Zasshi*, 15 : 514, 1941.

- Muazzan, M. G. and Ali, M. T. : Intestinal parasites in East Pakistan. *J. Pakistan Med. Ass.*, 11 : 185, 1961.
- Nakagawa, K. : The development of *Fasciolopsis buski* with special reference to the intermediate hosts and cercariae. *J. Form. Med. Ass.*, 211 : 553, 1920.
- Nakagawa, K. : On the life cycle of *Fasciolopsis buski*, Lankester. *The Kitasato Arch. Exptl. Med.*, 4 : 159, 1921.
- Nakagawa, K. : The development of *Fasciolopsis buski* (Lankester). *J. Parasit.*, 8 : 161, 1922.
- Nakagawa, K. : Studies on *Fasciolopsis buski*. *J. Form. Med. Ass.*, 258 : 241, 1925.
- Nakagawa, K. : An aspect of studies on *Paragonimus westermanii* and *Fasciolopsis buski* with reference to Karapai fluke and Rono fluke. *Tokyo Ijishinshi*, 75 : 29, 1958.
- Nakagawa, K. and Suzuki, S. : Investigations on the life history of *Fasciolopsis buski*. (Second Report). *J. Form. Med. Ass.*, 228 : 155, 1923.
- Nauck, E. G. and Yu, S. : Sur frage der verbreitung von wurm parasiten in Mittel-China. *Arch. Schiffs-u. Tropen-Hgy.*, 30 : 503, 1926.
- Neveu-Lemaire, M. : Notes de geographie medicale . Maladies propres au continent asiatique. *Geographie*, 38 : 433, 1922.
- Nicol, B. : *Fasciolopsis buski* in Natal. *J. Trop. Med. Hyg.*, 13 : 341, 1910.
- Nicoll, W. : The trematode parasites of North Queensland. *J. Parasit.*, 6 : 333, 1914.
- Odhner, T. : *Fasciolopsis buski* (Lank) (=Distomum crassum Cobb), ein bisher wenig bekannter parasit des memschen in Ostasien. *Centralbl. Bakteriol. 1. Abt. Orig.*, 31 : 573, 1902.
- Oi, T. : On the infection by *Fasciolopsis buski* (Lankester) among the Japanese and the infection route with the human body and the pig. *Tokyo Ijishinshi* (2396) : 2268-2272, *J. Form. Med. Ass.*, 1924.
- van Oije, P. : Korte bijdrage tot de kennis van het geslacht *Fasciolopsis*. *Vlaamsch. Geneesk. Tijdschr.*, 1 : 257, 1920.
- Onorato, A. R. and Stunkard, H. W. : The effect of certain enviromental factors on the development and hatching of the eggs of blood flukes. *Biol. Bull.*, 61 : 120, 1931.
- Penalver, L. M. and Aguilar, F. J., Arthes, S. F. and Aramburu, G. : Notas previas sobre trematodiasis en Guatemala . *Rev. Colegio Med. Guatemala*, 6: 101, 1955.

- von Querner, F. R. : Zur Histologie des exkretionsgefäß-systems digenetischer trematoden
I. Tiel. Ztschr. Parasitenk., Berlin, 1 : 489, 1929.
- Railliet, A. : Les helminths des animaux domestiques et de l'homme en Indochine., *Bull. Soc. Zool.*, 49 : 589, 1925.
- Railliet, A. and Henry, A. C. L. : Helminthes du porc recueillis par M. Bauche en Annam.
Bull. Soc. Path. Exot., 4 : 693, 1911.
- Kessang, A. A., Fischer, H. and Muchlis A. : The Indonesian veterinarian : his education, activities and problems. *Comm. Fac. Vet. Sci., Bogor Indonesia*, 3 : 55, 1959.
- Risquez, J. R. : Un cas de distomatose au Venezuela. *Rev. Med. et Hyg. Trop.*, 8 : 44, 1911.
- Robertson, R. C. : Division of pathological sciences. *Ann. Rep. Henry Lester Inst. Med. Res.* (1935) : 42, 1936.
- Robertson, R. C. : Division of pathological sciences. *Ann. Rep. Henry Lester Inst. Med. Res.* (1936) : 41, 1937.
- Robertson, R. C. : Division of pathological science. *Ann. Rep. Henry Lester Inst. Med. Res.* (1937-1938) : 38, 1939.
- Robertson, R. C. : Division of pathological science. *Ann. Rep. Henry Lester Inst. Med. Res.* (1939) : 31, 1940.
- Russell, H. K. and Scott, J. O. : Intestinal parasites among Melanesians. *U.S. Naval Med. Bull.*, 44 : 727, 1945.
- Sadun, E. H. and Maiphoom, C. : Studies on the epidemiology of the human intestinal fluke *Fasciolopsis buski* (Lankester) in Central Thailand. *Amer. J. Trop. Med. Hyg.*, 2 : 1070, 1953.
- Sandosham, A. A. : A check list of helminth parasites of man in Malaya with brief notes on their incidence. *Proc. Alumni. Ass. Malaya.*, 8 : 258, 1955.
- Sandosham, A. A. and bin Keling, M. N. : Animal parasites of animals which affect man in Malaysia. *Med. J. Malaya*, 22 : 16, 1967.
- Saovakontha, S., Charoenlarp, P., Radomyos, P. and Harinasuta, C. : A new endemic area of fasciolopsiasis in Thailand. *J. Med. Assoc. Thai* (Phaetaya Samakom Haeng Prathet Thai), 48 : 247, 1965.
- Satyā Prakash, G. J. N., Varmani, B. L. M. : A note on the prevalence of *Fasciolopsis buski* in Ghaziabad area (Uttar Pradesh - India). *Bull. Indian Soc. Mal. Com. Dis.*, 4 : 55, 1967.

- Schwartz, B. : The helminthological fauna of man in the Philippine Islands. *J. Parasit.*, 11 : 100, 1924.
- Schwartz, B. : Occurrence of *Fasciolopsis buski* in the Philippines. *J. Parasit.*, 11 : 93, 1924.
- Segal, D. B., Humphrey, J. M., Edwards, S. J. and Kirby, M. D. : Parasites of man and domestic animals in Vietnam, Thailand, Laos, and Cambodia. *Exper. Parasit.*, 23 : 412, 1968.
- Sekorrarithi, C. : Fasciolopsis. *Medical News, Bangkok*, 4 : 739, 1956.
- Semenov, G. M. and Garmatty, K. I. : Un premier cas de constatation en U.S.S.R. de l'invasion humaine par *Fasciolopsis buski* Lank. *Med. Mysl. Uzbekistana*, 6-7 : 49-57 : 130, 1927.
- Shah, A. *et al.* : Fasciolopsiasis in Bombay; A preliminary communication. *Ind. J. Med. Sci.*, 20 : 805, 1966.
- Shih, P. T. : The epidemiology of fasciolopsiasis in Sunwui, Kwangtung Province. *Nat. Med. J. China*, 243 : 454, 1957.
- Simmons, J. S., Whaune, T. F., Anderson, G. W. and Horack, H. M. : Global epidemiology. *A geography of disease and Sanitation* 1 : 504, J. B. Lippincott Co. Philadelphia, 1944.
- Skrjabin, K. I., Podiapolskais, V. P. and Statirova, N. A. : Contribution a la caracteristique de la fasciolopiose de l'intestine et sur son agent pathogene. *Russk. Zhurnal. Trop. Med.*, 7 : 262, 1929.
- Steen, H. B. : A case of *Fasciolopsis buski* (*Distoma buski* V. crassum and *Amphistoma hominis*). *Indian Med. Gaz.*, 42 : 376, 1907.
- Stoll, N. R. : This wormy world. *J. Parasit.* 33 : 1, 1947.
- Stoll, N. R. Cort, W. W. and Kwei, W. S. : Egg-worm correlations in cases of *Fasciolopsis buski* with additional data on the distribution of the parasite in China. *J. Parasit.*, 13 : 166, 1927.
- Sturton, S. D. : Oedema and ascites in Chinese patients. Being a dissertation for the degree of M. D. Cantab. *China Med. J.*, 45 : 584, 1931.
- Sugimoto, M. : Collection of the parasites (sic) of domestic animals by the veterinary students in our agri. expt. st. Taiwan Nojiho., 5 : 62, 1916.
- Sugimoto, M. : Catalogue of parasites of domestic animals in Formosa. *Dept. Agric. Govt. Res. Inst.* 1925., pp. 93, 1929.
- Suzuki, S. : Effects of certain physical and chemical influences upon the eggs of *Fasciolopsis buski*. *J. Form. Med. Ass.*, 225 : 539, 1922.

- Suzuki, S. : On the resistance of the encysted cercaria of *Fasciolopsis buski*. *J. Form. Med. Ass.*, 227 : 123, 1923.
- Suzuki, S. : On an inquiry into conditions of the spread of *Fasciolopsis buski*. *J. Form. Med. Ass.*, 234 : 257, 1924.
- Suzuki, S. : On human infection by *Fasciolopsis buski* in Formosa. *Tokyo Ijishinshi* (2339) : 2409, 1924.
- Suzuki, S. : Human cases of *Fasciolopsis buski* found in Formosan natives. *J. Form. Med. Ass.*, 238 : 95, 1925.
- Sweet, W. C. : *Fasciolopsis buski* infestation. *J. Am. Med. Ass.*, 76 : 1819, 1921.
- Teng, P., Li, P. H., Chin, C. M. and Fan, K. H. : Epidemiology of fasciolopsiasis among pupils of Weishan, ShanTung and the therapeutic value of *ARECA* nut as a vermifuge. *People's Health*, 1 : 807, 1959.
- Thapar, G. S. : Systematic survey of helminth parasites of domesticated animals in India. *Indian J. Vet. Sci. Animal Husb.*, 26 : 211, 1956.
- Tsang, P. Y. : General study on the *Fasciolopsis buski* in ShaoHsing Hsien, Hsieh Shan Hsien, and Huyang Hsien of Chekiang Province of China. *Nipponshokakibyō Gakkai Zasshi*, 33 : 46, 1934.
- Tubangui, M. A. : Trematode parasites of Philippine vertebrates, VI. Description of new species and classification. *Philippine J. Sci.*, 52 : 167, 1933.
- Tubangui, M. A. : *Schistosomiasis japonica* and other helminthic diseases in Philippines, *Proc. 4. Internat. Cong. Trop. Med. Malaria* (Washington, May 10-18), 2 : 1034, 1948.
- Varma, A. K. : Studies on the nature, incidence, distribution and control of nasal schistosomiasis and fasciolopsiasis in Bihar. *Indian J. Vet. Sci. Animal Husb.*, 24 : 11, 1954.
- Virenuvatti, V., Stitnimankarn, T. and Tansurat, P. : A fatal case of *Fasciolopsis buski* in Thailand. *Ann. Trop. Med. Parasit.*, 47 : 132, 1953.
- Vogel, H. : Beobachtungen über *Fasciolopsis*-infektion. *Arch. Schiffs-u. Tropen-Hyg.*, 40 : 181, 1936.
- Wallace, F. G. : A new intermediate host of *Fasciolopsis buski* (Lankester). *Lingnan Sci. J.*, 15 : 125, 1936.
- Wang, M. S. : Über die Krankheitserscheinungen bei Infektion mit *Fasciolopsis buski*. *Arch. Schiffs-u. Tropen-Hyg.*, 40 : 243; 295, 1936.

- Ward, H. B. : *Fasciolopsis buski*, *F. refectus* and related species in China. *Tr. Am. Micro. Soc.*, 29 : 5, 1909.
- Ward, H. B. : *Fasciolopsis* (Letter to editor dated July 7). *J. Trop. Med. Hyg.*, 24 : 222, 1921.
- Wenell, C. M. : *Fasciolopsis* endemic in Szech. *China Med. J.*, 38 : 613, 1924.
- Wu, J. Y. C. : Incidence of helminthic parasites with special reference to the epidemiology of *Fasciolopsis buski* in Shantung, Chekiang. *Chinese Med. J.*, 51 : 77, 1927.
- Wenzel, R. and Quinck, G. : Über die erwirkungsdauer (Festnahmepériode der parasitischen wurmer im winter). *Arch. Wissenschaft u. Prakt. Tierh.*, 75 : 236, 1902.
- Wu, K. : Deux nouvelles plantes peuvent transmettre le *Fasciolopsis buski* aux humains. *Ann. Parasit.*, 15 : 458, 1937.
- Wu, K. : *Fasciolopsis buski* infecting almost the entire population in the Su-chang district of Northern Chekiang. *China J.*, 26 : 158, 1937.
- Wu, K. : Helminthic fauna in vermines of the Hangchow area. *Peking Nat. Hist. Bull.*, 12 : 1, 1937.
- Wu, K. : Susceptibility of various mammals to experimental infection with *Fasciolopsis buski* (Trematoda: Fasciolidae). *Am. Trop. Med. Parasit.*, 31 : 361, 1937.
- Wu, K. : Two additional aquatic plants as carriers of *Fasciolopsis buski* in Chekiang Province. *Chinese Med. J.*, 51 : 734, 1937.
- Wu, K. : Some aspects of *Fasciolopsis buski*. *Far East Ass. Trop. Med. Congr.-Revd.*, 10 Cong. (Hanoi, Nov. 26-Dec. 2, 1938), 2 : 671, 1940.
- Wu, K. : Snails as carriers of human parasitic diseases in China. *China J.*, 34 : 173, 1941.
- Wu, K. : Susceptibility of rabbits to infection by *Fasciolopsis buski*. *Chinese Med. J.*, 53 : 32, 1944.
- Yokoyama, S. and Morishita, K. : Handbook of human parasitology (in Japanese). Tokyodoshoten, Tokyo, Japan, 1 : 529, 1931.
- Young, S. : The blood picture in human fasciolopsiasis. *J. Shanghai Sci. Inst.*, 1 : 177, 1925.
- Young, S. : Studies on the final host of *Fasciolopsis buski* and its development in the intestine of the pig. *J. Shanghai Sci. Inst.*, 2 : 225, 1936.
- Zo, H. : Studies on *Fasciolopsis buski* in China, Nihon Shyokakibyō Gakkai Zasshi, 33 : 46, 1934.

DISCUSSION

SANDESHAN: : So far in Malaysia we have had several adult specimens of *Fasciolopsis buski* sent from the mortuary both from Singapore and Federation of Malaysia (West Malaysia). Most of the persons who harboured the worms were Chinese, and many of them could have brought the infection from outside. However, some cases were supposed not to come from abroad. With regard to mis-diagnosis, you find in the reports from the clinical laboratories that "*Fasciola hepatica*" eggs were seen, but since *Fasciola hepatica* is very uncommon, and we have never seen, any specimens obtained from the postmortem, I assume that those would be *Fasciolopsis buski* and not the other way about.

Seo: I simply want to give information on the possible incidence of fasciolopsiasis in Korea. We have *Segmentina* snails in Korea and also some water chest-nut, its size being a little bit smaller than those in Taiwan and mainland China. Korean people do not have the habit of eating raw food. However, we have a great incidence of *Fasciola hepatica* in cattle. Recently we made faecal surveys among primary school children about 20,000 in number, but we did not find any single specimen with eggs like those of *Fasciola* or *Fasciolopsis buski*.

Measomug: Recently we just completed surveys in about 20 provinces in Central Thailand with special reference to *Fasciolopsis* infection. In those areas we found that approximately 100,000 individuals, we estimated, were infected with *Fasciolopsis buski*. However, the infection seemed to be somewhat different from other areas, i.e. the infection rates or the worm burdens in the people were somewhat lower than those reported by others workers. We very rarely got an individual with over 20-30 worms. Of course, we saw cases of young children with 80-90 worms, and those individuals also passed worms in their stools which is probably an indication of very heavy infection, and they were very sick. Recently a complete study on the clinical aspects of fasciolopsiasis using the D-xylose absorption screening test and the blood pictures seemed to indicate that there was no difference between the individuals infected with *Fasciolopsis* and those with negative faeces.

With regard to the incidence of infection, in one area over 70% of the people were found to be infected, which seemed to indicate that approximately 100% of the people were infected at one time or another during their life, because from one stool examination only over 70% of them were found to be positive and the majority was young children. From surveying the human population and the animals in several villages, we found that the infection rate in pigs was almost parallel to that in the people. For example in 3 villages we surveyed, the infection rates among man were very similar, i.e. about 10% and those among the pigs were about 8-10%, indicating that the eating habits are quite similar in many respects because neither one of them cook their vegetables. As far as the plants are involved, we took all the plants in the vicinity of the houses of known infected people and examined them; essentially all of the plants had infected snails on them. So I shall assume that all the plants are potentially capable of causing infection. However, there are only 2-3 species of the plants that the people normally eat raw; the most important ones are water caltrop and Morning Glory ("Pak-beong" in Thai). Those are eaten by most of the people. However, from case histories of individuals with fasciolopsiasis, there is no one common plant they eat, probably it is an indication that there is more than one plant involved in the transmission. From our work we found that the *Segmentina*

hemiphaeridia snail was naturally infected as well as *Segmentina trochoideus* and also we succeeded in experimentally infecting them. However, *Gyraulus convexiusculus* which is living in the same areas, we have never found infected, and also we could not infect it in the laboratory.

SORNMANI : I just would like to ask Dr. Manning, in which part of Thailand did you do the survey?

MANNING : The surveys were made in Central Thailand, limited to the provinces of Suphanburi, Ayuthaya, Angtong, Bangkok, Nakorn-pathom (all in the central plain). It also included Pha-yao district of North Thailand where there was a report of a focus *buski* infection. We did surveys twice in that area and failed to turn up any infection.

SORNMANI : I ask the question because it seems to me that it is difficult to differentiate the eggs of *Fasciolopsis buski* from those of *Echinostoma* flukes.

MANNING : It is not so difficult, if you have a micrometer on your microscope. The echinostome eggs are much smaller than those of *Fasciolopsis*.

SORNMANI : But it is better if you would confirm it by getting the adult worms.

MANNING : It is true. In clinical studies which we have completed and will be published soon, we treated many cases, and obtained the adult worms. First of all we made egg counts to find out how many eggs they had, and then found how many adult worms they had; thus we got the confirmation that the eggs were those of *Fasciolopsis buski*.

We suspected that perhaps the cattle would be a reservoir for *Fasciolopsis* and then we spent a great deal of time at the slaughter house examining them, and most of them had *Fasciola* worms but not *Fasciolopsis*. On the other hand in examinations of the people we have not yet found an individual infected with *Fasciola* either.

CHAMLONG : I would like to give more information on *Fasciolopsis buski* in Thailand. The first case of fasciolopsiasis, a Thai boy 6 years old, was reported in 1941. Later in 1951-1953 Sadun and Maiphoom discovered three endemic areas of *Fasciolopsis buski* in Central Thailand, i.e. Bang Khun Sri, Thonburi (18%), Pak-hai, Ayuthaya (15%) and Pra-pa-thon, Nakorn-pathom (1%). In these areas the people usually grew water-caltrop. They described that the infection was prevalent among the children because the children had a habit of biting and eating the raw fruit of the water-caltrop where *Fasciolopsis buski* metacercariae were located. Since then there was no work on fasciolopsiasis until 1964-1965 when the research team of the Faculty of Tropical Medicine discovered a new endemic area of this infection in Suphanburi province of Central Thailand (134 kilometers northwest of Bangkok). The infected area was confined to six villages of two districts of that province. The average prevalence of infection among the people was about 20%; there was not much difference in the rates of infection among different age groups, the lowest being 15% in 1-4 years age group and the highest 24% in 40-54 years. The severity of the infection was not high as only an average of 2,300 eggs per gram of faeces were found per individual. The local pigs were also found to harbour *Fasciolopsis buski*, the incidence of which being about 30%. The interesting thing

is that there was no water-caltrop in that area; only a plant called Morning Glory was found. Moreover, neither *Segmentina* nor *Gyraulus* snails were located in the water-beds, ponds and canals in that region. However, we succeeded in completion of the life cycle of this intestinal fluke in our laboratory using the pigs as experimental animals, and *Thochorbis trochoides* snails and Morning Glory vegetation as the intermediate hosts.

E. H. Kwo : I shall add a little information about *Fasciolopsis* in Indonesia. In all text books you will find they mention fasciolopsiasis in Indonesia, but when I was in Sumatra in 1962-1967, I was unable to find any human cases of fasciolopsiasis or any pigs infected with *Fasciolopsis* at all. I tried to find the literature concerning this infection but the only one which I could find was a publication before the Second World War; it was not a paper but just a list of the parasites found in the animals in Indonesia, mentioning about *Fasciolopsis buski* in pigs in Sumatra. I think from that it brought the authors to publish *Fasciolopsis* in the text books. Now whether or not *Fasciolopsis buski* would be found in man in Indonesia needs further observations and investigations.

MIYAZAKI : I would like to ask Dr. Cross the difference between the eggs of *Fasciolopsis buski* and *Fasciola hepatica*. Would it be difficult to differentiate them?

CROSS : It is very difficult.

MANNING : We tried to study the difference between those two parasites. We took fresh stool samples containing *Fasciola* eggs from water buffaloes and those from human cases containing *Fasciolopsis* eggs. We tried to use the criteria that had been used in the past to differentiate the two different eggs, i.e., the clustering of yolk material in the nucleus, the number of yolk cells and the size of germinal area. I, however, would not feel confident about it.

MIYAZAKI : In the northern part of Japan human infection with *Fasciola hepatica* is not so infrequent, but there is no fasciolopsiasis.

FAN : I would like to give brief information about *F. buski* in Taiwan.

In Taiwan fasciolopsiasis was found as an endemic disease located in certain areas especially in the southern part of the country. It was found mostly in pigs; human infections in general were not high. However, the infected people usually resided in houses situated near ponds (in the vicinity of the pig slaughter house) where the water caltrop and other edible water plants were cultivated. There were three important kinds of water plants in Taiwan, namely, water chest-nut, water caltrop and a kind of Chinese water plant; the last one is a vegetable commonly eaten among the Chinese families. In 1962-1964 I went to the endemic areas several times in order to examine the water plants in the ponds for *F. buski*. I found that our Chinese water plant named 'Kung-shin tsai' or *Ipomoea reptans* harboured metacercariae of *F. buski* on its surface as well as in the hollow of its stem.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
U. S. Naval Medical Research Unit No. 2 Box 14, APO San Francisco 96263		Unclassified	
3. REPORT TITLE		2b. GROUP	
Fasciolopsiasis in Southeast Asia and the Far East: A Review			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Technical Report			
5. AUTHOR(S) (First name, middle initial, last name)			
John H. Cross			
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS	
1969	23	209	
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO MR005.20-0098		NAMRU-2-TR-395	
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		MR005.20.01-0098.80	
10. DISTRIBUTION STATEMENT			
Distribution of this document is unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
Published in Proceedings of the Fourth Southeast Asian Seminar on Parasitology and Tropical Medicine, 1969		Bureau of Medicine and Surgery Department of the Navy Washington, D. C.	
13. ABSTRACT			
In 1947 it was estimated that over 10 million people in the Far East were infected with <u>Fasciolopsis buski</u>. Most of these infections were reported from mainland China. Although fasciolopsiasis is endemic in Taiwan, Thailand, Vietnam, Laos, East Pakistan and India, the distribution of infections in these countries is limited and the overall prevalence rates are considerably lower than those of mainland China. While human fasciolopsiasis has not been reported from Indonesia and Burma, pigs in these countries have been found infected with the parasite. Further studies should be carried out in these countries to determine whether or not the parasite is present in man. Studies are recommended in Cambodia where little information is available and in Korea in order to determine whether or not the parasite is endemic. Human infections have been reported from Japan, the Philippines, Malaysia, and a number of western countries, but it is probable these were from people who immigrated from endemic areas or where due to misidentification of the egg. cont'd/....			

DD FORM 1473 (PAGE 1)

1 NOV 61 161-857-6801

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

13. ABSTRACT

-Fasciolopsiasis seems to be restricted to areas where populations raise water plants, such as water caltrop, water chestnuts, water hyacinth and water bamboo; and to populations that commonly ingest the uncooked metacercariae-laden plants. In addition, in most but not all the endemic countries, pigs appear to be an important reservoir of infection and species of snails of the genera Segmentina, Hippeutis, and Planorbis, the intermediate hosts.

Investigations show infections are found most commonly in individuals and families that live in the immediate vicinity of water caltrop plantations and that prevalence rates are highest in children.

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
<u>Fasciolopsis buski</u>						
<u>South East Asia</u>						
<u>Far East</u>						
<u>Segmentina sp</u>						
<u>Hippeutis sp</u>						
<u>Planorbis sp</u>						
<u>Trapa bispinosa</u>						
<u>Ipomoea reptans</u>						
<u>Eliocharis tuberosa</u>						
<u>Eichornia crassipes</u>						
<u>Zizania aquatica</u>						
<u>Salvinia natans</u>						
<u>Lemna polyrhiza</u>						