

AD 715994

**ELECTRON DETACHMENT FROM F^- ION IN A
SHOCK-HEATED CESIUM FLUORIDE-ARGON MIXTURE**

A. Mandl, E. W. Evans and B. Kivel

AVCO EVERETT RESEARCH LABORATORY

AMP 301

December 1970

jointly sponsored by

**ADVANCED RESEARCH PROJECTS AGENCY
DEPARTMENT OF DEFENSE
ARPA Order #1092**

and

**SPACE AND MISSILE SYSTEMS ORGANIZATION
AIR FORCE SYSTEMS COMMAND
DEPUTY FOR RE-ENTRY SYSTEMS (SMY)
Norton Air Force Base, California 92409**



DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.

This document has been approved
for public release and sale; its
distribution is unlimited.

ELECTRON DETACHMENT FROM F^- ION IN A
SHOCK-HEATED CESIUM FLUORIDE-ARGON MIXTURE*†

by

A. Mandl, E. W. Evans†† and B. Kivel

December 1970

AVCO EVERETT RESEARCH LABORATORY
a division of
AVCO CORPORATION
Everett, Massachusetts

* This research was supported by the Advanced Research Projects Agency of the Department of Defense and Space and Missile Systems Organization, Air Force Systems Command and was monitored by Space and Missile Systems Organization, Air Force Systems Command under Contract F04701-70-C-0128.

† Submitted to Chemical Physical Letters

†† Present address: University of Missouri, Columbia Missouri

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.

FOREWORD

Distribution of this document is unlimited. This indicates document has been cleared for public release by competent authority.

"This research was supported by the Advanced Research Projects Agency of the Department of Defense and Space and Missile Systems Organization, Air Force Systems Command and was monitored by Space and Missile Systems Organization, Air Force Systems Command under Contract F04701-70-C-0128." The secondary report number as assigned by AERL is Avco Everett Research Laboratory AMP 301. The Air Force program monitor for this contract is Capt. R. W. Padfield, USAF, Project Officer, Environmental Technology Branch, RNSE.

Publication of this report does not constitute Air Force approval of the report's findings or conclusions; it is published only for the exchange and stimulation of ideas.

Capt. R. W. Padfield, USAF,
Project Officer
Environmental Technology Branch,
RNSE

ABSTRACT

An experimental determination has been made of the rate of electron detachment from F^- in shock-heated mixtures of cesium fluoride and argon. The detachment rate is found to increase exponentially with temperature, indicating an activation energy comparable to the electron affinity of F^- . The rate constants for the reactions $F^- + Ar \xrightarrow{k(Ar)} F + Ar + e$ and $F^- + Cs^+ \xrightarrow{k(Cs^+)} F + Cs^+ + e$ have been found to be $1.2 \times 10^{-11} e^{-40,000/T}$ cm^3/sec and $5.6 \times 10^{-9} e^{-40,000/T}$ cm^3/sec respectively, where T is in degrees Kelvin.

We have studied the collisional detachment rate of electrons from F^- in a shock tube environment following the nonequilibrium overshoot of ions in the dissociation of CsF in argon. F^- collisional detachment cross sections have been measured at energies above ~ 100 eV.¹ The cross sections measured were found to remain very high down to less than 100 eV in violation of the adiabatic criterion.² We have now extended the range of these cross-section measurements to much lower energies.

The extensive work of Berry, et al.,^{3,4} has shown that the small difference between ionization potential of the alkali atom and the electron affinity of the halide atom results in the crossing of the potential energy curve for the neutral atoms and the coulomb curve for the ions at large internuclear separation causing a violation in the non-crossing rule. The thermal dissociation of an alkali halide molecule leads to a nonequilibrium population of the primary dissociation products (Cs^+ and F^- in our case).

Thus, under the conditions of our experiment, the CsF dissociates in the shock tube almost entirely into Cs^+ and F^- by the following reaction



After the CsF dissociates, we follow the decay of the F^- . We also monitor the Cs^+ density and since we know the argon density from the initial gas pressure and the shock speed, we are


able to solve the equation for the logarithmic time derivative of the F^- density

$$\frac{1}{[F^-]} \frac{d[F^-]}{dt} = -k(\text{Ar}) [\text{Ar}] - k(\text{Cs}^+) [\text{Cs}^+] \quad (2)$$

for both rate constants $k(\text{Ar})$ and $k(\text{Cs}^+)$, where we assume that the major processes leading to the collisional detachment of F^- are⁵



and



CsF is introduced, with the argon, into our shock tube as a dust of very small granules. The CsF densities are typically 1% or less of the argon density. The mixture is heated by the shock wave causing the dust to evaporate and subsequently dissociate. The time duration of both evaporation and dissociation of the CsF has been seen to be short when compared with that of the F^- collisional detachment under our experimental conditions.

Observations of the F^- , Cs^+ and electrons have been made in both absorption and emission experiments. The various diagnostics used are summarized in Table I. It should be noted that several diagnostics have been used simultaneously in taking these measurements. In our most recent measurements, we have measured the F^- absorption signal at wavelengths between 3100 Å and 2100 Å since in this region the photoabsorption cross section for F^- is

TABLE I
SUMMARY OF VARIOUS MEASUREMENTS*

Measurement	Quantity Derived	Quantity Measured ‡	Relevant Equation	Wavelength of Measurement	Detection System
F ⁻ absorption	[F ⁻]	$I = I_0 e^{-[F^-] \sigma l}$	$[F^-] = [F^-]_0 e^{-k(T)[Ar]t}$	(a) 3440 Å (b) 3100 Å to 2100 Å	(a) 3 pass - 3440 Å filter - S11 photo-tube (b) 1 pass - double monochromator - solar blind photo-tube. Incident and reflected shock
Cs ⁺ emission free-bound	[Cs ⁺][e]	$I \propto [e][Cs^+]$	$Cs^+ = \text{const}$ $[e] = [F^-]_0 (1 - e^{-k(T)[Ar]t})$	4000 Å	4000 Å filter with S11 photo-tube
Bremsstrahlung emission †	[e]	$I \propto [e]$	$[e] = [F^-]_0 (1 - e^{-k(T)[Ar]t})$	2.5 μ	Liquid N ₂ cooled InSb photovoltaic element

*[] = density

[]₀ = initial density

† The relation for I holds if ion density is less than a few by 10⁻³ of neutral gas density.

‡ σl = absorption cross section per F⁻ ion multiplied by the length of the absorption path.

several times larger than at 3445 Å.^{6,7} The light source is a xenon flash lamp which is pulsed so that its maximum intensity is reached as the shock front passes the detector. The detector used is a double monochrometer with a solar blind phototube. Measurements were simultaneously made of the free-bound $\text{Cs}^+ + e$ emission in order to give us a measure of the Cs^+ concentration.

The results of these measurements are plotted in Fig. 1. We have rewritten Eq. (2) in terms of a total measured rate constant as

$$\frac{1}{[F^-]} \frac{d[F^-]}{dt} = k(T) [Ar] \quad (4)$$

where

$$k(T) = k(Ar) + k(\text{Cs}^+) \frac{[\text{Cs}^+]}{[Ar]} \quad (5)$$

and assuming the same T dependence for both reactions

$$= \left\{ a + \beta \frac{[\text{Cs}^+]}{[Ar]} \right\} e^{-40,000/T} \quad (6)$$

$$k(Ar) = a e^{-40,000/T} \quad (7a)$$

$$k(\text{Cs}^+) = \beta e^{-40,000/T} \quad (7b)$$

Thus, the intercept of Fig. 2 gives a and the slope β . If we substitute these values into Eqs. (7a) and (7b), we find

$$k(Ar) = 1.2 \times 10^{-11} e^{-40,000/T} \text{ cm}^3/\text{sec} \quad (8a)$$

and

$$k(\text{Cs}^+) = 5.6 \times 10^{-9} e^{-40,000/T} \text{ cm}^3/\text{sec} \quad (8b)$$

We have plotted our measured rate constant $k(Ar)$ as a function of $1/T$ in Fig. 2. It is seen that the best fit




Fig. 1 This is a plot of our measured rate constant $[k(T)]$ of Eq. 6], with the temperature dependence divided out as a function of the fractional concentration of Cs^+ to Ar. The y intercept gives the value of $k(T)/e^{-40,000/T}$ at zero $[\text{Cs}^+]$ [i. e., α of Eq. (7a)] whereas the slope determines β of Eq. (7b). The open circles are incident shock measurements and the closed circles are reflected shock measurements.





Fig. 2 The electron detachment rate of F^- assuming impact ionization by the atomic argon [$k(\text{Ar}) \text{ cm}^{-3}/\text{sec}$] as a function of $10^4 \text{ K}/T$. The effect of the Cs^+ (as determined from Fig. 1) has been subtracted. The solid line shows the dependence of a detachment rate having an activation energy equal to the electron affinity of F^- . The open circles are incident shock measurements and the closed circles are reflected shock measurements.

to these points gives a slope which is consistent with an activation energy equal to the electron affinity of fluorine.

We thus find ourselves in agreement with some of the results of the higher energy measurements, i. e. , that the collisional detachment cross sections remain larger than would be theoretically expected down to very low velocities.² We can also compare our results to those of Berry, et al.,³ who find for the reaction



that (in our notation)

$$k'(\text{Ar}) = 5.8 \times 10^{-11} e^{-39,000/T} \text{ cm}^3/\text{sec}, \quad (10)$$

in good agreement with our Eq. (8a).

ACKNOWLEDGMENTS

The authors are indebted to Dr. E. Sutton who proposed this experiment and guided the initial phases of the program and to R. Jonkman for much work performed in the initial phase. Significant technical help and guidance was received from K. Wray, Chairman of the Avco Everett Research Laboratory Atomic Physics Committee. We also acknowledge assistance from J. Keck, M. Camac and J. D. Teare. Numerical nonequilibrium calculations of shock front conditions were provided by G. E. Caledonia. The technical assistance received from J. W. Martz, Jr., in operation of the shock tube and other experimental apparatus was a most important contribution to this program.

REFERENCES

1. Hasted, J. B. , Proc. Roy. Soc. A222, 74 (1954).
2. Branscomb, L. M. , Adv. in Electronics & Electron Phys. , Vol. IX, Academic Press (1957).
3. Berry, R. S. , et al. , J. Chem. Phys. 49, 127 (1968).
4. Ewing, J. J. , Milstein, R. and Berry, R. S. , 7th Shock Tube Symp. , Toronto, Canada (1969).
5. Processes of the type $F^- + e \rightarrow F + 2e$, $F + F^- \rightarrow F_2 + e$ and $F + F^- \rightarrow F + F + e$ are not considered since if these processes were large, our resultant F^- decay curve would show an incubation time corresponding to the initially low electron density and fluorine density. This shape is not observed. The reaction $F^- + F^- \rightarrow F + F^- + e$ is assumed to be negligible when compared with (3b).
6. The results of the measurements are to be published elsewhere.
7. E. W. Evans and B. Kivel, Avco Everett Research Laboratory Research Note 827 (Avco Everett Research Laboratory, Everett, Mass. , 1969).

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Avco Everett Research Laboratory 2385 Revere Beach Parkway Everett, Massachusetts		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE Electron Detachment From F^- Ion in a Shock-Heated Cesium Fluoride-Argon Mixture			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) AMP 301			
5. AUTHOR(S) (Last name, first name, initial) A. Mandl, E. W. Evans and B. Kivel			
6. REPORT DATE December 1970		7a. TOTAL NO. OF PAGES 9	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO. F04701-70-C-0128		8b. ORIGINATOR'S REPORT NUMBER(S) AMP 301	
a. PROJECT NO.			
c.		8d. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		SAMSO-TR-70-101	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited. This indicates document has been cleared for public release by competent authority.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Advanced Research Projects Agency, Department of Defense, ARPA Order #1092 and SAMSO, AFSC, Deputy for Re-entry Systems (SMY), Norton Air Force Base, Cal. 92409	
13. ABSTRACT An experimental determination has been made of the rate of electron detachment from F^- in shock-heated mixtures of cesium fluoride and argon. The detachment rate is found to increase exponentially with temperature, indicating an activation energy comparable to the electron affinity of F^- . The rate constants for the reactions $F^- + Ar \xrightarrow{k(Ar)} F + Ar + e$ and $F^- + Cs \xrightarrow{k(Cs^+)} F + Cs^+ + e$ have been found to be $1.2 \times 10^{-11} e^{-40,000/T} \text{ cm}^3/\text{sec}$ and $5.6 \times 10^{-9} e^{-40,000/T} \text{ cm}^3/\text{sec}$ respectively, where T is in degrees Kelvin.			

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
1. Electron Detachment						
2. Negative Ion						
3. F ⁻ Ion						
4. Shock tube						
5. Cesium Fluoride						
6. Rate Constant						
7. Cs ⁺ ion						

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.