

THE EFFECT OF INTERMITTENT EXPOSURE TO 3% CO₂ ON
ACID-BASE BALANCE AND ELECTROLYTE EXCRETION


by

K. E. Schaefer
C. C. Morgan
A. A. Messier
and
M. J. Jacey

SUBMARINE MEDICAL RESEARCH LABORATORY
NAVAL SUBMARINE MEDICAL CENTER REPORT NO. 635

Bureau of Medicine and Surgery, Navy Department
Research Work Unit M4306.02-7050B.02

Reviewed and Approved by:



Charles F. Gell, M.D., D.Sc. (Med)
Scientific Director
SubMedResLab

Reviewed and Approved by:



J. D. Bloom, CDR MC USN
Officer-in-Charge
SubMedResLab

Approved and Released by:



J. E. Stark, CAPT MC USN
COMMANDING OFFICER
Naval Submarine Medical Center

This document has been approved for public release and sale;
its distribution is unlimited.

SUMMARY PAGE

THE PROBLEM

To determine the effects of intermittent exposure to increasing CO₂, rising at a constant rate from 0.03 to 3% CO₂ within a period of 15 hours followed by a nine-hour period of air breathing, on acid-base balance.

FINDINGS

Intermittent exposure of one subject for six days to increased CO₂ for fifteen hours per day produced an increase in hydrogen ion concentrations and PCO₂ in capillary blood during the air breathing periods of the fourth and fifth day. This CO₂ accumulation which could not be prevented by respiratory gas exchange was eliminated through renal mechanisms of increased hydrogen ion excretion during the fourth and fifth day of intermittent exposure. This renal mechanism was not found to operate under chronic exposure to 3% CO₂. The 17 hydroxysteroid excretion in the urine was not affected by intermittent exposure to CO₂ indicating that there was no significant stress effect.

APPLICATIONS

These findings are pertinent to submarine operations in snorkel type submarines in which CO₂ levels may be achieved intermittently for short periods of time.

ADMINISTRATIVE INFORMATION

This investigation was conducted as a part of Bureau of Medicine and Surgery Research Work Unit M4306.02-7050B -- The Effects of Intermittent CO₂ and Lowered Oxygen Levels on Diving and Submarine Personnel. The present report is No. 2 on this work unit. The manuscript was approved for publication on 16 July 1970, and designated as Submarine Medical Research Laboratory Report No. 635.

PUBLISHED BY THE NAVAL SUBMARINE MEDICAL CENTER

ABSTRACT

The effects of intermittent exposure to CO_2 on acid-base balance was investigated. One subject was exposed for six days to increasing CO_2 rising at a constant rate from 0.03 to 3.0% CO_2 within a period of 15 hours followed by a nine hour period of air breathing. To assess the acid-base parameters "arterialized" capillary blood was taken from the finger twice daily at 8 AM and 11 PM corresponding with the beginning and end of the intermittent exposure to CO_2 .

Urine specimens were collected at the same times daily while venous blood samples were obtained on alternate days.

Hydrogen ion concentrations and PCO_2 in the arterialized capillary blood were found increased during the air breathing periods of the fourth and fifth day but returned to normal during the sixth day. The elimination of this CO_2 accumulation in the blood was accomplished through renal mechanism showing a marked increase in urine volume, organic acids, titratable acidity, ammonia and hydrogen ion excretion during the fourth and fifth day. The stress effect produced by intermittent exposure to CO_2 in this subject must have been minimal since the 17-hydroxysteroid excretion did not change.

TABLE II. EFFECT OF INTERMITTENT EXPOSURE TO 3% CO₂
ON pH, P_{CO₂}, HCO₃ AND P_{O₂} OF VENOUS BLOOD

Condition		pH	P _{CO₂} mm Hg	HCO ₃ mM/L	P _{O₂} mm Hg
Control 8 AM	Mean	7.360	53.4	29.7	26.6
	SEM	-	-	-	-
	N	1	1	1	1
Control 11 PM	Mean	7.323	58.2	29.3	20.0
	SEM	-	-	-	-
	N	1	1	1	1
Intermittent Exposure to 3% CO ₂ -9 hrs. on air, 8 AM	Mean	7.319	57.3	28.9	22.2
	SEM	.006	1.7	.5	3.8
	N	3	3	3	3
Intermittent Exposure - 15 hrs. on 3% CO ₂ , 11 PM	Mean	7.295*	59.4	28.6	31.9
	SEM	.003	2.4	1.3	2.4
	N	3	3	3	3
Recovery on air 8 AM	Mean	7.338	59.5	31.6	21.4
	SEM	.018	.5	1.1	7.7
	N	2	2	2	2
Recovery on air 11 PM	Mean	7.342	60.8	32.5	21.6
	SEM	.005	1.8	.5	4.5
	N	2	2	2	2

* Significantly different from corresponding controls at the 5% level and better.

venous blood increased during CO₂ breathing which confirms the findings on alveolar oxygen tensions (Figure 3).

The responses of the renal functions primarily involved in acid-base regulations are shown in Figures 4 and 5. There is an immediate response to CO₂ breathing on the first day as shown in an increase in urine volume and excretion of organic acids, titratable acidity, and ammonia, which is followed by a decline in these parameters during the next two days. However, during the fourth and fifth day, which are marked by an increased CO₂ excretion and acid load during the air breathing period, there is a marked increase in urine volume, organic acids, titratable acidity, ammonia, and hydrogen ion excretion. During the second day of recovery an opposite




Fig. 3. Effect of intermittent exposure to CO₂ on PO₂ of arterialized capillary blood and venous blood. (Solid line: values obtained at 8 AM on air; Dotted line: values obtained at 11 PM after CO₂ exposure)




Fig. 4. Effect of intermittent exposure to CO₂ on urine volume and urinary excretion of organic acids. (Black blocks: 15 hour excretion from 8 AM - 11 PM; Stippled blocks: 9 hour excretion during night - breathing air)

trend can be noted which is characterized by a decreased excretion of ammonia and titratable acidity and a reduction in hydrogen ion excretion commensurate with a large increase in bicarbonate elimination.

Blood lactate and pyruvate and L/P ratio were not affected by intermittent exposure to CO₂ (Table III). The 17-hydroxysteroid excretion in the urine exhibited large diurnal variations which were not influenced by intermittent exposure to CO₂ (Table IV).

REFERENCES

1. Brackett, N. C., Jr., C. F. Wingo, O. Muren and J. T. Salano. Acid base response to chronic hypercapnia in man. N. Eng. J. Med. 280:124-130, 1969.
2. Bray's Clinical Laboratory Methods. Sixth Edition, D. V. Mosby Co., 1962, p. 73.
3. Clark, J.M., et al. Rate of Acclimatization to chronic hypercapnia in man. USAF School of Aerospace Medicine Environmental Systems Division, Brooks AFB, Texas.
4. Gambino, S. R. Comparisons of pH in human arterial, venous and capillary blood. Am. J. Clin. Path. 32:298, 1959.
5. Glatte, H. A., Jr., et al. Carbon dioxide tolerance studies. USAF School of Aerospace Medicine, Aerospace Medical Division (AFSC), Brooks AFB, Texas. Report SMA-TR-67-77.
6. Schaefer, K. E. Atmung und Saure-Basengleichgewicht bei langdauerndem Aufenthalt in 3% CO₂. Pfluegers Archiv. 251: 689-715, 1949.
7. Schaefer, K. E., G. Nichols, Jr. and C. R. Carey. Acid-base balance and blood and urine electrolytes of man during acclimatization to CO₂. Submarine Medical Research Laboratory. Report No. 425, March, 1964.
8. Schaefer, K. E., C. R. Carey and J. H. Dougherty, Jr. The effect of intermittent exposure to 3% CO₂ on respiration. Submarine Medical Research Laboratory Report No. 618, March, 1970.
9. Schwartz, W. B., N. C. Brackett, and J. J. Cohen. The response of extracellular hydrogen ion concentration of graded degrees of chronic hypercapnia. The physiological limits of the defense of the pH. J. Clin. Invest. 44:291, 1965.
10. Sullivan, W. J., and P. J. Dorman. Renal response to chronic respiratory acidosis. J. Clin. Invest. 34:268, 1955.
11. Weitzman, D. O. and J. A. S. Kinney and S. M. Luria. Effect on vision of repeated exposure to carbon dioxide. Submarine Medical Research Laboratory. Report No. 566, 14 February 1969.
12. Van Ypersele De Strihou, C., L. Brasseur and J. De Coninck. The "carbon dioxide response curve" for chronic hypercapnia in man. N. Eng. J. Med. 275: 117-122, 1966.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) NAVAL SUBMARINE MEDICAL CENTER, Submarine Medical Research Laboratory		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE THE EFFECT OF INTERMITTENT EXPOSURE TO 3% CO ₂ on ACID-BASE BALANCE AND ELECTROLYTE EXCRETION		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Interim Report		
5. AUTHOR(S) (First name, middle initial, last name) Karl E. SCHAEFER, C. C. MORGAN, Arthur A. MESSIER and Michael J. JACEY		
6. REPORT DATE 16 July 1970	7a. TOTAL NO. OF PAGES 8	7b. NO. OF REFS 12
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) SMRL Report No. 635	
b. PROJECT NO. M4306.02-705QB		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Naval Submarine Medical Center Box 600 Naval Submarine Base New London Groton, Connecticut 06340
13. ABSTRACT The effects of intermittent exposure to CO ₂ on acid-base balance was investigated. One subject was exposed for six days to increasing CO ₂ rising at a constant rate from 0.03 to 3.0% CO ₂ within a period of 15 hours followed by a nine hour period of air breathing. To assess the acid-base parameters "arterialized" capillary blood was taken from the finger twice daily at 8 AM and 11 PM corresponding with the beginning and end of the intermittent exposure to CO ₂ . Urine specimens were collected at the same times daily while venous blood samples were obtained on alternate days. Hydrogen ion concentrations and PCO ₂ in the arterialized capillary blood were found increased during the air breathing periods of the fourth and fifth day but returned to normal during the sixth day. The elimination of this CO ₂ accumulation in the blood was accomplished through renal mechanism showing a marked increase in urine volume, organic acids, titratable acidity, ammonia and hydrogen ion excretion during the fourth and fifth day. The stress effect produced by intermittent exposure to CO ₂ in this subject must have been minimal since the 17-hydroxysteroid excretion did not change.		

DD FORM 1473 (PAGE 1)

S/N 0101-807-6801

UNCLASSIFIED

Security Classification

3ND PPSO 13152

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Intermittent exposure to CO ₂						
Acid base balance						
Carbon dioxide toxicity						
Renal mechanism						

UNCLASSIFIED

Security Classification

