

ARL 71-0131
AUGUST 1971



Aerospace Research Laboratories

THE DISTRIBUTION OF A TRUNCATED LINEAR DIFFERENCE BETWEEN INDEPENDENT CHI-SQUARE VARIATES

DAVID A. HARVILLE

APPLIED MATHEMATICS RESEARCH LABORATORY


PROJECT No. 7071

Approved for public release; distribution unlimited.

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22151

AIR FORCE SYSTEMS COMMAND

United States Air Force



ARL 71-0131

**THE DISTRIBUTION OF A TRUNCATED
LINEAR DIFFERENCE BETWEEN
CHI-SQUARE VARIATES**

DAVID A. HARVILLE

APPLIED MATHEMATICS RESEARCH LABORATORY

AUGUST 1971

PROJECT No. 7071

Approved for public release; distribution unlimited

AEROSPACE RESEARCH LABORATORIES
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
WRIGHT-PATTERSON AIR FORCE BASE, OHIO

NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely related Government procurement operation, the United States Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

ACCESSION BY		
CPSTH	WHITE SECTION ☒	
DOC	DIFF SECTION ☐	
UNANNOUNCED	☐	
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
DUT.	AVAIL. and/or	SPECIAL
A		

Agencies of the Department of Defense, qualified contractors and other government agencies may obtain copies from the

Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314

This document has been released to the

CLEARINGHOUSE
U. S. Department of Commerce
Springfield, Virginia 22151

for sale to the public.

Copies of ARL Technical Documentary Reports should not be returned to Aerospace Research Laboratories unless return is required by security considerations, contractual obligations or notices on a specified document.

FOREWORD

This report was prepared by D. A. Harville of the Applied Mathematics Research Laboratory, Aerospace Research Laboratories, Wright-Patterson Air Force Base, Ohio. The research comprises part of Project 7071, "Research in Applied Mathematics".

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Aerospace Research Laboratories Applied Mathematics Research Laboratory Wright-Patterson AFB, Ohio 45433		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE The Distribution of a Truncated Linear Difference Between Independent Chi-square Variates		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Scientific Final		
5. AUTHOR(S) (First name, middle initial, last name) David A. Harville		
6. REPORT DATE August 1971	7a. TOTAL NO. OF PAGES 23	7b. NO. OF REFS 5
8a. CONTRACT OR GRANT NO. In-house research		8a. ORIGINATOR'S REPORT NUMBER(S)
b. PROJECT NO. 7071-00-10		
c. DoD Element 61102F		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
d. DoD Subelement 681304		ARL 71-0131
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.		
11. SUPPLEMENTARY NOTES TECH OTHER		12. SPONSORING MILITARY ACTIVITY Aerospace Research Laboratories (LB) Air Force Systems Command Wright-Patterson AFB, Ohio 45433
13. ABSTRACT A comprehensive account is given of the properties of the distribution of a doubly-truncated linear difference between two independent chi-square variates. Expressions for the probability density function, distribution function, moments, moment generating function, and characteristic function of the distribution are obtained, primarily by making use of known results on hypergeometric functions. Emphasis is placed on representations thought to be useful for computations. In general, simple expressions are possible only when at least one of the chi-squares has an even number of degrees of freedom. The results should prove useful in studying the properties of truncated quadratic estimators of variance components.		

DD FORM 1 NOV 65 1473

Unclassified

Security Classification

ABSTRACT

A comprehensive account is given of the properties of the distribution of a doubly-truncated linear difference between two independent chi-square variates. Expressions for the probability density function, distribution function, moments, moment generating function, and characteristic function of the distribution are obtained, primarily by making use of known results on hypergeometric functions. Emphasis is placed on representations thought to be useful for computations. In general, simple expressions are possible only when at least one of the chi-squares has an even number of degrees of freedom. The results should prove useful in studying the properties of truncated quadratic estimators of variance components.

Unclassified

Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Variance components Quadratic forms in normal variates Hypergeometric functions Moments Linear difference between chi-squares						

Unclassified

Security Classification

TABLE OF CONTENTS

Section	Page
1. Introduction	1
2. Notation and Preliminary Results	2
3. Evaluation of $g(r,s,a,b; x)$	5
4. Evaluation of $G(r,s,a,b; k,\theta,x)$	10
References	18

1. Introduction. Quadratic estimators of the variance components associated with random-effects models are in common use. If the experimental design is a balanced completely-nested one, if the usual independence and normality assumptions are appropriate, and if the estimators are those derived by analysis-of-variance techniques, then each of the estimators, save that of the residual variance, is distributed as a linear difference between two independent chi-square variates. More generally, any quadratic form in multivariate-normal data and in particular any quadratic variance-component estimator based on such data is either distributed as a linear difference between two independent chi-squares or, as Press [2] has shown, its distribution can be represented as a mixture of such distributions.

In cases where a quadratic estimator of a variance component can take on negative values, it is common practice to replace negative estimates by zero. If the data are multivariate-normal, the distribution of the modified estimator is identical to that of a truncated (at zero) linear difference between independent chi-squares or is a mixture of such truncated distributions.

It will be our purpose to investigate the distribution of a truncated linear difference between independent chi-square variates. Rather than assume that the truncation is from below and at zero, we will consider the more general situation where the truncation is either from above or below or both and at arbitrary values. In particular, we will obtain expressions

for the probability density function, distribution function, moments, moment generating function, and characteristic function of such a distribution. In the case of the probability density function, distribution function, and moments, emphasis will be on obtaining representations that appear to be convenient computational forms. Extensive use will be made of known results on hypergeometric functions.

Previously, Press [2] obtained the probability density function of a linear difference between independent chi-squares in terms of a confluent hypergeometric function of the second kind. Wang [5] found a simpler expression for the probability density function for the special case where the degrees of freedom associated with both chi-squares are even. She also gave, for that same special case, convenient expressions for the moments of that distribution, after its truncation from below at zero.

2. Notation and preliminary results. Define

$$X = \alpha W_1 - \beta W_2,$$

where α and β are positive constants, and W_1 and W_2 are independent chi-square variates having m and n degrees of freedom, respectively. Truncation of X from above and/or below yields the random variable

$$\begin{aligned} Y &= y_1 && \text{if } X > y_1, \\ &= X && \text{if } y_1 \geq X \geq y_0, \\ &= y_0 && \text{if } X < y_0, \end{aligned}$$

where y_0 and y_1 are known constants with $-\infty \leq y_0 < y_1 \leq \infty$.

A chi-square variate with r degrees of freedom has k th moment $2^k (r/2)_k$, where $(a)_0 = 1$ and $(a)_k = a(a+1)\dots(a+k-1)$, $k = 1, 2, \dots$; moment generating function

$$M(\theta) = (1 - 2\theta)^{-r/2}, \quad \theta < 1/2;$$

and characteristic function

$$C(\theta) = (1 - 2i\theta)^{-r/2}.$$

It follows that X has k th moment

$$(1) \quad (2\alpha)^k \sum_{j=0}^k \binom{k}{j} (-\zeta)^j (n/2)_j (m/2)_{k-j},$$

where $\zeta = \beta/\alpha$; moment generating function

$$(2) \quad M_X(\theta) = (1 - 2\alpha\theta)^{-m/2} (1 + 2\beta\theta)^{-n/2}, \quad (-2\beta)^{-1} < \theta < (2\alpha)^{-1};$$

and characteristic function

$$C_X(\theta) = (1 - 2\alpha i\theta)^{-m/2} (1 + 2\beta i\theta)^{-n/2}.$$

Take

$$U(c, d; x) = [1/\Gamma(c)] \int_0^\infty e^{-xt} t^{c-1} (1+t)^{d-c-1} dt, \quad c > 0, \quad x > 0,$$

to be a confluent hypergeometric function of the second kind, and put

$$(3) \quad g(r, s, a, b; t) = \frac{t^{(r+s-2)/2} e^{-t/(2a)} U[s/2, (r+s)/2; (a+b)t/(2ab)]}{2^{(r+s)/2} a^{r/2} b^{s/2} \Gamma(r/2)}$$

for $r = 1, 2, \dots$; $s = 1, 2, \dots$; $a > 0$, $b > 0$; and $t > 0$.

From [2], we have that the probability density function of X is

$$(4) \quad \begin{aligned} f_X(x) &= g(m, n, \alpha, \beta; x) && \text{if } x > 0, \\ &= g(n, m, \beta, \alpha; -x) && \text{if } x < 0, \\ &= \frac{\zeta^{m/2} \Gamma(m/2 + n/2 - 1)}{2^{(m+n)/2} \beta [(\zeta+1)/2]^{(m+n-2)/2} \Gamma(m/2) \Gamma(n/2)} && \text{if } x = 0. \end{aligned}$$

It will prove convenient to express the distribution functions of X and Y and the moments, moment generating function, and characteristic function of Y in terms of the function

$$(5) \quad \begin{aligned} G(r, s, a, b; k, z, u) &= \int_0^u t^k e^{zt} g(r, s, a, b; t) dt && \text{for } u > 0, \\ &= 0 && \text{for } u = 0, \end{aligned}$$

where $k = 0, 1, 2, \dots$

Note that, for $v < 0$,

$$(6) \quad \int_v^0 t^k e^{zt} g(s, r, b, a; -t) dt = (-1)^k G(s, r, b, a; k, -z, -v).$$

The distribution function of X is

$$\begin{aligned} F_X(x) &= G(m, n, \alpha, \beta; 0, 0, x) + G(n, m, \beta, \alpha; 0, 0, \infty) && \text{if } x \geq 0, \\ &= G(n, m, \beta, \alpha; 0, 0, \infty) - G(n, m, \beta, \alpha; 0, 0, -x) && \text{if } x \leq 0. \end{aligned}$$

Denote by $P_{r,s}(\cdot)$ the distribution function of an F random variable having r and s degrees of freedom. Note that

$$P_{m,n}[(n\beta)/(m\alpha)] = F_X(0) = G(n, m, \beta, \alpha; 0, 0, \infty).$$

The distribution function of Y is

$$\begin{aligned} F_Y(y) &= 1 && \text{if } y \geq y_1, \\ &= F_X(y) && \text{if } y_0 \leq y < y_1, \\ &= 0 && \text{if } y < y_0. \end{aligned}$$

If y_0 and y_1 are finite, $F_Y(\cdot)$ is discontinuous at those points. It makes a jump of $F_X(y_0)$ at y_0 and $1 - F_X(y_1)$ at y_1 . At other points, the probability density function of Y is

$$\begin{aligned} F'_Y(y) &= f_X(y) && \text{if } y_0 < y < y_1, \\ &= 0 && \text{if } y < y_0 \text{ or } y > y_1. \end{aligned}$$

Define

$$\begin{aligned} H(u_1, u_0; k, z) &= G(m, n, \alpha, \beta; k, z, u_1) - G(m, n, \alpha, \beta; k, z, u_0) && \text{if } 0 \leq u_0 < u_1, \\ &= G(m, n, \alpha, \beta; k, z, u_1) + (-1)^k G(n, m, \beta, \alpha; k, -z, -u_0) && \text{if } u_0 < 0 \leq u_1, \\ &= (-1)^k [G(n, m, \beta, \alpha; k, -z, -u_0) - G(n, m, \beta, \alpha; k, -z, -u_1)] && \text{if } u_0 < u_1 < 0; \end{aligned}$$

and

$$H^*(u_1, u_0; k, z) = H(u_1, u_0; k, z) + u_0^k e^{zu_0} F_X(u_0) + u_1^k e^{zu_1} [1 - F_X(u_1)]$$

for $-\infty < u_0 < u_1 < \infty$,

$$H^*(\infty, u_0; k, z) = H(\infty, u_0; k, z) + u_0^k e^{zu_0} F_X(u_0) \quad \text{for } -\infty < u_0,$$

$$H^*(u_1, -\infty; k, z) = H(u_1, -\infty; k, z) + u_1^k e^{zu_1} [1 - F_X(u_1)] \quad \text{for } u_1 < \infty,$$

and

$$H^*(\infty, -\infty; k, z) = H(\infty, -\infty; k, z),$$

with $0^0 = 1$.

The k th moment of Y is $H^*(y_1, y_0; k, 0)$, and its moment generating

function and characteristic function are

$$M_Y(\theta) = H^*(y_1, y_0; 0, \theta), \quad \theta > (-2\beta)^{-1} \text{ if } y_0 < 0 \text{ and } \theta < (2\alpha)^{-1} \text{ if } y_1 > 0,$$

and

$$C_Y(\theta) = H^*(y_1, y_0; 0, i\theta),$$

respectively.

It is clear that the problem of evaluating the probability density function of X or Y can be reduced to one of evaluating $g(r, s, a, b; t)$ for appropriate values of that function's parameters. Moreover, to evaluate the distribution function of X or Y and the moments and moment generating function of Y , it suffices to evaluate $G(r, s, a, b; k, \theta, u)$ for appropriate values of r, s, a , and b and for $k = 0, 1, 2, \dots, \theta$ real, and $u > 0$. For the most part, we will find it convenient to carry out the analysis for all such (k, θ, u) triplets, rather than to separately consider only those particular triplets that are relevant to the evaluation of the distribution function, the moments, and the moment generating function, respectively. Because, in (5), $t^k e^{zt}$ is 'absorbed' by $g(r, s, a, b; t)$, the more general approach is no more difficult. It has the advantage of avoiding repetition. Since the expressions for both the moment generating function and the moments are to be obtained directly, the derivatives at zero of expressions for the moment generating function can be compared with the corresponding expressions for the moments, so as to obtain a check on the correctness of the results.

Expressions for the characteristic function $C_Y(\theta)$ of Y will not be given explicitly; however, substitution of $i\theta$ for θ , in any of the expressions indicated for the moment generating function $M_Y(\theta)$, produces a valid representation for $C_Y(\theta)$, provided that representation is meaningful.

3. Evaluation of $g(r, s, a, b; x)$. Usual relationships. It is

easy to show that, for $c > 0$,

$$(7) \quad g(r, s, a, b; t) = c g(r, s, ca, cb; ct).$$

The recurrence relations for confluent hypergeometric functions given by (13.4.15) - (13.4.20) of Slater [4] yield

$$(8) \quad g(r, s, a, b; t) = (b/a)(r-2)^{-1} \{ [t(a^{-1}+b^{-1}) + s - r + 4] g(r-2, s+2, a, b; t) + (b/a)(s+2) g(r-4, s+4, a, b; t) \},$$

$$(9) \quad g(r, s, a, b; t) = b(a+b)^{-1}(r-2)^{-1} \{ [t(a^{-1}+b^{-1}) + s + r - 4] g(r-2, s, a, b; t) - (t/a) g(r-4, s, a, b; t) \},$$

$$(10) \quad g(r, s, a, b; t) = [a(r-2)]^{-1} [bs g(r-2, s+2, a, b; t) + t g(r-2, s, a, b; t)],$$

$$(11) \quad g(r, s, a, b; t) = (a+b)^{-1} [b g(r-2, s, a, b; t) + a g(r, s-2, a, b; t)],$$

$$(12) \quad g(r, s, a, b; t) = b(a+b)^{-1}(r-2)^{-1} \{ [t(a^{-1}+b^{-1}) + s] g(r-2, s, a, b; t) + (bs/a) g(r-4, s+2, a, b; t) \},$$

and

$$(13) \quad g(r, s, a, b; t) = (b/a)(r-2)^{-1} \{ [t(a^{-1}+b^{-1}) + s] g(r-2, s+2, a, b; t) - (t/a) g(r-4, s+2, a, b; t) \}.$$

The probability density function of a chi-square variate with r degrees of freedom is

$$p_r(t) = [2^{r/2} \Gamma(r/2)]^{-1} t^{(r/2)-1} e^{-t/2}, \quad 0 < t < \infty, \\ = 0, \quad \text{elsewhere.}$$

The domain of the function g can be extended by putting

$$g(r, 0, a, b; t) = (1/a) p_r(t/a)$$

for $r = 1, 2, \dots$; $a > 0$; $b > 0$; $t > 0$. The relationships (7) - (13) remain valid. Thus, tables of the chi-square density may be of use in evaluating the densities of X and Y .

Case (i): r an even integer. For $c > 0$ and $p = 1, 2, \dots$,

$$(14) \quad \begin{aligned} U(c, c+p; x) &= [1/\Gamma(c)] \sum_{j=0}^{p-1} \binom{p-1}{j} \int_0^\infty e^{-xt} t^{c+j-1} dt \\ &= \sum_{j=0}^{p-1} \binom{p-1}{j} (c)_j x^{-c-j}. \end{aligned}$$

Substitution of this expression into (3) gives

$$(15) \quad g(r, s, a, b; x) = [a^{r/2} \Gamma(r/2)]^{-1} [a/(a+b)]^{s/2} e^{-x/(2a)} \\ \cdot \sum_{j=0}^{r/2-1} \binom{r/2-1}{j} (s/2)_j [2b/(a+b)]^j (x/a)^{r/2-j-1},$$

for $r = 2, 4, 6, \dots$

Case (ii): s an even integer. Denote by $P_u(\cdot)$ the distribution function of a chi-square variate with u degrees of freedom. For $p = 1, 2, \dots$ and $2c = 1, 2, \dots$,

$$\begin{aligned} U(p, p+c; x) &= [1/\Gamma(p)] \int_1^\infty e^{-x(v-1)} (v-1)^{p-1} v^{c-1} dv \\ &= [1/\Gamma(p)] e^x \sum_{j=0}^{p-1} \binom{p-1}{j} (-1)^{p-j-1} \int_1^\infty e^{-xv} v^{c+j-1} dv \\ &= [1/\Gamma(p)] e^x \sum_{j=0}^{p-1} \binom{p-1}{j} (-1)^{p-j-1} x^{-c-j} \Gamma(c+j) \\ &\quad \cdot [1 - P_{2c+2j}(2x)]. \end{aligned}$$

Substituting this expression into (3), we find

$$\begin{aligned} g(r, s, a, b; x) &= [b^{s/2} \Gamma(s/2)]^{-1} [b/(a+b)]^{r/2} e^{x/(2b)} \\ &\quad \cdot \sum_{j=0}^{s/2-1} \binom{s/2-1}{j} (r/2)_j [2a/(a+b)]^j (-x/b)^{s/2-j-1} \\ &\quad \cdot \{1 - P_{r+2j}[x(a+b)/(ab)]\}, \end{aligned}$$

for $s = 2, 4, 6, \dots$

Case (iii): Both r and s are odd integers. Let

$$c_j(u, v) = \frac{\binom{u}{j}}{j! \binom{v}{j}}, \quad j = 1, 2, \dots$$

Take

$$\Phi(u, v; z) = \sum_{j=0}^{\infty} c_j(u, v) z^j$$

to be a confluent hypergeometric function of the first kind. This series is absolutely convergent for all values of u, v , and z , save $v = 0, -1, -2, \dots$

Its remainder after q terms will be denoted by the symbol

$$R_q(u, v; z) = \sum_{j=q}^{\infty} c_j(u, v) z^j.$$

The numerical evaluation of $U[s/2, (r+s)/2; (a+b)x/(2ab)]$ and consequently of $g(r, s, a, b; x)$ can be accomplished for $r = 1, 3, 5, \dots$, $s = 1, 3, 5, \dots$, by applying the well-known relationship

$$(16) \quad U(v, 1+p; t) = [(-1)^p p! \Gamma(v-p)]^{-1} L(v, 1+p; t) \\ + [(p-1)!/\Gamma(v)] \sum_{j=0}^{p-1} c_j(v-p, 1-p) t^{j-p},$$

for $p = 0, 1, 2, \dots$; where, taking $\psi(t) = \Gamma'(t)/\Gamma(t)$ to be the logarithmic derivative of the gamma function and $\gamma = -\psi(1) = 0.57721\ 56649\dots$ to be Euler's constant, $L(v, 1+p; t)$ is obtained by adding together the three convergent series

$$(17) \quad (-\gamma - \ln t) \phi(v, 1+p; t),$$

$$(18) \quad \sum_{j=0}^{\infty} c_j(v, 1+p) t^j [\psi(1+j+p) - \psi(v+j)],$$

and

$$(19) \quad \sum_{j=0}^{\infty} c_j(v, 1+p) t^j \sum_{u=1}^j u^{-1}.$$

Here, the meaningless sums $\sum_{j=0}^{-1}$ and $\sum_{u=1}^0$ are interpreted as zero.

The series (17), (18), and (19) can be approximated by their partial sums. We now show, for $0 < v < 1+p$ and $t > 0$, how to estimate the errors incurred in these approximations. These results, together with (16) and tables of the gamma function, allow us to calculate $U(v, 1+p; t)$, for such v, p , and t values, with known accuracy.

Denote by $[v]$ the largest integer that is less than v . By assumption $0 < v < 1+p$ and $t > 0$, so that

$$(20) \quad 0 < R_q(v, 1+p; t) < R_q(1+[v], 1+p; t).$$

Furthermore, since $\psi'(y) > 0$ and $\psi''(y) < 0$ for $y > 0$, we have

$$(21) \quad \begin{aligned} 0 &< \sum_{j=q}^{\infty} c_j(v, 1+p) t^j [\psi(1+j+p) - \psi(v+j)] \\ &< [\psi(1+q+p) - \psi(v+q)] R_q(v, 1+p; t) \end{aligned}$$

$$(22) \quad < [\psi(1+q+p) - \psi(v+q)] R_q(1+[v], 1+p; t).$$

Also, for $q \geq 1$,

$$(23) \quad \begin{aligned} 0 &< \sum_{j=q}^{\infty} c_j(v, 1+p) t^j \sum_{u=1}^j u^{-1} \\ &\leq \sum_{j=q}^{\infty} j c_j(v, 1+p) t^j \\ &= vt (1+p)^{-1} R_{q-1}(1+v, 2+p; t) \\ &\leq vt (1+p)^{-1} R_{q-1}(2+[v], 2+p; t). \end{aligned}$$

The recurrence relationship

$$(24) \quad w(w+1) \phi(y, w; t) - (w+1)(w+t) \phi(y, w+1; t) \\ + t(w-y+1) \phi(y, w+2; t) = 0;$$

together with the formulas

$$\phi(j, j; t) = e^t$$

and

$\phi(j, 1+j; t) = (-1)^j j! t^{-j} [1 - e^t e_{j-1}(-t)]$, $j = 1, 2, \dots$,
 where $e_q(t) = \sum_{j=0}^q t^j/j!$; can be used to construct convenient procedures for computing $R_q(1+[v], 1+p; t)$ and $R_{q-1}(2+[v], 2+p; t)$ and thus, in light of (20), (22), and (23), can be used for estimating errors resulting from the use of partial sums in place of the series (17), (18), and (19), and for determining the number of terms that need be included in each partial sum.

For many v , p , and t values, it may be advantageous to modify the outlined procedures for approximating the series (17), (18), and (19) with known accuracy. Existing tables, Kummer's transformations, recurrence relationships like (24), and/or other techniques from the theory of confluent hypergeometric functions (see e.g. [3] or [4]) can often be utilized to approximate $\phi(v, 1+p; t)$ and consequently the series (17) in more efficient fashion than the partial-sum approach described above. Our more efficient approximation for $\phi(v, 1+p; t)$ also yields $R_q(v, 1+p; t)$ which, upon applying (21), can also be used to estimate the error resulting from approximating (18) by one of its partial sums. Note that this error estimate is more precise than the one based on $R_q(1+[v], 1+p; t)$. Similarly, it may be possible to approximate $\phi(1+v, 2+p; t)$ and consequently $R_{q-1}(1+v, 2+p; t)$ to known accuracy with techniques not requiring the computation of $R_{q-1}(2+[v], 2+p; t)$; thus permitting more precise, and possibly easier, estimation of errors resulting from truncation of the series (19).

4. Evaluation of $G(r, s, a, b; k, \theta, x)$. Useful relationships.

Using (7), we find

$$G(r, s, a, b; k, \theta, x) = c^{-k} G(r, s, ca, cb; k, \theta/c, cx).$$

Upon substituting the expression (8) for g into the definition (5), we obtain

$$\begin{aligned} G(r, s, a, b; k, \theta, x) &= (b/a)(r-2)^{-1} \{ (a^{-1}+b^{-1}) G(r-2, s+2, a, b; k+1, \theta, x) \\ &\quad + (s-r+4) G(r-2, s+2, a, b; k, \theta, x) \\ &\quad + (b/a)(s+2) G(r-4, s+4, a, b; k, \theta, x) \}. \end{aligned}$$

Recurrence relations for G , corresponding to (9) - (13), can easily be obtained in the same way.

The domain of G can be extended by defining $G(r, 0, a, b; k, \theta, x)$ in terms of $g(r, 0, a, b; \cdot)$ through application of (5). Our recurrence relations for G remain valid. Thus, tables of the chi-square distribution function may prove useful in evaluating the distribution functions of X and Y and the moments of Y .

By making use of (1), (2), (4), and (6), we obtain

$$\begin{aligned} (25) \quad G(r, s, a, b; k, 0, \infty) &= (-1)^{k+1} G(s, r, b, a; k, 0, \infty) \\ &\quad + (2a)^k \sum_{j=0}^k \binom{k}{j} (-b/a)^j (s/2)_j (r/2)_{k-j} \end{aligned}$$

and

$$\begin{aligned} (26) \quad G(r, s, a, b; 0, \theta, \infty) &= (1 - 2a\theta)^{-r/2} (1 + 2b\theta)^{-s/2} \\ &\quad - G(s, r, b, a; 0, -\theta, \infty), \\ &\quad -1/(2b) < \theta < 1/(2a). \end{aligned}$$

Case (i): r an even integer. Using (15), we find

$$\begin{aligned} G(r, s, a, b; k, \theta, x) &= [a2^{r/2} \Gamma(r/2)]^{-1} [a/(a+b)]^{s/2} \\ &\quad \cdot \sum_{j=0}^{r/2-1} \binom{r/2-1}{j} (s/2)_j [2b/(a+b)]^j a^{-r/2+j+1} \\ &\quad \cdot \int_0^x t^{r/2+k-j-1} \exp[-t(1-2a\theta)/(2a)] dt \end{aligned}$$

$$\begin{aligned}
&= [2^{r/2} \Gamma(r/2)]^{-1} [a/(a+b)]^{s/2} a^k \\
&\quad \cdot \sum_{j=0}^{r/2-1} \binom{r/2-1}{j} (s/2)_j [2b/(a+b)]^j (1-2a\theta)^{-r/2-k+j} \\
&\quad \cdot \int_0^{x(1-2a\theta)/a} y^{r/2+k-j-1} e^{-y/2} dy \\
(27) \quad &= (2a)^k [a/(a+b)]^{s/2} \sum_{j=0}^{r/2-1} \frac{(s/2)_j (r/2-j)_k}{j!} \left(\frac{b}{a+b}\right)^j \\
&\quad \cdot (1-2a\theta)^{-r/2-k+j} P_{r+2k-2j}[x(1-2a\theta)/a];
\end{aligned}$$

for $r = 2, 4, 6, \dots$; $\theta < 1/(2a)$; $0 \leq x \leq \infty$.

In particular,

$$\begin{aligned}
(28) \quad G(r, s, a, b; k, \theta, \infty) &= (2a)^k [a/(a+b)]^{s/2} \sum_{j=0}^{r/2-1} \frac{(s/2)_j (r/2-j)_k}{j!} \\
&\quad \cdot \left(\frac{b}{a+b}\right)^j (1-2a\theta)^{-r/2-k+j};
\end{aligned}$$

for $r = 2, 4, 6, \dots$; $\theta < 1/(2a)$.

Applying the well-known relationship

$$P_{2v}(2x) = [\Gamma(v+1)]^{-1} x^v \phi(v, 1+v; -x)$$

to (27), we obtain, for $0 \leq x < \infty$, the alternate representation

$$\begin{aligned}
G(r, s, a, b; k, \theta, x) &= x^k [x/(2a)]^{r/2} [a/(a+b)]^{s/2} \\
&\quad \cdot \sum_{j=0}^{r/2-1} \frac{(s/2)_j}{j! (r/2-j-1)! (r/2+k-j)} \left(\frac{2ab}{x(a+b)}\right)^j \\
&\quad \cdot \phi[r/2+k-j, r/2+k-j+1; -x(1-2a\theta)/(2a)]; \\
&\quad r = 2, 4, 6, \dots; \quad \theta < 1/(2a).
\end{aligned}$$

Case (ii): s an even integer. By working with the joint probability distribution of the random variables W_1 and X , we obtain the representation

$$(29) \quad f_X(t) = (1/\beta) \int_{\max(0, t/\alpha)}^{\infty} p_m(u) p_n[(\alpha u - t)/\beta] du$$

for the probability density function of X .

Using (29), together with (4) and (6), we find; for $s = 2, 4, 6, \dots$;
 $-1/(2b) < \theta < 1/(2a)$; $0 < x < \infty$;

$$\begin{aligned}
G(r, s, a, b; k, \theta, x) &= (-1)^{k+1} G(s, r, b, a; k, -\theta, \infty) \\
&\quad + (1/b) \int_{-\infty}^x dt t^k e^{\theta t} \int_{\max(0, t/a)}^{\infty} p_r(u) p_s[(\alpha u - t)/b] du
\end{aligned}$$

$$\begin{aligned}
 &= (-1)^{k+1} G(s, r, b, a; k, -\theta, \infty) \\
 &\quad + a^k (a/b) \int_{-\infty}^{x/a} dv v^k e^{a\theta v} \int_{\max(0,v)}^{\infty} p_r(u) p_s[a(u-v)/b] du \\
 (30) \quad &= (-1)^{k+1} G(s, r, b, a; k, -\theta, \infty) \\
 &\quad + G_1(r, s, a, b; k, \theta, x) + G_2(r, s, a, b; k, \theta, x)
 \end{aligned}$$

where

$$\begin{aligned}
 G_1(r, s, a, b; k, \theta, x) &= a^k (a/b) \int_0^{x/a} du p_r(u) \int_{-\infty}^u v^k e^{a\theta v} p_s[a(u-v)/b] dv \\
 &= a^k \int_0^{x/a} du p_r(u) \int_0^{\infty} (u - by/a)^k e^{a\theta(u-by/a)} p_s(y) dy \\
 &= a^k \int_0^{x/a} du p_r(u) e^{a\theta u} \sum_{j=0}^k \binom{k}{j} u^j (-b/a)^{k-j} \int_0^{\infty} y^{k-j} e^{-b\theta y} p_s(y) dy \\
 &= (-2b)^k (1-2a\theta)^{-r/2} (1+2b\theta)^{-s/2-k} \\
 &\quad \cdot \sum_{j=0}^k \binom{k}{j} (r/2)_j (s/2)_{k-j} (-a/b)^j [(1+2b\theta)/(1-2a\theta)]^j P_{r+2j}[x(1-2a\theta)/a]
 \end{aligned}$$

and

$$\begin{aligned}
 G_2(r, s, a, b; k, \theta, x) &= a^k (a/b) \int_{x/a}^{\infty} du p_r(u) \int_{-\infty}^{x/a} v^k e^{a\theta v} p_s[a(u-v)/b] dv \\
 &= a^k (a/b)^{s/2} [2^{s/2} \Gamma(s/2)]^{-1} \int_{x/a}^{\infty} du p_r(u) e^{-au/(2b)} \\
 &\quad \cdot \sum_{j=0}^{s/2-1} \binom{s/2-1}{j} u^{s/2-j-1} (-1)^j \int_{-\infty}^{x/a} v^{k+j} e^{av(1+2b\theta)/(2b)} dv \\
 &= (-2b)^k [b/(a+b)]^{r/2} (1+2b\theta)^{-s/2-k} e^{x(1+2b\theta)/(2b)} \\
 &\quad \cdot \sum_{j=0}^{s/2-1} \frac{(r/2)_j (s/2-j)_k}{j!} \left(\frac{a}{a+b}\right)^j (1+2b\theta)^j \\
 &\quad \cdot e_{s/2+k-j-1}[-x(1+2b\theta)/(2b)] \{1 - P_{r+2j}[x(a+b)/(ab)]\}.
 \end{aligned}$$

Note that $G(s, r, b, a; k, -\theta, \infty)$, which appears on the right hand side of (30), can be evaluated from (28). Expression (28), together with (25) and (26), can also be used to evaluate $G(r, s, a, b; k, 0, \infty)$ and $G(r, s, a, b; 0, \theta, \infty)$ for $s = 2, 4, 6, \dots$

Case (iii): General case. Take

$$\begin{aligned}
 F(a, b, c; z) &= \frac{\Gamma(c)}{\Gamma(b) \Gamma(c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-tz)^{-a} dt, \\
 &\quad \operatorname{Re} c > \operatorname{Re} b > 0, \quad |\arg(1-z)| < \pi,
 \end{aligned}$$

to be the analytic continuation of the hypergeometric function.

For the Laplace transform of $y^{v-1} U(c, d; y)$, we have

$$\begin{aligned} \int_0^\infty e^{-zy} y^{v-1} U(c, d; y) dy \\ = \left\{ \frac{\Gamma(v) \Gamma(1+v-d)}{\Gamma(1+c+v-d)} \right\} F(v, 1+v-d, 1+c+v-d; 1-z) \end{aligned}$$

providing $\operatorname{Re} v > 0$, $\operatorname{Re} d < \operatorname{Re} v + 1$, $\operatorname{Re} z > 0$ (equation (3.2.51) of

Slater [3] and its extension by analytic continuation); so that

$$\begin{aligned} G(r, s, a, b; k, \theta, \infty) \\ = [z^{(r+s)/2} a^{r/2} b^{s/2} \Gamma(r/2)]^{-1} [2ab/(a+b)]^{r/2+s/2+k} \\ \cdot \int_0^\infty y^{r/2+s/2+k-1} \exp[-yb(1-2a\theta)/(a+b)] U(s/2, r/2+s/2; y) dy \\ = (2a)^k \left(\frac{a}{a+b}\right)^{s/2} \left(\frac{b}{a+b}\right)^{r/2+k} \frac{\Gamma(k+1) \Gamma(r/2+s/2+k)}{\Gamma(r/2) \Gamma(s/2+k+1)} \\ \cdot F[r/2+s/2+k, k+1, s/2+k+1; a(1+2b\theta)/(a+b)] \end{aligned}$$

for $\theta < 1/(2a)$.

By making use of linear transformation formulas for the hypergeometric function ((9.5.2) and (9.5.3) in Lebedev's book [1]), we obtain alternate representations

$$\begin{aligned} G(r, s, a, b; k, \theta, \infty) \\ = (2a)^k \left(\frac{a}{a+b}\right)^{s/2} \left(\frac{b}{a+b}\right)^{r/2-1} \frac{\Gamma(k+1) \Gamma(r/2+s/2+k)}{\Gamma(r/2) \Gamma(s/2+k+1)} (1-2a\theta)^{-k-1} \\ \cdot F[1-r/2, k+1, s/2+k+1; -(a/b)(1+2b\theta)/(1-2a\theta)], \end{aligned} \quad \theta < 1/(2a),$$

and

$$\begin{aligned} G(r, s, a, b; k, \theta, \infty) \\ = (2a)^k \left(\frac{a}{a+b}\right)^{s/2} \frac{\Gamma(k+1) \Gamma(r/2+s/2+k)}{\Gamma(r/2) \Gamma(s/2+k+1)} (1-2a\theta)^{-r/2-k} \\ \cdot F[1-r/2, s/2, s/2+k+1; a(1+2b\theta)/(a+b)], \end{aligned} \quad \theta < 1/(2a).$$

In particular, the representation (31) yields

$$\begin{aligned} G(r, s, a, b; k, 0, \infty) \\ = (2a)^k \left(\frac{a}{a+b}\right)^{s/2} \left(\frac{b}{a+b}\right)^{r/2-1} \frac{\Gamma(k+1) \Gamma(r/2+s/2+k)}{\Gamma(r/2) \Gamma(s/2+k+1)} \\ \cdot F(1-r/2, k+1, s/2+k+1; -a/b). \end{aligned} \quad (33)$$

This expression is an especially fruitful computational form.

For $r = 2, 4, 6, \dots$, it reduces to

$$(34) \quad G(r, s, a, b; k, 0, \infty) = (2a)^k \left(\frac{a}{a+b}\right)^{r/2+s/2-1} \cdot \sum_{j=0}^{r/2-1} \frac{(1-r/2-s/2-k)_j (r/2-j)_k}{j!} (-b/a)^j,$$

providing an alternative to (28) for computing purposes.

For $a/b \leq 1$ and $r = 1, 3, 5, \dots$, terms $(r-1)/2, (r+1)/2, (r+3)/2, \dots$ of the right hand side of

$$(35) \quad F(1-r/2, k+1, s/2+k+1; -a/b) = \sum_{j=0}^{\infty} \frac{(1-r/2)_j (k+1)_j}{j! (s/2+k+1)_j} (-a/b)^j$$

comprise a convergent alternating series, whose remainder is smaller in absolute value than the first neglected term and has the same sign; so that, by making use of the series representation (35), $G(r, s, a, b; k, 0, \infty)$ can be calculated to any desired accuracy.

In evaluating $G(r, s, a, b; k, 0, \infty)$ from (25) for $s = 2, 4, 6, \dots$, formula (34) can be used as an alternative to (28) for computing $G(s, r, b, a; k, 0, \infty)$. Similarly, if $a/b \geq 1$ and $s = 1, 3, 5, \dots$, $G(s, r, b, a; k, 0, \infty)$, which appears on the right hand side of the expression (25) for $G(r, s, a, b; k, 0, \infty)$, can be evaluated by making use of (31) and the series representation (35).

From (32), we obtain

$$G(r, s, a, b; k, 0, \infty) = (2a)^k (r/2+s/2)_k \sum_{j=0}^k \binom{k}{j} [(s/2)_j / (r/2+s/2)_j] \cdot (-1)^j (1+b/a)^j \{1 - P_{r,s+2j}[(b/a)(s+2j)/r]\},$$

so that this function can also be computed from tables of the F distribution function or the incomplete beta function.

For $0 < x < \infty$, we find

$$\begin{aligned} & 2^{(r+s)/2} a^{r/2} b^{s/2} \Gamma(r/2) \Gamma(s/2) G(r, s, a, b; k, \theta, x) \\ &= \int_0^\infty du u^{s/2-1} (1+u)^{r/2-1} \int_0^x t^{r/2+s/2+k-1} \exp\{-t[u(1+a/b)+1-2a\theta]/(2a)\} dt \end{aligned}$$

$$\begin{aligned}
&= (r/2+s/2+k)^{-1} x^{r/2+s/2+k} \exp[-x(1-2a\theta)/(2a)] \\
&\quad \cdot \int_0^\infty u^{s/2-1} (1+u)^{r/2-1} \exp[-ux(a+b)/(2ab)] \\
&\quad \cdot \Phi\{1, r/2+s/2+k+1; x[u(1+a/b)+1-2a\theta]/(2a)\} du \\
&= (r/2+s/2+k)^{-1} x^{r/2+s/2+k} \exp[-x(1-2a\theta)/(2a)] \\
&\quad \cdot \sum_{p=0}^\infty \frac{[x/(2a)]^p}{(\Gamma(r/2+s/2+k+1))_p} \sum_{j=0}^p \binom{p}{j} (1+a/b)^j (1-2a\theta)^{p-j} \\
&\quad \cdot \Gamma(s/2+j) U[s/2+j, r/2+s/2+j; x(a+b)/(2ab)] \\
&= x^{r/2+s/2+k} \exp[-x(1-2a\theta)/(2a)] \\
(36) \quad &\cdot \sum_{j=0}^\infty \frac{\Gamma(s/2+j)}{(\Gamma(r/2+s/2+k))_{j+1}} [x(a+b)/(2ab)]^j U[s/2+j, r/2+s/2+j; x(a+b)/(2ab)] \\
&\quad \cdot \Phi[j+1, r/2+s/2+k+j+1; x(1-2a\theta)/(2a)], \\
&\quad \theta < 1/(2a).
\end{aligned}$$

Using Kummer's transformation, we obtain an alternate representation

$$2^{(r+s)/2} a^{r/2} b^{s/2} \Gamma(r/2) \Gamma(s/2) G(r, s, a, b; k, \theta, x)$$

$$\begin{aligned}
&= x^{r/2+s/2+k} \sum_{j=0}^\infty \frac{\Gamma(s/2+j)}{(\Gamma(r/2+s/2+k))_{j+1}} [x(a+b)/(2ab)]^j \\
&\quad \cdot U[s/2+j, r/2+s/2+j; x(a+b)/(2ab)] \\
&\quad \cdot \Phi[r/2+s/2+k, r/2+s/2+k+j+1; -x(1-2a\theta)/(2a)], \\
&\quad 0 < x < \infty, \quad \theta < 1/(2a).
\end{aligned}$$

We now consider; for $r = 1, 3, 5, \dots$; $s = 1, 3, 5, \dots$; estimation of the error incurred in approximating the series appearing on the right hand side of (36) by a partial sum.

We will need the simple inequalities

$$\begin{aligned}
0 < \Gamma(c) U(c, d; t) < \Gamma(c-\delta) U(c-\delta, d; t), \\
t > 0, \quad c > c-\delta > 0,
\end{aligned}$$

and

$$\begin{aligned}
0 < \Phi(c, d; t) < e^t, \\
t > 0, \quad d > c > 0,
\end{aligned}$$

and formula (14).

Putting $r^* = r + 1$ and $s^* = s - 1$, we find that, for $r = 1, 3, 5, \dots$,
 $s = 1, 3, 5, \dots$, and for any positive integer v ,

$$\begin{aligned}
 (37) \quad & 0 < \sum_{j=v}^{\infty} [\Gamma(s/2+j)/(\Gamma(r/2+s/2+k))_{j+1}] [x(a+b)/(2ab)]^j \\
 & \quad \cdot U[s/2+j, r/2+s/2+j; x(a+b)/(2ab)] \Phi[j+1, r/2+s/2+k+j+1; x(1-2a\theta)/(2a)] \\
 & < \sum_{j=v}^{\infty} [\Gamma(s^*/2+j)/(\Gamma(r^*/2+s^*/2+k))_{j+1}] [x(a+b)/(2ab)]^j \\
 & \quad \cdot U[s^*/2+j, r^*/2+s^*/2+j; x(a+b)/(2ab)] \\
 & \quad \cdot \Phi[j+1, r^*/2+s^*/2+k+j+1; x(1-2a\theta)/(2a)] \\
 & < e^{x(1-2a\theta)/(2a)} \sum_{j=v}^{\infty} [\Gamma(s^*/2+j)/(\Gamma(r^*/2+s^*/2+k))_{j+1}] [x(a+b)/(2ab)]^j \\
 & \quad \cdot U[s^*/2+j, r^*/2+s^*/2+j; x(a+b)/(2ab)] \\
 & = [x(a+b)/(2ab)]^{-s^*/2} e^{x(1-2a\theta)/(2a)} \\
 & \quad \cdot \sum_{j=v}^{\infty} [(\Gamma(r^*/2+s^*/2+k))_{j+1}]^{-1} \sum_{p=0}^{r^*/2-1} \binom{r^*/2-1}{p} \Gamma(s^*/2+p+j) \\
 & \quad \quad \quad [x(a+b)/(2ab)]^{-p} \\
 (38) \quad & = [x(a+b)/(2ab)]^{-s^*/2} e^{x(1-2a\theta)/(2a)} \\
 & \quad \cdot \sum_{p=0}^{r^*/2-1} \binom{r^*/2-1}{p} [x(a+b)/(2ab)]^{-p} \\
 & \quad \cdot \left\{ \frac{\Gamma(s^*/2+p+1)}{(\Gamma(r^*/2+s^*/2+k))(\Gamma(r^*/2+k-p))} - \sum_{j=1}^{v-1} \frac{\Gamma(s^*/2+p+j)}{(\Gamma(r^*/2+s^*/2+k))_{j+1}} \right\}, \\
 & \quad \quad \quad 0 < x < \infty, \quad \theta < 1/(2a).
 \end{aligned}$$

For $r = 1, 3, 5, \dots$, $s = 3, 5, 7, \dots$, we note that the error
 bound (37) equals

$$\begin{aligned}
 (39) \quad & 2^{(r+s)/2} a^{r^*/2} b^{s^*/2} \Gamma(r^*/2) \Gamma(s^*/2) x^{-r/2-s/2-k} \\
 & \quad e^{x(1-2a\theta)/(2a)} G(r^*, s^*, a, b; k, \theta, x) \\
 & = \sum_{j=0}^{v-1} [\Gamma(s^*/2+j)/(\Gamma(r/2+s/2+k))_{j+1}] [x(a+b)/(2ab)]^j \\
 & \quad \cdot U[s^*/2+j, r/2+s/2+j; x(a+b)/(2ab)] \\
 & \quad \cdot \Phi[j+1, r/2+s/2+k+j+1; x(1-2a\theta)/(2a)].
 \end{aligned}$$

Here, $G(r^*, s^*, a, b; k, \theta, x)$ can be computed, for example, from (27);

$U[s^*/2+j, r/2+s/2+j; x(a+b)/(2ab)]$ can be calculated from (14); and

$\Phi[j+1, r/2+s/2+k+j+1; x(1-2a\theta)/(2a)]$ can be evaluated by using the recurrence
 relationship (24) together with its two accompanying formulas.

The error estimate (38) is less precise than (39); however it should prove easier to compute. Also, it can be calculated for $s = 1$.

To use (36) to approximate $G(r, s, a, b; k, \theta, x)$ for $r = 1, 3, 5, \dots$, $s = 1, 3, 5, \dots$, by replacing the infinite series with a partial sum, we must of course be able to compute the terms of the series. The evaluation of $\phi[j+1, r/2+s/2+k+j+1; x(1-2a\theta)/(2a)]$ can be accomplished as described above. A method for approximating $U[s/2+j, r/2+s/2+j; x(a+b)/(2ab)]$ to any desired accuracy can be found in Section 3 under Case (iii).

REFERENCES

- [1] LEBEDEV, N. N. (1965). Special Functions and Their Applications.
Prentice-Hall, Englewood Cliffs, New Jersey.
- [2] PRESS, S. J. (1966). Linear combinations of non-central chi-square
variates. Ann. Math. Statist. 37 480 - 487.
- [3] SLATER, L. J. (1960). Confluent Hypergeometric Functions.
Cambridge University Press, London.
- [4] SLATER, L. J. (1965). Confluent hypergeometric functions.
Handbook of Mathematical Functions (M. Abramowitz and
I. A. Stegun, eds.). Dover Publications, New York.
- [5] WANG, Y. Y. (1967). A comparison of several variance component
estimators. Biometrika 54 301 - 305.