

AROD-9559.1-C


AD 740178

University of Massachusetts
Department of Mechanical & Aerospace Engineering
Amherst, Massachusetts

Preliminary Experiments on the Gas-Phase Ignition Mechanism of
Ammonium Perchlorate

This work has been performed under
The Department of Army Research Grant
No. DA-ARO-31-124-71-G97 at
The Johns Hopkins University

Approved for public release; distribution
unlimited. The findings in this report are
not to be construed as an official Depart-
ment of the Army position, unless so desig-
nated by other authorized documents.



Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22151

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
The Johns Hopkins University		Unclassified	
		2b. GROUP	
		NA	
3. REPORT TITLE			
Preliminary Experiments on the Gas-Phase Ignition Mechanism of Ammonium Perchlorate			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Final Report: 1 March 1971 - 30 October 1971			
5. AUTHOR(S) (First name, middle initial, last name)			
Karl Jakus			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
1972		11	
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
DA-ARO-D-31-124-71-G97		NA	
b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
		9559.1-C	
c.			
d.			
10. DISTRIBUTION STATEMENT			
Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		U. S. Army Research Office-Durham Box CM, Duke Station Durham, North Carolina 27706	
13. ABSTRACT			
The results of the sublimation experiments that were performed and which were intended to be preparatory to the study of the gas-phase ignition mechanisms of Ammonium Perchlorate(A-F) are summarized in this report. These experiments were comprised of the study of the deposition rates from sublimating A-P as function of sublimation temperature, and the comparison of the mass spectra of the sublimating reagent grade A-P with that of already sublimed A-P. The results reinforce the previously reported observations that decomposition competes with sublimation in the temperature range of 100°C to 200°C. Furthermore, the results also indicate that the sublimed A-P is less stable than the reagent grade A-P, although the sublimation mass spectra from the two preparations are nearly indistinguishable.			
14. KEY WORDS			
Gas-Phase Ignition Mechanisms Ammonium Perchlorate			

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified

PRELIMINARY EXPERIMENTS ON THE GAS-PHASE IGNITION MECHANISM OF
AMMONIUM PERCHLORATE

Karl Jakus, University of Massachusetts

AROD Grant No. DA-ARO-31-124-71-G97

Abstract

The results of the sublimation experiments that were performed under the AROD Grant No. DA-ARO-31-124-71-G97 and which were intended to be preparatory to the study of the gas-phase ignition mechanisms of Ammonium Perchlorate(A-P) are summarized in this report. These experiments were comprised of the study of the deposition rates from sublimating A-P as function of sublimation temperature; and the comparison of the mass spectra of the sublimating reagent grade A-P with that of already sublimed A-P. The results reinforce the previously reported observations that decomposition competes with sublimation in the temperature range of 100°C to 200°C. Furthermore, the results also indicate that the sublimed A-P is less stable than the reagent grade A-P although the sublimation mass spectra from the two preparations are nearly indistinguishable.

FINAL REPORT

ARO-D Proposal Number: CRDARD-C-9559-C
Period Covered: 1 March 1971 - 31 August 1971
Title of Proposal: Study of Gas Phase Ignition Mechanisms with Skimming
Transient Mass Spectrometry
Grant Number: DA-ARO-D-31-124-71-G97
Name of Institution: The Johns Hopkins University
Author of Report: Karl Jakus

Scientific Personnel Supported by this Project:

1. Dr. Karl Jakus
Principal Investigator
1 March 1971 - 30 June 1971 : 25%
1 July 1971 - 31 August 1971: 100%
2. Stephen J. Thompson
Undergraduate at J.H.U.
410 hrs @ 2.50/hr.
3. Oliver Engel
Undergraduate at J.H.U.
86.5 hrs @ 2.50/hr.
4. Linda A. Ciotola
Undergraduate at J.H.U.
152 hrs @ 2.50/hr.
5. David R. Feineman
Undergraduate at J.H.U.
27 hrs @ 2.25/hr

Equipment Purchased
by this Project:

Bendix Model No. MA-3 Mass Spectroscope,
Serial No. S-12928
Total Cost: \$11,933.99

PRELIMINARY EXPERIMENTS ON THE GAS-PHASE IGNITION MECHANISM OF
AMMONIUM PERCHLORATE

Karl Jakus, University of Massachusetts

In May of 1971 Grant No. DA-AR0-31-124-71-G97 was awarded by the Department of Army to the Johns Hopkins University to study the gas phase ignition mechanisms of solid propellant oxydizers such as Ammonium Perchlorate (henceforth referred to as A-P). The primary goal of the research was charted to be the evaluation of the proposed "transient mass spectroscopy" in prospective of relevant propellant systems. Accordingly, the necessary apparatus has been constructed, a mass spectroscope has been selected, purchased and its performance in transient applications has been evaluated.

The technique under consideration entails the rapid vaporization in vacuum of a small sample of solid propellant to form an unsteadily expanding and simultaneously reacting gas cloud with an initial pressure and temperature corresponding to those likely to be encountered during actual propellant combustion. This gas cloud is then "skimmed" with a hypersonic orifice for mass spectroscopic analysis in a fashion similar to high pressure sampling. Previous theoretical¹ and experimental^{2,3} work on unsteady expansion of gas clouds as well as the extensive work on steady supersonic expansions⁴ serve as a foundation for the technique. The kind of flow field the "skimmer" orifice would experience in such an expansion process is illustrated in Figure 1 where the number flux and flow velocity in an expanding silver cloud are plotted against time.³ It is interesting to note that while the initial cloud density was approximately 5×10^{17} atoms/cm³ the density at the peak of the curve is only 1.5×10^{15} atoms/cm³ which represents, under isentropic conditions, a reduction in collision rate of 5×10^{-4} .

Such drastic reduction in collision rates may indeed be used to "freeze" the chemical reactions at various stages of development in a propellant gas cloud.

In the experimental program the gas phase decomposition of A-P was chosen as the first chemical system to be studied with the above "skimming" mass spectrometry. It was proposed that small samples of A-P would be deposited by sublimation on metallic filaments and subsequently, these samples would be vaporized rapidly by ohmic heating the filament. The preparation of these samples are essential to the study of the gas phase chemistry, consequently considerable attention has been devoted to the sublimation characteristics of A-P. The remainder of this report summarizes the results related to the sublimation portion of the program.

Two kinds of sublimation experiments were performed on A-P. First, the deposition rate of A-P onto a "filament" was measured and second, mass spectra were obtained from sublimating samples of reagent grade A-P as well as from samples of A-P that were collected as sublimate in previous experiments. Figure 2 shows the A-P deposition thicknesses as function of time and temperature that were obtained in vacuum on a 0.632" diameter nichrome wire. In the study related to the mass spectra, the sublimation and decomposition products emerging from a pyrex oven in vacuum were allowed to enter directly into the ionization chamber of a Bendix MA-3 time-of-flight mass spectroscope. The arrangement of the apparatus resembled those constructed by previous experimenters^{5,6}. Figure 3 shows, as function of temperature, the relative abundance of the predominant species present during the sublimation of reagent grade and sublimed A-P samples. Both of the samples exhibit an increase in the relative concentration of the decomposition products - O_2 , Cl_2 , NO_2 - and a decrease in the sublimation products

- HClO_4 , NH_3 - as the temperature increases from 100° to 200° C. Thus the previously reported observation that decomposition competes with sublimation at low temperatures is reinforced by the above results. Some differences exist, however, between the spectra of the two different A-P samples. Decomposition seems to overtake sublimation at a considerably lower temperature in the case of the sublimed A-P than it does in the case of the reagent grade A-P. This characteristic of the sublimed A-P corresponds to its previously reported⁷ shorter induction period. The following observations may be made concerning the results which are not illustrated on Figure 3. Both samples produced a spectrum in which the peaks 84 (HClO_3), 67 (ClO_2) and 51 (ClO) followed the parent peak of 100 (NClO_4) indicating that they, in fact, are ionization fragments of the perchloric acid. Furthermore, the peak 30 (NO) approximately followed the peak 46 (NO_2) as it was expected since NO is an ionization fragment of NO_2 ⁸. Figure 4 compares the concentration of O_2 (a decomposition product) and of HClO_4 (a sublimation product) of the herein discussed two samples with that of the experiments of Heath and Majer⁵. It is apparent from the graphs that Heath and Majer's results show marked similarity to the results obtained with the sublimed A-P sample. Figure 5 illustrates that, although the sublimed A-P sublimates itself its spectra show striking resemblance to the spectra of the decomposition products obtained in the event when all of the reagent grade A-P sample has vaporized out of the oven.

From the above results one may conclude that it is possible to deposit by sublimation small samples of A-P on "filaments" such that the chemical behavior of the sublimed A-P would be similar to A-P prepared by other methods although the sublimed compound seems to be relatively less stable.

BIBLIOGRAPHY

1. Mirels, H. and Mullen, J.F., "Expansion of Gas Clouds and Hypersonic Jets Bounded by a Vacuum", AIAA Journal, V. 1, No. 3, March 1963, 596-602.
2. Saltsburg, H., "Flash Evaporation", J. Chem. Phys., V. 42, No. 4, Feb. 1965, 1303-1309.
3. Jakus, K., "Study of the Transient Free Expansion of High Density Vapor Clouds", presented at the Annual Meeting of the Fluid Dynamics Division of the American Physical Society, Nov. 1970, Charlottesville, Virginia.
4. Chang, J. H., "Supersonic Molecular Beam Sampling System for Mass Spectrometric Studies of High Pressure Flow Systems", AIAA, Aerospace Science Meeting, 7th Paper No. 69-94, Jan. 1969.
5. Heath, G. A. and Majer, J. R., "Mass Spectrometric Study of the Thermal Decomposition of Ammonium Perchlorate", Trans. Faraday Soc., V. 60, 1964, 1783-1791.
6. Guillory, W. A., King, M. and Mack, J. L., "The Thermal Decomposition of Ammonium Chlorate", J. Phys. Chem., V. 73, No. 12, Dec. 1969, 4370-4374.
7. Hall, A. R. and Pearson, G. S., "Ammonium Perchlorate as an Oxidizer", Oxidation and Combustion Reviews, Elsevier, V. 3, 1968, 129-239.
8. Stenhagen, Abzahamsson and McLafferty, "Atlas of Mass Spectral Data", Interscience, 1969.




Figure 1. Aerodynamic Expansion of Silver Vapor (From Reference 3.)





Figure 2. Deposition of A-P onto a 0.032 inches Diameter Wire by Sublimation.





Figure 3. Relative Concentration of the Major Sublimation and Decomposition Products of A-P.







Figure 4. Relative Concentration of O_2 and of $HClO_4$ in the Mass Spectra of A-P.



SAMPLE: SUBLIMED A-P AT 165° C



SAMPLE: GASEOUS RESIDUE FROM REAGENT GRADE A-P AT 150°

Figure 5. Mass Spectra of Sublimed and Reagent Grade A-P Samples