


AD 740770

A FINAL REPORT TO
THE OFFICE OF NAVAL RESEARCH
on
INVESTIGATIONS OF THEORETICAL PROBLEMS
IN COSMICAL GAS DYNAMICS
Grant Number Nonr(G)-00005-66
Task Number NR 046-794

for the period
October 1, 1965 to December 31, 1970

February 10, 1971

Department of Astronomy and Astrophysics
University of Chicago
Chicago, Illinois 60637



Peter O. Vandervoort

Peter O. Vandervoort
Associate Professor of Astronomy
Principal Investigator

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22151

Reproduction in whole or in part is permitted for any purpose of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Department of Astronomy and Astrophysics The University of Chicago		2a. REPORT SECURITY CLASSIFICATION Unclassified 2b. GROUP
3. REPORT TITLE Theoretical Problems in Cosmical Gas Dynamics		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Final Report for the Period October 1, 1965 to December 31, 1970		
5. AUTHOR(S) (Last name, first name, initial) Vandervoort, Peter O.		
6. REPORT DATE February 10, 1971	7a. TOTAL NO. OF PAGES 9	7b. NO. OF REFS 13
8a. CONTRACT OR GRANT NO. NONR(G)-00005-66 b. PROJECT NO. NR 046-794 c. d.	9a. ORIGINATOR'S REPORT NUMBER(S) Final Report 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. AVAILABILITY/LIMITATION NOTICES This Document has been approved for Public Release and Sale; its distribution is unlimited.		
11. SUPPLEMENTARY NOTES None.	12. SPONSORING MILITARY ACTIVITY Geography Programs (Code 414) Office of Naval Research Arlington, Virginia 22217	
13. ABSTRACT This is the final report on investigations of theoretical problems in cosmical gas dynamics. An attempt has been made to derive a kinematical model of the Orion Nebula from observational models of the density distribution and data on the internal kinematics. Numerical methods have been developed for integrating the equations governing the dynamics of H II regions by the method of characteristics.		

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Gas Dynamics						
Interstellar Gas						
H II Regions						
Ionization Fronts						
Gaseous Nebulae						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report in d, y, month, year or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count shall follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, c, & d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, project number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER:** Enter the official report number by which the document will be identified and controlled by the originating activity. This number is unique to this report.

9b. **OTHER REPORT NUMBERS:** If the report has been identified by other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICE:** Enter any further dissemination of the report other than the

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

ABSTRACT

This is the final report on investigations of theoretical problems in cosmical gas dynamics. An attempt has been made to derive a kinematical model of the Orion Nebula from observational models of the density distribution and data on the internal kinematics. Numerical methods have been developed for integrating the equations governing the dynamics of H II regions by the method of characteristics.

Introduction

This is a Final Report to the Office of the Naval Research on work performed under Grant NONR(G)-00005-66 (Task Number NR 046-794) to the University of Chicago. Period of the work was October 1, 1965 to December 31, 1970. Work was interrupted during 1967-68, when the Principal Investigator (Peter O. Vandervoort) was on leave from the University of Chicago, and during 1969, as a result of service by the Principal Investigator on a University Disciplinary Committee.

Grant personnel were:

Peter O. Vandervoort, Principal Investigator
Stuart L. Mufson, Research Assistant

The work performed is described in the following two sections.

THE EXPANSION OF THE ORION NEBULAE

This investigation was an attempt to construct a kinematical model of the Orion Nebula on the basis of 1) Spherically symmetric models of the distribution of the electron density derived from optical and radio observations (Osterbrock and Flather 1959; Menon 1961) and 2) observational data on the internal kinematics of the nebula (Wilson, Munch, Flather, and Coffeen 1959). The nebula was treated as an expanding, spherically-symmetric system in which random motions of the gas are superimposed on the velocity of expansion. Under such conditions, the average radial velocity $\langle V(s) \rangle$ of the gas observed in a given spectral line along a line of sight at a projected distance S from the center of the nebulae is related to the velocity of expansion $u(r)$, where r is the true distance from the center, in the manner

$$\langle V(s) \rangle = \frac{\int \epsilon(r) u(r) \cos \theta e^{-\tau} dl}{\int \epsilon(r) e^{-\tau} dl} . \quad (1)$$

Here the integrations are performed along the line of sight through the nebulae, τ is the optical depth into the nebula, $\epsilon(r)$ is the volume emissivity in the spectral line, and the factor $\cos \theta$ projects the velocity of expansion

onto the line of sight. Values of $\langle V(s) \rangle$ were derived from the catalogue of Wilson, et al., while values of $\xi(r)$ were derived from the papers of Osterbrock and Flather and of Menon. Optical depths into the nebula were derived from results of an investigation by O'Dell and Hubbard (1955, private communication). The objective was then to find plausible functions $u(r)$ such that equation (1) would be satisfied over the range of s for which $\langle V(s) \rangle$ could be determined.

For the model envisaged, the expectation was that $u(r)$ should vanish at $r=0$, increase to a maximum value u_0 (say) at some distance $r=r_0$ (say), and fall to zero as $r \rightarrow \infty$. Dynamical considerations imply that u_0 should be of the order of the isothermal speed of sound (about 10 km-sec⁻¹) at most. It was found that equation (1) could be satisfied only if $u(r)$ attains a maximum value of the order of several hundred kilometers per second. This result is unacceptable because it violates the aforementioned dynamical considerations and because it implies that the dispersion of the radial velocities in the nebula should be much larger than is observed.

When this result was obtained, it was recognized that one possible source of error in the analysis lay in the assumption that the Orion Nebula cluster is at rest relative to the center of the Nebula. On this basis, $\langle V(s) \rangle$ had been determined relative to the cluster. This procedure would give erroneous

results for $u(\lambda)$ if the nebula were drifting relative to the cluster. Subsequently studies of the radio recombination lines of hydrogen revealed such a drift (Mezger and Hoglund 1967). Departures from spherical symmetry must be additional sources of error. Indeed, more recent investigations (e.g. Webster and Altenhoff 1970) now indicate that the Orion Nebula is much less regular and symmetric than was believed when this work was undertaken. For these reasons, it now appears that a reliable kinematical model cannot be derived along these lines.

NUMERICAL METHODS FOR THE INVESTIGATION
OF THE EVOLUTION OF H II REGIONS

It had been proposed under this Grant to investigate aspects of the early evolution of an H II region which had become of interest in earlier work (Vandervoort 1966). In particular, it was intended to consider further the effects of the finite time scale for the formation of the exciting star to include a more careful treatment of the ionization and heating of the gas in the formulation of the dynamics of the H II region. A central question was whether the early evolution of an H II region would be characterized by an ionization front of the R-type propagating directly into the undisturbed neutral gas or a front of the D-type following a shock into the undisturbed gas. And if the front were of the D-type, the question arose whether or not a rapid increase of the luminosity of the exciting star would induce a transition to a front of the R-type. This question had been raised earlier in an investigation of non-stationary ionization fronts (Vandervoort 1965 a,b).

It was clear at an early stage of the work that the development and use of numerical methods would be required. Accordingly, the thrust of the work shifted to the development of suitable numerical schemes for integrating the basic equations. At the time this work was undertaken, other investigators,

particularly Mathews (1965) and Lasker (1966), had investigated the evolution of H II regions by numerical integration of the hydrodynamic equations in a Lagrangian representation. It was thought that the present work might compliment those investigations if it were based on other numerical schemes. For this reason and because the Principal Investigator had had previous experience in the use of the method of characteristics (Courant and Friedrichs 1949), it was decided to develop numerical methods for integrating the basic equations by the method of characteristics.

In particular, such a scheme has been developed for the numerical integration of the equations governing the structure of a non-stationary ionization front (Vandervoort 1965a; all equation numbers in this section of this report refer to equations in that paper). Characteristic equations corresponding to equations (1), (2), (4), and (6) were derived in a conventional manner (Courant and Friedrichs 1949). These were replaced with a system of finite difference equations appropriate for a numerical integration by a "method of specified time intervals" (Lister 1960). At each time step in the integration, the flux must be determined by integrating equation (3); this was accomplished with the aid of a conventional scheme for numerical quadrature. The boundary conditions expressed by equations (8) - (10) were replaced with boundary conditions at the first and last points on the spatial grid.

A program for the execution of this method has been written, and it has been tested on the problem of a stationary ionization front for which an exact solution is known (see §IV in Vandervoort 1965a). Considerable experimentation was required to find satisfactory combinations of the number of points in the spatial grid and the size of the time step. The status of the work at the end of the grant period is that it has been possible to produce accurate solutions for the structures of stationary R-type fronts for intervals of time greater than the characteristic time scales during which elements of the neutral gas are over-run and completely ionized. These results encourage the belief that with further development these methods could be applied to non-stationary fronts and generalized for application to other aspects of the dynamics of H II regions. In view of the circumstance that such work would involve a considerable investment of computer time, for which funds are currently unavailable, work on this project was suspended at the end of the grant period.

REFERENCES

- Courant, R. and Friedrichs, K. O. 1948, Supersonic Flow and Shock Waves (New York: Interscience Publishers, Inc.), Chap. 2.
- Lasker, B. 1966, Ap. J., 143, 700.
- Lister, M. 1960, in Mathematical Methods for Digital Computers, ed. A. Ralston and H. S. Wilf (New York: John Wiley and Sons, Inc.), Chap. 15.
- Mathews, W. G. 1965, Ap. J. 142, 1120.
- Menon, T. K. 1961, Pub. N.R.A.O., 1, 1.
- Mezger, P. G. and Hoglund, B. 1967, Ap. J., 147, 490.
- O'Dell, C. R. and Hubbard, W. B. 1965, Ap. J., 142, 591.
- Osterbrock, D. E., and Flather, Edith 1959, Ap. J., 129, 26.
- Vandervoort, P. O. 1965a, Ap. J., 142, 507.
- _____. 1965b, ibid., 521.
- _____. 1966, ibid., 146, 104.
- Webster, W. J. and Altenhoff, W. J. 1970, Ap. Letters, 5, 233.
- Wilson, O. C., Munch, G., Flather, Edith, and Coffeen, Mary F. 1959, Ap. J. Suppl., 4, 199.