

AD-780 791

**EXPLICIT SOLUTIONS IN CONVEX GOAL
PROGRAMMING**

A. Charnes, et al

Carnegie-Mellon University

Prepared for:

Office of Naval Research

June 1973

DISTRIBUTED BY:

NTIS

**National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151**

**Best
Available
Copy**

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER MSRR No. 341		3. RECIPIENT'S CATALOG NUMBER AD-780 791	
4. TITLE (and Subtitle) Explicit Solutions in Convex Goal Programming		5. TYPE OF REPORT & PERIOD COVERED Technical Report-June 1973	
7. AUTHOR(s) A. Charnes D. Klingman W. W. Cooper R. J. Niehaus		6. PERFORMING ORG. REPORT NUMBER MSRR No. 341	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Graduate School of Industrial Administration Carnegie-Mellon University Pittsburgh, Pennsylvania 15213		8. CONTRACT OR GRANT NUMBER(s) N00014-67-A-0312-0007- N00014-67-A-0126-0008	
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research (Code 458) Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 047-048 NR 047-021	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1973	
		13. NUMBER OF PAGES 31	
		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) goal programming, linear programming, nonlinear programming, convex functional constrained regression, constrained generalized hypermedians, advanced starts, algorithms, explicit solution, manpower planning, organization design linear programming under uncertainty, multi-dimensional optimization, Markov processes			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Goal programming has now become an important tool in areas such as public management science. There is therefore a need for examining ways of securing improved computational efficiency, as is done in this paper, instead of resting only on the linear programming equivalences that were set forth when the original goal programming article was published in Vol. 1, No. 2 of <u>Management Science</u> . Based on lemmata which permit reduction of various im- portant classes of convex goal programming models to problems of full row			

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
U S Department of Commerce
Springfield VA 22151

rank-interval programming type, explicit solutions to convex goal programming problems are exhibited. Some of the equivalences herein established are also useful in their own right and for other classes of problems -- e.g., interval programming, as well as advanced start procedures and other such computational matters.

Unclassified

EXPLICIT SOLUTIONS IN CONVEX GOAL PROGRAMMING

by

A. Charnes*
W. W. Cooper**
D. Klingman*
R. J. Niehaus***


June, 1973



*University of Texas
**Carnegie-Mellon University, School of Urban and Public Affairs
***Office of Civilian Manpower Management, Department of the Navy

This report was prepared as part of the activities of the Management Sciences Research Group at Carnegie-Mellon University under Contract N00014-67-A-0312-0007 NR 047-048 with the U.S. Office of Naval Research. The research was also partly supported by ONR Contract No. NR 047-021, Task N00014-67-A-0126-0008, and the Center for Cybernetic Studies at the University of Texas. Reproduction in whole or in part is permitted for any purpose of the U.S. Government.

Management Sciences Research Group
Graduate School of Industrial Administration
Carnegie-Mellon University
Pittsburgh, Pennsylvania 15213



Background:

As material for possible historical (as well as motivational) interest, we may observe that Volume 1, No. 1 of Management Science contained an article entitled "The Stepping Stone Method for Explaining Linear Programming Calculations in Transportation Type Models." Volume 1, No. 2 contained an article entitled "Optimal Estimation of Executive Compensation by Linear Programming."¹ The purpose of these two papers was to help this new journal and the new society, TIMS, which published it get off to a start that would also provide a basis for further growth not only in themselves but also in a society (TIMS) which would, in turn, greatly enlarge the prospects for these and other related developments in the management sciences.

By emphasis at least, the first of these two articles was designed to appeal to immediate use and understanding. The emphasis in the second article was directed to longer range potentials for additional applications in the then new discipline of linear programming. Some of these applications included extensions to areas like "inequality constrained regression," "multi-dimensional objective optimizations" and their extensions to ordinal non-metric scaling, including non-Archimedean constructs.² New theory as well as new methodological possibilities were also naturally kept in mind, and exploited as their

1. See [2] and [3] in the bibliography appendix to this paper.

2. See [4].

potency for applications appeared to warrant. Thus, the transportation model in the first of these two articles was extended via a variety of new ideas which ranged from the development of the "poly - w method"³ to the use of model transformations and approximations. Ideas in the second article subsequently gave birth to attempts to exploit the use of linear programming models and methods to attack a variety of nonlinear problems.⁴ They also gave rise to the goal programming ideas which have now begun to be used in a variety of novel contexts. One such use, the OCMM⁵ series of models [12], has also involved a change in the state of modeling per se by joining the ideas of goal programming and Markov processes into a new approach for dealing with decision (Markov) processes.

As in these other developments, applications associated with the OCMM models have provided points which now make it possible to undertake further methodological and theoretical developments, as well as applications, that will provide still better bases for future extensions. This, in fact, is the point of the present paper, which will be followed by others, in which we shall undertake to join some of the preceding work (e. g., in model approximation and goal programming)

3. See [4].

4. See ~~Charnes and Lemke~~ [11], [13], [14].

5. OCMM (Office of Civilian Manpower Management, U. S. Navy) This series of models is described in [12].

not only with each other but also with presently disparate developments (e. g. , in linear programming under uncertainty) while we simultaneously attempt to provide solution procedures (including good starts) that can supply operational significance for the new problem areas that might then be addressed.

Introduction

Explicit solutions to linear programming or linear fractional programming problems presently exist only for special cases, the most general of which are the full row rank interval programming⁶ and the full row rank linear fractional programming.⁷

In some cases these models are of direct interest. This naturally includes those cases where problems can be modelled to fit the theoretical and computational requirements. It also includes cases for which model (structure) approximation and parameterization methods can yield either exact or approximate solutions to more complicated problems.⁸ Finally it includes cases in which advanced starts can be thus obtained, including ones which greatly accelerate the attainment of solutions--as when, e. g. , primal simplex technique applied to such starts have yielded great improvements in efficiency

6. See [1].

7. See [5].

8. See [6].

relative to alternative solution procedures that might otherwise have been required.⁹

There are at present no comparable classes of explicit solutions to "goal programming" models. The increased usage now being accorded to this type of model, however, would seem to justify such developments in order to anticipate benefits for this class which are analogous to those described in the preceding paragraph.

The present paper is directed toward developing such solution procedures for the general case of piecewise linear, but separable, "goal functionals." We define a separable functional as a sum of one-variable goal functionals. A one-variable goal functional, $f_j(x_j)$, is a function that is monotone decreasing for $x_j \leq g_j$ -- g_j = the stipulated j^{th} goal -- and monotone increasing for $x_j > g_j$. Note, however, that strict monotonicity is not required. In fact for the important "goal interval" type of model, the functional has a constant minimum over a whole interval.

The following examples are included in this class:

- (i) Absolute value functions, including those with asymmetric weights and multi-goal components.
- (ii) General convex piecewise linear functionals
- (iii) Goal interval functionals

9. See [10] and [8] ~~Charnes, Glover, Klingman, et. al.~~

- (iv) Hypermedian functionals and related functionals in extensions of ordinary goal programming.¹⁰

This list is not exhaustive, but it is nevertheless indicative of the many significant kinds of problems that are included for the development that we shall now provide as follows: First we shall state and prove a very general lemma which can be used to reduce and simplify such problems. This lemma will be presented in the next section--i.e., section 2--along with its application to a simple (but significant) class of cases. The lemma, we may note, is not restricted to the latter class of cases and is, indeed, established in far greater generality than required for any part of the present paper. In addition, we shall proceed to our results for obtaining explicit solutions via general separable convex piecewise linear functionals, which will be covered in section 3. In this section--i.e., section 3--we shall obtain a linear programming equivalent which will be extended in section 4 where procedures for obtaining explicit solutions will be delineated. Section 5 will then conclude this paper with a numerical example which will serve to illustrate some of the results achieved in the present paper and also indicate some of the possible extensions that will be covered in later papers in this series.

10. See [7] ~~Charnes, Cooper, Thompson~~ for exact definitions and elaborations.

2. A General Reduction Lemma:

The following lemma is established in far greater generality than we shall require for this paper. Indeed, no properties other than monotone decrease up to the goal and monotone increase to its right are stipulated. Thus, we let $f = \sum_j f_j(x_j)$ where $f_j(x_j)$ is monotone decreasing for $x_j \leq h_j$ and monotone increasing for $x_j \geq h_j$. Then we consider the following problem:

$$(1.1) \quad \text{Min. } f(x) = \sum_j f_j(x_j)$$

$$\text{subject to } \sum_j x_j \leq b_j$$

$$(1.2) \quad a_j \leq x_j \leq b_j$$

With this in mind we now develop

Lemma 1: If x satisfies (1.2) and for some j_0 , $x_{j_0} > h_{j_0}$, then $\bar{x}$ satisfies (1.2) and $f(\bar{x}) \leq f(x)$, where

$$\bar{x} = (\dots, \bar{x}_{j_0}, \dots),$$

$$\bar{x}_{j_0} = \begin{cases} x_{j_0}, & j \neq j_0 \\ \max(a_{j_0}, h_{j_0}), & j = j_0. \end{cases}$$

Proof: If x_{j_0} is replaced by $x_{j_0} - \Delta$, $0 \leq \Delta \leq \min.(x_{j_0} - h_{j_0}, x_{j_0} - a_{j_0})$, then

$$f_{j_0}(x_{j_0} - \Delta) \leq f_{j_0}(x_{j_0})$$

$$a_{j_0} \leq x_{j_0} - \Delta \leq b_{j_0}$$

$$\text{and } \sum_{j_0} x_j + x_{j_0} - \Delta \leq b_{j_0}, \text{ where } \hat{j}_0 \text{ signifies that the summation}$$

omits the variable with this index.

Thus, (1.2) is still satisfied. Also $f_{j_0}(x_{j_0})$ is monotone increasing for $h_{j_0} \leq x_{j_0}$. We therefore obtain a decrease (or at least no increase) by setting Δ at its upper limit $\bar{\Delta}$ and define $\bar{x}_{j_0} = x_{j_0} - \bar{\Delta} = \max(h_{j_0}, a_{j_0}) \leq x_{j_0}$. But $x_{j_0} > h_{j_0}$. Hence

$$f_{j_0}(\bar{x}_{j_0}) \leq f_{j_0}(x_{j_0})$$

and


$$\sum_j f_j(x_j) + f_{j_0}(\bar{x}_{j_0}) \leq f(x).$$

i.e.,


$$f(\bar{x}) \leq f(x),$$

and the lemma is thus established.


The lemma covers in particular nonlinear goal functionals of types such as those drawn in the following figures.



Non-Convex Functional
Figure 1a



Convex Functional
Figure 1b



Piecewise Linear Functional
Figure 1c

Note, further, also that the simple constraint $\sum x_j \leq b_0$ in the lemma may be replaced by any system of constraints such that decreasing any single variable preserves feasibility of the constraints. E. g., the system

$$5x_1 + 2x_2 + x_4 \leq 20$$

$$x_2 + 3x_3 + 5x_4 \leq 50$$

would be a valid type of replacement for the single constraint of the lemma.

See the illustration in Section 5.

We proceed first to consider, however, the particular case

$$(2.1) \quad \min \sum_j \mu_j |x_j - g_j|$$

subject to $\sum_j x_j \leq b_0$

$$(2.2) \quad a_j \leq x_j \leq b_j$$

where, without loss of generality, the constants μ_j are indexed so that $\mu_1 \geq \mu_2 \geq \dots \geq \mu_j \geq \dots \geq \mu_n \geq 0$. Because of Lemma 1, we can impose the additional condition

$$(3.0) \quad x_j \leq \max(g_j, a_j) \equiv b'_j.$$

Via (2.2), however, $a_j \leq x_j$ so that if $g_j \leq a_j$ we need only set $x_j = a_j$ and reduce the preceding problem to one in the remaining variables

$$(3.1) \quad \min \sum_j \mu_j |x_j - g_j|$$

subject to

$$(3.2) \quad \sum_j x_j \leq b'_0$$

$$a_j \leq x_j \leq b'_j$$

where b'_j is defined as in (3.0) and $a_j \leq g_j \leq b'_j$ with, again, $\mu_1 \geq \mu_2 \geq \dots \geq \mu_j \geq \dots \geq \mu_n \geq 0$, by renumbering, if required, and b'_0 is b_0 reduced by the $x_j = a_j$ values. Abusing notation, we replace n' by n too.

Because of Lemma 1 and the definition of b'_j , however, we now have $x_j - g_j \leq 0$ all j and this implies

$$|x_j - g_j| = g_j - x_j$$

Thus we may now replace the preceding problem by

$$(4.1) \quad \max \sum_j u_j x_j$$

$$\text{subject to } \sum_j x_j \leq b'_0$$

$$(4.2) \quad x_j \leq a_j.$$

This is an ordinary linear programming problem with a dual that may be written

$$(5.1) \quad \min v_0 b'_0 + \sum_j (v_j^+ b'_j - v_j^- a_j)$$

subject to

$$(5.2) \quad v_0 + v_j^+ - v_j^- = \mu_j$$

$$v_0, v_j^+, v_j^- \geq 0.$$

If $\sum_j a_j > b'_0$ then the constraints of (4.2) are evidently inconsistent since the a_j are lower bounds which the corresponding x_j must satisfy.

If we assume that $\sum_j a_j \leq b'_0$ then the optimum solution of (4.2) can be immediately written:

$$x_j^* = b'_j, \quad j=1, \dots, k-1$$

$$x_k^* = b'_0 - \sum_{j=1}^{k-1} b'_j - \sum_{j=k+1}^n a_j$$

$$x_j^* = a_j, \quad j = k+1, \dots, n$$

where k is the smallest positive integer such that

$$\sum_{j=1}^{k-1} b'_j \leq b'_0 - \sum_{j=k+1}^n a_j$$

and

$$\sum_{j=1}^k b'_j > b'_0 - \sum_{j=k+1}^n a_j$$

To see that these x^* choices are optimal we observe that they satisfy the constraints in (4.2) and produce $\sum_{j=1}^n \mu_j x_j^* = \sum_{j=1}^{k-1} \mu_j b_j' + \mu_k (b_0' - \sum_{j=1}^{k-1} b_j') - \sum_{j=k+1}^n a_j + \sum_{j=k+1}^n \mu_j a_j$ in the functional (4.1). Via regrouping¹¹ this becomes $\mu_k b_0' + \sum_{j=1}^{k-1} (\mu_j - \mu_k) b_j' + \sum_{j=k+1}^n (\mu_j - \mu_k) a_j$

Applying the "regrouping principle" (see [9]), we posit these coefficients as optimal dual variables and need only verify that they satisfy the dual constraints--viz.,

$$\begin{aligned} v_0^* &= \mu_k \\ v_j^{*+} &= \mu_j - \mu_k \geq 0, \quad j=1, \dots, k-1 \\ v_j^{*-} &= 0, \quad j=1, \dots, k-1 \\ v_k^{*+} &= v_k^{*-} = 0 \\ v_j^{*-} &= \mu_k - \mu_j \geq 0, \quad j=k+1, \dots, n \\ v_j^{*+} &= 0, \quad j=k+1, \dots, n. \end{aligned}$$

These values of v_0^* , v_j^{*+} , v_j^{*-} are non-negative as required. They also satisfy the other dual constraints in (5.2) since, as may be seen, these v^* choices give

$$\begin{aligned} \mu_k + \mu_j - \mu_k &= \mu_j & j=1, \dots, k-1 \\ \mu_k &= \mu_k \\ \mu_k + \mu_j - \mu_k &= \mu_j, & j=k+1, \dots, n \end{aligned}$$

upon substitution in (5.2).

11. I. e., we are here using a variant of the "regrouping principle" first set forth in [9] as a means of obtaining explicit solutions via the dual problem. See also [4].

This completes the proof that our explicit solution x^* is optimal. It also shows in full detail how we can use the lemma to obtain explicit solutions in this class of cases. The next section will extend this result to the case of general separable convex piecewise linear functionals.

5. The General Separable Convex Piecewise Linear Functional

The functional we are considering is separable--i.e., it is a sum of functions each involving only one variable. Hence, we will be able to apply Lemma 1 and perform a reduction on one variable at a time to obtain a linear programming format. The explicit solution will then be developed and exhibited in the section after this one.

The general (continuous) one variable case has a graph consisting of straight line segments with slopes that increase from left to right. See Figure 1c. As is well known classically, such a function can be written as

$$(6.1) \quad f^a(t) = \sum_j \mu_j |t - g_j| + pt + q$$

where the g_j , μ_j , p and q are given constants with $\mu_j \geq 0$, all j .

In this form we state the following:

Lemma 2: A necessary and sufficient condition for $f^a(t)$ with $\mu_j \geq 0$, $\forall_j$, to be a goal functional is that $\sum_j \mu_j \geq |p|$.

Consider then

$$f^a(t) = \sum_{j=1}^n \mu_j |t - g_j| + pt + q$$

and

$$f^s(t) = \sum_{j=1}^{n+1} k_j t_j + d, \quad k_j \leq k_{j+1}, \quad \forall j.$$

The graph of $f^s(t)$ is shown in Figure 2. Now consider

$$t_1 \leq g_1$$

$$0 \leq t_j \leq g_j - g_{j-1}$$

$$0 \leq t_{n+1}$$

$$\text{For } t \rightarrow \infty: \quad f^a(t) \sim \left(\sum_{j=1}^n \mu_j + p \right) t_{n+1}$$

$$f^s(t) \sim k_{n+1} t_{n+1}$$

Hence

$$p + \sum_{j=1}^n \mu_j = k_{n+1}$$

$$\text{For } t \rightarrow -\infty: \quad f^a(t) \sim \left(-\sum_{j=1}^n \mu_j + p \right) t.$$

$$f^s(t) \sim k_1 t_1$$

Hence

$$p - \sum_{j=1}^n \mu_j = k_1$$

Therefore

$$p = 1/2 (k_1 + k_{n+1}), \quad \sum_{j=1}^n \mu_j = 1/2 (k_{n+1} - k_1)$$

Yet, for $g_{j-1} \leq t \leq g_j$, we have $t = g_{j-1} + t_j$, so

$$f^a(t) = \sum_{r=1}^{j-1} \mu_r (g_{j-1} + t_j - g_r) + \sum_{r=j}^n \mu_r (g_r - g_j - t_j) + p(g_{j-1} + t_j) + q$$

$$f^S(t) = \sum_{r=1}^{j-1} k_r (g_r - g_{r-1}) + k_j t_j + d_1,$$

where $g_0 = 0$.

Thus

$$\begin{aligned} f^A(t) &= \left(\sum_{r=1}^{j-1} \mu_r - \sum_j \mu_r + p \right) t_j + \left(\sum_{r=1}^{j-1} \mu_r - \sum_j \mu_r + p \right) g_{j-1} + \\ &\quad + \sum_{r=1}^{j-1} \mu_r g_r + \sum_j \mu_r g_r + q \\ f^S(t) &= k_j t_j + \sum_{r=1}^{j-1} k_r (g_r - g_{r-1}) + d_1 \end{aligned}$$

For $t = t_1 \leq g_1$:

$$\begin{aligned} f^A(t) &= (p - \sum_1^n \mu_j) t_1 + \sum_1^n \mu_j + q \\ f^S(t) &= k_1 t_1 + d_1 \\ k_1 &= p - \sum_1^n \mu_j; \quad d_1 = q + \sum_1^n \mu_j \end{aligned}$$

From before:

$$k_{n+1} = p + \sum_1^n \mu_j \quad p = 1/2 (k_1 + k_{n+1}), \quad \sum_1^n \mu_j = 1/2 (k_{n+1} - k_1)$$

$$\text{For } t = g_{j-1} + t_j: \quad k_j = p + \sum_{r=1}^{j-1} \mu_r - \sum_j \mu_r$$

$$\text{For } t = g_{j-2} + t_{j-1}: \quad k_{j-1} = p + \sum_{r=1}^{j-2} \mu_r - \sum_{j-1}^n \mu_r$$

$$\text{Hence } k_j - k_{j-1} = 2\mu_{j-1}$$

Thus, given $k_1, \dots, k_{n+1}, g_1, \dots, g_n; d_1$ in the $f^S(t)$ form, we can determine the $f^A(t)$ form via

$$p = 1/2 (k_1 + k_{n+1})$$

$$g = d_1 - 1/2 (k_{n+1} - k_1)$$

$$\mu_j = 1/2 (k_{j+1} - k_j)$$

Conversely, given the $f^a(t)$ form with $\mu_1, \dots, \mu_n$; p, q and $\xi_1, \dots, \xi_n$, we can determine $d_1, k_1, \dots, k_{n+1}$ for the $f^s(t)$

form via

$$d_1 = q + \sum_{j=1}^n \mu_j$$

$$k_1 = p - \sum_{j=1}^n \mu_j$$

$$k_j = p + \sum_{s=1}^{j-1} \mu_s - \sum_{s=j}^n \mu_s = k_{j-1} + 2\mu_{j-1}$$




Figure 2

Because we are dealing with a piecewise linear and convex problem, we have been able to reduce the portion corresponding to one variable t to a separable linear form on the t_j with individual interval constraints. Going back to our total functional (which is the sum of functions of one variable each) and splitting each variable whose functional is nonlinear, into a sum, as above, we have thereby reduced the solution of the general convex problem to that of a linear program. The solution of the latter will be exhibited in the following section.

4.1 Explicit Solution

Because of the equivalence we have developed in the last section between functions in $f^a(t)$ and $f^s(t)$ format, we can evidently always write the general convex piecewise linear separable functional (with a finite number of pieces) as a linear functional subject to internal constraints on the "incremental" variables. We shall therefore take our candidate for explicit solution in the reduced (equivalent) form:

$$\begin{aligned}
 (8.1) \quad & \min k^{(1)T} x^{(1)} + k^{(2)T} x^{(2)} \\
 & d^{(1)T} x^{(1)} + d^{(2)T} x^{(2)} \leq b_0 \\
 & a^{(1)} \leq I x^{(1)} \leq b^{(1)} \\
 & a^{(2)} \leq A x^{(2)} \leq b^{(2)}
 \end{aligned}$$

in which we now use vector-matrix notation with

$$k^{(1)T} = (k_1^{(1)}, \dots, k_{n_1}^{(1)})$$

$$k^{(2)T} = (k_1^{(2)}, \dots, k_{n_2}^{(2)})$$

$$x^{(1)T} = (x_1^{(1)}, \dots, x_{n_1}^{(1)})$$

$$x^{(2)T} = (x_1^{(2)}, \dots, x_{n_2}^{(2)})$$

$$A = (a_{ij}) \text{ is of full row rank}$$

Note that if A is of full row rank, so is a matrix $\begin{pmatrix} I & A \end{pmatrix}$. Thus the above, which we have written to separate out that part of the constraints and functional which corresponds to the non-linear goals, can be rendered in the form

$$\begin{aligned} & \min k^T x \\ (8.2) \quad & d^T x \leq b_0 \\ & \bar{a} \leq \bar{A}x \leq \bar{b} \end{aligned}$$

where $\bar{A}$ is of full row rank. We can make the transformation $y = \bar{A}x$,

or $x = \bar{A}^\# y + (I - \bar{A}^\# \bar{A}) z$, where $\bar{A}^\#$ is a generalized inverse of $\bar{A}$ (e.g., a right inverse) and z is an arbitrary vector.

The problem may now be rewritten

$$(9.1) \quad \begin{aligned} & \min k^T \bar{A}^\# y + k^T (I - \bar{A}^\# \bar{A}) z \quad \text{with} \\ & d^T \bar{A}^\# y + d^T (I - \bar{A}^\# \bar{A}) z \leq b_0 \\ & \bar{a} \leq y \leq \bar{b} \end{aligned}$$

Now let

$$n_j = \begin{cases} d^T \bar{A}_j^\# y_j, & \text{if } d^T \bar{A}_j^\# \neq 0 \\ 0, & \text{if } d^T \bar{A}_j^\# = 0 \end{cases}$$

$$s_k = \begin{cases} d^T (I - \bar{A}^\# \bar{A})_k z_k, & \text{if } d^T (I - \bar{A}^\# \bar{A})_k \neq 0 \\ z_k, & \text{if } (I - \bar{A}^\# \bar{A})_k = 0 \end{cases}$$

Then the problem may be written

$$(9.2) \quad \begin{aligned} & \min \hat{k}^T n + \hat{k}^T s \\ & \text{e.t. } \hat{e}^T n + \hat{e}^T s \leq b_0 \\ & \hat{a} \leq n \leq \hat{b} \end{aligned}$$

where

$$(9.3) \quad \begin{aligned} \hat{e}_j &= \begin{cases} 1, & \text{if } d^T \bar{A}_j^\# \neq 0 \\ 0, & \text{if } d^T \bar{A}_j^\# = 0 \end{cases} \\ \hat{e}_k &= \begin{cases} 1, & \text{if } d^T (I - \bar{A}^\# \bar{A})_k \neq 0 \\ 0, & \text{if } (I - \bar{A}^\# \bar{A})_k = 0 \end{cases} \\ \hat{k}_j &= \begin{cases} k^T \bar{A}_j^\# / d^T \bar{A}_j^\#, & \text{if } d^T \bar{A}_j^\# \neq 0 \\ k^T \bar{A}_j^\#, & \text{if } d^T \bar{A}_j^\# = 0 \end{cases} \end{aligned}$$

$$\hat{k}_r = \begin{cases} k^T(I - \bar{A}^{\#}\bar{A})_r / d^T(I - \bar{A}^{\#}\bar{A})_r & \text{if } d^T(I - \bar{A}^{\#}\bar{A})_r \neq 0 \\ k^T(I - \bar{A}^{\#}\bar{A})_r & \text{if } d^T(I - \bar{A}^{\#}\bar{A})_r = 0 \end{cases}$$

$$\hat{a}_j = \begin{cases} d^T \bar{A}_j^{\#} a_j, & \text{if } d^T \bar{A}_j^{\#} > 0 \\ a_j, & \text{if } d^T \bar{A}_j^{\#} = 0 \\ d^T \bar{A}_j^{\#} b_j, & \text{if } d^T \bar{A}_j^{\#} < 0 \end{cases}$$

$$\hat{b}_j = \begin{cases} d^T \bar{A}_j^{\#} b_j, & \text{if } d^T \bar{A}_j^{\#} > 0 \\ b_j, & \text{if } d^T \bar{A}_j^{\#} = 0 \\ d^T \bar{A}_j^{\#} a_j, & \text{if } d^T \bar{A}_j^{\#} < 0 \end{cases}$$

Combining the components of n and s into a single vector

$\xi^T = (\xi_1, \dots, \xi_n)$ in the order of decrease of their functional coefficients

$\hat{k}_j$, $\hat{k}_r$, and adjoining $-\infty$, $+\infty$ lower and upper bounds for the ξ_r which are

S_r , s , the problem may be written in the form

$$(10.1) \quad \min \quad \tilde{k}^T \xi$$

$$\text{with } \tilde{c}^T \xi \leq b_0$$

$$\tilde{a} \leq \xi \leq \tilde{b}$$

where $\tilde{k}_1 \leq \tilde{k}_2 \leq \dots \leq \tilde{k}_n$, the $\tilde{a}_j$ correspond to $\hat{a}_j$'s or $-\infty$, the $\tilde{b}_j$ cor-

respond to $\hat{b}_j$'s or $+\infty$, the $\tilde{e}_j$ are the appropriate $\hat{e}_j$ or $\hat{e}_k$. For consistency we

must have $\sum_j^n \tilde{e}_j \tilde{a}_j \leq b$. Setting $\bar{\xi}_j = \tilde{a}_j + \tilde{e}_j$, the problem may be rewritten,

$$\begin{aligned} & \max. \quad \sum_j (-\tilde{k}_j) \bar{\xi}_j + \sum_j (-\tilde{k}_j) \tilde{a}_j \\ (10.2) \quad & \text{with} \quad \sum_j \tilde{e}_j \bar{\xi}_j \leq b_0 - \sum_j \tilde{e}_j \tilde{a}_j \equiv b'_0 \end{aligned}$$

$$\bar{\xi}_j \leq \tilde{b}_j - \tilde{a}_j \equiv \tilde{b}'_j$$

$$\bar{\xi}_j \geq 0.$$

This is a linear programming problem. It therefore has the following dual

$$\begin{aligned} & \min. \quad w_0 b'_0 + \sum_j w_j \tilde{b}'_j \\ (10.3) \quad & \text{with} \quad w_0 \tilde{e}_j + w_j \geq -\tilde{k}_j \\ & w_0, w_j \geq 0. \end{aligned}$$

We now introduce

$$\begin{aligned} (11) \quad J_0 & \equiv \{j: -\tilde{k}_j \geq 0 \text{ and } \tilde{e}_j = 0\} \\ J_1 & \equiv \{j: -\tilde{k}_j \geq 0 \text{ and } \tilde{e}_j = 1\} \end{aligned}$$

$$r = \max j \text{ such that } -\tilde{k}_j \geq 0, \tilde{e}_j = 1, \sum_{s=1}^r \tilde{e}_s \tilde{b}'_s \leq b'_0$$

and shall show that an optimal solution to the primal problem is

$$\begin{aligned} & \bar{\xi}_j^* = \tilde{b}'_j, \quad j \in J_0 \\ (12) \quad & \bar{\xi}_j^* = \tilde{b}'_j, \quad j \in J_1 \\ & \bar{\xi}_j^* = b_0 - \sum_{j \in J_1} \tilde{e}_j \tilde{b}'_j = b'_1 - \sum_{j \in J_1 \cup J_0} \tilde{e}_j \tilde{b}'_j \\ & \bar{\xi}_j^* = 0, \quad j \in \{r\} \cup J_0 \cup J_1 \end{aligned}$$

where $\{r\}$ is the singleton set with element r . Then

$$(13) \quad \sum_j -\tilde{k}_j \tilde{\xi}_j^* = \sum_{j_0} (-\tilde{k}_j) \tilde{b}_j + \sum_{j_1} (-\tilde{k}_j) \tilde{b}_j + (-\tilde{k}_r) (b_0' - \sum_{j_0 \cup j_1} \tilde{b}_j) \\ = (-\tilde{k}_r) b_0' + \sum_{j_0} [(-\tilde{k}_j) - \tilde{e}_j (-\tilde{k}_r)] \tilde{b}_j + \text{terms under } j_1.$$

We now employ the regrouping principle and posit the following values

for the dual variables: $w_0^* = -\tilde{k}_r$, $w_j^* = -\tilde{k}_j - e_j(-\tilde{k}_r)$, $j \in j_0 \cup j_1$ $w_j^* = 0$ otherwise

Noting that these are all non-negative, we check that these w^* satisfy the other dual constraints--viz.,

$$(14.1) \quad w_0^* \tilde{e}_j + w_j^* = (-\tilde{k}_r) \tilde{e}_j + [-\tilde{k}_j - \tilde{e}_j(-\tilde{k}_r)] = -\tilde{k}_j \text{ for } j \in j_0 \cup j_1$$

Thus the $j_0 \cup j_1$ dual constraints are satisfied. Next,

$$(14.2) \quad w_0^* \tilde{e}_j + w_j^* = (-\tilde{k}_r) \tilde{e}_j \text{ for } j \notin j_0 \cup j_1$$

We need to show here that $(-\tilde{k}_r) \tilde{e}_j \geq -\tilde{k}_j$. If $-\tilde{k}_j \geq 0$ then $\tilde{e}_j = 1$ and $j > r$ so

that $-\tilde{k}_r \geq -\tilde{k}_j$. If $-\tilde{k}_j < 0$, then since $(-\tilde{k}_r) \tilde{e}_j \geq 0$ we have $(-\tilde{k}_r) \tilde{e}_j \geq \tilde{k}_j$.

Thus all dual conditions are satisfied.

Passing back from $\tilde{\xi}$ to ξ , our optimal solution for (10.1), is therefore

$$\xi_j^* = \tilde{b}_j, j \in j_0 \cup j_1 \\ (15) \quad \xi_r^* = b_0 - \sum_{j_0 \cup j_1} \tilde{e}_j \tilde{b}_j - \sum_{j \notin j_0 \cup j_1} \tilde{e}_j \tilde{a}_j \\ \xi_j^* = \tilde{a}_j, j \notin j_0 \cup j_1$$

4.2 Further Computational Reduction:

For simplicity of exposition the general separable convex piecewise linear functional was transformed directly into linear interval programming format employing additional variables, one for each of the linear segments. Our fundamental lemma, however, makes it possible to reduce (sometimes very substantially) the number of segments which need to be considered and also reduces the number of variables and constraints which need to be introduced. Because of this lemma, we can restrict an original variable in the goal functional to be bounded above by the first g_j at which the goal functional minimum over $a \leq t \leq b$ is attained.

The use of this technique, together with other reductions that are possible for special goal programming functionals, will be presented in additional reports in this series together with applications involving various utilizations of advance starts. As a start toward that end we have supplied a numerical illustration which illustrates some of the preceding developments.

5. Numerical Illustration:

We now conclude the present paper with a numerical example which illustrates the remarks we have just made. For this purpose we adapt an example from the paper which initiated this series of studies

in manpower planning.

Applying the procedure of model approximation sketched in the next to last paragraph on page II-8 of [14] to a curtailment of that example we have as our problem

$$\min. |N_1(1) - 30| + |N_2(1) - 200| + |N_1(2) - 70| + |N_2(2) - 300|$$

subject to

$$0 \leq N_1(1) \leq \infty$$

$$0 \leq N_2(1) \leq \infty$$

$$(16) \quad 15N_1(1) + 13N_2(1) \leq 3,000$$

$$44 \leq N_1(2) \leq \infty$$

$$143 \leq N_2(2) \leq \infty$$

$$15N_1(2) + 13N_2(2) \leq 4,000$$

Here we have confined ourselves to a model involving only two types of manpower in a 2-period plan. Thus, the variable $N_i(t)$ refers to the amount of the i^{th} type of manpower, $i = 1, 2$, scheduled for recruitment in period $t = 1, 2$. The admissible values for these variables are in interval form except for the two budgetary constraints which are applicable at the indicated salary levels of 15 and 13(thousand dollars/annum) in each period. Essentially then this is all in the form of the expressions given in our General Lemma (see section 2, above) with the goals of 30, 200, 70 and 300 stipulated in the absolute value functional for (16). By our Lemma 1, the above example may be transformed into

maximize $N_1(1) + N_2(1) + N_1(2) + N_2(2)$
subject to

$$0 \leq N_1(1) \leq 30$$

$$0 \leq N_2(1) \leq 200$$

$$(17) \quad 15N_1(1) + 13N_2(1) \leq 3,000$$

$$44 \leq N_1(2) \leq 70$$

$$143 \leq N_2(2) \leq 300$$

$$15N_1(2) + 13N_2(2) \leq 4,000$$

We now have only an ordinary linear programming problem that involves no more variables than the nonlinear problem (16). Furthermore, (17) may be split into the following two separate subproblems.

maximize $N_1(1) + N_2(1)$ subject to

$$0 \leq N_1(1) \leq 30$$

$$(18.1) \quad 0 \leq N_2(1) \leq 200$$

$$15N_1(1) + 13N_2(1) \leq 3,000$$

and

maximize $N_1(2) + N_2(2)$ subject to

$$44 \leq N_1(2) \leq 70$$

$$143 \leq N_2(2) \leq 300$$

$$15N_1(2) + 13N_2(2) \leq 4,000$$

Without further ado we can then write our optimal solution for

(17), and hence (16), via (18.1) and (18.2) This gives

$$N_1^*(1) = 400/15, N_2^*(1) = 200$$

$$N_1^*(2) = 44, N_2^*(2) = 143 + \frac{1481}{13} = 257.$$

Evidently our lemma has produced a simplification so that achieving a solution to the nonlinear problem (16) involved practically nothing more than merely a solution by inspection of two much smaller linear problems. This suggests further possible ways in which the ideas contained in our lemma may be extended to other classes of problems. Such developments are best delayed, however, until they can be treated in their own right via other papers in this series.

BIBLIOGRAPHY

- [1] Ben-Israel, A., and A. Charnes "An Explicit Solution of a Special Class of Linear Programming Problems," Operations Research, Vol. 16, No. 6, Nov.-Dec. 1968, pp. 1166-1175.
- [2] Charnes, A. and W. W. Cooper, "The Stepping Stone Method for Explaining Linear Programming Calculations in Transportation Type Models," Management Science, Vol. 1, No. 1, October 1954, pp. 49-69.
- [3] _____, _____, and R. F. Froyen, "Optimal Estimation of Executive Compensation by Linear Programming," Management Science, Vol. 1, No. 2, January 1955, pp. 138-151.
- [4] _____, _____, Management Models and Industrial Applications of Linear Programming (New York: John Wiley & Sons, Inc., 1961).
- [5] _____, _____, "An Explicit General Solution in Linear Fractional Programming," Naval Research Logistics Quarterly, Vol. 20, No. 3, September 1973, pp. 449-467.
- [6] _____, _____, "A Strategy for Making Models by Linear Programming," R. Machol et. al., ed. Systems Engineering Handbook (New York: McGraw-Hill Book Co., Inc.).
- [7] _____, _____, and G. L. Thompson, "Constrained Generalized Medians and Hypermedians As Deterministic Equivalents for Two-Stage Linear Programs Under Uncertainty," Management Science, Vol. 12, No. 1, September 1965.
- [8] _____, F. Glover, D. Karney, D. Klingman and J. Stutz, "Past, Present, & Future of Development, Computational Comparison, and Use of Large Scale Transportation and Network Computer Codes," Research Report CS 131, Center for Cybernetic Studies, University of Texas at Austin, Texas, July 1973.
- [9] _____, W. W. Cooper, and B. Mellon, "A Model For Programming and Sensitivity Analysis in an Integrated Oil Company," Econometrica, Vol. 22, No. 4, April 1954.
- [10] Glover, F., D. Karney, D. Klingman, A. Napier "A Computational Study on Start Procedures, Basis Change Criteria, and Solution Algorithms for Transportation Problems," Research Report CS 93 (Austin, Texas: Center for Cybernetic Studies, The University of Texas Graduate School of Business, 1972).

- [11] Charnes, A., C. E. Lemke "Minimization of Non-linear Separable Convex Functionals," Naval Research Logistics Quarterly, Vol. 1, No. 4, December 1954.

- [12] _____, W. W. Cooper and R. J. Niehaus, Studies in Manpower Planning, (Washington: U.S. Navy, Office of Civilian Manpower Management, 1972).

- [13] Ijiri, Y. Management Goals and Accounting for Control (Chicago: Rand-McNally, Inc., 1964).

- [14] Lee, Sang M. Goal Programming for Decision Analysis, (Philadelphia: Auerbach Publishers, 1972).