

UNCLASSIFIED


AD NUMBER
AD825979
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; DEC 1967. Other requests shall be referred to Bureau of Medicine and Surgery, Washington, DC.
AUTHORITY
USNMRU No. 4 ltr, 14 May 1971

THIS PAGE IS UNCLASSIFIED

AD825979

NAVAL MEDICAL RESEARCH

UNIT No. 4



ALPHA-1 GLOBULINS IN THOSE SUSCEPTIBLE TO VIRAL ACUTE RESPIRATORY DISEASES

R. I. Lytle and M. J. McNamara

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of *Naval Medical Research*
Guantanamo, C.R. 60088

DDC
RECEIVED
DEC 1 1968
NEC

TO: [REDACTED]
FROM: [REDACTED]
SUBJECT: [REDACTED]

MR 005.20-0013

RU 66.28



December 1967

Best Available Copy

This document is for official use only and is not to be
transmitted to any other person or organization without prior approval of the
authorizing official.

NAVAL MEDICAL RESEARCH UNIT NO. 4
GREAT LAKES, ILLINOIS

11 December 1967

ALPHA-1 GLOBULINS IN THOSE SUSCEPTIBLE TO VIRAL
ACUTE RESPIRATORY DISEASES

By

R. I. LYTLER and M. J. MC NAMARA*

Research Project Report MR 005.20-0013, the Bureau of Medicine
and Surgery, Navy Department, Washington, D.C.

FOR OFFICIAL USE ONLY
MAY BE RELEASED ON: 11 June 1968

The opinions expressed herein are those of the authors and cannot be construed as reflecting the views of the Navy Department or the Naval Service at large. The use of commercially available products does not imply endorsement of these products or preference to other similar products on the market.

This document is for official use only until it has appeared in the open literature, or until 11 June 1968, whichever occurs earlier. It is not to be quoted for publication without the written permission of the authors. Quotation or reference to this report should cite the title of the report, authors, number of report and date of publication.

*Present address: University of Kentucky, Medical Center, Dept. Community Medicine, Lexington, Kentucky.

A high frequency of elevated initial serum alpha-1 globulin levels was noted among naval recruits who subsequently appeared to be more susceptible to acute viral respiratory diseases.

Correlating the biological indices with host resistance in disease patterns presents a real challenge. Qualitative and quantitative differences in susceptibility to infection also needs to be evaluated (1-3). During epidemiological surveys, individual differences were seen in which 50-60% of the recruits had an average of more than two discrete respiratory episodes, while 10-20% remained disease-free throughout the same training period (4). The study of the host is therefore of significant importance, since other extensive studies have failed to show why some recruits are more susceptible to respiratory infection than others (3-5).

The purpose of this paper is to determine the value of the initial serum alpha-1 globulin moiety in predicting susceptibility to respiratory infection.

MATERIALS AND METHODS

Population:

The study population consisted of two companies of recruits in basic training at the Great Lakes Naval Training Center. One company (A, 55 men) underwent training from November 1959 through January 1960. The other (B, 69 men) trained from July through September 1960. Recruits at this training center are predominantly 17-19-year olds from the eastern half of the United States.

Sampling procedures:

The men in both companies were first observed within 24-48 hours after their arrival, and were subsequently interviewed (MJM) at 0700 on Mondays, Wednesdays and Fridays, when oral temperatures and symptoms of respiratory disease were recorded. The

presence or absence of symptoms on non-interview days was also recorded.

A discrete respiratory disease episode was considered to have occurred if the symptoms of an individual differed in intensity, quality, or number from his usual pattern, and were present for at least two days. A patient with respiratory infection was considered febrile if the oral temperature was 100°F or above (2).

Collection of specimens:

At least three specimens were collected from every recruit who was observed for the entire training period, i.e., one within two days of his arrival, one during the middle (approximately 4th week) and, one just before completion of training. Specimens were also obtained at the onset of acute respiratory disease and three weeks convalescent. All were stored at -20°C until tested.

Laboratory procedures:

Serologic tests:

The following antigens were used in the complement fixation (CF) tests: influenza viruses - type A strain (A2/Japan/305/57) and type B strain (B/GL/1760/54). Parainfluenza viruses - types 1 (HA-2) and 3(HA-1)*. Adenovirus - types 3, 4, and 7 grown in HeLa cell cultures in this laboratory. Tests were performed by the modified Kolmer technique using two full units of complement and antigen (6).

Paper strip electrophoresis:

Serum protein distributions were determined on each recruit without knowledge of their respiratory disease pattern. Six lambda (.006 ml) of serum was applied at the center of 3X30.5 cm strip of Whatman No. 1 filter paper in a Durrum type cell. Specimens were

*Microbiological Associates

allowed to resolve for 16 hours in a barbitol buffer system (pH 8.6 at ionic strength .05 M) in a current of .1 ma (width of strip). Strips were then fixed and stained with bromphenol blue in accordance with procedures outlined elsewhere (7,8). Serum protein distributions were measured with the Spinco model R analytrol.

RESULTS

The results in Table 1 indicate that the average number of discrete respiratory episodes in the two companies of recruits was remarkably similar, therefore, no attempt has been made to present these data in reference to individual companies.

The distribution of serological antibody responses among the 119 recruits is presented in Table 2. One of the most striking observations to be made from these data is that very few (15.6%) failed to respond to at least one of the battery of antigens tested, while over 50% revealed significant antibody responses to 2 or more viral antigens.

The distribution of respiratory episodes and serological antibody rises in naval recruits is illustrated in Figure 1. Respiratory disease episodes (representing acquisition of new agents) were concentrated in the first 4 weeks of training with distinctive disease peaks at 10 and 25 days (Fig. 2). The serological antibody responses were also of a bimodal nature with an interval of 15 days between peaks which followed the respective episodes by 10 days. An additional flare-up was evident near the end of the recruit training period (60-65 days).

Figure 3 shows a comparison of the initial alpha-1 globulin levels of each individual with their subsequent number of respiratory episodes. Significantly elevated values (3.5 mg%) were found at a greater frequency among recruits who subsequently experienced more than 2 discrete episodes (see also lower half of Table 3). A more detailed investigation of recruits who experienced either single or multiple antibody responses is graphically compared to their alpha-1 globulin levels in Figure 4 --- difference being apparent not only within, but between groups

selected for single or multiple response. Initial serum alpha-1 globulin levels were not found to be significantly elevated in recruits who subsequently developed only antibody responses to either HA-1 or 1760 or to both. Evidence of antibody response to J-305, either alone or in combination with other antigens, was most commonly seen in recruits with significantly elevated alpha-1 globulin levels in their initial or subsequent serum specimens. A positive correlation, as shown in Figure 5, resulted from a comparison of the percentage of significantly elevated alpha-1 globulin with the number of antibody responses.

In Table 3 the mean alpha-1 globulin values and antibody responses are compared with the number of clinical episodes. In those men who had three or more respiratory episodes, initial alpha-1 globulin levels higher than 3.5 mg% are significantly more frequent ($p = <0.001$).

DISCUSSION

The frequency of discrete respiratory episodes among the military population described indicated that very few recruits completed training without experiencing a respiratory infection. In fact, an average of more than 2 discrete upper respiratory infections per man occurred during the Fall-Winter and Summer study period. A high incidence of viral respiratory infections was also indicated by the serological data. Over 50% of the men acquired responses to more than one respiratory antigen. Seventy-five percent of the individuals who experienced more than 2 different respiratory infections had significantly elevated initial serum alpha-1 globulin values. These results might have been even more significant if the nature of antibody response to influenza 1760 were known. An antibody response to influenza 1760 without clinical symptoms, and isolations during the across-the-board influenza vaccine program (although these companies were exempt) may indicate that antibody rises resulted from immunization. If so, then significantly elevated alpha-1 globulin levels might be used as an index in predicting susceptibility to natural respiratory diseases, as compared with those resulting from a vaccine response. To evaluate this, a knowledge of the role of

this parameter in antibody responses would be extremely important. Clinical symptoms, serological response and normal alpha-1 globulin levels may well indicate a difference in responsiveness to immunization rather than a natural episode. Statistically significant differences were found in both the epidemiological and serological response groups with respect to their alpha-1 globulin levels.

SUMMARY

The respiratory disease pattern observed in two companies of naval recruits during a 1959-1960 (Fall-Winter and Summer) epidemiological study, indicated an average of more than 2 discrete respiratory episodes per man. Over 50% demonstrated antibody responses to multiple antigens, while only 15.6% failed to show a serological rise to a viral respiratory antigen. Most of the new infections were acquired during the first half of the training period. There were two distinct peaks of clinical disease. The first occurred on the 10th day of training and the second on the 25th day. The peaks of clinical disease were followed in both instances by serological peaks 10 days later.

The alpha-1 globulin values were found to be significantly elevated in 40% of the recruits who subsequently demonstrated multiple serological responses. Seventy-five percent of the recruits considered to be susceptibles on the basis of demonstrating more than 2 discrete respiratory episodes had significantly elevated initial serum alpha-1 globulin levels. A highly significant relationship ($p = <0.001$) was found between number of discrete episodes and the alpha-1 globulin levels.

TABLE I

Respiratory Disease in Two Companies of Naval Recruits
during Nine Weeks of Training in the Winter of 1959-60
(Company A) and the Summer of 1960 (Company B)

Companies	Population	Total APD*	Average number ARD per man
A	50	117	2.3
B	69	151	2.2

*Acute respiratory disease

TABLE 2

The Distribution of Significant Serological Antibody Responses
to Seven Selected Acute Respiratory Viral Agents among
Naval Recruits during the 1959-1960 Epidemiological Study

Serological responses - 119 recruits	
Number of antigens	Percent recruits developing antibody rises
0	15.6
1	31.2
2	34.4
3	12.8
4	6.2

TABLE 3

Relationship of the Mean Alpha-1 Globulin among the Initial
Serum Specimens with the Antibody Responses and the
Number of Discrete Episodes

	Number of episodes*				
	0	1	2	3	4
Mean					
Alpha-1 globulin (mg%)	2.3	2.24	2.46	3.9**	4.25**
Antibody response (No. of antigens)	0	1.1	1.8	1.7	2.33
	Number of serological rises†				
	0	1	2	3	
Mean					
Alpha-1 globulin	2.28	2.79	3.56**	3.71**	
Episodes	.83	2.1	2.55	3.0	

*F 21.55 p = <0.001 significant difference between groups (episodes)

**Significantly elevated (3.5 mg%)

† F 5.17 p = <0.01 significant difference between groups (serological)




FIG. 1. Disease episodes and significant rises to selected respiratory viruses.




FIG. 2. Comparison of significant antibody response with that of respiratory episodes. Time in days relationship of respiratory episodes and development of viral antibodies.




FIG. 3. Initial serum alpha-1 globulin levels as related to respiratory episodes during the recruit training period.




FIG. 4. Percent of significantly increased alpha-1 globulins (above 3.5 mg%) among recruits who subsequently demonstrated either single or multiple antibody responses




FIG. 5. Positive correlation of antibody response with percent recruits having significantly elevated alpha-1 globulins.

REFERENCES

1. Lytle, R.I., Rytel, M. and Edwards, E.A.: Correlation of Immunological and Biological Factors in the Host with Susceptibility to Respiratory Infections. J. Infect. Dis. 116:67-74, 1966.
2. Lytle, R.I., Pierce, W.E. and Crawford, Y.E.: Multiple Infection in Acute Respiratory Disease. II. Gamma Globulin Disturbances in Dual Infections and Possible Significance in Acute Rheumatic Fever. J. Infect. Dis. 113:123-130, 1963.
3. Rytel, M. and Lytle, R.I.: Alterations in Humoral Factors in the Course of Acute Respiratory Infections. Naval Medical Research Unit No. 4, Project Report No. MF 022.03.07-4002, RU 66.14 to the Bureau of Medicine and Surgery, Navy Department, Washington, D.C. on 30 December 1966.
4. McNamara, M.J., Pierce, W., Crawford, Y.E. and Miller, L.F.: Patterns of Adenovirus Infection in the Respiratory Diseases of Naval Recruits. Amer. Rev. Resp. Dis. 86:485-497, 1962.
5. Stille, W.T., Pierce, W.E. and Crawford, Y.E.: Multiple Infections in Acute Respiratory Illness. J. Infect. Dis. 109:158-165, 1961.
6. Kolmer, J.A., Spaulding, E.N. and Robinson, H.S.: Approved Laboratory Techniques. Appleton-Century-Crofts, New York, 1950, p. 814.
7. Lytle, R.I., Rosenbaum, M.J., Miller, L.F. and Rosenthal, S.: A Protein Moiety Inhibitory to HeLa Growth in Sera of Burned Children. Fed. Proc. 19:73, 1960.
8. Lytle, R.I., Young, J. and Stille, W.T.: Standard Values and Variations of Paper Strip Electrophoresis. Naval Medical Research Unit No. 4, Project Report No. MR 005.12-1102.1 to the Bureau of Medicine and Surgery, Navy Department, Washington, D.C. on 2 May 1961.

ACKNOWLEDGMENTS

The authors are grateful to Captain L. F. Miller, MC USN, who encouraged this study and to Dr. Richard Winzler who provided advice and direction.

The authors also wish to thank Mrs. E. Hutchings for her faithful technical assistance, and Mrs. Lula Duggan for the expert preparation of the manuscript.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Naval Medical Research Unit No. 4 Great Lakes, Illinois 60088		Unclassified
		2b. GROUP
3. REPORT TITLE		
Alpha-1 Globulins in Those Susceptible to Viral Acute Respiratory Diseases		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
Final		
5. AUTHOR(S) (Last name, first name, initial)		
Lytle, R. I. and McNamara, M. J.		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
11 December 1967	15	8
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S)	
a. PROJECT NO. MR 005.20-0013	RU 66.28	
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. AVAILABILITY/LIMITATION NOTICES		
		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY	
	Bureau of Medicine and Surgery Navy Department Washington, D.C.	
13. ABSTRACT		
The respiratory disease pattern observed in two companies of naval recruits during a 1959-1960 (Fall-Winter and Summer) epidemiological survey indicated an average of more than 2 discrete respiratory episodes per man. Over 50% demonstrated antibody responses to multiple antigens, while only 15.6% failed to show a serological rise to a viral respiratory antigen. Most of the infections were acquired during the first half of the training period. There were 2 distinct peaks of clinical disease. The first occurred the 10th day of training, and the second on the 25th day. The peaks of clinical disease were followed on both instances by serological peaks 10 days later. The alpha-1 globulin values were found to be significantly elevated in 40% of the recruits who subsequently demonstrated multiple serological responses. Seventy-five percent of the recruits considered to be susceptibles on the basis of demonstrating more than 2 discrete respiratory episodes had significantly elevated initial serum alpha-1 globulin levels. A highly significant relationship ($p = <0.001$) was found between number of discrete episodes and the alpha-1 globulin levels.		

DD FORM 1473

Unclassified

Security Classification

Unclassified

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Viral etiology						
Upper respiratory disease						
Alpha-globulins						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch or service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8c. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms, or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical content. The assignment of links, roles, and weights is optional.