

UNCLASSIFIED

AD NUMBER
AD835290
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; MAR 1966. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701.
AUTHORITY
Fort Detrick/SMUFD ltr dtd 14 Feb 1972

THIS PAGE IS UNCLASSIFIED

AD835290

TRANSLATION NO. 1656

DATE: 31 March 1966

DDC AVAILABILITY NOTICE

Reproduction of this publication in whole or in part is prohibited. However, DDC is authorized to reproduce the publication for United States Government purposes.

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of


DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

attn: Tech. Release Sect 11D

DDC
JUL 10 1968

1

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

STUDIES ON THE BACTERIAL LEAF BLIGHT REGION OF RICE
PLANT FROM THE VIEWPOINT OF SOIL AND MANURE

I. ON THE OUTLINE OF SOILS

Kyushu Nogyo Kenkyu (Kyushu
Agricultural Research), No.15,
April 1955. Pages 97-100

M. Yoshino and M. Nakahara
of the Saga Prefecture
Agricultural Experiment
Station

I. INTRODUCTION

Heretofore in the usual locality where wet-land rice leaf blight occurs, flooded areas were considered the normal condition. However, above and beyond this, in Saga prefecture, there were some areas where the usual outbreak occur. Considering this in the category of "Poor Rice Land in the Broad Sense", the authors already classified them as standing water type poor rice fields (4).

In general, with this type of region even the yield of wet-land plants are unstable. Furthermore, the development of the winter crop is not good.

Accordingly, an investigation was made on the characteristics of the wet-land field soils as well as on the development of the wet-land rice plant and submitted as data for later improvements.

II. GENERAL CONDITIONS OF THE AREA

(1) Weather - Topography - Geology - Altitude:

a. Weather: Annual average temperature 16 - 17°, annual precipitation 1700 mm.

b. Topography: Flat land

c. Geological Condition: Fourth geological era new layer of river and ocean-bed alluvial soil

d. Altitude: 4 - 7.5 m

(2) Distribution:

The area is sporadically distributed mostly within the Saga plain at the foot of the mountain close to the strip of level land (Figure 1).




Figure 1. Distribution of Area

- 1) Saga
- 2) Chikugo River
- 3) Ariake Sea

III. SOIL

1. Cross-Sectional Characteristic of Soil

Details of the sectional configuration of the soils will be presented separately, but the characteristic of the soil configuration has been arranged in Table 1. (The soil characteristic map of Figure 2 has been omitted because of space requirement).

These in the category of standing water are divided into I, II, and III and these were compared with a reference field. According to this:

(1) The loam stratum which is primarily based on peat is 40 cm deep more or less. Furthermore, the loam-like stratum is approximately 20 cm deep. Moreover, disregarding the loam "grei" stratum, the recent ash "grei" stratum is quite deep.

(1) 区分	No.	(3) 場所	(12) 泥炭乃至黒泥層	(19) 灰層	(21) 地下水水位	土の性質	構造の発達	(32) 保水日数	(34) 地層
I	35	五時岡・五丁岡 (4)	36~56cm (13)(黒泥化)	56cm (20)以下	65cm	C(CL)	弱(24)	33日	(35) 河海成土
	50	牛神・乙神 (5)	41~55 (55~90) (黒泥化) 泥炭(14)	90~	62	CL	弱(25)	5	M(36)成
	11	三日月・佐藤前 (6)	41~70 (黒泥化)(15)	70~	60	C(CL)	弱(26)	6	M(37)成
	48	西海・横浜 (7)	42~80 (黒泥化)(16)	80~	60	C(CL)	弱(27)	5	(38) 河海成
II	61	兵庫・佛示野 (8)	16~26 (黒泥類似層)(17)	95~	53	CL(C)	弱(28)	5	(39) 河海成
	65	上峰・中村 (9)	25~80 (黒泥類似層)(18)	~	55	L	弱(29)	6	M(41)成
III	49	鹿島・世間 (10)	—	35~	35	CL	弱(30)	7	河海成(41)
(2) 参照	25	佐賀・神野 (11)(元農試)	—	95~	85	CL	弱(31)	3~4	河海成(42)

(43) 註: ここに黒泥類似層と仮称するものは、黒ぼくに由来するものとみられる黒色泥土層を意味する。

Table 1. Characteristics of the Soil Configuration

- | | |
|---|-------------------------------|
| 1) Subdivision | 22) Underground Water Level |
| 2) Control | 23) Development of Structure |
| 3) Location | 24) Weak |
| 4) Gochoda-Gotoda | 25) Slightly Weak |
| 5) Ushizu-Otsuryu | 26) Weak |
| 6) Mikkazuki-Sahatanami | 27) " |
| 7) Nishizato-Yokodake | 28) " |
| 8) Hyogo-Wakijino | 29) Slightly Weak |
| 9) Jomine-Nakamura | 30) Very minute |
| 10) Kashima-Seken | 31) Intermediate |
| 11) Saga-Kamiya(former Agriculture Sta.) | 32) Period of Water Retention |
| 12) Peat to Loam Strata | 33) Days |
| 13) (Loam Forming) | 34) Note |
| 14) (" ") Peat | 35) River-Sea Soil |
| 15) (" ") | 36) River Formed |
| 16) (" ") | 37) " " |
| 17) (Loam-like Stratum) | 38) River-Sea Soil |
| 18) (" ") | 39) " " |
| 19) Recent Ash "Grei" Stratum | 40) River Formed |
| 20) Less than 56 cm | 41) River-Sea Soil |
| 21) Ground water Level | 42) " " |
| 43) Note: That which is referred to as Loam-like, is the dark colored soil stratum. | |

(2) In the lower strata, the recent ash-grey layer appears more often. Moreover, the underground water is high, in general, attaining 50 - 60 cm in most places.

(3) The soil characteristic of the present area is mostly viscous. Moreover, the development of the plowed bed and structure is weak, the period of water retention is quite long and the water permeation is generally poor.

2. Physical Characteristic

(1) Mechanical Constitution - According to Table 2, all the dirt fractions contained large amounts of minute and fine sand; however, there are those which were more uniform. Accordingly, with regard to the mechanical constitution, it is rather difficult to note distinct characteristics.

(1) 区分	No.	層(2) 位	(3)ピペット法による%				(8)土 性 名
			細(4)砂	細(5)砂	微(6)砂	粘(7)土	
I	35	I (A)	13.46	18.24	44.23	24.07	Clay loam
		II (B)	6.71	16.34	45.89	31.06	Silty Clay
		III (G P m)	6.33	16.34	54.90	22.43	Silty Clay loam
		IV (G)	2.76	15.26	58.44	23.54	"
I	50	I (A)	11.59	46.39	25.60	16.43	Clay loam
		II (B ₁)	14.15	44.89	31.55	9.41	Loam
		III (B ₂)	24.62	45.08	21.92	8.38	Sandy loam
		IV (G M)	1.23	28.31	42.23	28.23	Light Clay
II	29	I (A ₁)	0.94	9.25	58.09	31.72	Silty Clay
		II (A ₂)	1.11	11.75	59.74	27.40	"
		III (B _m)	0.46	14.86	56.02	28.66	"
		IV (B G ₁)	0.45	9.88	58.52	31.15	"
III	49	I (A ₁)	0.85	25.49	55.92	17.74	Silty Clay loam
		II (A ₂)	1.40	15.55	55.91	27.14	Silty Clay
(9) 原 田	28	A ₁	18.49	12.46	47.06	21.99	Silty Clay loam
		A ₂	—	—	—	—	—
		B ₁	12.76	31.73	31.66	23.88	Clay loam
		B ₂	4.41	7.80	52.03	35.73	Silty Clay

Table 2. Mechanical Structure of Soil

- | | |
|------------------------|--------------------|
| 1) Subdivision | 6) Ultra-fine Sand |
| 2) Strata | 7) Clay |
| 3) % by Pipette Method | 8) Soil Type |
| 4) Coarse Sand | 9) Control Field |
| 5) Fine Sand | |

PRECEDING
PAGE BLANK

Table 4. Chemical Analysis of Soil

(1) 区分	No.	(2) 層位	(3) 全氮素%	(4) 腐植%	pH (H ₂ O)	(5) 置換石灰%	(6) 遊離鉄%	(7) NH ₄ 化成率
I	(8) 35 (三丁田)	A	0.195	4.23	5.6	0.45	1.19	7.8
		B	0.190	2.39	7.0	0.50	1.30	—
		GPm	0.293	6.03	6.3	0.55	—	8.1
		G	0.163	3.89	6.1	0.53	—	—
	(9) 50 (牛津)	A	0.185	2.58	5.3	0.19	1.93	11.3
		B ₁	0.069	1.49	6.4	0.22	1.43	—
		B ₂ Gm	0.057 0.332	1.35 8.94	6.7 6.4	0.18 0.38	—	4.3
	(10) ¹¹ — (三日月)	A	0.222	3.49	6.8	0.21	2.37	(11)X大
		B ₁	0.257	5.88	7.0	0.18	2.68	—
		B ₂	0.263	7.51	6.3	0.26	—	—
		Gm	0.303	9.17	6.5	0.43	—	2.5
	(12) ⁴⁸ — (西郷)	A	0.206	3.51	5.9	0.30	1.00	8.2
B ₁		0.077	1.37	6.3	0.34	0.95	—	
B ₂		0.095	2.03	6.6	0.35	—	—	
GPm		0.237	7.85	6.8	0.31	—	2.7	
(13) ⁶¹ — (兵庫)	A ₁	0.255	5.75	5.1	0.49	1.08	4.7	
	A ₂	0.213	7.02	6.0	0.53	1.11	—	
	Bm	0.360	13.31	5.7	0.51	—	2.4	
	G ₁	0.087	1.63	4.5	0.18	—	—	
II	(14) ⁶⁵ — (上峰)	A ₁	0.409	5.49	4.9	0.27	1.40	6.8
		B	0.174	1.53	6.5	0.27	0.96	—
		Gm	0.373	9.83	6.1	0.39	—	5.0
(15) ²⁹ — (兵庫)	A ₁	0.249	5.57	4.9	0.38	1.03	4.1	
	A ₂	0.183	4.79	5.7	0.49	1.26	—	
	B ₁	0.092	3.49	5.8	0.35	—	—	
III	(16) ⁴⁹ — (鹿島)	A ₁	0.273	4.29	6.1	0.46	1.58	8.5
		A ₂	0.255	3.96	6.4	0.49	1.47	8.0
(17)類 田	(18) ²⁵ — (佐賀)	A ₁	0.258	4.22	5.6	0.29	1.34	—
		A ₂	0.198	3.17	6.0	0.36	1.10	—
		B ₁	0.124	1.01	7.0	0.40	1.67	—
		B ₂	0.089	1.23	7.2	0.55	—	—

- | | |
|---|---------------------|
| 1) Subdivision | 10) (Mikkazuki) |
| 2) Strata | 11) Absent |
| 3) Total Nitrogen % | 12) (Nishizato) |
| 4) Humus % | 13) (Hyogo) |
| 5) Lime Replaced % | 14) (Jomine) |
| 6) Metallic Iron % | 15) (Hyogo) |
| 7) NH ₄ Transformation Index | 16) (Kashima) |
| 8) (Gotoda) | 17) Reference Field |
| 9) (Ushizu) | 18) (Saga) |

layer was generally high. For example, it compares with the NH_3 -transformation rate in rice fields with standing water at such places as Shioire and Acmine (2).

Although the chemical analysis was not complete, the results of determining the Eh (Acid reduction potential) of a portion of ground were as follows (Harvest soils during the first part of August):

No. 50-I indicated Eh 17.5 mv, No. 25-I (Reference) Eh 188 mv. Thus the Eh of the soil of the present area was extremely low compared to that of the reference.

Although a summary of the ground environment of the typical wet-land, where wet rice plant leaf blight outbreak occurred at Saga prefecture, has been presented above; it is hoped that future investigations can be made on the effect of soil dehydration, the texture of the humus, the relationship of potassium, nitrogen and of other inorganic components. Furthermore, using the investigation on cultivation as a basis, it is hoped that research can be conducted on the period of development and the permeation of nourishment for the wet rice plant.

REFERENCES

- (1) Yoshino, Koike: "Soil Structure and Development of Crops (I)", Kyushu Agricultural Research, No. 10 (1953).
- (2) S. Acmine: Effect of Culvert Drainage and Dehydration, p. 26 (1949).