

UNCLASSIFIED

AD NUMBER
AD853965
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAY 1969. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Release Branch, Attn: TID, Frederick, MD 21701.
AUTHORITY
BDRL D/A ltr, 29 Sep 1981

THIS PAGE IS UNCLASSIFIED

AD


TECHNICAL MANUSCRIPT 501

**ROOT PERMEABILITY AS AFFECTED
BY PICLORAM AND OTHER CHEMICALS**

AD853965

**Charles P. P. Reid
Woodland Hurtt**

MAY 1969



STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/TID. Frederick. Maryland 21701

**DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland**

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

ACCESSION NO.	
CFSTI	✓
DDC	✓
DDA/DMIC/2	
JUSTIFICATION	
BY	
DIST. NO.	
DATE	
2	

**DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701**

TECHNICAL MANUSCRIPT 501

**ROOT PERMEABILITY AS AFFECTED BY
PICLORAM AND OTHER CHEMICALS**

Charles P.P. Reid

Woodland Hurtt

**Plant Physiology Division
PLANT SCIENCES LABORATORIES**

Project 1B562602A061

May 1969

ABSTRACT

Because increased water deficit in the plant caused by changes in root permeability can result in transpiration reduction, the effects of picloram (4-amino-3,5,6-trichloropicolinic acid) and several other chemicals on root permeability were studied. Initially, effects on cell permeability were investigated by measuring the betacyanin efflux from red beet (Beta vulgaris L.) root sections. Picloram solutions ranging from 10^{-3} M to 10^{-6} M had no significant effect on betacyanin efflux when compared with controls. Similar results were found for 10^{-4} M and 10^{-5} M 2-methoxy-3,6-dichlorobenzoic acid (dicamba), 10^{-4} M 2,4-dichlorophenoxyacetic acid (2,4-D), and 10 ppm of ethylene. Compounds that caused significant pigment leakage were 10^{-4} M and 10^{-5} M phenylmercuric acetate, 10^{-3} M and 10^{-4} M 2,4-dinitrophenol, 10^{-3} M and 10^{-4} M 2,4,5-trichlorophenoxyacetic acid, and 10^{-3} M 2,4-D.

Further investigations were conducted with picloram applications to the roots of bean plants (Phaseolus vulgaris L. var. Black Valentine) grown in nutrient solution under controlled environmental conditions. Roots treated for 3 hours with 10^{-5} M picloram showed no significant electrolyte leakage as determined by conductivity measurements of the root bathing solution for 52 hours. When bean roots were treated for 3 hours with 10^{-5} M, 10^{-6} M, and 10^{-7} M picloram and then the plants detopped, increased stem exudation was observed. Xylem exudate of the treated plants also showed increased electrolytic conductivity. The increased exudation rate accompanied by increased conductivity indicates that picloram has little effect on root cell membrane integrity, does not act as a metabolic inhibitor, and in some way stimulates salt secretion into the xylem.

I. INTRODUCTION*

Picloram (4-amino-3,5,6-trichloropicolinic acid) is a highly biologically active herbicide with apparently unique properties. It causes distortions in growing tissue and other formative effects¹ and growth promotion in stem sections, similar to that caused by 2,4-dichlorophenoxyacetic acid (2,4-D) and other auxin-like growth regulators.^{2,3} Picloram has also been characterized as comparable to 2,4-D and 2,4,5-trichlorophenoxyacetic acid (2,4,5-T) in absorption and translocation.¹ To further elucidate the modes of action of picloram, studies have been conducted on its effect on foliar transpiration. Because increased water deficit in the plant caused by changes in root permeability can result in transpiration reduction, the effects of picloram and several other chemicals on root permeability were studied.

II. MATERIALS AND METHODS

A technique of Veldstra and Booij⁴ involving the measurement of betacyanin efflux from red beet root sections was used for investigating the possible effects of selected substances on cell permeability. Discs (10 mm by 3 mm) were cut from the roots of red beet (*Beta vulgaris* L.), rinsed several times in distilled water, and then retained in distilled water overnight at 8 C. After again rinsing, 10 discs were placed in Erlenmeyer flasks containing 50 ml of solution. Each of the following compounds was prepared in 0.5X Hoagland's nutrient solution and replicated three times: picloram, 10^{-3} M, 10^{-4} M, 10^{-5} M, and 10^{-6} M; 2,4-D, 10^{-3} M and 10^{-4} M; 2,4,5-T, 10^{-3} M and 10^{-4} M; 2-methoxy-3,6-dichlorobenzoic acid (dicamba), 10^{-4} M and 10^{-5} M; phenylmercuric acetate (PMA), 10^{-4} M and 10^{-5} M; 2,4-dinitrophenol (DNP), 10^{-3} M and 10^{-4} M; and ethylene, 10 ppm. All solutions were adjusted to pH 6.05 ± 0.12 .

Pigment release from the discs was determined by measuring the light absorbancy of the solution at 500 m μ with a Bausch & Lomb Spectronic 20 colorimeter and comparing the values with blanks of 0.5X Hoagland's solution. Measurements were made periodically up to 24 hours; the flasks were maintained at 25 C.

* This report should not be used as a literature citation in material to be published in the open literature. Readers interested in referencing the information contained herein should contact the senior author to ascertain when and where it may appear in citable form.

To study the effect of picloram on exudation rate, bean plants (*Phaseolus vulgaris* L. var. Black Valentine) were grown in nutrient culture in a controlled environment growth chamber with a 16-hour photoperiod provided by 1,400 ft-c of illumination at plant-top level. Temperature and relative humidity were maintained at 25 ± 1 C and $50 \pm 5\%$, respectively. When the first compound leaves were beginning to unfold, plant roots were treated for 3 hours with 300 ml of selected molarities of picloram in 0.5X Hoagland's nutrient solution. After treatment, each plant was transferred to a 250-ml Erlenmeyer flask containing aerated distilled water. Control plants were manipulated in the same manner as the treated plants except that picloram was omitted. Each series of picloram-treated plants and control plants consisted of 10 replications. To measure exudation rate, the top of each plant was removed 2 cm below the cotyledonary node immediately after the 3-hour treatment period, and a 0.2-ml graduated pipette was attached vertically to the cut stem by a short length of rubber tubing. At the start of each measurement period, the pipette was adjusted to the zero mark by injecting distilled water containing acid fuschin dye. The increase of volume in the pipette caused by xylem exudate was then recorded periodically and expressed as milliliters of exudate per hour.

Electrolyte conductivity of the root bathing solution was measured in 5-ml samples removed periodically from the flasks containing treated and control plants. Solution conductivity (μmhos at 25 C) was determined with an Industrial Instruments Inc. Conductivity Bridge Model RC16B2 using a 3-ml pipette cell with a 0.1 cell constant.

Conductivity of xylem exudate from several different experiments was also determined. Plants were treated as above but exudate was collected in 5-ml plastic containers attached to each stem. Eleven collections were made, each consisting of the pooled exudate from at least six plants. The conductivity of a particular exudate collection from plants treated with 10^{-5} M picloram was compared with the control plant exudate conductivity at the same collection period. A statistical analysis based on paired observations⁵ was then performed on all collections.

III. RESULTS AND DISCUSSION

The effects of picloram and the other compounds on cell permeability are shown in Figure 1. The degree of disturbance of the semipermeability of the membranes of the beet root cells is indicated by the absorbancy value of the bathing solution, with higher values demonstrating greater leakage of betacyanin from the vacuoles. The loss of pigment was greatest with 10^{-4} M FMA followed in descending order of loss by 10^{-3} M 2,4,5-T, 10^{-3} M DNP, 10^{-5} M FMA, 10^{-3} M 2,4-D, 10^{-4} M 2,4,5-T, and 10^{-4} M DNP. No significant loss of pigment over the 24-hour period was detected for 10^{-3} M, 10^{-4} M, 10^{-5} M, and 10^{-6} M picloram; 10^{-4} M 2,4-D; 10^{-4} M and 10^{-5} M dicamba; or 10 ppm of ethylene. The slight peak observed for ethylene at 2 hours is thought to be the result of a spurious reading caused by air bubbles in the solution.




FIGURE 1. Effects of Various Compounds on Pigment Efflux from Beet Root Discs at 25 C. Degree of efflux reflected as absorbancy value of solution. All values above 0.06 are significantly different from the control at 5% level.

In addition to picloram, 2,4-D, 2,4,5-T, and dicamba were tested because of their known effectiveness as herbicides. The responses observed with 2,4-D and 2,4,5-T are similar to those reported by Veldstra and Booij.⁴ They found that the chlorinated compound 2,4,6-trichlorophenoxyacetic acid interacted with beet membranes more strongly than 2,4-D. This stronger response was attributed to the increase of the lipophilic component of the molecule by the introduction of the added chlorine atom.

DNP, a metabolic inhibitor known to affect membrane permeability adversely, was included as a basis for comparison. Surprisingly, PMA proved to be the most effective compound in causing pigment leakage, even at a lower concentration than DNP. PMA has been studied extensively as an antitranspirant and fungicide.⁶ Siegenthaler and Packer⁷ found that PMA inhibits light-induced chloroplast swelling and also inhibits nicotinamide adenine dinucleotide phosphate reduction and photophosphorylation reactions. Perhaps the effect of PMA on regulation of chloroplast volume is also related to changes in membrane permeability. Ethylene, a gaseous plant growth regulator,⁸ has been considered to cause alteration to membrane permeability, although no such effect appeared under the particular experimental conditions used here.

It is particularly interesting that picloram had no appreciable effect on membrane permeability as compared with 2,4-D and 2,4,5-T, because picloram is often compared with these compounds as having similar growth regulation characteristics. Veldstra and Booij⁴ noted that the activity of compounds such as indoleacetic acid, naphthaleneacetic acid, and 2,4-D, as determined by the beet disc test, reflects very well their phytotoxicity as determined from studies on parts of plants or intact plants. Obviously, this is not the case for picloram.

Exudation rates resulting from treatment with picloram further support the thesis that picloram does not affect membrane permeability adversely. Increases in exudation rate were consistently observed for plants treated with 10^{-5} M, 10^{-6} M, and 10^{-7} M picloram (Fig. 2 to 4). Increases in exudation rate were significant 7 hours after the initiation of treatment for the 10^{-5} M and 10^{-6} M picloram-treated plants and remained significant throughout the experiments. Exudation rate of the 10^{-7} M picloram-treated plants was not significantly greater during the first 7 hours but was significant after 22 hours. Because exudation rate from detopped plants is primarily affected by salt transfer into the xylem,⁹⁻¹¹ it appears that picloram directly or indirectly causes an increased movement of salt into the xylem. In fact, statistically higher conductivities, indicating a greater salt concentration, occurred in the collected exudate from plants treated with 10^{-5} M picloram when compared with controls (Table 1).




FIGURE 2. Effect of 10^{-5} M Picloram on Exudation Rate of Detopped Plants. Three-hour picloram treatment on roots initiated at zero time. Vertical bars represent 95% confidence limits on the mean.




FIGURE 3. Effect of 10^{-6} M Picloram on Exudation Rate of Detopped Plants. Three-hour picloram treatment on roots initiated at zero time. Vertical bars represent 95% confidence limits on the mean.




FIGURE 4. Effect of 10^{-7} M Picloram on Exudation Rate of Detopped Plants. Three-hour picloram treatment on roots initiated at zero time. Vertical bars represent 95% confidence limits on the mean.

TABLE 1. COMPARISON OF XYLEM EXUDATE CONDUCTIVITY FROM
PICLORAM-TREATED PLANTS AND CONTROLS

Treatment	Conductivity, μmhos , for Indicated Exudate Collection Numbers/ 1 2 3 4 5 6 7 8 9 10 11											Mean
	1	2	3	4	5	6	7	8	9	10	11	
10^{-5} M Picloram	1,140	2,325	1,950	2,720	980	1,160	2,530	2,260	2,180	2,260	2,200	1,973
Control	600	1,725	1,725	1,910	995	745	2,250	2,250	2,120	2,080	1,860	1,660
Difference	540	600	225	810	(15)	415	280	10	60	180	340	313 ^b

a. Each figure represents the pooled exudate from at least six plants.

b. Statistically significant at 1% level using paired observations analysis.⁶

Salt moves into the xylem even though the roots are bathed in distilled water,^{9,11} as was done here. However, the exudation rate steadily declines with time (Fig. 2 to 4).

Wallace et al.¹¹ indicate that movement of salt into the xylem appears to be under metabolic control. Considering that inhibitors such as DNP and KCN decrease exudation rate,¹⁰ it seems very unlikely that picloram acts as a metabolic inhibitor in the root system.

The measurements of the conductivity of the root bathing solution of the plants treated with 10^{-5} M picloram indicated no statistically significant electrolyte loss as compared with controls (Fig. 5). The increase of exudation rate with no change in ion leakage from the roots seems to be in agreement with the results of Newman and Kramer.¹² In the reverse situation, they found that a decrease in root pressure exudation accompanied an increase of electrolyte leakage from roots immersed in decenylsuccinic acid, a metabolic inhibitor.




FIGURE 5. Conductivity of the Root Bathing Solution of Plants Treated for 3 Hours with 10^{-5} M Picloram. Vertical bars represent 95% confidence limits on the mean.

Although it is not known why exudation rate is increased by picloram, it might be explained as an auxin-like effect. It has been demonstrated that auxin in low concentrations causes increased exudation,¹³ and that auxin causes increased water uptake by plant tissues.¹⁴ However, it is generally agreed that auxin-induced water uptake is primarily related to effects on cell wall extensibility, which in turn permits osmotic water uptake, rather than effects on membrane permeability per se.¹⁴

The increased salt concentration found in the xylem exudate cannot be explained entirely on the basis of an auxin effect, and for this reason another mechanism appears to be involved other than wall extensibility, perhaps one affecting ion transport across membranes.

LITERATURE CITED

1. Hamaker, J.W.; Johnston, H.; Martin, R.J.; Redemann, C.T. 1963. A picolinic acid derivative: Plant growth regulator. *Science* 141:363.
2. Eisinger, W.; Morré, D.J.; Hess, C.E. 1966. Promotion of plant growth by Tordon herbicide. *Down to Earth* 21:8-10.
3. Fisher, D.A.; Bayer, D.E.; Weier, T.E. 1968. Morphological and anatomical effects of picloram on Phaseolus vulgaris. *Bot. Gaz.* 129:67-70.
4. Veldstra, H.; Booij, H.L. 1949. Researches on plant growth regulators: XVII. Structure and activity; on the mechanism of the action III. *Biochim. Biophys. Acta* 3:278-312.
5. Li, J.C.R. 1964. Statistical inference, Vol. I. Statistics, Inc., c/o Edwards Brothers, Inc., Ann Arbor, Michigan. 658 p.
6. Pallas, J.E., Jr.; Bertrand, A.R.; Harris, D.G.; Elkins, C.B., Jr.; Parks, C.L. 1965. Research in plant transpiration: 1962. U.S. Dep. Agr. Prod. Res. Rep. 87. 56 p.
7. Siegenthaler, P.A.; Packer, L. 1965. Light-dependent volume changes and reactions in chloroplasts: I. Action of alkenylsuccinic acids and phenylmercuric acetate and possible relation to mechanisms of stomatal control. *Plant Physiol.* 40:785-791.
8. Abeles, F.B.; Gahagan, H.E., III. 1968. Abscission: The role of ethylene, ethylene analogues, carbon dioxide, and oxygen. *Plant Physiol.* 43:1255-1258.
9. Arisz, W.H.; Helder, R.J.; Van Nie, R. 1951. Analysis of the exudation process in tomato plants. *J. Exp. Bot.* 2:257-297.
10. Vaadia, Y. 1960. Autonomic diurnal fluctuations in rate of exudation and root pressure of decapitated sunflower plants. *Physiol. Plant.* 13:701-717.
11. Wallace, A.; Ashcroft, R.T.; Lunt, O.R.; Joven, C.B.; Clark, J.A. 1966. Root pressure exudation in tobacco: I. Some general characteristics, p. 127-135. In A. Wallace (ed.) *Current topics in plant nutrition*. Edwards Brothers, Inc., Ann Arbor, Michigan.
12. Newman, E.I.; Kramer, P.J. 1966. Effects of decenylsuccinic acid on the permeability and growth of bean roots. *Plant Physiol.* 41:606-609.

13. Skoog, F.; Broyer, T.C.; Grossenbacher, K.A. 1938. Effects of auxin on rates, periodicity, and osmotic relations in exudation. *Amer. J. Bot.* 25:749-759.
14. Kozlowski, T.T. 1964. *Water metabolism in plants.* Harper and Row, N. Y. 227 p.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Department of the Army Fort Detrick, Frederick, Maryland, 21701		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE ROOT PERMEABILITY AS AFFECTED BY PICLORAM AND OTHER CHEMICALS		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name) Charles P.P. Reid Woodland (NMI) Hurtt		
6. REPORT DATE May 1969	7a. TOTAL NO. OF PAGES 16	7b. NO. OF REFS 14
8a. CONTRACT OR GRANT NO. A. PROJECT NO. 1B562602A061	8b. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 501	
9.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. DISTRIBUTION STATEMENT Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland, 21701	
13. ABSTRACT Because increased water deficit in the plant caused by changes in root permeability can result in transpiration reduction, the effects of picloram (4-amino-3,5,6-trichloropicolinic acid) and several other chemicals on root permeability were studied. Initially, effects on cell permeability were investigated by measuring the betacyanin efflux from red beet (<i>Beta vulgaris</i> L.) root sections. Picloram solutions ranging from 10^{-3} M to 10^{-6} M had no significant effect on betacyanin efflux when compared with controls. Similar results were found for 10^{-4} M and 10^{-5} M 2-methoxy-3,6-dichlorobenzoic acid (dicamba), 10^{-4} M 2,4-dichlorophenoxyacetic acid (2,4-D) and 10 ppm of ethylene. Compounds that caused significant pigment leakage were 10^{-4} M and 10^{-5} M phenylmercuric acetate, 10^{-4} M and 10^{-5} M 2,4-dinitrophenol, 10^{-4} M and 10^{-5} M 2,4,5-trichlorophenoxyacetic acid, and 10^{-4} M 2,4-D. Further investigations were conducted with picloram applications to the roots of bean plants (<i>Phaseolus vulgaris</i> L. var. Black Valentine) grown in nutrient solution under controlled environmental conditions. Roots treated for 3 hours with 10^{-4} M picloram showed no significant electrolyte leakage as determined by conductivity measurements of the root bathing solution for 52 hours. When bean roots were treated for 3 hours with 10^{-4} M, 10^{-5} M, and 10^{-6} M picloram and then the plants detopped, increased stem exudation was observed. Xylem exudate of the treated plants also showed increased electrolytic conductivity. The increased exudation rate accompanied by increased conductivity indicates that picloram has little effect on root cell membrane integrity, does not act as a metabolic inhibitor, and in some way stimulates salt secretion into the xylem. () ←		

DD FORM 1473 1 NOV 65 1 JAN 68, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified
Security Classification

Unclassified
Security Classification

14. Key Words
Picloran
Herbicides
Roots
Permeability
Transpiration
Plants (botany)

Unclassified
Security Classification