


UNCLASSIFIED

AD NUMBER
AD858521
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution: Further dissemination only as directed by Naval Oceanographic Office, Washington, DC 20390, JUN 1969, or higher DoD authority.
AUTHORITY
USNOO ltr, 25 Jun 1972

THIS PAGE IS UNCLASSIFIED

INFORMAL REPORT

MARINE BIOFOULING IN VIEQUES
SOUND, PUERTO RICO,
AN INTERIM REPORT
APRIL 1964 TO FEBRUARY 1969



JUNE 1969

This document has been approved for public
release and sale; its distribution is unlimited.

NAVAL OCEANOGRAPHIC OFFICE
WASHINGTON, D.C. 20390

Best Available Copy

20070119194

LIBRARY
JUL 80 1969
U.S. NATIONAL ACADEMY

INFORMAL REPORT

The Informal Report (IR) as produced at the Naval Oceanographic Office is a means for personnel to issue timely scientific and technical preliminary reports of their investigations. These are primarily informal documents used to report preliminary findings or useful byproducts of investigations and work to members of the scientific and industrial communities.

Informal Reports are assigned sequential numbers for each calendar year; the digits preceding the dash indicate the year.

The distribution made of this report is determined primarily by the author. Information concerning obtaining additional copies or being placed on a distribution list for all future Informal Reports in a given area of interest or specialty field, should be obtained from:

Distribution Control Department
Code 4420
Naval Oceanographic Office
Washington, D. C. 20390

ABSTRACT

In 1964, the U.S. Naval Oceanographic Office began a marine biofouling sampling program in Vieques Sound, Puerto Rico. The sampling program is continuing until May 1970. This program is one of a series of sampling programs to determine the character and extent of marine fouling communities in various marine biological provinces of the world.

The Vieques Sound program is being conducted by exposing test panels at 50- and 95-foot depths in 100 feet of water. From April 1964 to February 1969, 103 panels were recovered and analyzed for fouling and boring organisms.

Settlement of fouling larvae was found to occur throughout the year in Vieques Sound. The rate of fouling, however, was quite variable from month to month, apparently depending on whether water currents bring larvae of potential foulers from distant brood sites. A strong affinity exists between the fauna of the Vieques Sound fouling communities and the fauna of the South Florida/Bahamas region.

JOHN R. DePALMA
Nearshore Surveys Division
Oceanographic Surveys Department

This report has been reviewed and is approved for release as an UNCLASSIFIED Informal Report.


L. B. BERTHOLF

Director, Nearshore Surveys Division

ACKNOWLEDGMENTS

The author is indebted to Dr. Ruth D. Turner of the Harvard Museum of Comparative Zoology, Dr. Arnold Ross of the San Diego Museum of Natural History, Dr. Maximo Cerase-Vivas of the University of Puerto Rico, and Mr. Edward R. Long of NAVOCEANO for assisting with the identification of the various organisms.

The author also is indebted to the EOD divers from the U.S. Naval Station in San Juan who collected panels and maintained the test site with remarkable efficiency and a refreshing "CAN DO" attitude.

TABLE OF CONTENTS

	Page
I. INTRODUCTION.....	1
II. OBJECTIVES.....	1
III. METHODS OF COLLECTION AND ANALYSIS.....	1
IV. DISPOSITION OF DATA.....	2
V. DISCUSSION.....	2
VI. ADDITIONAL WORK NEEDED IN THE REGION.....	3
VII. BIBLIOGRAPHY.....	4

FIGURES

1. Location of Test Panels.....	5
2. EOD Diver Retrieving Test Panels.....	5
3. Radiograph of Test Panel Exposed in Vieques Sound for 4 Months at 95 Feet.....	6
4. Radiograph of Test Panel Exposed in Vieques Sound for 8 Months at 50 Feet.....	6
5. Radiograph of Test Panel Exposed in Vieques Sound for 8 Months at 95 Feet.....	7
6. Radiograph of Test Panel Exposed in Vieques Sound for 10 Months at 50 Feet.....	7
7. Radiograph of Test Panel Exposed in Vieques Sound for 10 Months at 95 Feet.....	8
8. Radiograph of Test Panel Exposed in Vieques Sound for 12 Months at 95 Feet.....	8
9. Growth Rates of Two Important Fouling Organisms from Vieques Sound.....	9

TABLES

I. Relative Abundance of Biofouling Organisms on Test Panels Exposed at Two Depths in Vieques Sound - April 1964 to February 1969.....	10
II. Dry Weight Production on Test Panels Exposed in Vieques Sound.....	11

I. INTRODUCTION

In April 1964, the U.S. Naval Oceanographic Office (NAVOCEANO) began a marine biofouling sampling program in Vieques Sound, Puerto Rico. The program is being conducted at one site, 18°18'N, 65°26'W, by exposing test panels at 50- and 95-foot depths in 100 feet of water (Fig. 1). From April 1964 to February 1969, 103 panels were recovered from the test site. This interim report presents the results from the analyses of these panels. The sampling program at Vieques Sound is continuing until May 1970, and a final report will be prepared when the program is completed.

Previous to this NAVOCEANO program, no detailed test panel studies of fouling communities (both foulers and borers) had been made in Puerto Rican waters. The William F. Clapp Laboratories (BUYARDS and DOCKS, 1951) exposed test panels in the harbors of San Juan and Roosevelt Roads in 1951, but these exposures were primarily to study marine borers. A series of moored mines exposed off San Juan during World War II were found to have accumulated up to 3 inches of unspecified fouling organisms after 6 months of exposure (Moritz, 1944).

II. OBJECTIVES

The NAVOCEANO fouling study in Vieques Sound is one of a series of programs to determine the character and extent of the biofouling communities in various faunal provinces; i.e., regions in which the animal inhabitants are more or less the same. The objectives of this study were as follows: (1) to identify the attaching organisms, (2) to ascertain if these organisms attach in sufficient numbers to affect military operations, and (3) to determine growth rates, seasonality, productivity, and patterns of distribution of coastal fouling communities. Data from this particular program will supplement published information and, together with data from other regions being studied, will eventually result in a worldwide atlas of biofouling conditions.

III. METHODS OF COLLECTION AND ANALYSIS

Biofouling organisms were collected on standard NAVOCEANO test panels (6 x 12-inch sections of wood and asbestos, attached back to back). Panels were exposed at intervals sufficient to yield monthly and longer term information. The longest exposure was for 17 months. The panels were planted and recovered from the test site by Explosive Ordnance Disposal (EOD) divers (Fig. 2) from the U.S. Naval Station at San Juan. Additionally, temperature measurements were made and water samples for salinity analysis were collected at each test depth and at the surface whenever test panels were removed. After recovery, the panels were preserved in alcohol and shipped to NAVOCEANO for analysis.

At NAVOCEANO, the analysis consisted of identifying the various organisms, determining the number and size of each, estimating the percent of surface covered by foulers, and estimating (by radiographic technique) the degree of wood damage by marine borers. The fouling material then was scraped cleanly from the panels and oven dried at 100°C until the weight was constant.

Maximum size measurements are used to calculate growth rates, which are an index of the well-being of various species. Growth rates also are useful in determining exposure time of recovered derelict objects. The dry weight of fouling per square meter of substrate provides a statistical measure of the biofouling productivity of the area.

IV. DISPOSITION OF DATA

All data are retained on file at NAVOCEANO. A reference collection of specimens and the dried biomass scrapings, radiographs, and data summary sheets are filed under Operation Numbers 924001 and 928032.

V. DISCUSSION

The waters of Vieques Sound are oceanic and tropical in character. The range of salinity values does not exceed 34.5 - 36.5 ‰, and the temperatures vary from a high of 84.5°F (29.2°C) in September to a low of 78.0°F (25.6°C) in February and March. Larvae of potential foulers showed no seasonal preference for settlement; attachment of one or more species occurred during all months of the year.

A total of 54 species of fouling organisms (foulers and borers) was collected at the Vieques Sound test site. Table I presents the relative abundance of biofouling organisms on the test panels. As expected, a strong affinity exists between the fauna of the Vieques Sound fouling communities and the fauna of the South Florida/Bahamas region (DePalma, 1969). More than two-thirds of the animal foulers collected at the Vieques Sound test site were common to both areas.

The rate of fouling (accumulation of organic material on the asbestos substrate per unit time) was light and extremely variable from month to month and from one year to the next (Table II), and wood damage by marine borers was also light, by nearshore tropical standards. Figures 3 through 8 are reproductions of radiographs of test panels showing wood damage by teredinid borers at each test level over a 12-month period. This normally light, but occasionally epidemic, settlement of fouling and boring organisms (see Figure 7) is probably related to the distance from the nearest land masses and potential brood sites (6 and 10 miles). At the Vieques Sound test site and in most other offshore areas, the fouling rate is dependent on the vagaries of water currents to deliver larvae of fouling and

boring organisms to the test surfaces. Once settled, the organisms grew rapidly (Fig. 9), indicating that the rate of fouling is not limited by deficiencies in the environment.

VI. ADDITIONAL WORK NEEDED IN THE REGION

Additional exposure sites are planned in Vieques Passage and at Roosevelt Roads Naval Station. Data from these sites will be useful to more clearly define the relationship between fouling rate and distance from shore.

VII. BIBLIOGRAPHY

- Bureau of Yards and Docks, 1951. Report on Marine Borers and Fouling Organisms in 56 Important Harbors. NAVDOCKS TP-Re-1. UNPUBLISHED MANUSCRIPT. Department of the Navy. Washington, D.C.
- DePalma, J. R., 1969. A Study of Deep Ocean Fouling, Straits of Florida and Tongue of the Ocean, 1961 to 1968. IR No. 69-22. UNPUBLISHED MANUSCRIPT. U.S. Naval Oceanographic Office. Washington, D.C.
- Moritz, C. E., 1944. Mine Warfare and Marine Fouling. Report No. 957. UNPUBLISHED MANUSCRIPT. U.S. Naval Ordnance Laboratory. Washington, D.C.




Figure 1. Location of Test Panels



Figure 2. EOD Diver Retrieving Test Panels

Figure 3. Radiograph of Test Panel Exposed in Vieques Sound for 4 Months at 95 Feet. Shells and pallets of four Lyrodus pedicellatus can be seen.




Figure 4. Radiograph of Test Panel Exposed in Vieques Sound for 8 Months at 50 Feet. Shells and pallets of (A) Teredora malleolus and (B) Bankia carinata can be seen.



Figure 5. Radiograph of Test Panel Exposed in Vieques Sound for 8 Months at 95 Feet. The tubes of teredine borers are lined with a calcareous substance which shows as a dark outline. The light area is the anterior or working end of the tube.




Figure 6. Radiograph of Test Panel Exposed in Vieques Sound for 10 Months at 50 Feet. Arrow shows shell in the boring position.



Figure 7. Radiograph of Test Panel Exposed in Vieques Sound for 10 Months at 95 Feet. Long tubes represent an early, light set of borers; short tubes are from a more recent, heavier set.




Figure 8. Radiograph of Test Panel Exposed in Vieques Sound for 12 Months at 95 Feet. Wood damage is so extensive that this panel can be easily broken with the fingers.




Figure 9. Growth Rates of Two Important Fouling Organisms from Vieques Sound

Table 1. Relative Abundance of Biofouling Organisms on Test Panels Exposed at Two Depths in Vieques Sound - April 1964 To February 1969

Organisms	Local Abundance	
	50-foot Panels	95-foot Panels
ALGAE		
<u>Pocockiella</u> sp.	XXX	XX
<u>Desmarestia</u> <u>lingulata</u>	X	-
<u>D. aculeata</u>	X	-
<u>Polysiphonia</u> <u>opaca</u>	XX	X
<u>Hymenena</u> <u>media</u>	X	-
<u>Dictyota</u> <u>dichotoma</u>	XX	XX
<u>Jania</u> <u>adherens</u>	X	-
<u>Antithamnion</u> <u>plumula</u>	X	-
<u>Enantiocladia</u> <u>duperreyi</u>	XX	-
<u>Corallina</u> <u>cubensis</u>	-	X
<u>Melobesia</u> eae	XXX	XX
<u>Chrysonephos</u> <u>lewisii</u>	XX	XX
PORIFERA		
<u>Sycon</u> sp.	X	XX
COELENTERATA		
<u>Eudendrium</u> <u>distichum</u>	-	XXX
<u>Tubulariidae</u>	XX	X
<u>Teleso</u> sp.	XX	XX
BRYOZOA		
<u>Thalamoporella</u> <u>distorta</u>	XXX	XXX
<u>Celleporaria</u> <u>albirostris</u>	X	X
<u>Schizoporella</u> <u>unicornis</u>	-	X
<u>Savignyella</u> <u>lafonti</u>	XX	XX
<u>Catenicella</u> <u>contei</u>	XX	XX
ANNELIDA		
<u>Hydroides</u> <u>parvus</u>	XX	XX
<u>H. norvegica</u>	X	XX
<u>H. dianthus</u>	-	X
<u>H. lunulifera</u>	X	-
<u>Pomatoceros</u> <u>triqueter</u>	-	X
<u>Crosslandii</u> <u>multispinosa</u>	-	X
<u>Spirobranchus</u> <u>giganteus</u>	-	X
<u>Vermiliopsis</u> <u>infundibulum</u>	X	-
<u>Pomatostegus</u> <u>latiscapus</u>	X	X
<u>Serpula</u> <u>vermicularis</u>	-	X
<u>Salmacina</u> <u>incrustans</u>	-	X

Table I. (Continued)

Organisms	Local Abundance	
	50-foot Panels	95-foot Panels
ARTHROPODA		
<u>Balanus tintinnabulum</u>	XX	X
<u>B. venustus niveus</u>	X	X
<u>B. trigonus</u>	XX	XX
<u>Tetraclita radiata</u>	XX	-
<u>Limnoria tripunctata</u>	-	X ¹
MOLLUSCA		
<u>Ostrea equestris</u>	XX	XXX
<u>O. frons</u>	XX	XX
<u>Chama macrophylla</u>	XX	XX
<u>Pycnodonta thomasi</u>	-	XX
<u>Musculus lateralis</u>	XX	XX
<u>Spondylus americanus</u>	X	XX
<u>Pinctada radiata</u>	XX	X
<u>Brevimalleus candeanus</u>	X	-
<u>Pteria colymbus</u>	XX	-
<u>Pinna carnea</u>	X	-
<u>Lyrodus pedicellatus</u>	XX ₂	XX ₂
<u>Bankia carinata</u>	XX ₂	XX ₂
<u>Teredora malleolus</u>	XX	XX ₂
<u>Teredothyra dominicensis</u>	-	XX ₂
<u>T. matocotana</u>	-	XX
CHORDATA (Tunicata)		
<u>Botrylloides sp.</u>	XX	XX
<u>Styela partita</u>	X	X

X - never exceeds 1 % coverage of cumulative panels or occurs rarely.

XX - less than 40 % coverage of cumulative panels.

XXX - 40 % coverage or greater on cumulative panels.

X¹ Marine borer. 10-25 % of 12-month wooden panel destroyed by this and other molluscan borers.

XX² Marine borer. 26-50 % of 12-month wooden panel destroyed by this and other molluscan borers.

Table II. Dry Weight Production (in gms/m²) on Test Panels Exposed in Vieques Sound

Months of Exposure	Apr. '64-Feb. '65		Dec. '65-May '67		Mar. '67-Jan. '68		Nov. '67-Nov. '68	
	50-foot	95-foot	50-foot	95-foot	50-foot	95-foot	50-foot	95-foot
1	<50	< 50	<50	<50	<50	56	-	-
2	<50	< 50	<50	<50	<50	<50	-	-
3	<50	< 50	188	61	<50	-	-	-
4	55	146	90	90	<50	99	-	-
5	281	331	524	128	-	-	-	-
6	319	223	1,094	727	155	686	-	-
7	-	-	-	-	-	-	-	-
8	-	-	-	-	439	236	-	-
9	-	-	2,025	-	-	-	-	-
10	655	571	-	1,112	517	1,361	-	-
11	-	-	-	-	-	-	-	-
12	-	-	2,014	-	-	-	697	2,227
14	-	-	2,509	1,462	-	-	-	-
15	-	-	1,892	-	-	-	-	-
17	-	-	2,183	1,163	-	-	-	-

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)

U.S. NAVAL OCEANOGRAPHIC OFFICE

2a. REPORT SECURITY CLASSIFICATION

Unclassified

2b. GROUP

3. REPORT TITLE

MARINE BIOFOULING IN VIEQUES SOUND, PUERTO RICO, AN INTERIM REPORT,
APRIL 1964 TO FEBRUARY 1969

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

Informal Report April 1964 to February 1969

5. AUTHOR(S) (First name, middle initial, last name)

JOHN R. DePALMA

6. REPORT DATE

June 1969

7a. TOTAL NO. OF PAGES

12

7b. NO. OF REFS

3

8a. CONTRACT OR GRANT NO.

b. PROJECT NO. 102-01

c.

d.

9a. ORIGINATOR'S REPORT NUMBER(S)

IR No. 69-50

9b. OTHER REPORT NO(S) (Any other numbers that may be assigned
this report)

10. DISTRIBUTION STATEMENT

This document has been approved for public release and sale; its distribution is
unlimited.

11. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

U.S. Naval Oceanographic Office

13. ABSTRACT

In 1964, the U.S. Naval Oceanographic Office began a marine biofouling sampling program in Vieques Sound, Puerto Rico. The sampling program is continuing until May 1970. This program is one of a series of sampling programs to determine the character and extent of marine fouling communities in various marine biological provinces of the world.

The Vieques Sound program is being conducted by exposing test panels at 50- and 95-foot depths in 100 feet of water. From April 1964 to February 1969, 103 panels were recovered and analyzed for fouling and boring organisms.

Settlement of fouling larvae was found to occur throughout the year in Vieques Sound. The rate of fouling, however, was quite variable from month to month, apparently depending on whether water currents bring larvae of potential foulers from distant brood sites. A strong affinity exists between the fauna of the Vieques Sound fouling communities and the fauna of the South Florida/Bahamas region.

UNCLASSIFIED

Security Classification

14.

KEY WORDS

LINK A

LINK B

LINK C

ROLE

WT

ROLE

WT

ROLE

WT

MARINE BIOFOULING
VIEQUES SOUND, PUERTO RICO

UNCLASSIFIED

Security Classification